

CMPE 322/327 - Theory of Computation

Week 5: State Minimization & Myhill-Nerode Relations

Burak Ekici

March 21-25, 2022

Outline

- 1 A Quick Recap
- 2 State Minimization
- 3 Myhill-Nerode Relations

Definition

regular expressions are restricted patterns which use only

$$a \in \Sigma \quad \varepsilon \quad \emptyset \quad \alpha + \beta \quad \alpha^* \quad \alpha\beta$$

Definition

regular expressions are restricted patterns which use only

$$a \in \Sigma \quad \varepsilon \quad \emptyset \quad \alpha + \beta \quad \alpha^* \quad \alpha\beta$$

Theorem

finite automata, patterns, and regular expressions are **equivalent**:

Definition

regular expressions are restricted patterns which use only

$$a \in \Sigma \quad \varepsilon \quad \emptyset \quad \alpha + \beta \quad \alpha^* \quad \alpha\beta$$

Theorem

finite automata, patterns, and regular expressions are **equivalent**:

- for all $A \subseteq \Sigma^*$
- ① A is regular
 - \iff ② $A = L(\alpha)$ for some pattern α
 - \iff ③ $A = L(\alpha)$ for some regular expression α

Definition

regular expressions are restricted patterns which use only

$$a \in \Sigma \quad \varepsilon \quad \emptyset \quad \alpha + \beta \quad \alpha^* \quad \alpha\beta$$

Theorem

finite automata, patterns, and regular expressions are **equivalent**:

- for all $A \subseteq \Sigma^*$
- ① A is regular
 - \iff ② $A = L(\alpha)$ for some pattern α
 - \iff ③ $A = L(\alpha)$ for some regular expression α

Proof.

- ③ \implies ② trivial (every regular expression is a pattern)

Definition

regular expressions are restricted patterns which use only

$$a \in \Sigma \quad \varepsilon \quad \emptyset \quad \alpha + \beta \quad \alpha^* \quad \alpha\beta$$

Theorem

finite automata, patterns, and regular expressions are **equivalent**:

- for all $A \subseteq \Sigma^*$
- ① A is regular
 - \iff ② $A = L(\alpha)$ for some pattern α
 - \iff ③ $A = L(\alpha)$ for some regular expression α

Proof.

- ③ \implies ② trivial (every regular expression is a pattern)
- ② \implies ① induction on α

Definition

regular expressions are restricted patterns which use only

$$a \in \Sigma \quad \varepsilon \quad \emptyset \quad \alpha + \beta \quad \alpha^* \quad \alpha\beta$$

Theorem

finite automata, patterns, and regular expressions are **equivalent**:

- for all $A \subseteq \Sigma^*$
- ① A is regular
 - \iff ② $A = L(\alpha)$ for some pattern α
 - \iff ③ $A = L(\alpha)$ for some regular expression α

Proof.

- ③ \implies ② trivial (every regular expression is a pattern)
- ② \implies ① induction on α
- ① \implies ③

Theorem

regular sets are effectively closed under homomorphic image and **preimage**

Theorem

regular sets are effectively closed under homomorphic image and **preimage**

Proof.

- DFA $M = (Q, \Gamma, \delta, s, F)$

Theorem

regular sets are effectively closed under homomorphic image and **preimage**

Proof.

- DFA $M = (Q, \Gamma, \delta, s, F)$
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$

Theorem

regular sets are effectively closed under homomorphic image and **preimage**

Proof.

- DFA $M = (Q, \Gamma, \delta, s, F)$
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$
- $h^{-1}(L(M)) = L(M')$ for DFA $M' = (Q, \Sigma, \delta', s, F)$ with $\delta'(q, a) := \widehat{\delta}(q, h(a))$

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Proof.

- regular expression α over Σ

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Proof.

- regular expression α over Σ
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Proof.

- regular expression α over Σ
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$
- $h(L(\alpha)) = L(\alpha')$ for regular expression α' defined inductively:

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Proof.

- regular expression α over Σ
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$
- $h(L(\alpha)) = L(\alpha')$ for regular expression α' defined inductively:

$$\mathbf{a'} = h(\mathbf{a}) \quad \text{for } \mathbf{a} \in \Sigma$$

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Proof.

- regular expression α over Σ
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$
- $h(L(\alpha)) = L(\alpha')$ for regular expression α' defined inductively:

$$\mathbf{a'} = h(\mathbf{a}) \quad \text{for } \mathbf{a} \in \Sigma$$

$$\boldsymbol{\epsilon'} = \boldsymbol{\epsilon}$$

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Proof.

- regular expression α over Σ
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$
- $h(L(\alpha)) = L(\alpha')$ for regular expression α' defined inductively:

$$\mathbf{a'} = h(\mathbf{a}) \quad \text{for } \mathbf{a} \in \Sigma$$

$$\epsilon' = \epsilon$$

$$\emptyset' = \emptyset$$

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Proof.

- regular expression α over Σ
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$
- $h(L(\alpha)) = L(\alpha')$ for regular expression α' defined inductively:

$$\mathbf{a'} = h(\mathbf{a}) \quad \text{for } \mathbf{a} \in \Sigma$$

$$\boldsymbol{\epsilon'} = \boldsymbol{\epsilon}$$

$$\boldsymbol{\emptyset'} = \boldsymbol{\emptyset}$$

$$(\beta + \gamma)' = \beta' + \gamma'$$

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Proof.

- regular expression α over Σ
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$
- $h(L(\alpha)) = L(\alpha')$ for regular expression α' defined inductively:

$$\mathbf{a'} = h(\mathbf{a}) \quad \text{for } \mathbf{a} \in \Sigma$$

$$\epsilon' = \epsilon$$

$$\emptyset' = \emptyset$$

$$(\beta + \gamma)' = \beta' + \gamma'$$

$$(\beta\gamma)' = \beta'\gamma'$$

Theorem

regular sets are effectively closed under **homomorphic image** and preimage

Proof.

- regular expression α over Σ
- homomorphism $h: \Sigma^* \rightarrow \Gamma^*$
- $h(L(\alpha)) = L(\alpha')$ for regular expression α' defined inductively:

$$\mathbf{a}' = h(\mathbf{a}) \quad \text{for } \mathbf{a} \in \Sigma$$

$$\epsilon' = \epsilon$$

$$\emptyset' = \emptyset$$

$$(\beta + \gamma)' = \beta' + \gamma'$$


$$(\beta\gamma)' = \beta'\gamma'$$

$$(\beta^*)' = (\beta')^*$$


Outline

- 1 A Quick Recap
- 2 State Minimization
- 3 Myhill-Nerode Relations


Example




Example



$A = \{1\}$ $C = \{1,3\}$ $E = \{1,3,4\}$
 $B = \{1,2\}$ $D = \{1,2,4\}$ $F = \{1,4\}$




Example




$A = \{1\}$ $C = \{1,3\}$ $E = \{1,3,4\}$

$B = \{1,2\}$ $D = \{1,2,4\}$ $F = \{1,4\}$




Example

DFA



Example

DFA




not **minimal**: states 2 and 3 can be collapsed




Example

DFA




Example

DFA




Example

DFA




not **minimal**: states 2 and 3 and states 4 and 5 can be collapsed



Example

DFA



Definitions

DFA $M = (Q, \Sigma, \delta, s, F)$

Definitions

DFA $M = (Q, \Sigma, \delta, s, F)$

- state p is inaccessible if $\widehat{\delta}(s, x) \neq p$ for all $x \in \Sigma^*$

Definitions

DFA $M = (Q, \Sigma, \delta, s, F)$

- state p is inaccessible if $\widehat{\delta}(s, x) \neq p$ for all $x \in \Sigma^*$
- states p and q are **distinguishable** if

$$\exists x \in \Sigma^*, (\widehat{\delta}(p, x) \in F \wedge \widehat{\delta}(q, x) \notin F) \vee (\widehat{\delta}(p, x) \notin F \wedge \widehat{\delta}(q, x) \in F)$$

Definitions

DFA $M = (Q, \Sigma, \delta, s, F)$

- state p is inaccessible if $\widehat{\delta}(s, x) \neq p$ for all $x \in \Sigma^*$
- states p and q are **distinguishable** if

$$\exists x \in \Sigma^*, (\widehat{\delta}(p, x) \in F \wedge \widehat{\delta}(q, x) \notin F) \vee (\widehat{\delta}(p, x) \notin F \wedge \widehat{\delta}(q, x) \in F)$$

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

Definitions

DFA $M = (Q, \Sigma, \delta, s, F)$

- state p is inaccessible if $\widehat{\delta}(s, x) \neq p$ for all $x \in \Sigma^*$
- states p and q are **distinguishable** if

$$\exists x \in \Sigma^*, (\widehat{\delta}(p, x) \in F \wedge \widehat{\delta}(q, x) \notin F) \vee (\widehat{\delta}(p, x) \notin F \wedge \widehat{\delta}(q, x) \in F)$$

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

- 1 remove inaccessible states

Definitions

DFA $M = (Q, \Sigma, \delta, s, F)$

- state p is inaccessible if $\widehat{\delta}(s, x) \neq p$ for all $x \in \Sigma^*$
- states p and q are **distinguishable** if

$$\exists x \in \Sigma^*, (\widehat{\delta}(p, x) \in F \wedge \widehat{\delta}(q, x) \notin F) \vee (\widehat{\delta}(p, x) \notin F \wedge \widehat{\delta}(q, x) \in F)$$

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

- ① remove inaccessible states
- ② for every two different states, determine whether they are distinguishable (**marking**)

Definitions

DFA $M = (Q, \Sigma, \delta, s, F)$

- state p is inaccessible if $\widehat{\delta}(s, x) \neq p$ for all $x \in \Sigma^*$
- states p and q are **distinguishable** if

$$\exists x \in \Sigma^*, (\widehat{\delta}(p, x) \in F \wedge \widehat{\delta}(q, x) \notin F) \vee (\widehat{\delta}(p, x) \notin F \wedge \widehat{\delta}(q, x) \in F)$$

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

- ① remove inaccessible states
- ② for every two different states, determine whether they are distinguishable (**marking**)
- ③ **collapse** indistinguishable states

Marking Algorithm

given DFA $M = (Q, \Sigma, \delta, s, F)$ without inaccessible states

① tabulate all unordered pairs $\{p, q\}$ with $p, q \in Q$, initially unmarked

Marking Algorithm

given DFA $M = (Q, \Sigma, \delta, s, F)$ without inaccessible states

- ① tabulate all unordered pairs $\{p, q\}$ with $p, q \in Q$, initially unmarked
- ② mark $\{p, q\}$ if $p \in F$ and $q \notin F$ or vice versa

Marking Algorithm

given DFA $M = (Q, \Sigma, \delta, s, F)$ without inaccessible states

- ① tabulate all unordered pairs $\{p, q\}$ with $p, q \in Q$, initially unmarked
- ② mark $\{p, q\}$ if $p \in F$ and $q \notin F$ or vice versa
- ③ repeat until no change:

mark $\{p, q\}$ if $\{\delta(p, a), \delta(q, a)\}$ is marked for some $a \in \Sigma$

Marking Algorithm

given DFA $M = (Q, \Sigma, \delta, s, F)$ without inaccessible states

- ① tabulate all unordered pairs $\{p, q\}$ with $p, q \in Q$, initially unmarked
- ② mark $\{p, q\}$ if $p \in F$ and $q \notin F$ or vice versa
- ③ repeat until no change:

mark $\{p, q\}$ if $\{\delta(p, a), \delta(q, a)\}$ is marked for some $a \in \Sigma$

Notation

$p \approx q \iff$ states p and q are indistinguishable

Marking Algorithm

given DFA $M = (Q, \Sigma, \delta, s, F)$ without inaccessible states

- ① tabulate all unordered pairs $\{p, q\}$ with $p, q \in Q$, initially unmarked
- ② mark $\{p, q\}$ if $p \in F$ and $q \notin F$ or vice versa
- ③ repeat until no change:

mark $\{p, q\}$ if $\{\delta(p, a), \delta(q, a)\}$ is marked for some $a \in \Sigma$


Notation

$p \approx q \iff$ states p and q are indistinguishable


Lemma

$p \approx q \iff \{p, q\}$ is unmarked

Example




Example



1
 ✓ 2
 ✓ 3
 ✓ ✓ 4
 ✓ ✓ 5
 ✓ ✓ ✓ 6

① final/non-final states are distinguishable


Example



1
 ✓ 2
 ✓ 3
 ✓ ✓ 4
 ✓ ✓ 5
 ✓ ✓ ✓ 6

- ① final/non-final states are distinguishable
- ② $\{2, 6\} \xrightarrow{a} \{4, 6\}$ $\{3, 6\} \xrightarrow{a} \{5, 6\}$


Example



1
 ✓ 2
 ✓ 3
 ✓ ✓ ✓ 4
 ✓ ✓ ✓ 5
 ✓ ✓ ✓ ✓ ✓ 6


- ① final/non-final states are distinguishable
- ② $\{2, 6\} \xrightarrow{a} \{4, 6\}$ $\{3, 6\} \xrightarrow{a} \{5, 6\}$
- ③ $\{1, 4\} \xrightarrow{a} \{2, 6\}$ $\{1, 5\} \xrightarrow{a} \{2, 6\}$

Example



1
 ✓ 2
 ✓ 3
 ✓ ✓ ✓ 4
 ✓ ✓ ✓ 5
 ✓ ✓ ✓ ✓ ✓ 6

- ① final/non-final states are distinguishable
- ② $\{2, 6\} \xrightarrow{a} \{4, 6\}$ $\{3, 6\} \xrightarrow{a} \{5, 6\}$
- ③ $\{1, 4\} \xrightarrow{a} \{2, 6\}$ $\{1, 5\} \xrightarrow{a} \{2, 6\}$



Definition

states p and q of DFA $M = (Q, \Sigma, \delta, s, F)$ are **indistinguishable** ($p \approx q$) if

$$\forall x \in \Sigma^*, \hat{\delta}(p, x) \in F \iff \hat{\delta}(q, x) \in F$$

Definition

states p and q of DFA $M = (Q, \Sigma, \delta, s, F)$ are **indistinguishable** ($p \approx q$) if

$$\forall x \in \Sigma^*, \widehat{\delta}(p, x) \in F \iff \widehat{\delta}(q, x) \in F$$

Lemma

\approx is **equivalence relation** on Q

Definition

states p and q of DFA $M = (Q, \Sigma, \delta, s, F)$ are **indistinguishable** ($p \approx q$) if

$$\forall x \in \Sigma^*, \widehat{\delta}(p, x) \in F \iff \widehat{\delta}(q, x) \in F$$

Lemma

\approx is **equivalence relation** on Q

$$\textcircled{1} \forall p \in Q \quad p \approx p \quad (\text{reflexivity})$$

Definition

states p and q of DFA $M = (Q, \Sigma, \delta, s, F)$ are **indistinguishable** ($p \approx q$) if

$$\forall x \in \Sigma^*, \widehat{\delta}(p, x) \in F \iff \widehat{\delta}(q, x) \in F$$

Lemma

\approx is **equivalence relation** on Q

- ① $\forall p \in Q \quad p \approx p$ (reflexivity)
- ② $\forall p, q \in Q \quad p \approx q \implies q \approx p$ (symmetry)

Definition

states p and q of DFA $M = (Q, \Sigma, \delta, s, F)$ are **indistinguishable** ($p \approx q$) if

$$\forall x \in \Sigma^*, \widehat{\delta}(p, x) \in F \iff \widehat{\delta}(q, x) \in F$$

Lemma

\approx is **equivalence relation** on Q

- ① $\forall p \in Q \quad p \approx p$ (reflexivity)
- ② $\forall p, q \in Q \quad p \approx q \implies q \approx p$ (symmetry)
- ③ $\forall p, q, r \in Q \quad p \approx q \wedge q \approx r \implies p \approx r$ (transitivity)

Definition

states p and q of DFA $M = (Q, \Sigma, \delta, s, F)$ are **indistinguishable** ($p \approx q$) if

$$\forall x \in \Sigma^*, \widehat{\delta}(p, x) \in F \iff \widehat{\delta}(q, x) \in F$$

Lemma

\approx is **equivalence relation** on Q

- ① $\forall p \in Q \quad p \approx p$ (reflexivity)
- ② $\forall p, q \in Q \quad p \approx q \implies q \approx p$ (symmetry)
- ③ $\forall p, q, r \in Q \quad p \approx q \wedge q \approx r \implies p \approx r$ (transitivity)

Notation

$[p]_{\approx} := \{q \in Q \mid p \approx q\}$ denotes **equivalence class** of p

Definition (Collapsing Indistinguishable States)

DFA M/\approx is defined as $(Q', \Sigma, \delta', s', F')$ with

Definition (Collapsing Indistinguishable States)

DFA M/\approx is defined as $(Q', \Sigma, \delta', s', F')$ with

- $Q' := \{[p]_{\approx} \mid p \in Q\}$

Definition (Collapsing Indistinguishable States)

DFA M/\approx is defined as $(Q', \Sigma, \delta', s', F')$ with

- $Q' := \{[p]_{\approx} \mid p \in Q\}$
- $\delta'([p]_{\approx}, a) := [\delta(p, a)]_{\approx}$ well defined: $p \approx q \implies \delta(p, a) \approx \delta(q, a)$

Definition (Collapsing Indistinguishable States)

DFA M/\approx is defined as $(Q', \Sigma, \delta', s', F')$ with

- $Q' := \{[p]_{\approx} \mid p \in Q\}$
- $\delta'([p]_{\approx}, a) := [\delta(p, a)]_{\approx}$ well defined: $p \approx q \implies \delta(p, a) \approx \delta(q, a)$
- $s' := [s]_{\approx}$

Definition (Collapsing Indistinguishable States)

DFA M/\approx is defined as $(Q', \Sigma, \delta', s', F')$ with

- $Q' := \{[p]_{\approx} \mid p \in Q\}$
- $\delta'([p]_{\approx}, a) := [\delta(p, a)]_{\approx}$ well defined: $p \approx q \implies \delta(p, a) \approx \delta(q, a)$
- $s' := [s]_{\approx}$
- $F' := \{[p]_{\approx} \mid p \in F\}$

Definition (Collapsing Indistinguishable States)

DFA M/\approx is defined as $(Q', \Sigma, \delta', s', F')$ with

- $Q' := \{[p]_{\approx} \mid p \in Q\}$
- $\delta'([p]_{\approx}, a) := [\delta(p, a)]_{\approx}$ well defined: $p \approx q \implies \delta(p, a) \approx \delta(q, a)$
- $s' := [s]_{\approx}$
- $F' := \{[p]_{\approx} \mid p \in F\}$

Lemma

- ① $\widehat{\delta'}([p]_{\approx}, x) = [\widehat{\delta}(p, x)]_{\approx}$ for all $x \in \Sigma^*$

Definition (Collapsing Indistinguishable States)

DFA M/\approx is defined as $(Q', \Sigma, \delta', s', F')$ with

- $Q' := \{[p]_{\approx} \mid p \in Q\}$
- $\delta'([p]_{\approx}, a) := [\delta(p, a)]_{\approx}$ well defined: $p \approx q \implies \delta(p, a) \approx \delta(q, a)$
- $s' := [s]_{\approx}$
- $F' := \{[p]_{\approx} \mid p \in F\}$

Lemma

- 1 $\widehat{\delta'}([p]_{\approx}, x) = [\widehat{\delta}(p, x)]_{\approx}$ for all $x \in \Sigma^*$
- 2 $p \in F \iff [p]_{\approx} \in F'$

for all $p \in Q$

Theorem

$$L(M/\approx) = L(M)$$

Theorem

$$L(M/\approx) = L(M)$$

Proof.

$$x \in L(M/\approx) \iff \widehat{\delta'}([s]_{\approx}, x) \in F'$$

Theorem

$$L(M/\approx) = L(M)$$

Proof.

$$\begin{aligned} x \in L(M/\approx) &\iff \widehat{\delta'}([s]_{\approx}, x) \in F' \\ &\iff [\widehat{\delta}(s, x)]_{\approx} \in F' \end{aligned}$$

Theorem

$$L(M/\approx) = L(M)$$

Proof.

$$\begin{aligned} x \in L(M/\approx) &\iff \widehat{\delta}'([s]_{\approx}, x) \in F' \\ &\iff [\widehat{\delta}(s, x)]_{\approx} \in F' \\ &\iff \widehat{\delta}(s, x) \in F \end{aligned}$$

Theorem

$$L(M/\approx) = L(M)$$

Proof.

$$\begin{aligned}
 x \in L(M/\approx) &\iff \widehat{\delta'}([s]_{\approx}, x) \in F' \\
 &\iff [\widehat{\delta}(s, x)]_{\approx} \in F' \\
 &\iff \widehat{\delta}(s, x) \in F \\
 &\iff x \in L(M)
 \end{aligned}$$

Question

is M/\approx minimum-state DFA for $L(M)$?

Question

is M/\approx minimum-state DFA for $L(M)$?

Lemma

M/\approx cannot be collapsed further

Question

is M/\approx minimum-state DFA for $L(M)$?

Lemma

M/\approx cannot be collapsed further

Proof.

$$[p]_{\approx} \approx [q]_{\approx} \iff \forall x \in \Sigma^* \quad (\widehat{\delta}'([p]_{\approx}, x) \in F' \iff \widehat{\delta}'([q]_{\approx}, x) \in F')$$



Question

is M/\approx minimum-state DFA for $L(M)$?

Lemma

M/\approx cannot be collapsed further

Proof.

$$\begin{aligned}
 [p]_{\approx} \approx [q]_{\approx} &\iff \forall x \in \Sigma^* \quad (\widehat{\delta}'([p]_{\approx}, x) \in F' \iff \widehat{\delta}'([q]_{\approx}, x) \in F') \\
 &\iff \forall x \in \Sigma^* \quad ([\widehat{\delta}(p, x)]_{\approx} \in F' \iff [\widehat{\delta}(q, x)]_{\approx} \in F')
 \end{aligned}$$



Question

is M/\approx minimum-state DFA for $L(M)$?

Lemma

M/\approx cannot be collapsed further

Proof.

$$\begin{aligned}
 [p]_{\approx} \approx [q]_{\approx} &\iff \forall x \in \Sigma^* \quad (\widehat{\delta}'([p]_{\approx}, x) \in F' \iff \widehat{\delta}'([q]_{\approx}, x) \in F') \\
 &\iff \forall x \in \Sigma^* \quad ([\widehat{\delta}(p, x)]_{\approx} \in F' \iff [\widehat{\delta}(q, x)]_{\approx} \in F') \\
 &\iff \forall x \in \Sigma^* \quad (\widehat{\delta}(p, x) \in F \iff \widehat{\delta}(q, x) \in F)
 \end{aligned}$$



Question

is M/\approx minimum-state DFA for $L(M)$?

Lemma

M/\approx cannot be collapsed further

Proof.

$$\begin{aligned} [p]_{\approx} \approx [q]_{\approx} &\iff \forall x \in \Sigma^* \quad (\widehat{\delta}'([p]_{\approx}, x) \in F' \iff \widehat{\delta}'([q]_{\approx}, x) \in F') \\ &\iff \forall x \in \Sigma^* \quad ([\widehat{\delta}(p, x)]_{\approx} \in F' \iff [\widehat{\delta}(q, x)]_{\approx} \in F') \\ &\iff \forall x \in \Sigma^* \quad (\widehat{\delta}(p, x) \in F \iff \widehat{\delta}(q, x) \in F) \\ &\iff \forall x \in \Sigma^* \quad p \approx q \end{aligned}$$



Question

is M/\approx minimum-state DFA for $L(M)$?

Lemma


M/\approx cannot be collapsed further

Proof.


$$\begin{aligned}
 [p]_{\approx} \approx [q]_{\approx} &\iff \forall x \in \Sigma^* \quad (\widehat{\delta}'([p]_{\approx}, x) \in F' \iff \widehat{\delta}'([q]_{\approx}, x) \in F') \\
 &\iff \forall x \in \Sigma^* \quad ([\widehat{\delta}(p, x)]_{\approx} \in F' \iff [\widehat{\delta}(q, x)]_{\approx} \in F') \\
 &\iff \forall x \in \Sigma^* \quad (\widehat{\delta}(p, x) \in F \iff \widehat{\delta}(q, x) \in F) \\
 &\iff \forall x \in \Sigma^* \quad p \approx q \\
 &\iff \forall x \in \Sigma^* \quad [p]_{\approx} = [q]_{\approx}
 \end{aligned}$$





Example




Example



Example




Example




a
 b
 c
 ✓✓✓ d
 ✓ e
 ✓ f
 ✓✓✓ ✓✓ g
 ✓✓✓✓✓✓✓ h
 ✓ ✓✓✓ i
 ✓✓✓ ✓✓✓ j
 ✓✓✓✓✓✓✓✓ k
 ✓✓✓✓✓✓✓✓✓✓ l

Example




a
 b
 ✓✓ c
 ✓✓✓ d
 ✓✓ e
 ✓✓ ✓✓ f
 ✓✓✓ ✓✓ g
 ✓✓✓✓✓ h
 ✓✓ ✓✓ ✓✓ i
 ✓✓✓ ✓✓ ✓✓ j
 ✓✓✓✓✓✓✓ k
 ✓✓✓✓✓✓✓✓✓ l

Example




a
 ✓ b
 ✓✓ c
 ✓✓✓ d
 ✓ ✓✓ e
 ✓✓ ✓✓ f
 ✓✓✓ ✓✓ g
 ✓✓✓✓✓✓✓ h
 ✓✓✓✓✓✓✓ i
 ✓✓✓✓✓✓✓ j
 ✓✓✓✓✓✓✓ k
 ✓✓✓✓✓✓✓✓ l

Example



a
 ✓ b
 ✓ ✓ c
 ✓ ✓ ✓ d
 ✓ ✓ ✓ ✓ e
 ✓ ✓ ✓ ✓ f
 ✓ ✓ ✓ ✓ g
 ✓ ✓ ✓ ✓ h
 ✓ ✓ ✓ ✓ i
 ✓ ✓ ✓ ✓ j
 ✓ ✓ ✓ ✓ k
 ✓ ✓ ✓ ✓ l

Example



a
 ✓ b
 ✓ ✓ c
 ✓ ✓ ✓ d
 ✓ ✓ ✓ ✓ e
 ✓ ✓ ✓ ✓ ✓ f
 ✓ ✓ ✓ ✓ ✓ g
 ✓ ✓ ✓ ✓ ✓ h
 ✓ ✓ ✓ ✓ ✓ i
 ✓ ✓ ✓ ✓ ✓ j
 ✓ ✓ ✓ ✓ ✓ k
 ✓ ✓ ✓ ✓ ✓ l
 states d, g and h, k can be merged

Outline

- 1 A Quick Recap
- 2 State Minimization
- 3 Myhill-Nerode Relations**

Definition

Myhill-Nerode relation \equiv for $L \subseteq \Sigma^*$ is an equivalence relation that

Definition

Myhill-Nerode relation \equiv for $L \subseteq \Sigma^*$ is an equivalence relation that

- is **right congruent**: $\forall x, y \in \Sigma^* \quad x \equiv y \implies \forall a \in \Sigma \quad xa \equiv ya$

Definition

Myhill-Nerode relation \equiv for $L \subseteq \Sigma^*$ is an equivalence relation that

- is right congruent: $\forall x, y \in \Sigma^* \quad x \equiv y \implies \forall a \in \Sigma \quad xa \equiv ya$
- **refines** L : $\forall x, y \in \Sigma^* \quad x \equiv y \implies \text{either } x, y \in L \text{ or } x, y \notin L$

Definition

Myhill-Nerode relation \equiv for $L \subseteq \Sigma^*$ is an equivalence relation that

- is right congruent: $\forall x, y \in \Sigma^* \quad x \equiv y \implies \forall a \in \Sigma \quad xa \equiv ya$
- refines L : $\forall x, y \in \Sigma^* \quad x \equiv y \implies \text{either } x, y \in L \text{ or } x, y \notin L$
- is of **finite index**: \equiv has finitely many equivalence classes

Definition

equivalence relation \equiv_M on Σ^* for DFA $M = (Q, \Sigma, \delta, s, F)$ is defined as follows:

$$x \equiv_M y \iff \widehat{\delta}(s, x) = \widehat{\delta}(s, y)$$

Definition

equivalence relation \equiv_M on Σ^* for DFA $M = (Q, \Sigma, \delta, s, F)$ is defined as follows:

$$x \equiv_M y \iff \widehat{\delta}(s, x) = \widehat{\delta}(s, y)$$

Lemma

- \equiv_M is **right congruent**: $\forall x, y \in \Sigma^* \quad x \equiv_M y \implies \forall a \in \Sigma \quad xa \equiv_M ya$

Definition

equivalence relation \equiv_M on Σ^* for DFA $M = (Q, \Sigma, \delta, s, F)$ is defined as follows:

$$x \equiv_M y \iff \widehat{\delta}(s, x) = \widehat{\delta}(s, y)$$

Lemma

- \equiv_M is right congruent: $\forall x, y \in \Sigma^* \quad x \equiv_M y \implies \forall a \in \Sigma \quad xa \equiv_M ya$
- \equiv_M **refines** $L(M)$: $\forall x, y \in \Sigma^* \quad x \equiv_M y \implies \text{either } x, y \in L(M) \text{ or } x, y \notin L(M)$

Definition

equivalence relation \equiv_M on Σ^* for DFA $M = (Q, \Sigma, \delta, s, F)$ is defined as follows:

$$x \equiv_M y \iff \widehat{\delta}(s, x) = \widehat{\delta}(s, y)$$

Lemma

- \equiv_M is right congruent: $\forall x, y \in \Sigma^* \quad x \equiv_M y \implies \forall a \in \Sigma \quad xa \equiv_M ya$
- \equiv_M refines $L(M)$: $\forall x, y \in \Sigma^* \quad x \equiv_M y \implies \text{either } x, y \in L(M) \text{ or } x, y \notin L(M)$
- \equiv_M is of **finite index**: \equiv_M has finitely many equivalence classes

Definition

equivalence relation \equiv_M on Σ^* for DFA $M = (Q, \Sigma, \delta, s, F)$ is defined as follows:

$$x \equiv_M y \iff \widehat{\delta}(s, x) = \widehat{\delta}(s, y)$$


Lemma

- \equiv_M is right congruent: $\forall x, y \in \Sigma^* \quad x \equiv_M y \implies \forall a \in \Sigma \quad xa \equiv_M ya$
- \equiv_M refines $L(M)$: $\forall x, y \in \Sigma^* \quad x \equiv_M y \implies \text{either } x, y \in L(M) \text{ or } x, y \notin L(M)$
- \equiv_M is of finite index: \equiv_M has finitely many equivalence classes


Corollary

\equiv_M is Myhill-Nerode relation for $L(M)$


Example





Example





\equiv_M




Example




Example




Example




Example





Example





Example



Example



Example



M

start \rightarrow 1 \xrightarrow{ab} \rightarrow \rightarrow \rightarrow 6 \xrightarrow{ab} 6

\equiv_M

ϵ a aa baa
 b bb ab $aabba$
 ba $bbba$ \dots

Definition

given Myhill-Nerode relation \equiv for set $L \subseteq \Sigma^*$, DFA M_{\equiv} is defined as $(Q, \Sigma, \delta, s, F)$ with

- $Q := \{[x]_{\equiv} \mid x \in \Sigma^*\}$

Definition

given Myhill-Nerode relation \equiv for set $L \subseteq \Sigma^*$, DFA M_{\equiv} is defined as $(Q, \Sigma, \delta, s, F)$ with

- $Q := \{[x]_{\equiv} \mid x \in \Sigma^*\}$
- $\delta([x]_{\equiv}, a) := [xa]_{\equiv}$ well-defined: $x \equiv y \implies xa \equiv ya$

Definition

given Myhill-Nerode relation \equiv for set $L \subseteq \Sigma^*$, DFA M_{\equiv} is defined as $(Q, \Sigma, \delta, s, F)$ with

- $Q := \{[x]_{\equiv} \mid x \in \Sigma^*\}$
- $\delta([x]_{\equiv}, a) := [xa]_{\equiv}$ well-defined: $x \equiv y \implies xa \equiv ya$
- $s := [\epsilon]_{\equiv}$

Definition

given Myhill-Nerode relation \equiv for set $L \subseteq \Sigma^*$, DFA M_{\equiv} is defined as $(Q, \Sigma, \delta, s, F)$ with

- $Q := \{[x]_{\equiv} \mid x \in \Sigma^*\}$
- $\delta([x]_{\equiv}, a) := [xa]_{\equiv}$ well-defined: $x \equiv y \implies xa \equiv ya$
- $s := [\epsilon]_{\equiv}$
- $F := \{[x]_{\equiv} \mid x \in L\}$

Definition

given Myhill-Nerode relation \equiv for set $L \subseteq \Sigma^*$, DFA M_{\equiv} is defined as $(Q, \Sigma, \delta, s, F)$ with

- $Q := \{[x]_{\equiv} \mid x \in \Sigma^*\}$
- $\delta([x]_{\equiv}, a) := [xa]_{\equiv}$ well-defined: $x \equiv y \implies xa \equiv ya$
- $s := [\epsilon]_{\equiv}$
- $F := \{[x]_{\equiv} \mid x \in L\}$

Lemma

① $\widehat{\delta}([x]_{\equiv}, y) = [xy]_{\equiv}$ for all $y \in \Sigma^*$

for all $x \in \Sigma^*$

Definition

given Myhill-Nerode relation \equiv for set $L \subseteq \Sigma^*$, DFA M_{\equiv} is defined as $(Q, \Sigma, \delta, s, F)$ with

- $Q := \{[x]_{\equiv} \mid x \in \Sigma^*\}$
- $\delta([x]_{\equiv}, a) := [xa]_{\equiv}$ well-defined: $x \equiv y \implies xa \equiv ya$
- $s := [\epsilon]_{\equiv}$
- $F := \{[x]_{\equiv} \mid x \in L\}$

Lemma

① $\widehat{\delta}([x]_{\equiv}, y) = [xy]_{\equiv}$ for all $y \in \Sigma^*$

② $x \in L \iff [x]_{\equiv} \in F$

for all $x \in \Sigma^*$

Theorem

$$L(M_{\equiv}) = L$$

Theorem

$$L(M_{\equiv}) = L$$

Proof.

$$x \in L(M_{\equiv}) \iff \hat{\delta}([\varepsilon]_{\equiv}, x) \in F$$



Theorem

$$L(M_{\equiv}) = L$$

Proof.

$$\begin{aligned} x \in L(M_{\equiv}) &\iff \hat{\delta}([\varepsilon]_{\equiv}, x) \in F \\ &\iff [x]_{\equiv} \in F \end{aligned}$$



Theorem

$$L(M_{\equiv}) = L$$

Proof.

$$\begin{aligned} x \in L(M_{\equiv}) &\iff \hat{\delta}([\varepsilon]_{\equiv}, x) \in F \\ &\iff [x]_{\equiv} \in F \\ &\iff x \in L \end{aligned}$$



Theorem

$$L(M_{\equiv}) = L$$

Proof.

$$\begin{aligned} x \in L(M_{\equiv}) &\iff \widehat{\delta}([\varepsilon]_{\equiv}, x) \in F \\ &\iff [x]_{\equiv} \in F \\ &\iff x \in L \end{aligned}$$



Corollary

if L admits Myhill-Nerode relation then L is regular

Theorem

two mappings (for $L \subseteq \Sigma^*$)

- $D \mapsto \equiv_D$ from DFAs for L to Myhill-Nerode relations for L
- $\approx \mapsto M_\approx$ from Myhill-Nerode relations for L to DFAs for L

are each others **inverse** (up to isomorphism of automata):

Theorem

two mappings (for $L \subseteq \Sigma^*$)

- $D \mapsto \equiv_D$ from DFAs for L to Myhill-Nerode relations for L
- $\approx \mapsto M_\approx$ from Myhill-Nerode relations for L to DFAs for L

are each others **inverse** (up to isomorphism of automata):

$$\equiv_D \quad \forall \text{ DFA } D \text{ without inaccessible states}$$

Theorem

two mappings (for $L \subseteq \Sigma^*$)

- $D \mapsto \equiv_D$ from DFAs for L to Myhill-Nerode relations for L
- $\approx \mapsto M_{\approx}$ from Myhill-Nerode relations for L to DFAs for L

are each others **inverse** (up to isomorphism of automata):

- $M_{(\equiv_D)}$ \forall DFA D without inaccessible states

Theorem

two mappings (for $L \subseteq \Sigma^*$)

- $D \mapsto \equiv_D$ from DFAs for L to Myhill-Nerode relations for L
- $\approx \mapsto M_{\approx}$ from Myhill-Nerode relations for L to DFAs for L

are each others **inverse** (up to isomorphism of automata):

- $M_{(\equiv_D)} \simeq D \quad \forall \text{ DFA } D \text{ without inaccessible states}$

Theorem

two mappings (for $L \subseteq \Sigma^*$)

- $D \mapsto \equiv_D$ from DFAs for L to Myhill-Nerode relations for L
- $\approx \mapsto M_{\approx}$ from Myhill-Nerode relations for L to DFAs for L

are each others **inverse** (up to isomorphism of automata):

- $M_{(\equiv_D)} \simeq D$ \forall DFA D without inaccessible states
 M_{\approx} \forall Myhill-Nerode relation \approx

Theorem

two mappings (for $L \subseteq \Sigma^*$)

- $D \mapsto \equiv_D$ from DFAs for L to Myhill-Nerode relations for L
- $\approx \mapsto M_{\approx}$ from Myhill-Nerode relations for L to DFAs for L

are each others **inverse** (up to isomorphism of automata):

- $M_{(\equiv_D)} \simeq D$ \forall DFA D without inaccessible states
- $\equiv_{(M_{\approx})}$ \forall Myhill-Nerode relation \approx


Theorem

two mappings (for $L \subseteq \Sigma^*$)


- $D \mapsto \equiv_D$ from DFAs for L to Myhill-Nerode relations for L
- $\approx \mapsto M_{\approx}$ from Myhill-Nerode relations for L to DFAs for L

are each others **inverse** (up to isomorphism of automata):

- $M_{(\equiv_D)} \simeq D$ \forall DFA D without inaccessible states
- $\equiv_{(M_{\approx})} = \approx$ \forall Myhill-Nerode relation \approx



DFAs for L



Myhill-Nerode relations for L

DFAs with inaccessible states

DFAs for L


Myhill-Nerode relations for L


DFAs with inaccessible states


D


DFAs for L


Myhill-Nerode relations for L

















Definition

for any set $L \subseteq \Sigma^*$, equivalence relation \equiv_L on Σ^* is defined as follows:

$$x \equiv_L y \iff \forall z \in \Sigma^*, (xz \in L \iff yz \in L)$$

Definition

for any set $L \subseteq \Sigma^*$, equivalence relation \equiv_L on Σ^* is defined as follows:

$$x \equiv_L y \iff \forall z \in \Sigma^*, (xz \in L \iff yz \in L)$$

Lemma

for any set $L \subseteq \Sigma^*$, \equiv_L is **coarsest** right congruent refinement of L :

Definition

for any set $L \subseteq \Sigma^*$, equivalence relation \equiv_L on Σ^* is defined as follows:

$$x \equiv_L y \iff \forall z \in \Sigma^*, (xz \in L \iff yz \in L)$$

Lemma

for any set $L \subseteq \Sigma^*$, \equiv_L is **coarsest** right congruent refinement of L :

if \sim is right congruent equivalence relation refining L then

$$\forall x, y \in \Sigma^*, x \sim y \implies x \equiv_L y$$

Definition

for any set $L \subseteq \Sigma^*$, equivalence relation \equiv_L on Σ^* is defined as follows:

$$x \equiv_L y \iff \forall z \in \Sigma^*, (xz \in L \iff yz \in L)$$

Lemma

for any set $L \subseteq \Sigma^*$, \equiv_L is **coarsest** right congruent refinement of L :

if \sim is right congruent equivalence relation refining L then

$$\forall x, y \in \Sigma^*, x \sim y \implies x \equiv_L y$$

\equiv_L has fewest equivalence classes

Theorem (Myhill-Nerode)

following statements are equivalent for any set $L \subseteq \Sigma^*$:

- L is regular
- L admits Myhill-Nerode relation

Theorem (Myhill-Nerode)

following statements are equivalent for any set $L \subseteq \Sigma^*$:

- L is regular
- L admits Myhill-Nerode relation
- \equiv_L is of finite index

Theorem (Myhill-Nerode)

following statements are equivalent for any set $L \subseteq \Sigma^*$:

- L is regular
- L admits Myhill-Nerode relation
- \equiv_L is of finite index

Corollary

for every regular set L , $M_{(\equiv_L)}$ is minimum-state DFA for L

Theorem (Myhill-Nerode)

following statements are equivalent for any set $L \subseteq \Sigma^*$:

- L is regular
- L admits Myhill-Nerode relation
- \equiv_L is of finite index

Corollary

for every regular set L , $M_{(\equiv_L)}$ is minimum-state DFA for L

Theorem

for every DFA M , $M/\approx \simeq M_{\equiv_L}$

Example

- 1 $A := \{a^n b^n \mid n \geq 0\}$ is not regular

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular
because \equiv_A has infinitely many equivalence classes

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular
because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j$$

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular
because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j \quad (a^i b^i \in A \text{ and } a^j b^i \notin A)$$

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular
because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j \quad (a^i b^i \in A \text{ and } a^j b^i \notin A)$$

- ② $B := \{a^{2^n} \mid n \geq 0\}$ is not regular

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular
because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j \quad (a^i b^i \in A \text{ and } a^j b^i \notin A)$$

- ② $B := \{a^{2^n} \mid n \geq 0\}$ is not regular
because \equiv_B has infinitely many equivalence classes

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular
because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j \quad (a^i b^i \in A \text{ and } a^j b^i \notin A)$$

- ② $B := \{a^{2^n} \mid n \geq 0\}$ is not regular
because \equiv_B has infinitely many equivalence classes

$$i < j \implies a^{2^i} \not\equiv_B a^{2^j}$$

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular
because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j \quad (a^i b^i \in A \text{ and } a^j b^i \notin A)$$

- ② $B := \{a^{2^n} \mid n \geq 0\}$ is not regular
because \equiv_B has infinitely many equivalence classes

$$i < j \implies a^{2^i} \not\equiv_B a^{2^j} \quad (a^{2^i} a^{2^i} = a^{2^{i+1}} \in B \text{ and } a^{2^j} a^{2^i} \notin B)$$

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular

because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j \quad (a^i b^i \in A \text{ and } a^j b^i \notin A)$$

- ② $B := \{a^{2^n} \mid n \geq 0\}$ is not regular

because \equiv_B has infinitely many equivalence classes

$$i < j \implies a^{2^i} \not\equiv_B a^{2^j} \quad (a^{2^i} a^{2^i} = a^{2^{i+1}} \in B \text{ and } a^{2^j} a^{2^i} \notin B)$$

- ③ $C := \{a^{n!} \mid n \geq 0\}$ is not regular

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular
because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j \quad (a^i b^i \in A \text{ and } a^j b^i \notin A)$$

- ② $B := \{a^{2^n} \mid n \geq 0\}$ is not regular
because \equiv_B has infinitely many equivalence classes

$$i < j \implies a^{2^i} \not\equiv_B a^{2^j} \quad (a^{2^i} a^{2^i} = a^{2^{i+1}} \in B \text{ and } a^{2^j} a^{2^i} \notin B)$$

- ③ $C := \{a^{n!} \mid n \geq 0\}$ is not regular
because \equiv_C has infinitely many equivalence classes

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular

because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j \quad (a^i b^i \in A \text{ and } a^j b^i \notin A)$$

- ② $B := \{a^{2^n} \mid n \geq 0\}$ is not regular

because \equiv_B has infinitely many equivalence classes

$$i < j \implies a^{2^i} \not\equiv_B a^{2^j} \quad (a^{2^i} a^{2^i} = a^{2^{i+1}} \in B \text{ and } a^{2^j} a^{2^i} \notin B)$$

- ③ $C := \{a^{n!} \mid n \geq 0\}$ is not regular

because \equiv_C has infinitely many equivalence classes

$$i < j \implies a^{i!} \not\equiv_C a^{j!}$$

Example

- ① $A := \{a^n b^n \mid n \geq 0\}$ is not regular

because \equiv_A has infinitely many equivalence classes

$$i \neq j \implies a^i \not\equiv_A a^j \quad (a^i b^i \in A \text{ and } a^j b^i \notin A)$$

- ② $B := \{a^{2^n} \mid n \geq 0\}$ is not regular

because \equiv_B has infinitely many equivalence classes

$$i < j \implies a^{2^i} \not\equiv_B a^{2^j} \quad (a^{2^i} a^{2^i} = a^{2^{i+1}} \in B \text{ and } a^{2^j} a^{2^i} \notin B)$$

- ③ $C := \{a^{n!} \mid n \geq 0\}$ is not regular

because \equiv_C has infinitely many equivalence classes

$$i < j \implies a^{i!} \not\equiv_C a^{j!} \quad (a^{i!} a^{i!i} = a^{(i+1)!} \in C \text{ and } a^{j!} a^{i!i} \notin C)$$

Example

- ④ $D := \{a^p \mid p \text{ is prime}\}$ is not regular
because \equiv_D has infinitely many equivalence classes

$$i < j \text{ and } i, j \text{ are primes} \implies a^i \not\equiv_D a^j$$

Example

- ④ $D := \{a^p \mid p \text{ is prime}\}$ is not regular
because \equiv_D has infinitely many equivalence classes

$$i < j \text{ and } i, j \text{ are primes} \implies a^i \not\equiv_D a^j$$

- suppose $a^i \equiv_D a^j$ and let $k = j - i$

Example

- ④ $D := \{a^p \mid p \text{ is prime}\}$ is not regular
because \equiv_D has infinitely many equivalence classes

$$i < j \text{ and } i, j \text{ are primes} \implies a^i \not\equiv_D a^j$$

- suppose $a^i \equiv_D a^j$ and let $k = j - i$
- $a^i \equiv_D a^j = a^i a^k \equiv_D a^j a^k$

Example

- ④ $D := \{a^p \mid p \text{ is prime}\}$ is not regular
because \equiv_D has infinitely many equivalence classes

$$i < j \text{ and } i, j \text{ are primes} \implies a^i \not\equiv_D a^j$$

- suppose $a^i \equiv_D a^j$ and let $k = j - i$
- $a^i \equiv_D a^j = a^i a^k \equiv_D a^j a^k \equiv_D a^j a^k a^k$

Example

- ④ $D := \{a^p \mid p \text{ is prime}\}$ is not regular
because \equiv_D has infinitely many equivalence classes

$$i < j \text{ and } i, j \text{ are primes} \implies a^i \not\equiv_D a^j$$

- suppose $a^i \equiv_D a^j$ and let $k = j - i$
- $a^i \equiv_D a^j = a^i a^k \equiv_D a^j a^k \equiv_D a^j a^k a^k = a^j a^{2k}$

Example

- ④ $D := \{a^p \mid p \text{ is prime}\}$ is not regular
because \equiv_D has infinitely many equivalence classes

$$i < j \text{ and } i, j \text{ are primes} \implies a^i \not\equiv_D a^j$$

- suppose $a^i \equiv_D a^j$ and let $k = j - i$
- $a^i \equiv_D a^j = a^i a^k \equiv_D a^j a^k \equiv_D a^j a^k a^k = a^j a^{2k} \equiv_D \dots \equiv_D a^j a^{jk} = a^{j(k+1)}$

Example

- ④ $D := \{a^p \mid p \text{ is prime}\}$ is not regular
because \equiv_D has infinitely many equivalence classes

$$i < j \text{ and } i, j \text{ are primes} \implies a^i \not\equiv_D a^j$$

- suppose $a^i \equiv_D a^j$ and let $k = j - i$
- $a^i \equiv_D a^j = a^i a^k \equiv_D a^j a^k \equiv_D a^j a^k a^k = a^j a^{2k} \equiv_D \dots \equiv_D a^j a^{jk} = a^{j(k+1)}$
- $a^i \in D$ and $a^{j(k+1)} \notin D$

Example

- ④ $D := \{a^p \mid p \text{ is prime}\}$ is not regular
because \equiv_D has infinitely many equivalence classes

$$i < j \text{ and } i, j \text{ are primes} \implies a^i \not\equiv_D a^j$$

- suppose $a^i \equiv_D a^j$ and let $k = j - i$
- $a^i \equiv_D a^j = a^i a^k \equiv_D a^j a^k \equiv_D a^j a^k a^k = a^j a^{2k} \equiv_D \dots \equiv_D a^j a^{jk} = a^{j(k+1)}$
- $a^i \in D$ and $a^{j(k+1)} \notin D$
- \equiv_D does not refine D

Thanks! & Questions?