

# CMPE 322/327 - Theory of Computation

## Week 2: Deterministic Finite Automata & Closure Properties

Burak Ekici

February 28 - March 4, 2022

# Outline

1 A Quick Recap

2 Chomsky Hierarchy

3 Deterministic Finite State Automata

4 Closure Properties

## Definitions

- alphabet is finite set; its elements are called symbols or letters

## Definitions

- alphabet is finite set; its elements are called symbols or letters
- string over alphabet  $\Sigma$  is finite sequence of elements of  $\Sigma$

## Example

strings over  $\Sigma = \{0, 1\}$  : 0 0110

## Definitions

- alphabet is finite set; its elements are called symbols or letters
- string over alphabet  $\Sigma$  is finite sequence of elements of  $\Sigma$
- length  $|x|$  of string  $x$  is number of symbols in  $x$

## Example

strings over  $\Sigma = \{0, 1\}$  : 0 0110

## Definitions

- alphabet is finite set; its elements are called symbols or letters
- string over alphabet  $\Sigma$  is finite sequence of elements of  $\Sigma$
- length  $|x|$  of string  $x$  is number of symbols in  $x$
- empty string is unique string of length 0 and denoted by  $\epsilon$

## Example

strings over  $\Sigma = \{0, 1\}$  : 0 0110

## Definitions

- alphabet is finite set; its elements are called symbols or letters
- string over alphabet  $\Sigma$  is finite sequence of elements of  $\Sigma$
- length  $|x|$  of string  $x$  is number of symbols in  $x$
- empty string is unique string of length 0 and denoted by  $\epsilon$
- $\Sigma^*$  is set of all strings over  $\Sigma$  (  $\emptyset^* = \{\epsilon\}$  )

## Example

strings over  $\Sigma = \{0, 1\}$  : 0 0110

## Definitions

- alphabet is finite set; its elements are called symbols or letters
- string over alphabet  $\Sigma$  is finite sequence of elements of  $\Sigma$
- length  $|x|$  of string  $x$  is number of symbols in  $x$
- empty string is unique string of length 0 and denoted by  $\epsilon$
- $\Sigma^*$  is set of all strings over  $\Sigma$  (  $\emptyset^* = \{\epsilon\}$  )
- language over  $\Sigma$  is subset of  $\Sigma^*$

## Example

strings over  $\Sigma = \{0, 1\}$  : 0 0110

languages over  $\Sigma$ :

- $\{\epsilon, 0, 1, 00, 01, 10, 11\}$  (all strings having at most two symbols)



## Definitions

- alphabet is finite set; its elements are called symbols or letters
- string over alphabet  $\Sigma$  is finite sequence of elements of  $\Sigma$
- length  $|x|$  of string  $x$  is number of symbols in  $x$
- empty string is unique string of length 0 and denoted by  $\epsilon$
- $\Sigma^*$  is set of all strings over  $\Sigma$  (  $\emptyset^* = \{\epsilon\}$  )
- language over  $\Sigma$  is subset of  $\Sigma^*$

## Example

strings over  $\Sigma = \{0, 1\}$  : 0 0110

languages over  $\Sigma$ :

- $\{\epsilon, 0, 1, 00, 01, 10, 11\}$  (all strings having at most two symbols)
- $\{x \mid x \text{ is valid program in some machine language}\}$

## Definitions

- string concatenation  $x, y \in \Sigma^* \implies xy \in \Sigma^*$  is associative:

$$(xy)z = x(yz) \quad \forall x, y, z \in \Sigma^*$$

## Definitions

- string concatenation  $x, y \in \Sigma^* \implies xy \in \Sigma^*$  is associative:

$$(xy)z = x(yz) \quad \forall x, y, z \in \Sigma^*$$

- empty string is identity for concatenation:

$$\varepsilon x = x\varepsilon \quad \forall x \in \Sigma^*$$

## Definitions

- string concatenation  $x, y \in \Sigma^* \implies xy \in \Sigma^*$  is associative:

$$(xy)z = x(yz) \quad \forall x, y, z \in \Sigma^*$$

- empty string is identity for concatenation:

$$\varepsilon x = x\varepsilon \quad \forall x \in \Sigma^*$$

- $x$  is substring (prefix, suffix) of  $y$  if  $y = uxv$  ( $y = xv, y = ux$ )

## Definitions

- string concatenation  $x, y \in \Sigma^* \implies xy \in \Sigma^*$  is associative:

$$(xy)z = x(yz) \quad \forall x, y, z \in \Sigma^*$$

- empty string is identity for concatenation:

$$\varepsilon x = x\varepsilon \quad \forall x \in \Sigma^*$$

- $x$  is substring (prefix, suffix) of  $y$  if  $y = uxv$  ( $y = xv, y = ux$ )

- $x^n$  ( $x \in \Sigma^*, n \in \mathbb{N}$ ):

$$\begin{aligned} x^0 &= \varepsilon \\ x^{n+1} &= x^n x \end{aligned}$$

## Definitions

- string concatenation  $x, y \in \Sigma^* \implies xy \in \Sigma^*$  is associative:

$$(xy)z = x(yz) \quad \forall x, y, z \in \Sigma^*$$

- empty string is identity for concatenation:

$$\varepsilon x = x\varepsilon \quad \forall x \in \Sigma^*$$

- $x$  is substring (prefix, suffix) of  $y$  if  $y = uxv$  ( $y = xv, y = ux$ )
- $x^n$  ( $x \in \Sigma^*, n \in \mathbb{N}$ ):

$$\begin{aligned} x^0 &= \varepsilon \\ x^{n+1} &= x^n x \end{aligned}$$

- $\#a(x)$  ( $a \in \Sigma, x \in \Sigma^*$ ) denotes number of  $a$ 's in  $x$

## Definitions ( $A, B \subseteq \Sigma^*$ )

① union

$$A \cup B := \{x \mid x \in A \text{ or } x \in B\}$$

## Definitions ( $A, B \subseteq \Sigma^*$ )

- 1 union  $A \cup B := \{x \mid x \in A \text{ or } x \in B\}$
- 2 intersection  $A \cap B := \{x \mid x \in A \text{ and } x \in B\}$



## Definitions ( $A, B \subseteq \Sigma^*$ )

- |                |  |
|----------------|--|
| ① union        | $A \cup B := \{x \mid x \in A \text{ or } x \in B\}$           |
| ② intersection | $A \cap B := \{x \mid x \in A \text{ and } x \in B\}$          |
| ③ complement   | $\sim A := \Sigma^* - A := \{x \in \Sigma^* \mid x \notin A\}$ |

## Definitions ( $A, B \subseteq \Sigma^*$ )

- |                     |  |
|---------------------|--|
| ① union             | $A \cup B := \{x \mid x \in A \text{ or } x \in B\}$           |
| ② intersection      | $A \cap B := \{x \mid x \in A \text{ and } x \in B\}$          |
| ③ complement        | $\sim A := \Sigma^* - A := \{x \in \Sigma^* \mid x \notin A\}$ |
| ④ set concatenation | $AB := \{xy \mid x \in A \text{ and } y \in B\}$               |

## Definitions ( $A, B \subseteq \Sigma^*$ )

- |                                   |  |
|-----------------------------------|--|
| ① union                           | $A \cup B := \{x \mid x \in A \text{ or } x \in B\}$           |
| ② intersection                    | $A \cap B := \{x \mid x \in A \text{ and } x \in B\}$          |
| ③ complement                      | $\sim A := \Sigma^* - A := \{x \in \Sigma^* \mid x \notin A\}$ |
| ④ set concatenation               | $AB := \{xy \mid x \in A \text{ and } y \in B\}$               |
| ⑤ powers $A^n (n \in \mathbb{N})$ | $A^0 = \{\varepsilon\} \quad A^{n+1} = AA^n$                   |

Definitions ( $A, B \subseteq \Sigma^*$ )

- ① union  $A \cup B := \{x \mid x \in A \text{ or } x \in B\}$
- ② intersection  $A \cap B := \{x \mid x \in A \text{ and } x \in B\}$
- ③ complement  $\sim A := \Sigma^* - A := \{x \in \Sigma^* \mid x \notin A\}$
- ④ set concatenation  $AB := \{xy \mid x \in A \text{ and } y \in B\}$
- ⑤ powers  $A^n (n \in \mathbb{N})$   $A^0 = \{\varepsilon\}$   $A^{n+1} = AA^n$
- ⑥ asterate  $A^*$  is union of all finite powers of  $A$

$$A^* := \bigcup_{n \geq 0} A^n = A^0 \cup A^1 \cup A^2 \cup A^3 \cup \dots = \{x_1 x_2 \dots x_n \text{ and } x_i \in A \text{ for all } 1 \leq i \leq n\}$$

## Definitions ( $A, B \subseteq \Sigma^*$ )

- ① union  $A \cup B := \{x \mid x \in A \text{ or } x \in B\}$
- ② intersection  $A \cap B := \{x \mid x \in A \text{ and } x \in B\}$
- ③ complement  $\sim A := \Sigma^* - A := \{x \in \Sigma^* \mid x \notin A\}$
- ④ set concatenation  $AB := \{xy \mid x \in A \text{ and } y \in B\}$
- ⑤ powers  $A^n (n \in \mathbb{N})$   $A^0 = \{\varepsilon\}$   $A^{n+1} = AA^n$
- ⑥ asterate  $A^*$  is union of all finite powers of  $A$

$$A^* := \bigcup_{n \geq 0} A^n = A^0 \cup A^1 \cup A^2 \cup A^3 \cup \dots = \{x_1 x_2 \dots x_n \text{ and } x_i \in A \text{ for all } 1 \leq i \leq n\}$$

- ⑦ plus  $A^+$  is union of all finite powers of  $A$  except  $\varepsilon$

$$A^+ = AA^* := \bigcup_{n \geq 1} A^n$$

## Definitions ( $A, B \subseteq \Sigma^*$ )

- ① union  $A \cup B := \{x \mid x \in A \text{ or } x \in B\}$
- ② intersection  $A \cap B := \{x \mid x \in A \text{ and } x \in B\}$
- ③ complement  $\sim A := \Sigma^* - A := \{x \in \Sigma^* \mid x \notin A\}$
- ④ set concatenation  $AB := \{xy \mid x \in A \text{ and } y \in B\}$
- ⑤ powers  $A^n (n \in \mathbb{N})$   $A^0 = \{\varepsilon\}$   $A^{n+1} = AA^n$
- ⑥ asterate  $A^*$  is union of all finite powers of  $A$

$$A^* := \bigcup_{n \geq 0} A^n = A^0 \cup A^1 \cup A^2 \cup A^3 \cup \dots = \{x_1 x_2 \dots x_n \text{ and } x_i \in A \text{ for all } 1 \leq i \leq n\}$$

- ⑦ plus  $A^+$  is union of all finite powers of  $A$  except  $\varepsilon$

$$A^+ = AA^* := \bigcup_{n \geq 1} A^n$$

- ⑧ power set  $2^A := \{Q \mid Q \subseteq A\}$

## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$

## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$
- prefixes of 011: 0, 01, 011,  $\epsilon$



## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$
- prefixes of 011: 0, 01, 011,  $\epsilon$
- suffixes of 011: 1, 11, 011,  $\epsilon$

## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$
- prefixes of 011: 0, 01, 011,  $\epsilon$
- suffixes of 011: 1, 11, 011,  $\epsilon$
- $(011)^3 = 011011011 \neq 011^3$

## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$
- prefixes of 011: 0, 01, 011,  $\epsilon$
- suffixes of 011: 1, 11, 011,  $\epsilon$
- $(011)^3 = 011011011 \neq 011^3$
- $\#1(011011011) = 6$      $\#0(\epsilon) = 0$

## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$
- prefixes of 011: 0, 01, 011,  $\epsilon$
- suffixes of 011: 1, 11, 011,  $\epsilon$
- $(011)^3 = 011011011 \neq 011^3$
- $\#1(011011011) = 6$      $\#0(\epsilon) = 0$
- $\{0, 10, 111\}\{1, 11\} = \{01, 101, 1111, 011, 1011, 11111\}$

## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$
- prefixes of 011: 0, 01, 011,  $\epsilon$
- suffixes of 011: 1, 11, 011,  $\epsilon$
- $(011)^3 = 011011011 \neq 011^3$
- $\#1(011011011) = 6$      $\#0(\epsilon) = 0$
- $\{0, 10, 111\}\{1, 11\} = \{01, 101, 1111, 011, 1011, 11111\}$
- $\{0, 01, 111\}\{1, 11\} = \{01, 011, 1111, 0111, 11111\}$

## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$
- prefixes of 011: 0, 01, 011,  $\epsilon$
- suffixes of 011: 1, 11, 011,  $\epsilon$
- $(011)^3 = 011011011 \neq 011^3$
- $\#1(011011011) = 6$      $\#0(\epsilon) = 0$
- $\{0, 10, 111\}\{1, 11\} = \{01, 101, 1111, 011, 1011, 11111\}$
- $\{0, 01, 111\}\{1, 11\} = \{01, 011, 1111, 0111, 11111\}$
- $\{1, 01\}^3 = \{111, 0111, 1011, 01011, 1101, 01101, 10101, 010101\}$

## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$
- prefixes of 011: 0, 01, 011,  $\epsilon$
- suffixes of 011: 1, 11, 011,  $\epsilon$
- $(011)^3 = 011011011 \neq 011^3$
- $\#1(011011011) = 6$      $\#0(\epsilon) = 0$
- $\{0, 10, 111\}\{1, 11\} = \{01, 101, 1111, 011, 1011, 11111\}$
- $\{0, 01, 111\}\{1, 11\} = \{01, 011, 1111, 0111, 11111\}$
- $\{1, 01\}^3 = \{111, 0111, 1011, 01011, 1101, 01101, 10101, 010101\}$
- $\{1, 01\}^* = \{\epsilon, 1, 01, 11, 011, 101, 0101, 111, 0111, 1011, 01011, \dots\}$

## Example

- substrings of 011: 0, 1, 01, 11, 011,  $\epsilon$
- prefixes of 011: 0, 01, 011,  $\epsilon$
- suffixes of 011: 1, 11, 011,  $\epsilon$
- $(011)^3 = 011011011 \neq 011^3$
- $\#1(011011011) = 6$      $\#0(\epsilon) = 0$
- $\{0, 10, 111\}\{1, 11\} = \{01, 101, 1111, 011, 1011, 11111\}$
- $\{0, 01, 111\}\{1, 11\} = \{01, 011, 1111, 0111, 11111\}$
- $\{1, 01\}^3 = \{111, 0111, 1011, 01011, 1101, 01101, 10101, 010101\}$
- $\{1, 01\}^* = \{\epsilon, 1, 01, 11, 011, 101, 0101, 111, 0111, 1011, 01011, \dots\}$
- $2^{\{1, 01\}} = \{\emptyset, \{1\}, \{01\}, \{1, 01\}\}$



## Some Useful Properties

- $\{\epsilon\}A = A\{\epsilon\} = A$

## Some Useful Properties

- $\{\epsilon\}A = A\{\epsilon\} = A$
- $\emptyset A = A\emptyset = \emptyset$

## Some Useful Properties

- $\{\epsilon\}A = A\{\epsilon\} = A$
- $\emptyset A = A\emptyset = \emptyset$
- $\sim (A \cup B) = (\sim A) \cap (\sim B)$

## Some Useful Properties

- $\{\epsilon\}A = A\{\epsilon\} = A$
- $\emptyset A = A\emptyset = \emptyset$
- $\sim (A \cup B) = (\sim A) \cap (\sim B)$
- $\sim (A \cap B) = (\sim A) \cup (\sim B)$

## Some Useful Properties

- $\{\epsilon\}A = A\{\epsilon\} = A$
- $\emptyset A = A\emptyset = \emptyset$
- $\sim (A \cup B) = (\sim A) \cap (\sim B)$
- $\sim (A \cap B) = (\sim A) \cup (\sim B)$
- $A^{m+n} = A^m A^n$

## Some Useful Properties

- $\{\epsilon\}A = A\{\epsilon\} = A$
- $\emptyset A = A\emptyset = \emptyset$
- $\sim (A \cup B) = (\sim A) \cap (\sim B)$
- $\sim (A \cap B) = (\sim A) \cup (\sim B)$
- $A^{m+n} = A^m A^n$
- $A^* A^* = A^*$

## Some Useful Properties

- $\{\epsilon\}A = A\{\epsilon\} = A$
- $\emptyset A = A\emptyset = \emptyset$
- $\sim (A \cup B) = (\sim A) \cap (\sim B)$
- $\sim (A \cap B) = (\sim A) \cup (\sim B)$
- $A^{m+n} = A^m A^n$
- $A^* A^* = A^*$
- $A^{**} = A^*$

## Some Useful Properties

- $\{\epsilon\}A = A\{\epsilon\} = A$
- $\emptyset A = A\emptyset = \emptyset$
- $\sim (A \cup B) = (\sim A) \cap (\sim B)$
- $\sim (A \cap B) = (\sim A) \cup (\sim B)$
- $A^{m+n} = A^m A^n$
- $A^* A^* = A^*$
- $A^{**} = A^*$
- $A^* = \{\epsilon\} \cup AA^* = \{\epsilon\} \cup A^* A$






## Some Useful Properties



- $\{\epsilon\}A = A\{\epsilon\} = A$
- $\emptyset A = A\emptyset = \emptyset$
- $\sim(A \cup B) = (\sim A) \cap (\sim B)$
- $\sim(A \cap B) = (\sim A) \cup (\sim B)$
- $A^{m+n} = A^m A^n$
- $A^* A^* = A^*$
- $A^{**} = A^*$
- $A^* = \{\epsilon\} \cup AA^* = \{\epsilon\} \cup A^* A$
- $\emptyset^* = \{\epsilon\}$


# Outline


- 1 A Quick Recap
- 2 Chomsky Hierarchy
- 3 Deterministic Finite State Automata
- 4 Closure Properties
















# Outline

- 1 A Quick Recap
- 2 Chomsky Hierarchy
- 3 Deterministic Finite State Automata**
- 4 Closure Properties

## Definitions

- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with

## Definitions

- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with
  - 1  $Q$  : finite set of states

## Definitions


- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with
  - 1  $Q$  : finite set of states
  - 2  $\Sigma$  : input alphabet

## Definitions

- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with
  - ①  $Q$ : finite set of states
  - ②  $\Sigma$ : input alphabet
  - ③  $\delta: Q \times \Sigma \rightarrow Q$ : transition function


## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



①  $Q = \{1, 2, 3, 4\}$

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




①  $Q = \{1, 2, 3, 4\}$

②  $\Sigma = \{a, b\}$



## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

## Definitions


- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with
  - ①  $Q$ : finite set of states
  - ②  $\Sigma$ : input alphabet
  - ③  $\delta: Q \times \Sigma \rightarrow Q$ : transition function
  - ④  $s \in Q$ : start state

## Definitions

- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with
  - ①  $Q$ : finite set of states
  - ②  $\Sigma$ : input alphabet
  - ③  $\delta: Q \times \Sigma \rightarrow Q$ : transition function
  - ④  $s \in Q$ : start state
  - ⑤  $F \subseteq Q$ : final (accept) states

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	$a$	$b$
1	2	1
2	2	3
3	4	1
4	4	4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	$a$	$b$
1	2	1
2	2	3
3	4	1
4	4	4



## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

1    1    2    3    4    4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	$a$	$b$
1	2	1
2	2	3
3	4	1
4	4	4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	$a$	$b$
1	2	1
2	2	3
3	4	1
4	4	4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	$a$	$b$
1	2	1
2	2	3
3	4	1
4	4	4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

b a b a a  
1 1 2 3 4 4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	$a$	$b$
1	2	1
2	2	3
3	4	1
4	4	4

1    1    2    3    4    4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	$a$	$b$
1	2	1
2	2	3
3	4	1
4	4	4



## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

$b$     $a$     $b$     $a$     $a$   
 1   1   2   3   4   4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	$a$	$b$
1	2	1
2	2	3
3	4	1
4	4	4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

1 1 2 3 4 4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$


$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

b a b a a

1 1 2 3 4 4

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	$a$	$b$
1	2	1
2	2	3
3	4	1
4	4	4

## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$



## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

	b	a	b	a	a
1	1	2	3	4	4
	a	a	b	b	b
1	2	2	3	1	1

## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$


$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

	b	a	b	a	a
1	1	2	3	4	4
	a	a	b	b	b
1	2	2	3	1	1



## Example (DFAs → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

## Definitions

- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with

- ①  $Q$  : finite set of states
- ②  $\Sigma$  : input alphabet
- ③  $\delta : Q \times \Sigma \rightarrow Q$  : transition function
- ④  $s \in Q$  : start state
- ⑤  $F \subseteq Q$  : final (accept) states

- $\hat{\delta} : Q \times \Sigma^* \rightarrow Q$  is inductively defined by

$$\hat{\delta}(q, \varepsilon) := q$$

$$\hat{\delta}(q, xa) := \delta(\hat{\delta}(q, x), a)$$

### Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$$\delta(\widehat{\delta}(q_0, abbaa), b)$$

first recursive call

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$$\delta(\widehat{\delta}(q_0, abbaa), b)$$

first recursive call

$$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$$

second recursive call

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$$\delta(\widehat{\delta}(q_0, abbaa), b)$$

first recursive call

$$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$$

second recursive call

$$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$$

third recursive call

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$$\delta(\widehat{\delta}(q_0, abbaa), b)$$

first recursive call

$$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$$

second recursive call

$$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$$

third recursive call

$$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$$

fourth recursive call



## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$\delta(\widehat{\delta}(q_0, abbaa), b)$

first recursive call

$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$

second recursive call

$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$

third recursive call

$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$

fourth recursive call

$\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, a), b), b), a), a), b)$

fifth recursive call

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$\delta(\widehat{\delta}(q_0, abbaa), b)$	first recursive call
$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$	second recursive call
$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$	third recursive call
$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$	fourth recursive call
$\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, a), b), b), a), a), b)$	fifth recursive call
$\delta(\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, \epsilon), a), b), b), a), a), b)$	sixth recursive call

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$\delta(\widehat{\delta}(q_0, abbaa), b)$

first recursive call

$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$

second recursive call

$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$

third recursive call

$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$

fourth recursive call

$\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, a), b), b), a), a), b)$

fifth recursive call

$\delta(\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, \epsilon), a), b), b), a), a), b)$

sixth recursive call

$\delta(\delta(\delta(\delta(\delta(\delta(\delta(q_0, a), b), b), a), a), b)$

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$\delta(\widehat{\delta}(q_0, abbaa), b)$	first recursive call
$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$	second recursive call
$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$	third recursive call
$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$	fourth recursive call
$\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, a), b), b), a), a), b)$	fifth recursive call
$\delta(\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, \epsilon), a), b), b), a), a), b)$	sixth recursive call
$\delta(\delta(\delta(\delta(\delta(q_0, a), b), b), a), a), b)$	
$\delta(\delta(\delta(\delta(q_1, b), b), a), a), b)$	assuming $\delta(q_0, a) = q_1$

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$\delta(\widehat{\delta}(q_0, abbaa), b)$

first recursive call

$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$

second recursive call

$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$

third recursive call

$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$

fourth recursive call

$\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, a), b), b), a), a), b)$

fifth recursive call

$\delta(\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, \epsilon), a), b), b), a), a), b)$

sixth recursive call

$\delta(\delta(\delta(\delta(\delta(\delta(q_0, a), b), b), a), a), b)$

$\delta(\delta(\delta(\delta(q_1, b), b), a), a), b)$

assuming  $\delta(q_0, a) = q_1$

$\delta(\delta(\delta(q_2, b), a), a), b)$

assuming  $\delta(q_1, b) = q_2$

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$\delta(\widehat{\delta}(q_0, abbaa), b)$

first recursive call

$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$

second recursive call

$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$

third recursive call

$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$

fourth recursive call

$\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, a), b), b), a), a), b)$

fifth recursive call

$\delta(\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, \epsilon), a), b), b), a), a), b)$

sixth recursive call

$\delta(\delta(\delta(\delta(\delta(\delta(q_0, a), b), b), a), a), b)$

$\delta(\delta(\delta(\delta(q_1, b), b), a), a), b)$

assuming  $\delta(q_0, a) = q_1$

$\delta(\delta(\delta(q_2, b), a), a), b)$

assuming  $\delta(q_1, b) = q_2$

$\delta(\delta(\delta(q_3, a), a), b)$

assuming  $\delta(q_2, b) = q_3$

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$\delta(\widehat{\delta}(q_0, abbaa), b)$	first recursive call
$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$	second recursive call
$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$	third recursive call
$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$	fourth recursive call
$\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, a), b), b), a), a), b)$	fifth recursive call
$\delta(\delta(\delta(\delta(\delta(\delta(q_0, \epsilon), a), b), b), a), a), b)$	sixth recursive call
$\delta(\delta(\delta(\delta(\delta(q_0, a), b), b), a), a), b)$	
$\delta(\delta(\delta(\delta(q_1, b), b), a), a), b)$	assuming $\delta(q_0, a) = q_1$
$\delta(\delta(\delta(\delta(q_2, b), a), a), b)$	assuming $\delta(q_1, b) = q_2$
$\delta(\delta(\delta(q_3, a), a), b)$	assuming $\delta(q_2, b) = q_3$
$\delta(\delta(q_4, a), b)$	assuming $\delta(q_3, a) = q_4$

## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$\delta(\widehat{\delta}(q_0, abbaa), b)$	first recursive call
$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$	second recursive call
$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$	third recursive call
$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$	fourth recursive call
$\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, a), b), b), a), a), b)$	fifth recursive call
$\delta(\delta(\delta(\delta(\delta(\delta(q_0, \epsilon), a), b), b), a), a), b)$	sixth recursive call
$\delta(\delta(\delta(\delta(\delta(q_0, a), b), b), a), a), b)$	
$\delta(\delta(\delta(\delta(q_1, b), b), a), a), b)$	assuming $\delta(q_0, a) = q_1$
$\delta(\delta(\delta(\delta(q_2, b), a), a), b)$	assuming $\delta(q_1, b) = q_2$
$\delta(\delta(\delta(q_3, a), a), b)$	assuming $\delta(q_2, b) = q_3$
$\delta(\delta(q_4, a), b)$	assuming $\delta(q_3, a) = q_4$
$\delta(q_5, b)$	assuming $\delta(q_4, a) = q_5$



## Example (Unfolding of the multistep function $\widehat{\delta}$ )

Let  $x = abbaab$  over the alphabet  $\Sigma = \{a, b\}$

$\delta(\widehat{\delta}(q_0, abbaa), b)$	first recursive call
$\delta(\delta(\widehat{\delta}(q_0, abba), a), b)$	second recursive call
$\delta(\delta(\delta(\widehat{\delta}(q_0, abb), a), a), b)$	third recursive call
$\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, ab), b), a), a), b)$	fourth recursive call
$\delta(\delta(\delta(\delta(\delta(\widehat{\delta}(q_0, a), b), b), a), a), b)$	fifth recursive call
$\delta(\delta(\delta(\delta(\delta(\delta(q_0, \epsilon), a), b), b), a), a), b)$	sixth recursive call
$\delta(\delta(\delta(\delta(\delta(q_0, a), b), b), a), a), b)$	
$\delta(\delta(\delta(\delta(q_1, b), b), a), a), b)$	assuming $\delta(q_0, a) = q_1$
$\delta(\delta(\delta(\delta(q_2, b), a), a), b)$	assuming $\delta(q_1, b) = q_2$
$\delta(\delta(\delta(q_3, a), a), b)$	assuming $\delta(q_2, b) = q_3$
$\delta(\delta(q_4, a), b)$	assuming $\delta(q_3, a) = q_4$
$\delta(q_5, b)$	assuming $\delta(q_4, a) = q_5$
$q_6$	assuming $\delta(q_5, b) = q_6$

## Definitions

- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with

- ①  $Q$  : finite set of states
- ②  $\Sigma$  : input alphabet
- ③  $\delta : Q \times \Sigma \rightarrow Q$  : transition function
- ④  $s \in Q$  : start state
- ⑤  $F \subseteq Q$  : final (accept) states

- $\hat{\delta} : Q \times \Sigma^* \rightarrow Q$  is inductively defined by

$$\hat{\delta}(q, \varepsilon) := q$$

$$\hat{\delta}(q, xa) := \delta(\hat{\delta}(q, x), a)$$

- string  $x \in \Sigma^*$  is **accepted** by  $M$  if  $\hat{\delta}(s, x) \in F$

## Definitions

- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with

- ①  $Q$  : finite set of states
- ②  $\Sigma$  : input alphabet
- ③  $\delta : Q \times \Sigma \rightarrow Q$  : transition function
- ④  $s \in Q$  : start state
- ⑤  $F \subseteq Q$  : final (accept) states

- $\hat{\delta} : Q \times \Sigma^* \rightarrow Q$  is inductively defined by


$$\hat{\delta}(q, \varepsilon) := q$$

$$\hat{\delta}(q, xa) := \delta(\hat{\delta}(q, x), a)$$

- string  $x \in \Sigma^*$  is **accepted** by  $M$  if  $\hat{\delta}(s, x) \in F$
- string  $x \in \Sigma^*$  is **rejected** by  $M$  if  $\hat{\delta}(s, x) \notin F$

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

	b	a	b	a	a	$\in L(M)$
1	1	2	3	4	4	
	a	a	b	b	b	$\notin L(M)$
1	2	2	3	1	1	

## Definitions

- **deterministic finite automaton (DFA)** is quintuple  $M = (Q, \Sigma, \delta, s, F)$  with

- ①  $Q$  : finite set of states
- ②  $\Sigma$  : input alphabet
- ③  $\delta : Q \times \Sigma \rightarrow Q$  : transition function
- ④  $s \in Q$  : start state
- ⑤  $F \subseteq Q$  : final (accept) states

- $\hat{\delta} : Q \times \Sigma^* \rightarrow Q$  is inductively defined by


$$\hat{\delta}(q, \varepsilon) := q$$

$$\hat{\delta}(q, xa) := \delta(\hat{\delta}(q, x), a)$$

- string  $x \in \Sigma^*$  is **accepted** by  $M$  if  $\hat{\delta}(s, x) \in F$
- string  $x \in \Sigma^*$  is **rejected** by  $M$  if  $\hat{\delta}(s, x) \notin F$
- language accepted by  $M$  is given by  $L(M) := \{x \mid \hat{\delta}(s, x) \in F\}$

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$

$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4


	b	a	b	a	a	$\in L(M)$
1	1	2	3	4	4	
	a	a	b	b	b	$\notin L(M)$
1	2	2	3	1	1	

$L(M) := \{x \mid$

$\}$

## Example (DFAs → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3, 4\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{4\}$


$\delta$	a	b
1	2	1
2	2	3
3	4	1
4	4	4

	b	a	b	a	a	$\in L(M)$
1	1	2	3	4	4	
	a	a	b	b	b	$\notin L(M)$
1	2	2	3	1	1	

$L(M) := \{x \mid x \text{ contains } aba \text{ as substring}\}$

## Example (DFA $\rightarrow$ Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$





## Example (DFA $\rightarrow$ Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



①  $Q = \{1, 2, 3\}$

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




①  $Q = \{1, 2, 3\}$

②  $\Sigma = \{a, b\}$

## Example (DFA $\rightarrow$ Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$

$\delta$	$a$	$b$
1	1	2
2	3	3
3	3	3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

## Example (DFA $\rightarrow$ Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

## Example (DFA $\rightarrow$ Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

a a a a b  
1 1 1 1 1 2

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$


$\delta$	a	b
1	1	2
2	3	3
3	3	3

a a a a b  
1 1 1 1 1 2



## Example (DFA $\rightarrow$ Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$


$\delta$	a	b
1	1	2
2	3	3
3	3	3

a    a    a    a    b

1    1    1    1    1    2

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

a a a a b  
1 1 1 1 1 2

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

a a a a b  
1 1 1 1 1 2

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

## Example (DFA $\rightarrow$ Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

## Example (DFA → Regular Sets)


$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

a a a a b  
1 1 1 1 1 2

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$


$\delta$	a	b
1	1	2
2	3	3
3	3	3

a a a a b  
1 1 1 1 1 2



## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

a a a a b  
1 1 1 1 1 2

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

a a a a b  
1 1 1 1 1 2

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	1	2
	a	a	b	b	a	
1	1	1	1	2	3	3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

$a \quad a \quad a \quad a \quad b \quad \in L(M)$   
 1    1    1    1    1    2  
 $a \quad a \quad b \quad b \quad a$   
 1    1    1    2    3    3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	1	2
2	a	a	b	b	a	
3	1	1	1	2	3	3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

$a \quad a \quad a \quad a \quad b \quad \in L(M)$   
 1    1    1    1    1    2  
 $a \quad a \quad b \quad b \quad a$   
 1    1    2    3    3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$


$\delta$	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	1	2
2	a	a	b	b	a	
3	1	1	1	2	3	3



## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

$a \quad a \quad a \quad a \quad b \quad \in L(M)$   
 1    1    1    1    1    2  
 $a \quad a \quad b \quad b \quad a$   
 1    1    1    2    3    3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	1	2
	a	a	b	b	a	
1	1	1	1	2	3	3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

$a \quad a \quad a \quad a \quad b \quad \in L(M)$   
 1    1    1    1    1    2  
 $a \quad a \quad b \quad b \quad a$   
 1    1    1    2    3    3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	1	2
	a	a	b	b	a	
1	1	1	2	3	3	

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

$a \quad a \quad a \quad a \quad b \quad \in L(M)$   
 1    1    1    1    1    2  
 $a \quad a \quad b \quad b \quad a$   
 1    1    1    2    3    3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	1	2
	a	a	b	b	a	
1	1	1	1	2	3	3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$


$\delta$	a	b
1	1	2
2	3	3
3	3	3

a   a   a   a   b    $\in L(M)$

1	1	1	1	1	2
	a	a	b	b	a
1	1	1	2	3	3

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$


$\delta$	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	1	2
	a	a	b	b	a	$\notin L(M)$
1	1	1	2	3	3	



## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$


$\delta$	$a$	$b$
1	1	2
2	3	3
3	3	3

	$a$	$a$	$a$	$a$	$b$	$\in L(M)$
1	1	1	1	1	1	2
	$a$	$a$	$b$	$b$	$a$	$\notin L(M)$
1	1	1	1	2	3	3

$L(M) := \{x \mid \dots\}$

## Example (DFA → Regular Sets)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{2\}$

$\delta$	a	b
1	1	2
2	3	3
3	3	3

	a	a	a	a	b	$\in L(M)$
1	1	1	1	1	1	2
	a	a	b	b	a	$\notin L(M)$
1	1	1	2	3	3	

$$L(M) := \{x \mid x = a^n b, n \geq 0\}$$

## Theorem

The DFA  $M$  is correct with respect to predefined specs.

## Theorem

The DFA  $M$  is correct with respect to predefined specs. Namely,  $M$  accepts every string of the form  $a^n b$  s.t.  $n \in \mathbb{N}$ , rejecting all others.

## Theorem

The DFA  $M$  is correct with respect to predefined specs. Namely,  $M$  accepts every string of the form  $a^n b$  s.t.  $n \in \mathbb{N}$ , rejecting all others.

Formally: 
$$\hat{\delta}(1, x) = \begin{cases} 1 & \iff x \in L(a^*) \\ 2 & \iff x \in L(a^* b) \\ 3 & \iff x \in L(a^* b(a + b)^+) \end{cases}$$

## Theorem

The DFA  $M$  is correct with respect to predefined specs. Namely,  $M$  accepts every string of the form  $a^n b$  s.t.  $n \in \mathbb{N}$ , rejecting all others.

Formally: 
$$\hat{\delta}(1, x) = \begin{cases} 1 & \iff x \in L(a^*) \\ 2 & \iff x \in L(a^* b) \\ 3 & \iff x \in L(a^* b(a + b)^+) \end{cases}$$

## Proof.

We argue by mathematical induction on the length of  $x$ .

## Theorem

The DFA  $M$  is correct with respect to predefined specs. Namely,  $M$  accepts every string of the form  $a^n b$  s.t.  $n \in \mathbb{N}$ , rejecting all others.

Formally: 
$$\hat{\delta}(1, x) = \begin{cases} 1 & \iff x \in L(a^*) \\ 2 & \iff x \in L(a^* b) \\ 3 & \iff x \in L(a^* b(a+b)^+) \end{cases}$$

## Proof.

We argue by mathematical induction on the length of  $x$ .

① Base Case:  $|x| = 0 \iff x = \epsilon \quad \hat{\delta}(1, \epsilon) = 1 \iff \epsilon \in L(a^*)$

## Theorem

The DFA  $M$  is correct with respect to predefined specs. Namely,  $M$  accepts every string of the form  $a^n b$  s.t.  $n \in \mathbb{N}$ , rejecting all others.

Formally: 
$$\hat{\delta}(1, x) = \begin{cases} 1 & \iff x \in L(a^*) \\ 2 & \iff x \in L(a^* b) \\ 3 & \iff x \in L(a^* b(a+b)^+) \end{cases}$$

## Proof.

We argue by mathematical induction on the length of  $x$ .

- ① Base Case:  $|x| = 0 \iff x = \epsilon \quad \hat{\delta}(1, \epsilon) = 1 \iff \epsilon \in L(a^*)$
- ② Step Case:      Given IH :  $M$  is correct on every  $x \in \Sigma^*$       such that  $|x| = k$  with  $k \geq 0$   
                     Show :  $M$  is correct on every  $xy$  for all  $y \in \Sigma = \{a, b\}$       such that  $|xy| = k + 1$



## Theorem

The DFA  $M$  is correct with respect to predefined specs. Namely,  $M$  accepts every string of the form  $a^n b$  s.t.  $n \in \mathbb{N}$ , rejecting all others.

Formally: 
$$\hat{\delta}(1, x) = \begin{cases} 1 & \iff x \in L(a^*) \\ 2 & \iff x \in L(a^* b) \\ 3 & \iff x \in L(a^* b(a+b)^+) \end{cases}$$

## Proof.

We argue by mathematical induction on the length of  $x$ .

① Base Case:  $|x| = 0 \iff x = \varepsilon \quad \hat{\delta}(1, \varepsilon) = 1 \iff \varepsilon \in L(a^*)$

② Step Case: Given IH :  $M$  is correct on every  $x \in \Sigma^*$  such that  $|x| = k$  with  $k \geq 0$   
Show :  $M$  is correct on every  $xy$  for all  $y \in \Sigma = \{a, b\}$  such that  $|xy| = k + 1$

IH : 
$$\hat{\delta}(1, x) = \begin{cases} 1 & \iff x \in L(a^*) \\ 2 & \iff x \in L(a^* b) \\ 3 & \iff x \in L(a^* b(a+b)^+) \end{cases}$$

Formally:

Show : 
$$\hat{\delta}(1, xy) = \begin{cases} 1 & \iff xy \in L(a^*) \\ 2 & \iff xy \in L(a^* b) \\ 3 & \iff xy \in L(a^* b(a+b)^+) \end{cases}$$

## Proof. (cont'd)

①  $\widehat{\delta}(1, x) = 1$  and  $y = a$

## Proof. (cont'd)

$$\begin{array}{llll} \textcircled{1} & \widehat{\delta}(1, x) = 1 & \text{and} & y = a \\ & \widehat{\delta}(1, x) = 1 & \iff & x \in L(a^*) \text{ (by IH)} \end{array} \qquad \delta(\widehat{\delta}(1, x), a) = 1 \iff xa \in L(a^*)$$

## Proof. (cont'd)

$$\textcircled{1} \quad \begin{array}{ll} \widehat{\delta}(1, x) = 1 & \text{and} \quad y = a \\ \widehat{\delta}(1, x) = 1 & \iff x \in L(a^*) \text{ (by IH)} \end{array} \quad \delta(\widehat{\delta}(1, x), a) = 1 \iff xa \in L(a^*)$$

$$\textcircled{2} \quad \widehat{\delta}(1, x) = 1 \quad \text{and} \quad y = b$$

## Proof. (cont'd)

- ①  $\hat{\delta}(1, x) = 1$  and  $y = a$   
 $\hat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\hat{\delta}(1, x), a) = 1 \iff xa \in L(a^*)$
- ②  $\hat{\delta}(1, x) = 1$  and  $y = b$   
 $\hat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\hat{\delta}(1, x), b) = 2 \iff xb \in L(a^*b)$

## Proof. (cont'd)

- ①  $\hat{\delta}(1, x) = 1$  and  $y = a$   
 $\hat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\hat{\delta}(1, x), a) = 1 \iff xa \in L(a^*)$
- ②  $\hat{\delta}(1, x) = 1$  and  $y = b$   
 $\hat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\hat{\delta}(1, x), b) = 2 \iff xb \in L(a^*b)$
- ③  $\hat{\delta}(1, x) = 2$  and  $y = a$

## Proof. (cont'd)

- |   |                          |        |                 |         |                                     |        |                         |
|---|--------------------------|--------|-----------------|---------|-------------------------------------|--------|-------------------------|
| ① | $\hat{\delta}(1, x) = 1$ | and    | $y = a$         |         | $\delta(\hat{\delta}(1, x), a) = 1$ | $\iff$ | $xa \in L(a^*)$         |
|   | $\hat{\delta}(1, x) = 1$ | $\iff$ | $x \in L(a^*)$  | (by IH) |                                     |        |                         |
| ② | $\hat{\delta}(1, x) = 1$ | and    | $y = b$         |         | $\delta(\hat{\delta}(1, x), b) = 2$ | $\iff$ | $xb \in L(a^*b)$        |
|   | $\hat{\delta}(1, x) = 1$ | $\iff$ | $x \in L(a^*)$  | (by IH) |                                     |        |                         |
| ③ | $\hat{\delta}(1, x) = 2$ | and    | $y = a$         |         | $\delta(\hat{\delta}(1, x), a) = 3$ | $\iff$ | $xa \in L(a^*b(a+b)^+)$ |
|   | $\hat{\delta}(1, x) = 2$ | $\iff$ | $x \in L(a^*b)$ | (by IH) |                                     |        |                         |

## Proof. (cont'd)

- ①  $\widehat{\delta}(1, x) = 1$  and  $y = a$   
 $\widehat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)

$\delta(\widehat{\delta}(1, x), a) = 1 \iff xa \in L(a^*)$
- ②  $\widehat{\delta}(1, x) = 1$  and  $y = b$   
 $\widehat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)

$\delta(\widehat{\delta}(1, x), b) = 2 \iff xb \in L(a^*b)$
- ③  $\widehat{\delta}(1, x) = 2$  and  $y = a$   
 $\widehat{\delta}(1, x) = 2 \iff x \in L(a^*b)$  (by IH)

$\delta(\widehat{\delta}(1, x), a) = 3 \iff xa \in L(a^*b(a+b)^+)$
- ④  $\widehat{\delta}(1, x) = 2$  and  $y = b$



# Proof. (cont'd)

- |   |                          |        |                 |         |                                     |        |                         |
|---|--------------------------|--------|-----------------|---------|-------------------------------------|--------|-------------------------|
| ① | $\hat{\delta}(1, x) = 1$ | and    | $y = a$         |         | $\delta(\hat{\delta}(1, x), a) = 1$ | $\iff$ | $xa \in L(a^*)$         |
|   | $\hat{\delta}(1, x) = 1$ | $\iff$ | $x \in L(a^*)$  | (by IH) |                                     |        |                         |
| ② | $\hat{\delta}(1, x) = 1$ | and    | $y = b$         |         | $\delta(\hat{\delta}(1, x), b) = 2$ | $\iff$ | $xb \in L(a^*b)$        |
|   | $\hat{\delta}(1, x) = 1$ | $\iff$ | $x \in L(a^*)$  | (by IH) |                                     |        |                         |
| ③ | $\hat{\delta}(1, x) = 2$ | and    | $y = a$         |         | $\delta(\hat{\delta}(1, x), a) = 3$ | $\iff$ | $xa \in L(a^*b(a+b)^+)$ |
|   | $\hat{\delta}(1, x) = 2$ | $\iff$ | $x \in L(a^*b)$ | (by IH) |                                     |        |                         |
| ④ | $\hat{\delta}(1, x) = 2$ | and    | $y = b$         |         | $\delta(\hat{\delta}(1, x), b) = 3$ | $\iff$ | $xb \in L(a^*b(a+b)^+)$ |
|   | $\hat{\delta}(1, x) = 2$ | $\iff$ | $x \in L(a^*b)$ | (by IH) |                                     |        |                         |

## Proof. (cont'd)

- ①  $\hat{\delta}(1, x) = 1$  and  $y = a$   
 $\hat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\hat{\delta}(1, x), a) = 1 \iff xa \in L(a^*)$
- ②  $\hat{\delta}(1, x) = 1$  and  $y = b$   
 $\hat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\hat{\delta}(1, x), b) = 2 \iff xb \in L(a^*b)$
- ③  $\hat{\delta}(1, x) = 2$  and  $y = a$   
 $\hat{\delta}(1, x) = 2 \iff x \in L(a^*b)$  (by IH)  $\delta(\hat{\delta}(1, x), a) = 3 \iff xa \in L(a^*b(a+b)^+)$
- ④  $\hat{\delta}(1, x) = 2$  and  $y = b$   
 $\hat{\delta}(1, x) = 2 \iff x \in L(a^*b)$  (by IH)  $\delta(\hat{\delta}(1, x), b) = 3 \iff xb \in L(a^*b(a+b)^+)$
- ⑤  $\hat{\delta}(1, x) = 3$  and  $y = a$

## Proof. (cont'd)

- ①  $\widehat{\delta}(1, x) = 1$  and  $y = a$   
 $\widehat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\widehat{\delta}(1, x), a) = 1 \iff xa \in L(a^*)$
- ②  $\widehat{\delta}(1, x) = 1$  and  $y = b$   
 $\widehat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\widehat{\delta}(1, x), b) = 2 \iff xb \in L(a^*b)$
- ③  $\widehat{\delta}(1, x) = 2$  and  $y = a$   
 $\widehat{\delta}(1, x) = 2 \iff x \in L(a^*b)$  (by IH)  $\delta(\widehat{\delta}(1, x), a) = 3 \iff xa \in L(a^*b(a+b)^+)$
- ④  $\widehat{\delta}(1, x) = 2$  and  $y = b$   
 $\widehat{\delta}(1, x) = 2 \iff x \in L(a^*b)$  (by IH)  $\delta(\widehat{\delta}(1, x), b) = 3 \iff xb \in L(a^*b(a+b)^+)$
- ⑤  $\widehat{\delta}(1, x) = 3$  and  $y = a$   
 $\widehat{\delta}(1, x) = 3 \iff x \in L(a^*b(a+b)^+)$  (by IH)  $\delta(\widehat{\delta}(1, x), a) = 3 \iff xa \in L(a^*b(a+b)^+)$

## Proof. (cont'd)

- ①  $\widehat{\delta}(1, x) = 1$  and  $y = a$   
 $\widehat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\widehat{\delta}(1, x), a) = 1 \iff xa \in L(a^*)$
- ②  $\widehat{\delta}(1, x) = 1$  and  $y = b$   
 $\widehat{\delta}(1, x) = 1 \iff x \in L(a^*)$  (by IH)  $\delta(\widehat{\delta}(1, x), b) = 2 \iff xb \in L(a^*b)$
- ③  $\widehat{\delta}(1, x) = 2$  and  $y = a$   
 $\widehat{\delta}(1, x) = 2 \iff x \in L(a^*b)$  (by IH)  $\delta(\widehat{\delta}(1, x), a) = 3 \iff xa \in L(a^*b(a+b)^+)$
- ④  $\widehat{\delta}(1, x) = 2$  and  $y = b$   
 $\widehat{\delta}(1, x) = 2 \iff x \in L(a^*b)$  (by IH)  $\delta(\widehat{\delta}(1, x), b) = 3 \iff xb \in L(a^*b(a+b)^+)$
- ⑤  $\widehat{\delta}(1, x) = 3$  and  $y = a$   
 $\widehat{\delta}(1, x) = 3 \iff x \in L(a^*b(a+b)^+)$  (by IH)  $\delta(\widehat{\delta}(1, x), a) = 3 \iff xa \in L(a^*b(a+b)^+)$
- ⑥  $\widehat{\delta}(1, x) = 3$  and  $y = b$

## Proof. (cont'd)

- |   |  |     |         |  |
|---|--|-----|---------|--|
| ① | $\hat{\delta}(1, x) = 1$<br>$\hat{\delta}(1, x) = 1 \iff x \in L(a^*)$ (by IH)         | and | $y = a$ | $\delta(\hat{\delta}(1, x), a) = 1 \iff xa \in L(a^*)$         |
| ② | $\hat{\delta}(1, x) = 1$<br>$\hat{\delta}(1, x) = 1 \iff x \in L(a^*)$ (by IH)         | and | $y = b$ | $\delta(\hat{\delta}(1, x), b) = 2 \iff xb \in L(a^*b)$        |
| ③ | $\hat{\delta}(1, x) = 2$<br>$\hat{\delta}(1, x) = 2 \iff x \in L(a^*b)$ (by IH)        | and | $y = a$ | $\delta(\hat{\delta}(1, x), a) = 3 \iff xa \in L(a^*b(a+b)^+)$ |
| ④ | $\hat{\delta}(1, x) = 2$<br>$\hat{\delta}(1, x) = 2 \iff x \in L(a^*b)$ (by IH)        | and | $y = b$ | $\delta(\hat{\delta}(1, x), b) = 3 \iff xb \in L(a^*b(a+b)^+)$ |
| ⑤ | $\hat{\delta}(1, x) = 3$<br>$\hat{\delta}(1, x) = 3 \iff x \in L(a^*b(a+b)^+)$ (by IH) | and | $y = a$ | $\delta(\hat{\delta}(1, x), a) = 3 \iff xa \in L(a^*b(a+b)^+)$ |
| ⑥ | $\hat{\delta}(1, x) = 3$<br>$\hat{\delta}(1, x) = 3 \iff x \in L(a^*b(a+b)^+)$ (by IH) | and | $y = b$ | $\delta(\hat{\delta}(1, x), b) = 3 \iff xb \in L(a^*b(a+b)^+)$ |



## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

②  $\Sigma = \{a, b\}$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

①  $Q =$

②  $\Sigma = \{a, b\}$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

①  $Q =$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$



## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

①  $Q =$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

④  $s =$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

①  $Q =$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

④  $s =$

⑤  $F =$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

$M \text{ start} \rightarrow \textcircled{1}$

①  $Q =$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

④  $s =$

⑤  $F =$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

$M \text{ start} \rightarrow \textcircled{1} \qquad \textcircled{2}$

①  $Q =$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$


④  $s =$

⑤  $F =$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



①  $Q =$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$


④  $s =$

⑤  $F =$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$



## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



①  $Q = \{1, 2\}$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$


④  $s = 1$

⑤  $F = \{1\}$

$\delta$	$a$	$b$
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$$M = (Q, \Sigma, \delta, s, F)$$


①  $Q = \{1, 2\}$

②  $\Sigma = \{a, b\}$

$$\textcircled{3} \delta : Q \times \Sigma \rightarrow Q$$

④  $s = 1$

⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

a	b	a	b	a	
1	1	2	2	1	1

$$L(M) := \{x \mid x \text{ contains even number of } b\text{'s over } \Sigma\}$$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	<span style="color: blue;">a</span>	<span style="color: blue;">b</span>
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	<span style="color: blue;">a</span>	<span style="color: blue;">b</span>
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



a b a b a

1 1 2 2 1 1


- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	a	b
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	<span style="color: blue;">a</span>	<span style="color: blue;">b</span>
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



①  $Q = \{1, 2\}$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

④  $s = 1$

⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$



## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

a   b   a   b   a  
 1   1   2   2   1   1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	a	b
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	<span style="color: blue;">a</span>	<span style="color: blue;">b</span>
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	a	b
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	a	b
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

a    b    a    b    a  
 1    1    2    2    1    1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



a b a b a ∈ L(M)

1 1 2 2 1 1


- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	a	b
1	1	2
2	2	1

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



①  $Q = \{1, 2\}$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

④  $s = 1$

⑤  $F = \{1\}$

$\delta$	a	b
1	1	2
2	2	1


	a	b	a	b	a	
	$\in L(M)$					
1	1	2	2	1	1	
	b	a	a	a	a	
1	2	2	2	2	2	

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$



## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



①  $Q = \{1, 2\}$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

④  $s = 1$

⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



①  $Q = \{1, 2\}$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

④  $s = 1$

⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



①  $Q = \{1, 2\}$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

④  $s = 1$

⑤  $F = \{1\}$

$\delta$	a	b
1	1	2
2	2	1


$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$



## Example (Regular Sets → DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



①  $Q = \{1, 2\}$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$

④  $s = 1$

⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$


$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
 1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a$   
 1    2    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{1\}$

$\delta$	a	b
1	1	2
2	2	1

$a \quad b \quad a \quad b \quad a \quad \in L(M)$   
           1    2    2    1    1  
 $b \quad a \quad a \quad a \quad a \quad \notin L(M)$   
           1    2    2    2    2

$L(M) := \{x \mid x \text{ contains even number of } bs \text{ over } \Sigma\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

②  $\Sigma = \{a, b\}$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

①  $Q =$

②  $\Sigma = \{a, b\}$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$



## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$

- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$


$M$  start  $\rightarrow$  (1)

- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



①  $Q =$

②  $\Sigma = \{a, b\}$

③  $\delta : Q \times \Sigma \rightarrow Q$


④  $s =$

⑤  $F =$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$$



## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q =$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s =$
- ⑤  $F =$

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$




- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$

$\delta$	a	b
1	1	2
2	1	3
3	3	3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b   a   b   b   a  
 1   2   1   2   3   3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b a b b a  
 1 2 1 2 3 3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b   a   b   b   a  
 1   2   1   2   3   3

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b   a   b   b   a  
 1   2   1   2   3   3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$

$\delta$	a	b
1	1	2
2	1	3
3	3	3


b   a   b   b   a  
 1   2   1   2   3   3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$$



## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b   a   b   b   a  
 1   2   1   2   3   3

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b   a   b   b   a  
 1   2   1   2   3   3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b   a   b   b   a  
 1   2   1   2   3   3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b   a   b   b   a  
 1   2   1   2   3   3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b   a   b   b   a  
 1   2   1   2   3   3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

1    2    1    2    3    3

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

b   a   b   b   a  
 1   2   1   2   3   3

$$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$

$\delta$	a	b
1	1	2
2	1	3
3	3	3


$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring}\}$



## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
     a    a    b    a    b  
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$

$\delta$	a	b
1	1	2
2	1	3
3	3	3


$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$



## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
   a    a    b    a    b  
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta: Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$


$\delta$	a	b
1	1	2
2	1	3
3	3	3

$b \quad a \quad b \quad b \quad a \quad \in L(M)$   
 1    2    1    2    3    3  
 $a \quad a \quad b \quad a \quad b$   
 1    1    1    2    1    2

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

## Example (Regular Sets $\rightarrow$ DFA)

$M = (Q, \Sigma, \delta, s, F)$



- ①  $Q = \{1, 2, 3\}$
- ②  $\Sigma = \{a, b\}$
- ③  $\delta : Q \times \Sigma \rightarrow Q$
- ④  $s = 1$
- ⑤  $F = \{3\}$

$\delta$	a	b
1	1	2
2	1	3
3	3	3

	b	a	b	b	a	$\in L(M)$
1	2	1	2	3	3	
	a	a	b	a	b	$\notin L(M)$
1	1	1	2	1	2	

$L(M) := \{x \mid x \text{ contains } bb \text{ as substring} \}$

# Outline

- 1 A Quick Recap
- 2 Chomsky Hierarchy
- 3 Deterministic Finite State Automata
- 4 Closure Properties

## Theorem

regular sets are **effectively closed** under **intersection**

## Theorem

regular sets are **effectively closed** under **intersection**

Proof. (closure under intersection)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$



## Theorem

regular sets are **effectively closed** under **intersection**

Proof. (closure under intersection)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cap B := L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$

## Theorem

regular sets are **effectively closed** under **intersection**

Proof. (closure under intersection)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cap B := L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 := Q_1 \times Q_2 = \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$

## Theorem

regular sets are **effectively closed** under **intersection**

Proof. (closure under intersection)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cap B := L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 := Q_1 \times Q_2 = \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 := F_1 \times F_2$

## Theorem

regular sets are **effectively closed** under **intersection**

### Proof. (closure under intersection)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cap B := L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 := Q_1 \times Q_2 = \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 := F_1 \times F_2$
  - ③  $s_3 := (s_1, s_2)$

## Theorem

regular sets are **effectively closed** under **intersection**

### Proof. (closure under intersection)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cap B := L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 := Q_1 \times Q_2 = \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 := F_1 \times F_2$
  - ③  $s_3 := (s_1, s_2)$
  - ④  $\delta_3((p, q), a) := (\delta_1(p, a), \delta_2(q, a)) \quad \forall p \in Q_1, \forall q \in Q_2, \forall a \in \Sigma$

## Theorem

regular sets are **effectively closed** under **intersection**

### Proof. (closure under intersection)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cap B := L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 := Q_1 \times Q_2 = \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 := F_1 \times F_2$
  - ③  $s_3 := (s_1, s_2)$
  - ④  $\delta_3((p, q), a) := (\delta_1(p, a), \delta_2(q, a)) \quad \forall p \in Q_1, \forall q \in Q_2, \forall a \in \Sigma$
- claim:  $\widehat{\delta_3}((p, q), x) = (\widehat{\delta_1}(p, x), \widehat{\delta_2}(q, x)) \quad \forall x \in \Sigma^*$

## Theorem

regular sets are **effectively closed** under **intersection**

### Proof. (closure under intersection)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cap B := L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 := Q_1 \times Q_2 = \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 := F_1 \times F_2$
  - ③  $s_3 := (s_1, s_2)$
  - ④  $\delta_3((p, q), a) := (\delta_1(p, a), \delta_2(q, a)) \quad \forall p \in Q_1, \forall q \in Q_2, \forall a \in \Sigma$
- claim:  $\widehat{\delta_3}((p, q), x) = (\widehat{\delta_1}(p, x), \widehat{\delta_2}(q, x)) \quad \forall x \in \Sigma^*$   
 proof: induction on  $|x|$  next slide

## proof of the claim

claim:  $\widehat{\delta_3}((p, q), x) = (\widehat{\delta_1}(p, x), \widehat{\delta_2}(q, x)) \quad \forall x \in \Sigma^*$



## proof of the claim

claim:  $\widehat{\delta}_3((p, q), x) = (\widehat{\delta}_1(p, x), \widehat{\delta}_2(q, x)) \quad \forall x \in \Sigma^*$

- base case:  $|x| = 0$  thus  $x = \epsilon$

$$\widehat{\delta}_3((p, q), \epsilon) = (p, q) = (\widehat{\delta}_1(p, \epsilon), \widehat{\delta}_2(q, \epsilon))$$

## proof of the claim

claim:  $\widehat{\delta}_3((p, q), x) = (\widehat{\delta}_1(p, x), \widehat{\delta}_2(q, x)) \quad \forall x \in \Sigma^*$

- base case:  $|x| = 0$  thus  $x = \varepsilon$

$$\widehat{\delta}_3((p, q), \varepsilon) = (p, q) = (\widehat{\delta}_1(p, \varepsilon), \widehat{\delta}_2(q, \varepsilon))$$

- step case:  $|x| > 0$  thus  $x = ya$  s.t.  $|y| = |x| - 1$  with IH:  $\widehat{\delta}_3((p, q), y) = (\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y))$

## proof of the claim

claim:  $\widehat{\delta}_3((p, q), x) = (\widehat{\delta}_1(p, x), \widehat{\delta}_2(q, x)) \quad \forall x \in \Sigma^*$

- base case:  $|x| = 0$  thus  $x = \varepsilon$

$$\widehat{\delta}_3((p, q), \varepsilon) = (p, q) = (\widehat{\delta}_1(p, \varepsilon), \widehat{\delta}_2(q, \varepsilon))$$

- step case:  $|x| > 0$  thus  $x = ya$  s.t.  $|y| = |x| - 1$  with IH:  $\widehat{\delta}_3((p, q), y) = (\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y))$

$$\widehat{\delta}_3((p, q), ya) = \delta_3(\widehat{\delta}_3((p, q), y), a) \quad \text{(by definition of } \widehat{\delta}_3)$$

## proof of the claim

claim:  $\widehat{\delta}_3((p, q), x) = (\widehat{\delta}_1(p, x), \widehat{\delta}_2(q, x)) \quad \forall x \in \Sigma^*$

- base case:  $|x| = 0$  thus  $x = \varepsilon$

$$\widehat{\delta}_3((p, q), \varepsilon) = (p, q) = (\widehat{\delta}_1(p, \varepsilon), \widehat{\delta}_2(q, \varepsilon))$$

- step case:  $|x| > 0$  thus  $x = ya$  s.t.  $|y| = |x| - 1$  with IH:  $\widehat{\delta}_3((p, q), y) = (\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y))$

$$\begin{aligned} \widehat{\delta}_3((p, q), ya) &= \delta_3(\widehat{\delta}_3((p, q), y), a) && \text{(by definition of } \widehat{\delta}_3) \\ &= \delta_3((\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y)), a) && \text{(by induction hypothesis IH)} \end{aligned}$$

## proof of the claim

claim:  $\widehat{\delta}_3((p, q), x) = (\widehat{\delta}_1(p, x), \widehat{\delta}_2(q, x)) \quad \forall x \in \Sigma^*$

- base case:  $|x| = 0$  thus  $x = \varepsilon$

$$\widehat{\delta}_3((p, q), \varepsilon) = (p, q) = (\widehat{\delta}_1(p, \varepsilon), \widehat{\delta}_2(q, \varepsilon))$$

- step case:  $|x| > 0$  thus  $x = ya$  s.t.  $|y| = |x| - 1$  with IH:  $\widehat{\delta}_3((p, q), y) = (\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y))$

$$\begin{aligned} \widehat{\delta}_3((p, q), ya) &= \delta_3(\widehat{\delta}_3((p, q), y), a) && \text{(by definition of } \widehat{\delta}_3) \\ &= \delta_3((\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y)), a) && \text{(by induction hypothesis IH)} \\ &= (\delta_1(\widehat{\delta}_1(p, y), a), \delta_2(\widehat{\delta}_2(q, y), a)) && \text{(by definition of } \delta_3) \end{aligned}$$

## proof of the claim

claim:  $\widehat{\delta}_3((p, q), x) = (\widehat{\delta}_1(p, x), \widehat{\delta}_2(q, x)) \quad \forall x \in \Sigma^*$

- base case:  $|x| = 0$  thus  $x = \varepsilon$

$$\widehat{\delta}_3((p, q), \varepsilon) = (p, q) = (\widehat{\delta}_1(p, \varepsilon), \widehat{\delta}_2(q, \varepsilon))$$

- step case:  $|x| > 0$  thus  $x = ya$  s.t.  $|y| = |x| - 1$  with IH:  $\widehat{\delta}_3((p, q), y) = (\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y))$

$$\begin{aligned} \widehat{\delta}_3((p, q), ya) &= \delta_3(\widehat{\delta}_3((p, q), y), a) && \text{(by definition of } \widehat{\delta}_3) \\ &= \delta_3((\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y)), a) && \text{(by induction hypothesis IH)} \\ &= (\delta_1(\widehat{\delta}_1(p, y), a), \delta_2(\widehat{\delta}_2(q, y), a)) && \text{(by definition of } \delta_3) \\ &= (\widehat{\delta}_1(p, ya), \widehat{\delta}_2(q, ya)) && \text{(by definitions of } \widehat{\delta}_1 \text{ and } \widehat{\delta}_2) \end{aligned}$$

## proof of the claim

claim:  $\widehat{\delta}_3((p, q), x) = (\widehat{\delta}_1(p, x), \widehat{\delta}_2(q, x)) \quad \forall x \in \Sigma^*$

- base case:  $|x| = 0$  thus  $x = \varepsilon$

$$\widehat{\delta}_3((p, q), \varepsilon) = (p, q) = (\widehat{\delta}_1(p, \varepsilon), \widehat{\delta}_2(q, \varepsilon))$$

- step case:  $|x| > 0$  thus  $x = ya$  s.t.  $|y| = |x| - 1$  with IH:  $\widehat{\delta}_3((p, q), y) = (\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y))$

$$\begin{aligned} \widehat{\delta}_3((p, q), ya) &= \delta_3(\widehat{\delta}_3((p, q), y), a) && \text{(by definition of } \widehat{\delta}_3) \\ &= \delta_3((\widehat{\delta}_1(p, y), \widehat{\delta}_2(q, y)), a) && \text{(by induction hypothesis IH)} \\ &= (\delta_1(\widehat{\delta}_1(p, y), a), \delta_2(\widehat{\delta}_2(q, y), a)) && \text{(by definition of } \delta_3) \\ &= (\widehat{\delta}_1(p, ya), \widehat{\delta}_2(q, ya)) && \text{(by definitions of } \widehat{\delta}_1 \text{ and } \widehat{\delta}_2) \\ &= (\widehat{\delta}_1(p, x), \widehat{\delta}_2(q, x)) \end{aligned}$$



## Proof. (closure under intersection (cont'd))

statement:  $L(M_3) = L(M_1) \cap L(M_2)$



### Proof. (closure under intersection (cont'd))

statement:  $L(M_3) = L(M_1) \cap L(M_2)$

$$\forall x \in \Sigma^*, x \in L(M_3) \iff \widehat{\delta_3}(s_3, x) \in F_3 \quad \text{(by definition of acceptance)}$$

## Proof. (closure under intersection (cont'd))

statement:  $L(M_3) = L(M_1) \cap L(M_2)$

$$\begin{aligned} \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 && \text{(by definition of acceptance)} \\ &\iff \widehat{\delta}_3((s_1, s_2), x) \in F_1 \times F_2 && \text{(by definition of } s_3 \text{ and } F_3) \end{aligned}$$

## Proof. (closure under intersection (cont'd))

statement:  $L(M_3) = L(M_1) \cap L(M_2)$

$$\begin{aligned}
 \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 && \text{(by definition of acceptance)} \\
 &\iff \widehat{\delta}_3((s_1, s_2), x) \in F_1 \times F_2 && \text{(by definition of } s_3 \text{ and } F_3) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in F_1 \times F_2 && \text{(by claim proven in slide 21)}
 \end{aligned}$$

## Proof. (closure under intersection (cont'd))

statement:  $L(M_3) = L(M_1) \cap L(M_2)$

$$\begin{aligned}
 \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 && \text{(by definition of acceptance)} \\
 &\iff \widehat{\delta}_3((s_1, s_2), x) \in F_1 \times F_2 && \text{(by definition of } s_3 \text{ and } F_3) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in F_1 \times F_2 && \text{(by claim proven in slide 21)} \\
 &\iff \widehat{\delta}_1(s_1, x) \in F_1 \text{ and } \widehat{\delta}_2(s_2, x) \in F_2 && \text{(by definition of product)}
 \end{aligned}$$

**Proof. (closure under intersection (cont'd))**statement:  $L(M_3) = L(M_1) \cap L(M_2)$ 

$$\begin{aligned} \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 && \text{(by definition of acceptance)} \\ &\iff \widehat{\delta}_3((s_1, s_2), x) \in F_1 \times F_2 && \text{(by definition of } s_3 \text{ and } F_3) \\ &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in F_1 \times F_2 && \text{(by claim proven in slide 21)} \\ &\iff \widehat{\delta}_1(s_1, x) \in F_1 \text{ and } \widehat{\delta}_2(s_2, x) \in F_2 && \text{(by definition of product)} \\ &\iff x \in L(M_1) \text{ and } x \in L(M_2) && \text{(by definition of acceptance)} \end{aligned}$$



## Proof. (closure under intersection (cont'd))

statement:  $L(M_3) = L(M_1) \cap L(M_2)$ 

$$\begin{aligned} \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 && \text{(by definition of acceptance)} \\ &\iff \widehat{\delta}_3((s_1, s_2), x) \in F_1 \times F_2 && \text{(by definition of } s_3 \text{ and } F_3) \\ &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in F_1 \times F_2 && \text{(by claim proven in slide 21)} \\ &\iff \widehat{\delta}_1(s_1, x) \in F_1 \text{ and } \widehat{\delta}_2(s_2, x) \in F_2 && \text{(by definition of product)} \\ &\iff x \in L(M_1) \text{ and } x \in L(M_2) && \text{(by definition of acceptance)} \\ &\iff x \in L(M_1) \cap L(M_2) && \text{(by definition of intersection)} \end{aligned}$$






## Example (intersection)



$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$

## Example (intersection)






$$A = (1, 1)$$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$






## Example (intersection)



$$A = (1, 1)$$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$   
 $E = (1, 2)$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$   
 $E = (1, 2)$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$

## Example (intersection)






$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$   
 $E = (1, 2)$   
 $F = (2, 3)$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$






## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$   
 $E = (1, 2)$   
 $F = (2, 3)$

$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$   
 $E = (1, 2)$   
 $F = (2, 3)$

$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$   
 $E = (1, 2)$   
 $F = (2, 3)$

$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$   
 $E = (1, 2)$   
 $F = (2, 3)$

$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$




## Example (intersection)



$A = (1, 1)$   
 $B = (2, 2)$   
 $C = (2, 1)$   
 $D = (1, 3)$   
 $E = (1, 2)$   
 $F = (2, 3)$

$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$

## Example (intersection)



- $A = (1, 1)$
- $B = (2, 2)$
- $C = (2, 1)$
- $D = (1, 3)$
- $E = (1, 2)$
- $F = (2, 3)$

$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$

## Theorem

regular sets are **effectively closed** under **complement**

## Theorem

regular sets are **effectively closed** under **complement**

Proof. (closure under complement)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$



## Theorem

regular sets are **effectively closed** under **complement**

Proof. (closure under complement)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $\sim A := \Sigma^* - A$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$

## Theorem

regular sets are **effectively closed** under **complement**

Proof. (closure under complement)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $\sim A := \Sigma^* - A$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$

$$\textcircled{1} Q_2 := Q_1$$

## Theorem

regular sets are **effectively closed** under **complement**

Proof. (closure under complement)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $\sim A := \Sigma^* - A$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$

$$\textcircled{1} Q_2 := Q_1$$

$$\textcircled{2} F_2 := Q_1 - F_1$$

## Theorem

regular sets are **effectively closed** under **complement**

### Proof. (closure under complement)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $\sim A := \Sigma^* - A$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$

- ①  $Q_2 := Q_1$
- ②  $F_2 := Q_1 - F_1$
- ③  $s_2 := s_1$

## Theorem

regular sets are **effectively closed** under **complement**

### Proof. (closure under complement)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $\sim A := \Sigma^* - A$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$

- 1  $Q_2 := Q_1$
- 2  $F_2 := Q_1 - F_1$
- 3  $s_2 := s_1$
- 4  $\delta_2(p, a) := \delta_1(p, a) \quad \forall p \in Q_1, \forall a \in \Sigma$

## Theorem

regular sets are **effectively closed** under **complement**

### Proof. (closure under complement)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $\sim A := \Sigma^* - A$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$ 
  - 1  $Q_2 := Q_1$
  - 2  $F_2 := Q_1 - F_1$
  - 3  $s_2 := s_1$
  - 4  $\delta_2(p, a) := \delta_1(p, a) \quad \forall p \in Q_1, \forall a \in \Sigma$
- obvious claim:  $\widehat{\delta_2}(p, x) = \widehat{\delta_1}(p, x) \quad \forall x \in \Sigma^*$

## Proof. (closure under complement (cont'd))

statement:  $L(M_2) = \Sigma^* - L(M_1)$

### Proof. (closure under complement (cont'd))

statement:  $L(M_2) = \Sigma^* - L(M_1)$

$$\forall x \in \Sigma^*, x \in L(M_2) \iff \widehat{\delta_2}(s_2, x) \in F_2 \quad \text{(by definition of acceptance)}$$



## Proof. (closure under complement (cont'd))

statement:  $L(M_2) = \Sigma^* - L(M_1)$

$$\begin{aligned} \forall x \in \Sigma^*, x \in L(M_2) &\iff \widehat{\delta_2}(s_2, x) \in F_2 && \text{(by definition of acceptance)} \\ &\iff \widehat{\delta_1}(s_2, x) \notin F_1 && \text{(by the obvious claim in slide 24)} \end{aligned}$$

## Proof. (closure under complement (cont'd))

statement:  $L(M_2) = \Sigma^* - L(M_1)$

$$\begin{aligned}
 \forall x \in \Sigma^*, x \in L(M_2) &\iff \widehat{\delta}_2(s_2, x) \in F_2 && \text{(by definition of acceptance)} \\
 &\iff \widehat{\delta}_1(s_2, x) \in F_2 && \text{(by the obvious claim in slide 24)} \\
 &\iff \widehat{\delta}_1(s_1, x) \in Q_1 - F_1 && \text{(by definitions of } s_2 \text{ and } F_2)
 \end{aligned}$$

## Proof. (closure under complement (cont'd))

statement:  $L(M_2) = \Sigma^* - L(M_1)$

$$\begin{aligned}
 \forall x \in \Sigma^*, x \in L(M_2) &\iff \widehat{\delta}_2(s_2, x) \in F_2 && \text{(by definition of acceptance)} \\
 &\iff \widehat{\delta}_1(s_2, x) \in F_2 && \text{(by the obvious claim in slide 24)} \\
 &\iff \widehat{\delta}_1(s_1, x) \in Q_1 - F_1 && \text{(by definitions of } s_2 \text{ and } F_2) \\
 &\iff \widehat{\delta}_1(s_1, x) \in Q_1 \text{ and } \widehat{\delta}_1(s_1, x) \notin F_1 && \text{(by definition of set difference)}
 \end{aligned}$$



## Proof. (closure under complement (cont'd))

statement:  $L(M_2) = \Sigma^* - L(M_1)$




$$\begin{array}{llll}
 \forall x \in \Sigma^*, x \in L(M_2) & \iff & \widehat{\delta}_2(s_2, x) \in F_2 & \text{(by definition of acceptance)} \\
 & \iff & \widehat{\delta}_1(s_2, x) \in F_2 & \text{(by the obvious claim in slide 24)} \\
 & \iff & \widehat{\delta}_1(s_1, x) \in Q_1 - F_1 & \text{(by definitions of } s_2 \text{ and } F_2) \\
 & \iff & \widehat{\delta}_1(s_1, x) \in Q_1 \text{ and } \widehat{\delta}_1(s_1, x) \notin F_1 & \text{(by definition of set difference)} \\
 & \iff & x \notin L(M_1) & \text{(by definition of acceptance)}
 \end{array}$$

□

## Example (complement)







## Example (complement)



$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$

## Example (complement)



$\sim L(M_3) := \{x \mid x \text{ contains odd number of } bs \text{ or no } bb \text{ as substring}\}$

## Theorem

regular sets are **effectively closed** under **union**



## Theorem

regular sets are **effectively closed** under **union**

Proof. (closure under union)

$$A \cup B = \sim ((\sim A) \cap (\sim B))$$

## Theorem

regular sets are **effectively closed** under **union**

## Theorem

regular sets are **effectively closed** under **union**

Proof. (closure under union – explicit construction)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$

## Theorem

regular sets are **effectively closed** under **union**

Proof. (closure under union – explicit construction)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cup B = L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$

## Theorem

regular sets are **effectively closed** under **union**

Proof. (closure under union – explicit construction)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cup B = L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 = Q_1 \times Q_2 := \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$

## Theorem

regular sets are **effectively closed** under **union**

Proof. (closure under union – explicit construction)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cup B = L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 = Q_1 \times Q_2 \quad := \quad \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 \quad := \quad (F_1 \times Q_2) \cup (Q_1 \times F_2)$

## Theorem

regular sets are **effectively closed** under **union**

Proof. (closure under union – explicit construction)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$
- $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cup B = L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 = Q_1 \times Q_2 \quad := \quad \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 \quad := \quad (F_1 \times Q_2) \cup (Q_1 \times F_2)$
  - ③  $s_3 \quad := \quad (s_1, s_2)$

## Theorem

regular sets are **effectively closed** under **union**

Proof. (closure under union – explicit construction)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cup B = L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 = Q_1 \times Q_2 := \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 := (F_1 \times Q_2) \cup (Q_1 \times F_2)$
  - ③  $s_3 := (s_1, s_2)$
  - ④  $\delta_3((p, q), a) := (\delta_1(p, a), \delta_2(q, a)) \quad \forall p \in Q_1, \forall q \in Q_2, \forall a \in \Sigma$



## Theorem

regular sets are **effectively closed** under **union**

Proof. (closure under union – explicit construction)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cup B = L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 = Q_1 \times Q_2 \quad := \quad \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 \quad := \quad (F_1 \times Q_2) \cup (Q_1 \times F_2)$
  - ③  $s_3 \quad := \quad (s_1, s_2)$
  - ④  $\delta_3((p, q), a) \quad := \quad (\delta_1(p, a), \delta_2(q, a)) \quad \forall p \in Q_1, \forall q \in Q_2, \forall a \in \Sigma$
- claim:  $\widehat{\delta_3}((p, q), x) = (\widehat{\delta_1}(p, x), \widehat{\delta_2}(q, x)) \quad \forall x \in \Sigma^*$

## Theorem

regular sets are **effectively closed** under **union**

Proof. (closure under union – explicit construction)

- $A = L(M_1)$  for DFA  $M_1 = (Q_1, \Sigma, \delta_1, s_1, F_1)$   
 $B = L(M_2)$  for DFA  $M_2 = (Q_2, \Sigma, \delta_2, s_2, F_2)$
- $A \cup B = L(M_3)$  for DFA  $M_3 = (Q_3, \Sigma, \delta_3, s_3, F_3)$ 
  - ①  $Q_3 = Q_1 \times Q_2 \quad := \quad \{(p, q) \mid p \in Q_1 \text{ and } q \in Q_2\}$
  - ②  $F_3 \quad := \quad (F_1 \times Q_2) \cup (Q_1 \times F_2)$
  - ③  $s_3 \quad := \quad (s_1, s_2)$
  - ④  $\delta_3((p, q), a) \quad := \quad (\delta_1(p, a), \delta_2(q, a)) \quad \forall p \in Q_1, \forall q \in Q_2, \forall a \in \Sigma$
- claim:  $\widehat{\delta_3}((p, q), x) = (\widehat{\delta_1}(p, x), \widehat{\delta_2}(q, x)) \quad \forall x \in \Sigma^*$
- proof: induction on  $|x|$  – skipped (follows exact same steps with that is given at slide #21)

## Proof. (closure under union – explicit construction (cont'd))

statement:  $L(M_3) = L(M_1) \cup L(M_2)$

### Proof. (closure under union – explicit construction (cont'd))

statement:  $L(M_3) = L(M_1) \cup L(M_2)$

$\forall x \in \Sigma^*, x \in L(M_3) \iff \widehat{\delta}_3(s_3, x) \in F_3$

## Proof. (closure under union – explicit construction (cont'd))

statement:  $L(M_3) = L(M_1) \cup L(M_2)$

$$\begin{aligned} \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 \\ &\iff \widehat{\delta}_3((s_1, s_2), x) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \end{aligned}$$

**Proof. (closure under union – explicit construction (cont'd))**

statement:  $L(M_3) = L(M_1) \cup L(M_2)$

$$\begin{aligned}\forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 \\ &\iff \widehat{\delta}_3((s_1, s_2), x) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \\ &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (F_1 \times Q_2) \cup (Q_1 \times F_2)\end{aligned}$$

## Proof. (closure under union – explicit construction (cont'd))

statement:  $L(M_3) = L(M_1) \cup L(M_2)$

$$\begin{aligned}
 \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 \\
 &\iff \widehat{\delta}_3((s_1, s_2), x) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (F_1 \times Q_2) \text{ or } (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (Q_1 \times F_2)
 \end{aligned}$$

## Proof. (closure under union – explicit construction (cont'd))

statement:  $L(M_3) = L(M_1) \cup L(M_2)$

$$\begin{aligned}
 \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 \\
 &\iff \widehat{\delta}_3((s_1, s_2), x) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (F_1 \times Q_2) \text{ or } (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (Q_1 \times F_2) \\
 &\iff \left( \widehat{\delta}_1(s_1, x) \in F_1 \text{ and } \widehat{\delta}_2(s_2, x) \in Q_2 \right) \text{ or } \left( \widehat{\delta}_1(s_1, x) \in Q_1 \text{ and } \widehat{\delta}_2(s_2, x) \in F_2 \right)
 \end{aligned}$$



## Proof. (closure under union – explicit construction (cont'd))

statement:  $L(M_3) = L(M_1) \cup L(M_2)$

$$\begin{aligned}
 \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 \\
 &\iff \widehat{\delta}_3((s_1, s_2), x) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (F_1 \times Q_2) \text{ or } (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (Q_1 \times F_2) \\
 &\iff \left( \widehat{\delta}_1(s_1, x) \in F_1 \text{ and } \widehat{\delta}_2(s_2, x) \in Q_2 \right) \text{ or } \left( \widehat{\delta}_1(s_1, x) \in Q_1 \text{ and } \widehat{\delta}_2(s_2, x) \in F_2 \right) \\
 &\iff x \in L(M_1) \text{ or } x \in L(M_2)
 \end{aligned}$$



# Proof. (closure under union – explicit construction (cont'd))

statement:  $L(M_3) = L(M_1) \cup L(M_2)$




$$\begin{aligned}
 \forall x \in \Sigma^*, x \in L(M_3) &\iff \widehat{\delta}_3(s_3, x) \in F_3 \\
 &\iff \widehat{\delta}_3((s_1, s_2), x) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (F_1 \times Q_2) \cup (Q_1 \times F_2) \\
 &\iff (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (F_1 \times Q_2) \text{ or } (\widehat{\delta}_1(s_1, x), \widehat{\delta}_2(s_2, x)) \in (Q_1 \times F_2) \\
 &\iff \left( \widehat{\delta}_1(s_1, x) \in F_1 \text{ and } \widehat{\delta}_2(s_2, x) \in Q_2 \right) \text{ or } \left( \widehat{\delta}_1(s_1, x) \in Q_1 \text{ and } \widehat{\delta}_2(s_2, x) \in F_2 \right) \\
 &\iff x \in L(M_1) \text{ or } x \in L(M_2) \\
 &\iff x \in L(M_1) \cup L(M_2)
 \end{aligned}$$



## Example (union)







## Example (union)



$$L(M_1) \cap L(M_2) = L(M_3) := \{x \mid x \text{ contains even number of } bs \text{ and } bb \text{ as substring}\}$$

## Example (union)



$$L(M_1) \cup L(M_2) = L(M_4) := \{x \mid x \text{ contains even number of } bs \text{ or } bb \text{ as substring}\}$$

Thanks! & Questions?