

# CMPE 322/327 - Theory of Computation

## Week 9: Chomsky Normal Form & Pumping Lemma – CKY Algorithm

Burak Ekici

April 18-22, 2022

# Outline

- 1 A Quick Recap
- 2 Chomsky Normal Form
- 3 Pumping Lemma
- 4 CKY Algorithm

## New Algorithm for Testing Equivalence of NFAs

## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence


## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)

## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)
- first: DFAs (by example)


### Example



## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)
- first: DFAs (by example)


### Example



## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)
- first: DFAs (by example)

### Example






## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)
- first: DFAs (by example)


### Example



## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)
- first: DFAs (by example)


### Example



## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)
- first: DFAs (by example)

### Example




$1 \sim 4$      $1 \uparrow$      $4 \uparrow$   
 $2 \sim 5$      $2 \downarrow$      $5 \downarrow$   
 $3 \sim 4 ?$

## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)
- first: DFAs (by example)

### Example




1 ~ 4	1↑	4↑
2 ~ 5	2↓	5↓
3 ~ 4 ?	3↑	4↑

## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)
- first: DFAs (by example)

### Example




1 ~ 4	1↑	4↑
2 ~ 5	2↓	5↓
3 ~ 4	3↑	4↑
2 ~ 5		

## New Algorithm for Testing Equivalence of NFAs

- bisimulation up to congruence
- developed by F. Bonchi and D. Pous (POPL 2013, CACM 2015)
- first: DFAs (by example)

### Example



$1 \sim 4$

$2 \sim 5$

$3 \sim 4$

$2 \sim 5$

$1 \uparrow \quad 4 \uparrow$

$2 \downarrow \quad 5 \downarrow$

$3 \uparrow \quad 4 \uparrow$

$\sim$  is **bisimulation**

## Remark

bisimulation relates states with same observable behaviour

## Remark

bisimulation relates states with same observable behaviour

## Definition

- **bisimulation** is binary relation  $R$  on states  $Q$  of DFA  $M = (Q, \Sigma, \delta, s, F)$  such that for all  $pRq$ 
  - 1  $p \in F \iff q \in F$



## Remark

bisimulation relates states with same observable behaviour

## Definition

- **bisimulation** is binary relation  $R$  on states  $Q$  of DFA  $M = (Q, \Sigma, \delta, s, F)$  such that for all  $p R q$ 
  - ①  $p \in F \iff q \in F$
  - ②  $\delta(p, a) R \delta(q, a)$  for all  $a \in \Sigma$

## Remark

bisimulation relates states with same observable behaviour

## Definition

- **bisimulation** is binary relation  $R$  on states  $Q$  of DFA  $M = (Q, \Sigma, \delta, s, F)$  such that for all  $p R q$ 
  - ①  $p \in F \iff q \in F$
  - ②  $\delta(p, a) R \delta(q, a)$  for all  $a \in \Sigma$
- states  $p$  and  $q$  are **bisimilar** ( $p \sim q$ ) if  $p R q$  for some bisimulation  $R$

## Remark

bisimulation relates states with same observable behaviour

## Definition

- **bisimulation** is binary relation  $R$  on states  $Q$  of DFA  $M = (Q, \Sigma, \delta, s, F)$  such that for all  $p R q$ 
  - ①  $p \in F \iff q \in F$
  - ②  $\delta(p, a) R \delta(q, a)$  for all  $a \in \Sigma$
- states  $p$  and  $q$  are **bisimilar** ( $p \sim q$ ) if  $p R q$  for some bisimulation  $R$

## Example

- $=$  (identity relation)

## Remark

bisimulation relates states with same observable behaviour

## Definition

- **bisimulation** is binary relation  $R$  on states  $Q$  of DFA  $M = (Q, \Sigma, \delta, s, F)$  such that for all  $p R q$ 
  - ①  $p \in F \iff q \in F$
  - ②  $\delta(p, a) R \delta(q, a)$  for all  $a \in \Sigma$
- states  $p$  and  $q$  are **bisimilar** ( $p \sim q$ ) if  $p R q$  for some bisimulation  $R$

## Example

- $=$  (identity relation)
- $\approx$  (indistinguishability relation of lecture 5)

## Remark

bisimulation relates states with same observable behaviour

## Definition

- **bisimulation** is binary relation  $R$  on states  $Q$  of DFA  $M = (Q, \Sigma, \delta, s, F)$  such that for all  $p R q$ 
  - ①  $p \in F \iff q \in F$
  - ②  $\delta(p, a) R \delta(q, a)$  for all  $a \in \Sigma$
- states  $p$  and  $q$  are **bisimilar** ( $p \sim q$ ) if  $p R q$  for some bisimulation  $R$

## Remarks

- bisimilarity is equivalence relation

## Remark

bisimulation relates states with same observable behaviour

## Definition

- **bisimulation** is binary relation  $R$  on states  $Q$  of DFA  $M = (Q, \Sigma, \delta, s, F)$  such that for all  $p R q$ 
  - ①  $p \in F \iff q \in F$
  - ②  $\delta(p, a) R \delta(q, a)$  for all  $a \in \Sigma$
- states  $p$  and  $q$  are **bisimilar** ( $p \sim q$ ) if  $p R q$  for some bisimulation  $R$

## Remarks

- bisimilarity is equivalence relation
- $L(M, p) := \{x \in \Sigma^* \mid \hat{\delta}(p, x) \in F\}$

## Remark

bisimulation relates states with same observable behaviour


## Definition

- **bisimulation** is binary relation  $R$  on states  $Q$  of DFA  $M = (Q, \Sigma, \delta, s, F)$  such that for all  $p R q$ 
  - ①  $p \in F \iff q \in F$
  - ②  $\delta(p, a) R \delta(q, a)$  for all  $a \in \Sigma$
- states  $p$  and  $q$  are **bisimilar** ( $p \sim q$ ) if  $p R q$  for some bisimulation  $R$

## Remarks


- bisimilarity is equivalence relation
- $L(M, p) := \{x \in \Sigma^* \mid \hat{\delta}(p, x) \in F\}$
- $p \sim q \iff L(M, p) \sim L(M, q)$

## Example






## Example




## Example




$a \sim f$


## Example




## Example




## Example




## Example



## Example




## Example







## Example




## Example




## Example




## Example




## Example




## Example



## Example




## Example







## Example




## Example




## Example



## Example



## Example



$a \sim f$   
 $c \sim i$   
 $e \sim h$

$b \sim g$   
 $d \sim j$   
 $a \sim i$

$c \sim h$   
 $e \sim g$   
 $b \sim j$


$d \sim i$   
 $a \sim h$   
 $c \sim g$

$a$   
 $e \sim j$   
 $b \sim i$   
 $d \sim h$

$a \sim g$   
 $c \sim j$   
 $e \sim i$

$b \sim h$   
 $d \sim g$

## Example



$a \sim f$

$b \sim g$

$c \sim h$

$d \sim i$

$a$

$e \sim j$

$a \sim g$

$b \sim h$

$c \sim i$

$d \sim j$

$e \sim g$

$a \sim h$

$b \sim i$

$c \sim j$

$d \sim g$

$e \sim h$

$a \sim i$

$b \sim j$

$c \sim g$

$d \sim h$


$e \sim i$

$a \sim j$

bisimulation

21 comparisons

## Example



$a \sim f$

$b \sim g$

$c \sim h$

$d \sim i$

$a$

$e \sim j$

$a \sim g$

$b \sim h$

$c \sim i$

$d \sim j$

$e \sim g$

$a \sim h$

$b \sim i$

$c \sim j$

$d \sim g$

$e \sim h$

$a \sim i$

$b \sim j$

$c \sim g$

$d \sim h$

$e \sim i$


$a \sim j$

bisimulation

$a \sim j$  follows from  $a \sim g, c \sim g, c \sim j$

21 comparisons

## Example



$a \sim f$

$b \sim g$

$c \sim h$

$d \sim i$

$a$   
 $e \sim j$

$a \sim g$

$b \sim h$

$c \sim i$

$d \sim j$

$e \sim g$

$a \sim h$

$b \sim i$

$c \sim j$

$d \sim g$

$e \sim h$

$a \sim i$

$b \sim j$

$c \sim g$

$d \sim h$

$e \sim i$


bisimulation

$e \sim i$  follows from  $d \sim i, d \sim j, e \sim j$

20 comparisons



## Example




$a \sim f$	$b \sim g$	$c \sim h$	$d \sim i$	$e \sim j$	$a \sim g$	$b \sim h$
$c \sim i$	$d \sim j$	$e \sim g$	$a \sim h$	$b \sim i$	$c \sim j$	$d \sim g$
$e \sim h$	$a \sim i$	$b \sim j$	$c \sim g$	$d \sim h$		

bisimulation

$d \sim h$  follows from  $c \sim h, c \sim i, d \sim i$

19 comparisons

## Example




$a \sim f$	$b \sim g$	$c \sim h$	$d \sim i$	$e \sim j$	$a \sim g$	$b \sim h$
$c \sim i$	$d \sim j$	$e \sim g$	$a \sim h$	$b \sim i$	$c \sim j$	$d \sim g$
$e \sim h$	$a \sim i$	$b \sim j$	$c \sim g$			

bisimulation

$c \sim g$  follows from  $b \sim g, b \sim h, c \sim h$

18 comparisons

## Example




$a \sim f$	$b \sim g$	$c \sim h$	$d \sim i$	$e \sim j$	$a \sim g$	$b \sim h$
$c \sim i$	$d \sim j$	$e \sim g$	$a \sim h$	$b \sim i$	$c \sim j$	$d \sim g$
$e \sim h$	$a \sim i$	$b \sim j$				

bisimulation

$b \sim j$  follows from  $b \sim g, e \sim g, e \sim j$

17 comparisons

## Example




$a \sim f$	$b \sim g$	$c \sim h$	$d \sim i$	$e \sim j$	$a \sim g$	$b \sim h$
$c \sim i$	$d \sim j$	$e \sim g$	$a \sim h$	$b \sim i$	$c \sim j$	$d \sim g$
$e \sim h$	$a \sim i$					

bisimulation

$a \sim i$  follows from  $a \sim g, d \sim g, d \sim i$

16 comparisons

## Example




$a \sim f$	$b \sim g$	$c \sim h$	$d \sim i$	$e \sim j$	$a \sim g$	$b \sim h$
$c \sim i$	$d \sim j$	$e \sim g$	$a \sim h$	$b \sim i$	$c \sim j$	$d \sim g$
$e \sim h$						

bisimulation

$e \sim h$  follows from  $c \sim h, c \sim j, e \sim j$

15 comparisons

## Example




$a \sim f$	$b \sim g$	$c \sim h$	$d \sim i$	$e \sim j$	$a \sim g$	$b \sim h$
$c \sim i$	$d \sim j$	$e \sim g$	$a \sim h$	$b \sim i$	$c \sim j$	$d \sim g$

bisimulation

$d \sim g$  follows from  $b \sim g, b \sim i, d \sim i$

14 comparisons

## Example




bisimulation

$c \sim j$  follows from  $c \sim i, d \sim i, d \sim j$

13 comparisons

## Example




bisimulation

$b \sim i$  follows from  $b \sim h, c \sim h, c \sim i$

12 comparisons



## Example




bisimulation

$a \sim h$  follows from  $a \sim g, b \sim g, b \sim h$

11 comparisons

## Example




bisimulation

$e \sim g$  follows from  $b \sim g, b \sim h, c \sim h, c \sim i, d \sim i, d \sim j, e \sim j$

10 comparisons

## Example




bisimulation **up to equivalence**

9 instead of 21 comparisons



for NFAs on-the-fly determinization is incorporated

## Example



for NFAs on-the-fly determinization is incorporated




## Example



a  
- - -  
d




for NFAs on-the-fly determinization is incorporated

Example





for NFAs on-the-fly determinization is incorporated

## Example



for NFAs on-the-fly determinization is incorporated



## Example







for NFAs on-the-fly determinization is incorporated

## Example





for NFAs on-the-fly determinization is incorporated

## Example




for NFAs on-the-fly determinization is incorporated

## Example




for NFAs on-the-fly determinization is incorporated

## Example




$$a \sim d \wedge c \sim e + f \\ \Rightarrow a + c \sim d + e + f$$






for NFAs on-the-fly determinization is incorporated


## Example



$$a \sim d \wedge c \sim e + f \\ \Rightarrow a + c \sim d + e + f$$







## Definitions

- **context-free grammar (CFG)** is quadruple  $G = (N, \Sigma, P, S)$  with



## Definitions

- **context-free grammar (CFG)** is quadruple  $G = (N, \Sigma, P, S)$  with
  - 1  $N$  : finite set of **nonterminals**

## Definitions

- **context-free grammar (CFG)** is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of **terminals**, disjoint from  $N$

## Definitions

- **context-free grammar (CFG)** is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of **productions** of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$

## Definitions

- **context-free grammar (CFG)** is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of productions of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$
  - ④  $S \in N$ : **start symbol**

## Definitions

- context-free grammar (CFG) is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of productions of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$
  - ④  $S \in N$ : start symbol
- **one step derivation** relation  $\xrightarrow[G]{1}$  on  $(N \cup \Sigma)^*$ :  $\beta A \gamma \xrightarrow[G]{1} \beta \alpha \gamma$  if  $A \rightarrow \alpha \in P$  and  $\beta, \gamma \in (N \cup \Sigma)^*$

## Definitions

- context-free grammar (CFG) is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of productions of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$
  - ④  $S \in N$ : start symbol
- one step derivation relation  $\xrightarrow{1}_G$  on  $(N \cup \Sigma)^*$ :  $\beta A \gamma \xrightarrow{1}_G \beta \alpha \gamma$  if  $A \rightarrow \alpha \in P$  and  $\beta, \gamma \in (N \cup \Sigma)^*$
- $\xrightarrow{n}_G = (\xrightarrow{1}_G)^n \quad \forall n \geq 0$

## Definitions

- context-free grammar (CFG) is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of productions of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$
  - ④  $S \in N$ : start symbol
- one step derivation relation  $\xrightarrow[G]{1}$  on  $(N \cup \Sigma)^*$ :  $\beta A \gamma \xrightarrow[G]{1} \beta \alpha \gamma$  if  $A \rightarrow \alpha \in P$  and  $\beta, \gamma \in (N \cup \Sigma)^*$
- $\xrightarrow[G]{n} = (\xrightarrow[G]{1})^n \quad \forall n \geq 0 \quad \xrightarrow[G]{*} = \bigcup_{n \geq 0} \xrightarrow[G]{n}$

## Definitions

- context-free grammar (CFG) is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of productions of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$
  - ④  $S \in N$ : start symbol
- one step derivation relation  $\xrightarrow{1}_G$  on  $(N \cup \Sigma)^*$ :  $\beta A \gamma \xrightarrow{1}_G \beta \alpha \gamma$  if  $A \rightarrow \alpha \in P$  and  $\beta, \gamma \in (N \cup \Sigma)^*$
- $\xrightarrow{n}_G = (\xrightarrow{1}_G)^n \quad \forall n \geq 0 \quad \xrightarrow{*}_G = \bigcup_{n \geq 0} \xrightarrow{n}_G$
- members of the set  $(N \cup \Sigma)^*$  are called **strings**



## Definitions

- context-free grammar (CFG) is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of productions of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$
  - ④  $S \in N$ : start symbol
- one step derivation relation  $\xrightarrow{1}_G$  on  $(N \cup \Sigma)^*$ :  $\beta A \gamma \xrightarrow{1}_G \beta \alpha \gamma$  if  $A \rightarrow \alpha \in P$  and  $\beta, \gamma \in (N \cup \Sigma)^*$
- $\xrightarrow{n}_G = (\xrightarrow{1}_G)^n \quad \forall n \geq 0 \quad \xrightarrow{*}_G = \bigcup_{n \geq 0} \xrightarrow{n}_G$
- members of the set  $(N \cup \Sigma)^*$  are called strings
- string  $s$  is called a **sentential form** if  $S \xrightarrow{*}_G s$  (derivable from the start symbol  $S$ )

## Definitions

- context-free grammar (CFG) is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of productions of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$
  - ④  $S \in N$ : start symbol
- one step derivation relation  $\xrightarrow{1}_G$  on  $(N \cup \Sigma)^*$ :  $\beta A \gamma \xrightarrow{1}_G \beta \alpha \gamma$  if  $A \rightarrow \alpha \in P$  and  $\beta, \gamma \in (N \cup \Sigma)^*$
- $\xrightarrow{n}_G = (\xrightarrow{1}_G)^n \quad \forall n \geq 0 \quad \xrightarrow{*}_G = \bigcup_{n \geq 0} \xrightarrow{n}_G$
- members of the set  $(N \cup \Sigma)^*$  are called strings
- string  $s$  is called a sentential form if  $S \xrightarrow{*}_G s$  (derivable from the start symbol  $S$ )
- sentential form  $x$  is called a **sentence** if  $x \in \Sigma^*$  (consisting terminal symbols only)

## Definitions


- context-free grammar (CFG) is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of productions of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$
  - ④  $S \in N$ : start symbol
- one step derivation relation  $\xrightarrow{1}_G$  on  $(N \cup \Sigma)^*$ :  $\beta A \gamma \xrightarrow{1}_G \beta \alpha \gamma$  if  $A \rightarrow \alpha \in P$  and  $\beta, \gamma \in (N \cup \Sigma)^*$
- $\xrightarrow{n}_G = (\xrightarrow{1}_G)^n \quad \forall n \geq 0 \quad \xrightarrow{*}_G = \bigcup_{n \geq 0} \xrightarrow{n}_G$
- members of the set  $(N \cup \Sigma)^*$  are called strings
- string  $s$  is called a sentential form if  $S \xrightarrow{*}_G s$  (derivable from the start symbol  $S$ )
- sentential form  $x$  is called a sentence if  $x \in \Sigma^*$  (consisting terminal symbols only)
- language generated by  $G$ :  $L(G) = \{x \in \Sigma^* \mid S \xrightarrow{*}_G x\}$

## Definitions

- context-free grammar (CFG) is quadruple  $G = (N, \Sigma, P, S)$  with
  - ①  $N$ : finite set of nonterminals
  - ②  $\Sigma$ : finite set of terminals, disjoint from  $N$
  - ③  $P$ : finite set of productions of the form  $A \rightarrow \alpha$  with  $A \in N$  and  $\alpha \in (N \cup \Sigma)^*$
  - ④  $S \in N$ : start symbol
- one step derivation relation  $\xrightarrow{1}_G$  on  $(N \cup \Sigma)^*$ :  $\beta A \gamma \xrightarrow{1}_G \beta \alpha \gamma$  if  $A \rightarrow \alpha \in P$  and  $\beta, \gamma \in (N \cup \Sigma)^*$
- $\xrightarrow{n}_G = (\xrightarrow{1}_G)^n \quad \forall n \geq 0 \quad \xrightarrow{*}_G = \bigcup_{n \geq 0} \xrightarrow{n}_G$
- members of the set  $(N \cup \Sigma)^*$  are called strings
- string  $s$  is called a sentential form if  $S \xrightarrow{*}_G s$  (derivable from the start symbol  $S$ )
- sentential form  $x$  is called a sentence if  $x \in \Sigma^*$  (consisting terminal symbols only)
- language generated by  $G$ :  $L(G) = \{x \in \Sigma^* \mid S \xrightarrow{*}_G x\}$
- set  $B \subseteq \Sigma^*$  is **context-free** if  $B = L(G)$  for some CFG  $G$


## Example

DFA  $M$



## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$


$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$

$4 \rightarrow a4 \mid b4 \mid \epsilon$

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$


$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$

$4 \rightarrow a4 \mid b4 \mid \epsilon$

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$


$3 \rightarrow a4 \mid b1$

$4 \rightarrow a4 \mid b4 \mid \epsilon$



## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$


$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$

$4 \rightarrow a4 \mid b4 \mid \epsilon$

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$


$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$

$4 \rightarrow a4 \mid b4 \mid \epsilon$

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$

$4 \rightarrow a4 \mid b4 \mid \epsilon$

DFA


1

CFG

1

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$


$4 \rightarrow a4 \mid b4 \mid \epsilon$

DFA  $1 \xrightarrow{b} 1$

CFG  $1 \rightarrow b1$

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$


$4 \rightarrow a4 \mid b4 \mid \epsilon$

DFA  $1 \xrightarrow{b} 1 \xrightarrow{a} 2$

CFG  $1 \rightarrow b1 \rightarrow ba2$

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$


$4 \rightarrow a4 \mid b4 \mid \epsilon$

DFA  $b_1 a_1 b_2$

CFG  $1 \rightarrow b1 \rightarrow ba2 \rightarrow bab3$

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$


$4 \rightarrow a4 \mid b4 \mid \epsilon$

DFA  $1 \xrightarrow{b} 1 \xrightarrow{a} 2 \xrightarrow{b} 3 \xrightarrow{a} 4$

CFG  $1 \rightarrow b1 \rightarrow ba2 \rightarrow bab3 \rightarrow baba4$

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$

$4 \rightarrow a4 \mid b4 \mid \epsilon$


DFA     $b \ a \ b \ a \ a$   
 1   1   2   3   4   4

CFG     $1 \rightarrow b1 \rightarrow ba2 \rightarrow bab3 \rightarrow baba4 \rightarrow baba a4$



## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$


$4 \rightarrow a4 \mid b4 \mid \epsilon$

DFA     $b \ a \ b \ a \ a$   
 1   1   2   3   4   4

CFG     $1 \rightarrow b1 \rightarrow ba2 \rightarrow bab3 \rightarrow baba4 \rightarrow babaa4 \rightarrow babaa$

## Example

DFA  $M$



CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$


$4 \rightarrow a4 \mid b4 \mid \epsilon$

DFA  $\underset{1}{b} \underset{1}{a} \underset{2}{b} \underset{3}{a} \underset{4}{a} \in L(M)$

CFG  $1 \rightarrow b1 \rightarrow ba2 \rightarrow bab3 \rightarrow baba4 \rightarrow babaa4 \rightarrow babaa \in L(G_M)$

## Example

DFA  $M$



strongly right-linear CFG  $G_M$

$1 \rightarrow a2 \mid b1$

$2 \rightarrow a2 \mid b3$

$3 \rightarrow a4 \mid b1$

$4 \rightarrow a4 \mid b4 \mid \epsilon$

DFA  $\underset{1}{b} \underset{1}{a} \underset{2}{b} \underset{3}{a} \underset{4}{a} \in L(M)$

CFG  $1 \rightarrow b1 \rightarrow ba2 \rightarrow bab3 \rightarrow baba4 \rightarrow babaa4 \rightarrow babaa \in L(G_M)$

## Definition

CFG  $G = (N, \Sigma, P, S)$  is **strongly right-linear** if

$$\alpha = aB \in \Sigma N \quad \text{or} \quad \alpha = \varepsilon$$

for all  $A \rightarrow \alpha$  in  $P$

## Definition

CFG  $G = (N, \Sigma, P, S)$  is **strongly right-linear** if

$$\alpha = aB \in \Sigma N \quad \text{or} \quad \alpha = \varepsilon$$

for all  $A \rightarrow \alpha$  in  $P$

## Lemma

$L$  is **regular**  $\iff L$  is generated by **strongly right-linear CFG**

## Definition

CFG  $G = (N, \Sigma, P, S)$  is **strongly right-linear** if

$$\alpha = aB \in \Sigma N \quad \text{or} \quad \alpha = \varepsilon$$

for all  $A \rightarrow \alpha$  in  $P$

## Lemma

$L$  is regular  $\iff L$  is generated by strongly right-linear CFG

## Proof. ( $\implies$ )

- DFA  $M = (Q, \Sigma, \delta, s, F)$
- $L(M) = L(G_M)$  for strongly right-linear CFG  $G_M = \{Q, \Sigma, P, s\}$  with
$$P = \{p \rightarrow aq \mid \delta(p, a) = q\} \cup \{q \rightarrow \varepsilon \mid q \in F\}$$

## Definition

CFG  $G = (N, \Sigma, P, S)$  is **strongly right-linear** if

$$\alpha = aB \in \Sigma N \quad \text{or} \quad \alpha = \varepsilon$$

for all  $A \rightarrow \alpha$  in  $P$

## Lemma

$L$  is regular  $\iff L$  is generated by strongly right-linear CFG

## Proof. ( $\Leftarrow$ )

- strongly right-linear CFG  $G = (N, \Sigma, P, S)$
- $L(G) = L(M_G)$  for NFA  $M_G = (N, \Delta, \{S\}, F)$  with

$$\Delta(A, a) = \{B \mid A \rightarrow aB \in P\} \quad \text{and} \quad F = \{A \mid A \rightarrow \varepsilon \in P\}$$

## Definition

CFG  $G = (N, \Sigma, P, S)$  is **strongly right-linear** if

$$\alpha = aB \in \Sigma N \quad \text{or} \quad \alpha = \varepsilon$$

for all  $A \rightarrow \alpha$  in  $P$

## Lemma

$L$  is regular  $\iff L$  is generated by strongly right-linear CFG

## Corollary

every regular set is context-free




## Example

CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

①  $S \xrightarrow[G]{1} [S]$

parse tree




## Example

CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow[G]{1} [\textcolor{red}{S}] \xrightarrow[G]{1} [[S]]$$

parse tree




## Example

CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow[G]{1} [S] \xrightarrow[G]{1} [[S]] \xrightarrow[G]{1} [[[ ]]]$$

parse tree





## Example

CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow[G]{1} [S] \xrightarrow[G]{1} [[S]] \xrightarrow[G]{1} [[[ ]]$$

$$\textcircled{2} \quad S \xrightarrow[G]{1} [S]$$

parse trees





## Example

CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow[G]{1} [S] \xrightarrow[G]{1} [[S]] \xrightarrow[G]{1} [[[ ]]$$

$$\textcircled{2} \quad S \xrightarrow[G]{1} [S] \xrightarrow[G]{1} [SS]$$

parse trees





## Example

CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[[ ]]$$

$$\textcircled{2} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S]$$

parse trees





## Example

CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow{G} [S] \xrightarrow{G} [[S]] \xrightarrow{G} [[[ ]]$$

$$\textcircled{2} \quad S \xrightarrow{G} [S] \xrightarrow{G} [SS] \xrightarrow{G} [S] \xrightarrow{G} [[S]]$$

parse trees





## Example

CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow{G} [S] \xrightarrow{G} [[S]] \xrightarrow{G} [[[ ]]$$

$$\textcircled{2} \quad S \xrightarrow{G} [S] \xrightarrow{G} [SS] \xrightarrow{G} [S] \xrightarrow{G} [[S]] \xrightarrow{G} [[[ ]]$$

parse trees





## Example



CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[[ ]]$$

$$\textcircled{2} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[[ ]]$$

$$\textcircled{3} \quad S \xrightarrow{1_G} [S]$$

parse trees



## Example




CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[[ ]]$$

$$\textcircled{2} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[[ ]]$$

$$\textcircled{3} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS]$$

parse trees



## Example




CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[[ ]]$$

$$\textcircled{2} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[[ ]]$$

$$\textcircled{3} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [S\textcolor{red}{S}] \xrightarrow{1_G} [S[S]]$$

parse trees



## Example




CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[[ ]]$$

$$\textcircled{2} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[[ ]]$$

$$\textcircled{3} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S[S]] \xrightarrow{1_G} [S[ ]]$$

parse trees



## Example




CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[]]$$

$$\textcircled{2} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[]]$$

$$\textcircled{3} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S[S]] \xrightarrow{1_G} [S[[]]] \xrightarrow{1_G} [[]]$$

parse trees



## Example




CFG  $G$ :  $S \rightarrow [S] \mid SS \mid \epsilon$

$$\textcircled{1} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[]]$$

$$\textcircled{2} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S] \xrightarrow{1_G} [[S]] \xrightarrow{1_G} [[]]$$

$$\textcircled{3} \quad S \xrightarrow{1_G} [S] \xrightarrow{1_G} [SS] \xrightarrow{1_G} [S[S]] \xrightarrow{1_G} [S[]] \xrightarrow{1_G} [[]]$$

parse trees



## Definition

- CFG is **ambiguous** if some string has different parse trees

## Example

- CFG  $G$  :  $S \rightarrow S \times S \mid S + S \mid \text{int}$

- $G$  is ambiguous

with  $G$      $7 + 5 \times 2$  could be parsed as     $7 + (5 \times 2)$  and  $(7 + 5) \times 2$

## Example

- CFG  $G$  :  $S \rightarrow S \times S \mid S + S \mid \textit{int}$
- CFG  $G'$  :  
 $S \rightarrow S + T \mid T$   
 $T \rightarrow T \times U \mid U$   
 $U \rightarrow \textit{int} \mid (S)$

- $G$  is ambiguous

with  $G$      $7 + 5 \times 2$  could be parsed as     $7 + (5 \times 2)$  and  $(7 + 5) \times 2$



## Example


• CFG  $G$  :  $S \rightarrow S \times S \mid S + S \mid \text{int}$

• CFG  $G'$  :  $S \rightarrow S + T \mid T$   
 $T \rightarrow T \times U \mid U$   
 $U \rightarrow \text{int} \mid (S)$

•  $G$  is ambiguous       $G'$  is unambiguous       $L(G) = L(G')$


with  $G$        $7 + 5 \times 2$  could be parsed as       $7 + (5 \times 2)$  and  $(7 + 5) \times 2$

with  $G'$        $7 + 5 \times 2$  could only be parsed as       $7 + (5 \times 2)$



## Example

- CFG  $G$  :  $S \rightarrow S \times S \mid S + S \mid \text{int}$
- CFG  $G'$  :  $S \rightarrow S + T \mid T$   
 $T \rightarrow T \times U \mid U$   
 $U \rightarrow \text{int} \mid (S)$
- $G$  is ambiguous       $G'$  is unambiguous       $L(G) = L(G')$   
     with  $G$      $7 + 5 \times 2$  could be parsed as       $7 + (5 \times 2)$  and  $(7 + 5) \times 2$   
     with  $G'$      $7 + 5 \times 2$  could only be parsed as       $7 + (5 \times 2)$
- (if applicable) one way to remove ambiguity is to benefit from precedence of operators



## Remark

there exist context-free sets without unambiguous grammars

## Remark

there exist context-free sets without unambiguous grammars ( **inherently ambiguous** )

## Remark

there exist context-free sets without unambiguous grammars (inherently ambiguous)

## Example

$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$  is context-free and inherently ambiguous

## Remark

there exist context-free sets without unambiguous grammars (inherently ambiguous)

## Example

$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$  is context-free and inherently ambiguous

$A = \{a^i b^i c^k\} \cup \{a^i b^j c^j\}$

## Remark

there exist context-free sets without unambiguous grammars (inherently ambiguous)

## Example

$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$  is context-free and inherently ambiguous

$A = \{a^i b^i c^k\} \cup \{a^i b^j c^j\}$

let  $A = L(G)$  such that  $G$ :

## Remark

there exist context-free sets without unambiguous grammars (inherently ambiguous)

## Example

$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$  is context-free and inherently ambiguous

$A = \{a^i b^i c^k\} \cup \{a^i b^j c^j\}$

let  $A = L(G)$  such that  $G$ :

$$\begin{array}{ll} S \rightarrow T \mid W \\ T \rightarrow UV & W \rightarrow XY \\ U \rightarrow aUb \mid \varepsilon & X \rightarrow aX \mid \varepsilon \\ V \rightarrow cV \mid \varepsilon & Y \rightarrow bYc \mid \varepsilon \end{array}$$



## Remark

there exist context-free sets without unambiguous grammars (inherently ambiguous)

## Example

$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$  is context-free and inherently ambiguous

$A = \{a^i b^i c^k\} \cup \{a^i b^j c^j\}$

let  $A = L(G)$  such that  $G$ :

$$\begin{array}{ll} S \rightarrow T \mid W \\ T \rightarrow UV & W \rightarrow XY \\ U \rightarrow aUb \mid \varepsilon & X \rightarrow aX \mid \varepsilon \\ V \rightarrow cV \mid \varepsilon & Y \rightarrow bYc \mid \varepsilon \end{array}$$

the union we used has a non-empty intersection, where letters  $a$ ,  $b$  and  $c$  all are in equal number

## Remark

there exist context-free sets without unambiguous grammars (inherently ambiguous)

## Example

$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$  is context-free and inherently ambiguous

$A = \{a^i b^i c^k\} \cup \{a^i b^j c^j\}$

let  $A = L(G)$  such that  $G$ :

$$\begin{array}{ll} S \rightarrow T \mid W \\ T \rightarrow UV & W \rightarrow XY \\ U \rightarrow aUb \mid \varepsilon & X \rightarrow aX \mid \varepsilon \\ V \rightarrow cV \mid \varepsilon & Y \rightarrow bYc \mid \varepsilon \end{array}$$

the union we used has a non-empty intersection, where letters  $a$ ,  $b$  and  $c$  all are in equal number

## Lemma

there is no CFG  $G'$  such that  $L(G')$  is unambiguous with  $L(G') = A$

## Remark

there exist context-free sets without unambiguous grammars (inherently ambiguous)

## Example

$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$  is context-free and inherently ambiguous

$A = \{a^i b^i c^k\} \cup \{a^i b^j c^j\}$

let  $A = L(G)$  such that  $G$ :

$$\begin{array}{ll} S \rightarrow T \mid W \\ T \rightarrow UV & W \rightarrow XY \\ U \rightarrow aUb \mid \varepsilon & X \rightarrow aX \mid \varepsilon \\ V \rightarrow cV \mid \varepsilon & Y \rightarrow bYc \mid \varepsilon \end{array}$$

the union we used has a non-empty intersection, where letters  $a$ ,  $b$  and  $c$  all are in equal number

## Lemma

there is no CFG  $G'$  such that  $L(G')$  is unambiguous with  $L(G') = A$

## Remark

- 1 given an **ambiguous** CFG  $G$ , the language  $L(G)$  **may or may not be ambiguous**

## Remark

there exist context-free sets without unambiguous grammars (inherently ambiguous)

## Example

$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$  is context-free and inherently ambiguous

$A = \{a^i b^i c^k\} \cup \{a^i b^j c^j\}$

let  $A = L(G)$  such that  $G$ :

$$\begin{array}{ll}
 S \rightarrow T \mid W & \\
 T \rightarrow UV & W \rightarrow XY \\
 U \rightarrow aUb \mid \varepsilon & X \rightarrow aX \mid \varepsilon \\
 V \rightarrow cV \mid \varepsilon & Y \rightarrow bYc \mid \varepsilon
 \end{array}$$

the union we used has a non-empty intersection, where letters  $a$ ,  $b$  and  $c$  all are in equal number

## Lemma

there is no CFG  $G'$  such that  $L(G')$  is unambiguous with  $L(G') = A$

## Remark

- 1 given an ambiguous CFG  $G$ , the language  $L(G)$  may or may not be ambiguous
- 2 there is **no algorithm** to convert ambiguous CFG to unambiguous CFG

## Remark

there exist context-free sets without unambiguous grammars (inherently ambiguous)

## Example

$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$  is context-free and inherently ambiguous

$A = \{a^i b^i c^k\} \cup \{a^i b^j c^j\}$

let  $A = L(G)$  such that  $G$ :

$$\begin{array}{lcl}
 & S \rightarrow T \mid W & \\
 T \rightarrow UV & & W \rightarrow XY \\
 U \rightarrow aUb \mid \varepsilon & & X \rightarrow aX \mid \varepsilon \\
 V \rightarrow cV \mid \varepsilon & & Y \rightarrow bYc \mid \varepsilon
 \end{array}$$

the union we used has a non-empty intersection, where letters  $a$ ,  $b$  and  $c$  all are in equal number

## Lemma

there is no CFG  $G'$  such that  $L(G')$  is unambiguous with  $L(G') = A$

## Remark

- 1 given an ambiguous CFG  $G$ , the language  $L(G)$  may or may not be ambiguous
- 2 there is no algorithm to convert ambiguous CFG to unambiguous CFG
- 3 unambiguous context free languages can be **parsed** by **deterministic** push down automata

# Outline

- 1 A Quick Recap
- 2 Chomsky Normal Form
- 3 Pumping Lemma
- 4 CKY Algorithm

## Definitions

CFG  $G = (N, \Sigma, P, S)$  is in

- **Chomsky normal form** if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = BC \in N^2$  or  $\alpha = a \in \Sigma$

## Definitions

CFG  $G = (N, \Sigma, P, S)$  is in

- Chomsky normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = BC \in N^2$  or  $\alpha = a \in \Sigma$
- **Greibach normal form** if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = aB_1 \cdots B_n \in \Sigma N^*$



## Definitions

CFG  $G = (N, \Sigma, P, S)$  is in

- Chomsky normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = BC \in N^2$  or  $\alpha = a \in \Sigma$
- Greibach normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = aB_1 \cdots B_n \in \Sigma N^*$



## Definitions

CFG  $G = (N, \Sigma, P, S)$  is in

- Chomsky normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = BC \in N^2$  or  $\alpha = a \in \Sigma$
- Greibach normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = aB_1 \cdots B_n \in \Sigma N^*$



## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in

- 1 Greibach normal form
- 2 Chomsky normal form

## Definitions

CFG  $G = (N, \Sigma, P, S)$  is in

- Chomsky normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = BC \in N^2$  or  $\alpha = a \in \Sigma$
- Greibach normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = aB_1 \cdots B_n \in \Sigma N^*$



## Theorem

every context-free set **not containing  $\epsilon$**  is generated by CFG in

- ① Greibach normal form
- ② Chomsky normal form

## Definitions

CFG  $G = (N, \Sigma, P, S)$  is in

- Chomsky normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = BC \in N^2$  or  $\alpha = a \in \Sigma$
- Greibach normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = aB_1 \cdots B_n \in \Sigma N^*$



## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in

- ① Greibach normal form
- ② Chomsky normal form

## Application

membership  $x \in L(G)$  is easily decidable for CFG  $G$  in Greibach normal form

## Definitions

CFG  $G = (N, \Sigma, P, S)$  is in

- Chomsky normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = BC \in N^2$  or  $\alpha = a \in \Sigma$
- Greibach normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = aB_1 \cdots B_n \in \Sigma N^*$



## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in

- ① Greibach normal form
- ② Chomsky normal form

## Application

membership  $x \in L(G)$  is easily decidable for CFG  $G$  in Greibach normal form:  
*generate all derivations of length  $|x|$  and test whether one of them ends in  $x$*

## Definitions

CFG  $G = (N, \Sigma, P, S)$  is in

- Chomsky normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = BC \in N^2$  or  $\alpha = a \in \Sigma$
- Greibach normal form if for all  $A \rightarrow \alpha$  in  $P$   $\alpha = aB_1 \cdots B_n \in \Sigma N^*$



## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in

- ① Greibach normal form
- ② Chomsky normal form

## Application

membership  $x \in L(G)$  is easily decidable for CFG  $G$  in Chomsky normal form:

*generate all derivations of length  $2|x| - 1$  and test whether one of them ends in  $x$*

## Definitions

- $\epsilon$ -production:  $A \rightarrow \epsilon$

## Definitions

- $\epsilon$ -production:  $A \rightarrow \epsilon$
- **unit production**:  $A \rightarrow B$



## Definitions

- $\epsilon$ -production:  $A \rightarrow \epsilon$
- unit production:  $A \rightarrow B$

## Lemma

every context-free set is generated by CFG without unit productions

## Definitions

- $\epsilon$ -production:  $A \rightarrow \epsilon$
- unit production:  $A \rightarrow B$

## Lemma

every context-free set is generated by CFG without unit productions

## Proof.

- $\hat{P}$  is smallest set containing  $P$  such that

$$A \rightarrow B \in \hat{P} \wedge B \rightarrow \alpha \in \hat{P} \implies A \rightarrow \alpha \in \hat{P}$$

## Definitions

- $\epsilon$ -production:  $A \rightarrow \epsilon$
- unit production:  $A \rightarrow B$

## Lemma

every context-free set is generated by CFG without unit productions

## Proof.

- $\hat{P}$  is smallest set containing  $P$  such that

$$A \rightarrow B \in \hat{P} \wedge B \rightarrow \alpha \in \hat{P} \implies A \rightarrow \alpha \in \hat{P}$$

- remove all unit productions from  $\hat{P}$

## Definitions

- $\varepsilon$ -production:  $A \rightarrow \varepsilon$
- unit production:  $A \rightarrow B$

## Lemma

every context-free set not containing  $\varepsilon$  is generated by CFG without  $\varepsilon$ -productions

## Proof.

- $\hat{P}$  is smallest set containing  $P$  such that

$$A \rightarrow \alpha B \beta \in \hat{P} \quad \wedge \quad B \rightarrow \varepsilon \in \hat{P} \quad \implies \quad A \rightarrow \alpha \beta \in \hat{P}$$

- remove all  $\varepsilon$ -productions from  $\hat{P}$

## Definitions

- $\epsilon$ -production:  $A \rightarrow \epsilon$
- unit production:  $A \rightarrow B$

## Lemma

every context-free set not containing  $\epsilon$  is generated by CFG without  $\epsilon$  and unit productions

## Proof.

- $\hat{P}$  is smallest set containing  $P$  such that

$$\begin{aligned} A \rightarrow B \in \hat{P} \wedge B \rightarrow \alpha \in \hat{P} &\implies A \rightarrow \alpha \in \hat{P} \\ A \rightarrow \alpha B \beta \in \hat{P} \wedge B \rightarrow \epsilon \in \hat{P} &\implies A \rightarrow \alpha \beta \in \hat{P} \end{aligned}$$

- remove all  $\epsilon$  and unit productions from  $\hat{P}$

## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in Chomsky normal form

## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in Chomsky normal form

## Proof.

- remove  $\varepsilon$  and unit productions

## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in Chomsky normal form

## Proof.

- remove  $\varepsilon$  and unit productions
- introduce  $A_a \rightarrow a$  for every  $a \in \Sigma$

## Example

$$S \rightarrow SS$$

$$S \rightarrow [S]$$

$$S \rightarrow []$$



## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in Chomsky normal form

## Proof.

- remove  $\varepsilon$  and unit productions
- introduce  $A_a \rightarrow a$  for every  $a \in \Sigma$

## Example

$S \rightarrow SS$   
 $A \rightarrow [$

$S \rightarrow ASB$   
 $B \rightarrow ]$

$S \rightarrow AB$

## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in Chomsky normal form

## Proof.

- remove  $\varepsilon$  and unit productions
- introduce  $A_a \rightarrow a$  for every  $a \in \Sigma$
- split long right-hand sides of productions

## Example

$$S \rightarrow SS$$

$$A \rightarrow [$$

$$S \rightarrow ASB$$

$$B \rightarrow ]$$

$$S \rightarrow AB$$

## Theorem

every context-free set not containing  $\varepsilon$  is generated by CFG in Chomsky normal form

## Proof.

- remove  $\varepsilon$  and unit productions
- introduce  $A_a \rightarrow a$  for every  $a \in \Sigma$
- split long right-hand sides of productions

## Example

$$\begin{aligned} S &\rightarrow SS \\ A &\rightarrow [ \end{aligned}$$
$$\begin{aligned} S &\rightarrow AC \\ B &\rightarrow ] \end{aligned}$$
$$\begin{aligned} S &\rightarrow AB \\ C &\rightarrow SB \end{aligned}$$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \varepsilon$      $Y \rightarrow bXY \mid \varepsilon$      $Z \rightarrow a$   
remove  $\varepsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$

$X \rightarrow Z \mid \varepsilon$

$Y \rightarrow bXY \mid \varepsilon$      $Z \rightarrow a$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$

$X \rightarrow Z \mid \epsilon$

$Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$   
 $S \rightarrow bS$

$X \rightarrow Z \mid \epsilon$

$Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$   
 $S \rightarrow bS \mid Yb$

$X \rightarrow Z \mid \epsilon$

$Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$   
 $S \rightarrow bS \mid Yb \mid YZ$

$X \rightarrow Z \mid \epsilon$

$Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$



## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$   
 $S \rightarrow bS \mid Yb \mid YZ$

$X \rightarrow Z \mid \epsilon$   
 $Y \rightarrow bY$

$Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$   
 $S \rightarrow bS \mid Yb \mid YZ$

$X \rightarrow Z \mid \epsilon$   
 $Y \rightarrow bY$

$Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$$\begin{aligned} S &\rightarrow XbS \mid XYb \mid YXZ \\ S &\rightarrow bS \mid Yb \mid YZ \mid Xb \end{aligned}$$
$$\begin{aligned} X &\rightarrow Z \mid \epsilon \\ Y &\rightarrow bY \end{aligned}$$
$$Y \rightarrow bXY \mid \epsilon \quad Z \rightarrow a$$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$

$X \rightarrow Z \mid \epsilon$

$Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$

$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ$

$Y \rightarrow bY$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$

$X \rightarrow Z \mid \epsilon$

$Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$

$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ$

$Y \rightarrow bY \mid bX$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$	$X \rightarrow Z \mid \epsilon$	$Y \rightarrow bXY \mid \epsilon$	$Z \rightarrow a$
$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b$	$Y \rightarrow bY \mid bX$		

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$	$X \rightarrow Z \mid \epsilon$	$Y \rightarrow bXY \mid \epsilon$	$Z \rightarrow a$
$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b \mid Z$	$Y \rightarrow bY \mid bX$		

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$	$X \rightarrow Z \mid \epsilon$	$Y \rightarrow bXY \mid \epsilon$	$Z \rightarrow a$
$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b \mid Z$	$Y \rightarrow bY \mid bX \mid b$		



## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and **unit** productions

$S \rightarrow XbS \mid XYb \mid YXZ$	$X \rightarrow Z \mid \epsilon$	$Y \rightarrow bXY \mid \epsilon$	$Z \rightarrow a$
$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b \mid Z$	$Y \rightarrow bY \mid bX \mid b$		

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \epsilon$      $Y \rightarrow bXY \mid \epsilon$      $Z \rightarrow a$   
remove  $\epsilon$  and **unit** productions

$S \rightarrow XbS \mid XYb \mid YXZ$	$X \rightarrow Z \mid \epsilon$	$Y \rightarrow bXY \mid \epsilon$	$Z \rightarrow a$
$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b \mid Z$	$Y \rightarrow bY \mid bX \mid b$	$X \rightarrow a$	

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \varepsilon$      $Y \rightarrow bXY \mid \varepsilon$      $Z \rightarrow a$   
remove  $\varepsilon$  and **unit** productions

$S \rightarrow XbS \mid XYb \mid YXZ$	$X \rightarrow Z \mid \varepsilon$	$Y \rightarrow bXY \mid \varepsilon$	$Z \rightarrow a$
$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b \mid Z$	$Y \rightarrow bY \mid bX \mid b$	$X \rightarrow a$	

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \varepsilon$      $Y \rightarrow bXY \mid \varepsilon$      $Z \rightarrow a$   
remove  $\varepsilon$  and **unit** productions

$S \rightarrow XbS \mid XYb \mid YXZ$	$X \rightarrow Z \mid \varepsilon$	$Y \rightarrow bXY \mid \varepsilon$	$Z \rightarrow a$
$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b \mid Z$	$Y \rightarrow bY \mid bX \mid b$	$X \rightarrow a$	$S \rightarrow a$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \varepsilon$      $Y \rightarrow bXY \mid \varepsilon$      $Z \rightarrow a$   
remove  $\varepsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$

$Y \rightarrow bXY$

$Z \rightarrow a$

$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b$

$Y \rightarrow bY \mid bX \mid b$

$X \rightarrow a$

$S \rightarrow a$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \varepsilon$      $Y \rightarrow bXY \mid \varepsilon$      $Z \rightarrow a$

remove  $\varepsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$

$Y \rightarrow bXY$

$Z \rightarrow a$

$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b$

$Y \rightarrow bY \mid bX \mid b$

$X \rightarrow a$

$S \rightarrow a$

introduce new non-terminals

$B \rightarrow b$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \varepsilon$      $Y \rightarrow bXY \mid \varepsilon$      $Z \rightarrow a$   
remove  $\varepsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$	$Y \rightarrow bXY$	$Z \rightarrow a$
$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b$	$Y \rightarrow bY \mid bX \mid b$	$X \rightarrow a$ $S \rightarrow a$

introduce new non-terminals

$B \rightarrow b$	$S \rightarrow XBS \mid XYB \mid YXZ \mid BS \mid YB \mid YZ \mid XB \mid XZ \mid b \mid a$
$X \rightarrow a$	$Y \rightarrow BXY \mid BY \mid BX \mid b$ $Z \rightarrow a$

## Example

CFG  $G$ :  $S \rightarrow XbS \mid XYb \mid YXZ$      $X \rightarrow Z \mid \varepsilon$      $Y \rightarrow bXY \mid \varepsilon$      $Z \rightarrow a$   
remove  $\varepsilon$  and unit productions

$S \rightarrow XbS \mid XYb \mid YXZ$		$Y \rightarrow bXY$	$Z \rightarrow a$
$S \rightarrow bS \mid Yb \mid YZ \mid Xb \mid XZ \mid b$	$Y \rightarrow bY \mid bX \mid b$	$X \rightarrow a$	$S \rightarrow a$

introduce new non-terminals

$B \rightarrow b$	$S \rightarrow XBS \mid XbB \mid YXZ \mid BS \mid YB \mid YZ \mid XB \mid XZ \mid b \mid a$
$X \rightarrow a$	$Y \rightarrow BXY \mid BY \mid BX \mid b$ $Z \rightarrow a$

split long right-hand sides

$B \rightarrow b$	$S \rightarrow TS \mid UB \mid VZ \mid BS \mid YB \mid YZ \mid XB \mid XZ \mid b \mid a$
$X \rightarrow a$	$Y \rightarrow BU \mid BY \mid BX \mid b$ $Z \rightarrow a$
$T \rightarrow XB$	$U \rightarrow XY$ $V \rightarrow YX$



# Outline

- 1 A Quick Recap
- 2 Chomsky Normal Form
- 3 Pumping Lemma**
- 4 CKY Algorithm

## Pumping Lemma

$$A \text{ is context-free} \implies \left\{ \begin{array}{ll} \exists k & \\ \forall z \in A & \text{with } |z| \geq k \\ \exists u, v, w, x, y & \text{with } \begin{cases} z = uvwxy \\ |vwx| \leq k \\ vx \neq \varepsilon \end{cases} \\ \forall i \geq 0 & uv^iwx^iy \in A \end{array} \right.$$

## Pumping Lemma

$$A \text{ is context-free} \implies \left\{ \begin{array}{ll} \exists k & \\ \forall z \in A & \text{with } |z| \geq k \\ \exists u, v, w, x, y & \text{with } \begin{cases} z = uvwxy \\ |vwx| \leq k \\ vx \neq \varepsilon \end{cases} \\ \forall i \geq 0 & uv^iwx^iy \in A \end{array} \right.$$

## Proof. (Idea)


take  $k = 2^{n+1}$  where  $n$  is number of nonterminals of any CFG in Chomsky normal form that accepts  $A - \{\varepsilon\}$

## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$

## Example


$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G: S \rightarrow AC \mid AB \quad C \rightarrow SB \quad A \rightarrow a \quad B \rightarrow b$



parse tree for *aaaabbbb*

## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$




parse tree for *aaaabbbb*

long string  $\Rightarrow$  long path in parse tree

## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$




parse tree for  $aaaabbbb$

long string  $\Rightarrow$  long path in parse tree

(at depth  $m$  at most  $2^m$  symbols)

## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$



parse tree for  $aaaabbbb$

long string  $\implies$  long path in parse tree


(at depth  $m$  at most  $2^m$  symbols)

consider longest path



## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$



parse tree for *aaaabbbb*

long string  $\implies$  long path in parse tree


(at depth  $m$  at most  $2^m$  symbols)

consider longest path

look for **repetition of nonterminals near bottom**

## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$



parse tree for *aaaabbbb*

long string  $\implies$  long path in parse tree

(at depth  $m$  at most  $2^m$  symbols)


consider longest path

look for repetition of nonterminals near bottom

determine  $u, v, w, x, y$

## Example

$L = \{a^i b^j \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$



parse tree for *aaaabbbb*

long string  $\Rightarrow$  long path in parse tree

(at depth  $m$  at most  $2^m$  symbols)

consider longest path


look for repetition of nonterminals near bottom

determine  $u, v, w, x, y$

$$u = aa \quad v = a \quad w = ab \quad x = b \quad y = bb$$

## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$



parse tree for  $aaaabbbb$

long string  $\implies$  long path in parse tree

(at depth  $m$  at most  $2^m$  symbols)

consider longest path


look for repetition of nonterminals near bottom

determine  $u, v, w, x, y$

$u = aa$   $v = a$   $w = ab$   $x = b$   $y = bb$

## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$



parse tree for  $aaaabbbb$

long string  $\implies$  long path in parse tree

(at depth  $m$  at most  $2^m$  symbols)

consider longest path


look for repetition of nonterminals near bottom

determine  $u, v, w, x, y$

$u = aa$   $v = a$   $w = ab$   $x = b$   $y = bb$

## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$



parse tree for  $aaaabbbb$

long string  $\implies$  long path in parse tree

(at depth  $m$  at most  $2^m$  symbols)

consider longest path


look for repetition of nonterminals near bottom

determine  $u, v, w, x, y$

$u = aa$   $v = a$   $w = ab$   $x = b$   $y = bb$

## Example

$L = \{a^i b^i \mid i > 0\}$  Chomsky normal form  $G$ :  $S \rightarrow AC \mid AB$   $C \rightarrow SB$   $A \rightarrow a$   $B \rightarrow b$



parse tree for *aaaabbbb*

long string  $\implies$  long path in parse tree


(at depth  $m$  at most  $2^m$  symbols)

consider longest path


look for repetition of nonterminals near bottom


determine  $u, v, w, x, y$


$u = aa$   $v = a$   $w = ab$   $x = b$   $y = bb$





















$$S \xrightarrow[G]{*} uAy \quad A \xrightarrow[G]{*} vAx \quad A \xrightarrow[G]{*} w$$






$$S \xrightarrow[G]{*} uAy \quad A \xrightarrow[G]{*} vAx \quad A \xrightarrow[G]{*} w$$

$i = 0: uwy$



$$S \xrightarrow{*}_G uAy \quad A \xrightarrow{*}_G vAx \quad A \xrightarrow{*}_G w$$



$$S \xrightarrow{*}_G uAy \quad A \xrightarrow{*}_G vAx \quad A \xrightarrow{*}_G w$$



## Pumping Lemma (Contrapositive)

$$\left. \begin{array}{l} \forall k \\ \exists z \in A \quad \text{with } |z| \geq k \\ \forall u, v, w, x, y \quad \text{with } \begin{cases} z = uvwxy \\ |vwx| \leq k \\ vx \neq \varepsilon \end{cases} \\ \exists i \geq 0 \quad \text{with } uv^iwx^iy \notin A \end{array} \right\} \Rightarrow A \text{ is not context-free}$$

## Example

$A = \{a^i b^j c^i \mid i \geq 0\}$  is not context-free

## Example

$A = \{a^i b^j c^i \mid i \geq 0\}$  is not context-free

- choose  $z = a^k b^k c^k$

## Example

$A = \{a^i b^j c^i \mid i \geq 0\}$  is not context-free

- choose  $z = a^k b^k c^k$       check:  $z \in A \quad |z| = 3k \geq k$

## Example

$A = \{a^i b^j c^i \mid i \geq 0\}$  is not context-free

- choose  $z = a^k b^k c^k$       check:  $z \in A \quad |z| = 3k \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$

## Example

$A = \{a^i b^i c^i \mid i \geq 0\}$  is not context-free

- choose  $z = a^k b^k c^k$       check:  $z \in A \quad |z| = 3k \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- choose  $i = 0$

## Example

$A = \{a^i b^i c^i \mid i \geq 0\}$  is not context-free

- choose  $z = a^k b^k c^k$       check:  $z \in A \quad |z| = 3k \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- choose  $i = 0$
- $vwx$  cannot contain both  $a$ 's and  $c$ 's

## Example

$A = \{a^i b^i c^i \mid i \geq 0\}$  is not context-free

- choose  $z = a^k b^k c^k$       check:  $z \in A \quad |z| = 3k \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- choose  $i = 0$
- $vwx$  cannot contain both  $a$ 's and  $c$ 's
  - $vwx$  has no  $a$ 's:  $uv^i wx^i y$  has more  $a$ 's than  $b$ 's or  $c$ 's



## Example

$A = \{a^i b^i c^i \mid i \geq 0\}$  is not context-free

- choose  $z = a^k b^k c^k$       check:  $z \in A \quad |z| = 3k \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- choose  $i = 0$
- $vwx$  cannot contain both  $a$ 's and  $c$ 's
  - ①  $vwx$  has no  $a$ 's:  $uv^i wx^i y$  has more  $a$ 's than  $b$ 's or  $c$ 's
  - ②  $vwx$  has no  $c$ 's:  $uv^i wx^i y$  has more  $c$ 's than  $b$ 's or  $a$ 's

## Example

$B = \{a^i b^j c^k \mid i < j < k\}$  is not context-free

## Example

$B = \{a^i b^j c^k \mid i < j < k\}$  is not context-free

- choose

$$z = a^k b^{k+1} c^{k+2}$$

## Example

$B = \{a^i b^j c^k \mid i < j < k\}$  is not context-free

- choose  $z = a^k b^{k+1} c^{k+2}$       check:  $z \in B \quad |z| = 3k + 3 \geq k$

## Example

$B = \{a^i b^j c^k \mid i < j < k\}$  is not context-free

- choose  $z = a^k b^{k+1} c^{k+2}$       check:  $z \in B \quad |z| = 3k + 3 \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$

## Example

$B = \{a^i b^j c^k \mid i < j < k\}$  is not context-free

- choose  $z = a^k b^{k+1} c^{k+2}$       check:  $z \in B \quad |z| = 3k + 3 \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- $vwx$  cannot contain both  $a$ 's and  $c$ 's

## Example

$B = \{a^i b^j c^k \mid i < j < k\}$  is not context-free

- choose  $z = a^k b^{k+1} c^{k+2}$       check:  $z \in B \quad |z| = 3k + 3 \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- $vwx$  cannot contain both  $a$ 's and  $c$ 's

①  $vwx$  has no  $a$ 's

choose  $i = 0$       check:  $uv^i wx^i y \notin B$

## Example

$B = \{a^i b^j c^k \mid i < j < k\}$  is not context-free

- choose  $z = a^k b^{k+1} c^{k+2}$       check:  $z \in B \quad |z| = 3k + 3 \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- $vwx$  cannot contain both  $a$ 's and  $c$ 's

①  $vwx$  has no  $a$ 's

choose  $i = 0$       check:  $uv^i wx^i y \notin B$

②  $vwx$  has no  $c$ 's

choose  $i = 2$       check:  $uv^i wx^i y \notin B$



## Example

$C = \{a^p \mid p \text{ is prime}\}$  is not context-free

## Example

$C = \{a^p \mid p \text{ is prime}\}$  is not context-free

- choose  $z = a^p$  where  $p$  is any prime larger than  $k$

## Example

$C = \{a^p \mid p \text{ is prime}\}$  is not context-free

- choose  $z = a^p$  where  $p$  is any prime larger than  $k$       check:  $z \in C \quad |z| = p \geq k$

## Example

$C = \{a^p \mid p \text{ is prime}\}$  is not context-free

- choose  $z = a^p$  where  $p$  is any prime larger than  $k$       check:  $z \in C \quad |z| = p \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$

## Example

$C = \{a^p \mid p \text{ is prime}\}$  is not context-free

- choose  $z = a^p$  where  $p$  is any prime larger than  $k$       check:  $z \in C \quad |z| = p \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- choose  $i = p + 1$       check:  $uv^iwx^iy \notin C$

## Example

$C = \{a^p \mid p \text{ is prime}\}$  is not context-free

- choose  $z = a^p$  where  $p$  is any prime larger than  $k$       check:  $z \in C \quad |z| = p \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- choose  $i = p + 1$       check:  $uv^iwx^iy \notin C$   
 $|uv^iwx^iy| = p + p|vx|$

## Example

$C = \{a^p \mid p \text{ is prime}\}$  is not context-free

- choose  $z = a^p$  where  $p$  is any prime larger than  $k$       check:  $z \in C \quad |z| = p \geq k$
- split:  $z = uvwxy$  with  $|vwx| \leq k$  and  $vx \neq \varepsilon$
- choose  $i = p + 1$       check:  $uv^iwx^iy \notin C$   
 $|uv^iwx^iy| = p + p|vx| = p(1 + |vx|)$  is not prime

# Outline

- 1 A Quick Recap
- 2 Chomsky Normal Form
- 3 Pumping Lemma
- 4 CKY Algorithm**



## Theorem

given CFG  $G = (N, \Sigma, P, S)$  and string  $x \in \Sigma^*$ , it is decidable whether  $x \in L(G)$

## Theorem

given CFG  $G = (N, \Sigma, P, S)$  and string  $x \in \Sigma^*$ , it is decidable whether  $x \in L(G)$

## Proof.

efficient and elegant algorithm: Cocke Kasami Younger (CKY)

- convert  $G$  into Chomsky normal form

## Theorem

given CFG  $G = (N, \Sigma, P, S)$  and string  $x \in \Sigma^*$ , it is decidable whether  $x \in L(G)$

## Proof.

efficient and elegant algorithm: Cocke Kasami Younger (CKY)

- convert  $G$  into Chomsky normal form
- for all  $0 \leq i < j \leq |x|$ 
  - $x_{ij}$  is substring of  $x$  of length  $j - i$  starting at position  $i$

## Theorem

given CFG  $G = (N, \Sigma, P, S)$  and string  $x \in \Sigma^*$ , it is decidable whether  $x \in L(G)$

## Proof.

efficient and elegant algorithm: Cocke Kasami Younger (CKY)

- convert  $G$  into Chomsky normal form
- for all  $0 \leq i < j \leq |x|$ 
  - $x_{ij}$  is substring of  $x$  of length  $j - i$  starting at position  $i$
  - $T_{ij} = \{A \in N \mid A \xrightarrow[G]{*} x_{ij}\}$

## Theorem

given CFG  $G = (N, \Sigma, P, S)$  and string  $x \in \Sigma^*$ , it is decidable whether  $x \in L(G)$

## Proof.

efficient and elegant algorithm: Cocke Kasami Younger (CKY)

- convert  $G$  into Chomsky normal form
- for all  $0 \leq i < j \leq |x|$ 
  - $x_{ij}$  is substring of  $x$  of length  $j - i$  starting at position  $i$
  - $T_{ij} = \{A \in N \mid A \xrightarrow[G]{*} x_{ij}\}$

compute  $T_{ij}$  by induction on  $j - i$

## Theorem

given CFG  $G = (N, \Sigma, P, S)$  and string  $x \in \Sigma^*$ , it is decidable whether  $x \in L(G)$

## Proof.

efficient and elegant algorithm: Cocke Kasami Younger (CKY)


- convert  $G$  into Chomsky normal form
- for all  $0 \leq i < j \leq |x|$ 
  - $x_{ij}$  is substring of  $x$  of length  $j - i$  starting at position  $i$
  - $T_{ij} = \{A \in N \mid A \xrightarrow{*}_G x_{ij}\}$

compute  $T_{ij}$  by induction on  $j - i$

- $x \in L(G) \iff S \in T_{0|x|}$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$




$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$



$x = baaba$


$$T_{ij} = \{X \in N \mid X \xrightarrow{*}_G x_{ij}\}$$

$$T_{01} = \{X \in N \mid X \xrightarrow{*}_G b\}$$



## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$



$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow{*}_G x_{ij}\}$$

$$\begin{aligned} T_{01} &= \{X \in N \mid X \xrightarrow{*}_G b\} \\ &= \{X \in N \mid X \rightarrow b\} \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0				
{B}	1			
		2		
			3	
			{B}	4
				5

$$x = baaba$$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned} T_{34} &= T_{01} = \{X \in N \mid X \xrightarrow[G]{*} b\} \\ &= \{X \in N \mid X \rightarrow b\} \\ &= \{B\} \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
	?	2			
		?	3		
			{B}	4	
				?	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned} T_{12} = T_{23} = T_{45} &= \{X \in N \mid X \xrightarrow[G]{*} a\} \\ &= \{X \in N \mid X \rightarrow a\} \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
	{A, C}	2			
		{A, C}	3		
			{B}	4	
				{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{12} = T_{23} = T_{45} &= \{X \in N \mid X \xrightarrow[G]{*} a\} \\
 &= \{X \in N \mid X \rightarrow a\} \\
 &= \{A, C\}
 \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
?	{A, C}	2			
		{A, C}	3		
			{B}	4	
				{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow_G^* x_{ij}\}$$

$$T_{02} = \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{01} \text{ and } Z \in T_{12}\}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
?	{A, C}	2			
		{A, C}	3		
			{B}	4	
				{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned} T_{02} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{01} \text{ and } Z \in T_{12}\} \\ &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{A, C\}\} \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
?	{A, C}	2			
		{A, C}	3		
			{B}	4	
				{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{02} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{01} \text{ and } Z \in T_{12}\} \\
 &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{A, C\}\} \\
 &= \{X \in N \mid X \rightarrow BA \in P \text{ or } X \rightarrow BC \in P\}
 \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S,A}	{A,C}	2			
		{A,C}	3		
			{B}	4	
				{A,C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{02} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{01} \text{ and } Z \in T_{12}\} \\
 &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{A, C\}\} \\
 &= \{X \in N \mid X \rightarrow BA \in P \text{ or } X \rightarrow BC \in P\} \\
 &= \{S, A\}
 \end{aligned}$$



## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
	?	{A, C}	3		
			{B}	4	
				{A, C}	5

$$x = baaba$$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{13} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{12} \text{ and } Z \in T_{23}\} \\
&= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{A, C\}\}
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
	{B}	{A, C}	3		
			{B}	4	
				{A, C}	5

$$x = baaba$$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{13} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{12} \text{ and } Z \in T_{23}\} \\
 &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{A, C\}\} \\
 &= \{B\}
 \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
	{B}	{A, C}	3		
		?	{B}	4	
				{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned} T_{24} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{23} \text{ and } Z \in T_{34}\} \\ &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{B\}\} \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
	{B}	{A, C}	3		
		{S, C}	{B}	4	
				{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{24} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{23} \text{ and } Z \in T_{34}\} \\
 &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{B\}\} \\
 &= \{S, C\}
 \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
	{B}	{A, C}	3		
		{S, C}	{B}	4	
			?	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned} T_{35} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{34} \text{ and } Z \in T_{45}\} \\ &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{A, C\}\} \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
	{B}	{A, C}	3		
		{S, C}	{B}	4	
			{S, A}	{A, C}	5

$$x = baaba$$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{35} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{34} \text{ and } Z \in T_{45}\} \\
&= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{A, C\}\} \\
&= \{S, A\}
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
?	{B}	{A, C}	3		
		{S, C}	{B}	4	
			{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$T_{03} = \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{01} \text{ and } Z \in T_{13}\}$$

$$\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{02} \text{ and } Z \in T_{23}\}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
?	{B}	{A, C}	3		
		{S, C}	{B}	4	
			{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{03} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{01} \text{ and } Z \in T_{13}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{02} \text{ and } Z \in T_{23}\} \\
 &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{B\}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{S, A\} \text{ and } Z \in \{A, C\}\}
 \end{aligned}$$



## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
		{S, C}	{B}	4	
			{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{03} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{01} \text{ and } Z \in T_{13}\} \\
&\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{02} \text{ and } Z \in T_{23}\} \\
&= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{B\}\} \\
&\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{S, A\} \text{ and } Z \in \{A, C\}\} \\
&= \emptyset
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
	?	{S, C}	{B}	4	
			{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$T_{14} = \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{12} \text{ and } Z \in T_{24}\} \\ \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{13} \text{ and } Z \in T_{34}\}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
	?	{S, C}	{B}	4	
			{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{14} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{12} \text{ and } Z \in T_{24}\} \\
&\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{13} \text{ and } Z \in T_{34}\} \\
&= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{S, C\}\} \\
&\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{B\}\}
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
	{B}	{S, C}	{B}	4	
			{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{14} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{12} \text{ and } Z \in T_{24}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{13} \text{ and } Z \in T_{34}\} \\
 &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{S, C\}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{B\}\} \\
 &= \{B\}
 \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
	{B}	{S, C}	{B}	4	
		?	{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$T_{25} = \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{23} \text{ and } Z \in T_{35}\} \\ \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{24} \text{ and } Z \in T_{45}\}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
	{B}	{S, C}	{B}	4	
		?	{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{25} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{23} \text{ and } Z \in T_{35}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{24} \text{ and } Z \in T_{45}\} \\
 &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{S, A\}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{S, C\} \text{ and } Z \in \{A, C\}\}
 \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
	{B}	{S, C}	{B}	4	
		{B}	{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{25} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{23} \text{ and } Z \in T_{35}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{24} \text{ and } Z \in T_{45}\} \\
 &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{S, A\}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{S, C\} \text{ and } Z \in \{A, C\}\} \\
 &= \{B\}
 \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0				
{B}	1			
{S, A}	{A, C}	2		
$\emptyset$	{B}	{A, C}	3	
?	{B}	{S, C}	{B}	4
		{B}	{S, A}	{A, C} 5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned} T_{04} = & \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{01} \text{ and } Z \in T_{14}\} \\ & \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{02} \text{ and } Z \in T_{24}\} \\ & \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{03} \text{ and } Z \in T_{34}\} \end{aligned}$$



## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
?	{B}	{S, C}	{B}	4	
		{B}	{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{04} = & \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{B\}\} \\
 & \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{S, A\} \text{ and } Z \in \{S, C\}\} \\
 & \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \emptyset \text{ and } Z \in \{B\}\}
 \end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
∅	{B}	{S, C}	{B}	4	
		{B}	{S, A}	{A, C}	5

$$x = baaba$$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{04} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{B\}\} \\
&\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{S, A\} \text{ and } Z \in \{S, C\}\} \\
&\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \emptyset \text{ and } Z \in \{B\}\} \\
&= \emptyset
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
∅	{B}	{S, C}	{B}	4	
	?	{B}	{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{15} = & \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{12} \text{ and } Z \in T_{25}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{13} \text{ and } Z \in T_{35}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{14} \text{ and } Z \in T_{45}\}
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
∅	{B}	{S, C}	{B}	4	
	?	{B}	{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{15} = & \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{B\}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{S, A\}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{A, C\}\}
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
∅	{B}	{S, C}	{B}	4	
	{S, A, C}	{B}	{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{15} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{A, C\} \text{ and } Z \in \{B\}\} \\
&\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{S, A\}\} \\
&\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{A, C\}\} \\
&= \{S, A, C\}
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC$   $A \rightarrow BA \mid a$   $B \rightarrow CC \mid b$   $C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
∅	{B}	{S, C}	{B}	4	
?	{S, A, C}	{B}	{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{05} = & \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{01} \text{ and } Z \in T_{15}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{02} \text{ and } Z \in T_{25}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{03} \text{ and } Z \in T_{35}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in T_{04} \text{ and } Z \in T_{45}\}
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
∅	{B}	{S, C}	{B}	4	
?	{S, A, C}	{B}	{S, A}	{A, C}	5

$x = baaba$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
T_{05} = & \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{S, A, C\}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{S, A\} \text{ and } Z \in \{B\}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \emptyset \text{ and } Z \in \{S, A\}\} \\
& \cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \emptyset \text{ and } Z \in \{A, C\}\}
\end{aligned}$$

## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
∅	{B}	{S, C}	{B}	4	
{S, A, C}	{S, A, C}	{B}	{S, A}	{A, C}	5

$x = \text{baaba}$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$\begin{aligned}
 T_{05} &= \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{B\} \text{ and } Z \in \{S, A, C\}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \{S, A\} \text{ and } Z \in \{B\}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \emptyset \text{ and } Z \in \{S, A\}\} \\
 &\cup \{X \in N \mid X \rightarrow YZ \in P \text{ with } Y \in \emptyset \text{ and } Z \in \{A, C\}\} \\
 &= \{S, A, C\}
 \end{aligned}$$



## Example

CFG  $G$  in Chomsky normal form:  $S \rightarrow AB \mid BC \quad A \rightarrow BA \mid a \quad B \rightarrow CC \mid b \quad C \rightarrow AB \mid a$

0					
{B}	1				
{S, A}	{A, C}	2			
∅	{B}	{A, C}	3		
∅	{B}	{S, C}	{B}	4	
{S, A, C}	{S, A, C}	{B}	{S, A}	{A, C}	5

$$x = baaba$$

$$T_{ij} = \{X \in N \mid X \xrightarrow[G]{*} x_{ij}\}$$

$$S \in T_{05} \implies x \in L(G)$$

Thanks! & Questions?