

CMPE 322/327 - Theory of Computation

Week 7: An Overview for the Midterm

Burak Ekici

April 4-8, 2022

Outline

- 1 Midterm
- 2 Finite Automata
- 3 Regular Expressions

Logistics & Questions

- on April the 10th in between 13h00 – 15h30 (UTC+3 İstanbul Time)

Logistics & Questions

- on April the 10th in between 13h00 – 15h30 (UTC+3 İstanbul Time)
- **three questions** (with sub-items):

Logistics & Questions

- on April the 10th in between 13h00 – 15h30 (UTC+3 İstanbul Time)
- **three questions** (with sub-items):
 - **designing DFAs recognizing certain regular languages** (no correctness proof is necessary, however provide a few sentences of explanation)

Logistics & Questions

- on April the 10th in between 13h00 – 15h30 (UTC+3 İstanbul Time)
- **three questions** (with sub-items):
 - designing DFAs recognizing certain regular languages (no correctness proof is necessary, however provide a few sentences of explanation)
 - $NFA_{\epsilon} \xrightarrow{\epsilon\text{-elimination}} NFA \xrightarrow{\text{subset construction}} DFA \xrightarrow{\text{minimization}} \text{minimal DFA}$ (with all intermediate steps clearly stated)

Logistics & Questions

- on April the 10th in between 13h00 – 15h30 (UTC+3 İstanbul Time)
- **three questions** (with sub-items):
 - designing DFAs recognizing certain regular languages (no correctness proof is necessary, however provide a few sentences of explanation)
 - $\text{NFA}_\epsilon \xrightarrow{\epsilon\text{-elimination}} \text{NFA} \xrightarrow{\text{subset construction}} \text{DFA} \xrightarrow{\text{minimization}} \text{minimal DFA}$ (with all intermediate steps clearly stated)
 - $\text{NFA}_\epsilon \rightarrow$ **regular expressions** (with all recursive calls to the algorithm (definition) clearly traced and computed)

Topics Covered Thus Far


deterministic finite state machines (DFA), nondeterminism in finite state machines (NFA), closure properties of regular sets, ε -transitions (NFA_ε), homomorphisms, minimization, Myhill-Nerode relations, regular expressions, derivatives, Kleene algebra


Topics Appear in the Midterm


deterministic finite state machines (DFA), nondeterminism in finite state machines (NFA), closure properties of regular sets, ϵ -transitions (NFA_ϵ), homomorphisms, minimization, Myhill-Nerode relations, regular expressions, derivatives, Kleene algebra


Outline



- 1 Midterm
- 2 Finite Automata
- 3 Regular Expressions











Topics Appear in the Midterm (Finite Automata)

deterministic finite state machines (DFA),

Topics Appear in the Midterm (Finite Automata)

deterministic finite state machines (DFA), **nondeterminism in finite state machines (NFA)**,

Topics Appear in the Midterm (Finite Automata)

deterministic finite state machines (DFA), nondeterminism in finite state machines (NFA), closure properties of regular sets,

Topics Appear in the Midterm (Finite Automata)

deterministic finite state machines (DFA), nondeterminism in finite state machines (NFA), closure properties of regular sets, ϵ -transitions (NFA_ϵ),

Topics Appear in the Midterm (Finite Automata)

deterministic finite state machines (DFA), nondeterminism in finite state machines (NFA), closure properties of regular sets, ϵ -transitions (NFA_ϵ), **minimization**

Definition

set $A \subseteq \Sigma^*$ is **regular** if $A = L(M)$ for some DFA M

Definition

set $A \subseteq \Sigma^*$ is regular if $A = L(M)$ for some DFA M

Theorem

regular sets are **effectively** closed under **intersection**,

Definition

set $A \subseteq \Sigma^*$ is regular if $A = L(M)$ for some DFA M

Theorem

regular sets are **effectively** closed under intersection, **complement**,

Definition

set $A \subseteq \Sigma^*$ is regular if $A = L(M)$ for some DFA M

Theorem

regular sets are **effectively** closed under intersection, complement, **union**,

Definition

set $A \subseteq \Sigma^*$ is regular if $A = L(M)$ for some DFA M

Theorem

regular sets are **effectively** closed under intersection, complement, union, **concatenation**,

Definition

set $A \subseteq \Sigma^*$ is regular if $A = L(M)$ for some DFA M

Theorem

regular sets are **effectively** closed under intersection, complement, union, concatenation, **asterate**,

Definition

set $A \subseteq \Sigma^*$ is regular if $A = L(M)$ for some DFA M

Theorem

regular sets are **effectively** closed under intersection, complement, union, concatenation, asterate, homomorphic image

Definition

set $A \subseteq \Sigma^*$ is regular if $A = L(M)$ for some DFA M

Theorem

regular sets are **effectively** closed under intersection, complement, union, concatenation, asterate, homomorphic image and homomorphic preimage

Definition

set $A \subseteq \Sigma^*$ is regular if $A = L(M)$ for some DFA M

Theorem

regular sets are effectively closed under intersection, complement, union, concatenation, asterate, homomorphic image and homomorphic preimage

Theorem

every set accepted by NFA is regular

Definition

set $A \subseteq \Sigma^*$ is regular if $A = L(M)$ for some DFA M

Theorem

regular sets are effectively closed under intersection, complement, union, concatenation, asterate, homomorphic image and homomorphic preimage

Theorem

every set accepted by NFA is regular

Theorem

every set accepted by NFA_{ϵ} is regular

Effective Computation (ε -elimination)

- $\text{NFA}_\varepsilon N_1 = (Q, \Sigma, \varepsilon, \Delta_1, S, F_1)$

Effective Computation (ε -elimination)

- $\text{NFA}_\varepsilon N_1 = (Q, \Sigma, \varepsilon, \Delta_1, S, F_1)$
- $L(N_1) = L(N_2)$ for NFA $N_2 = (Q, \Sigma, \Delta_2, S, F_2)$ with


Effective Computation (ε -elimination)

- $\text{NFA}_\varepsilon N_1 = (Q, \Sigma, \varepsilon, \Delta_1, S, F_1)$
- $L(N_1) = L(N_2)$ for NFA $N_2 = (Q, \Sigma, \Delta_2, S, F_2)$ with
 - ① $\Delta_2(q, a) := \widehat{\Delta}_1(\{q\}, a) \quad \forall q \in Q \ \forall a \in \Sigma$

Effective Computation (ε -elimination)

- $\text{NFA}_\varepsilon N_1 = (Q, \Sigma, \varepsilon, \Delta_1, S, F_1)$
- $L(N_1) = L(N_2)$ for $\text{NFA } N_2 = (Q, \Sigma, \Delta_2, S, F_2)$ with
 - ① $\Delta_2(q, a) := \widehat{\Delta}_1(\{q\}, a) \quad \forall q \in Q \ \forall a \in \Sigma$
 - ② $F_2 := \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$


Example (ϵ -elimination)



Example (ϵ -elimination)

$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	$\{1\}$	\emptyset	\emptyset	$\{2\}$
2	\emptyset	$\{2\}$	\emptyset	$\{3\}$
3	\emptyset	\emptyset	$\{3\}$	\emptyset




Example (ϵ -elimination)

$\text{NFA}_\epsilon N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$\text{NFA } N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\epsilon(\{q\}) \cap F_1 \neq \emptyset\}$



Example (ϵ -elimination)


$\text{NFA}_\epsilon N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$\text{NFA } N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\epsilon(\{q\}) \cap F_1 \neq \emptyset\}$

- | Δ | a | b | c |
|----------|-----|-----|-----|
| | | | |



Example (ϵ -elimination)


$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

- | Δ | a | b | c |
|----------|-----|-----|-----|
|----------|-----|-----|-----|



$$\Delta_2(1, a) = \widehat{\Delta}_1(\{1\}, a) = \cup \{C_{\epsilon}(\Delta_1(q, a)) \mid q \in \widehat{\Delta}_1(\{1\}, \epsilon)\}$$

Example (ϵ -elimination)


$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

- | Δ | a | b | c |
|----------|-----|-----|-----|
|----------|-----|-----|-----|



$$\begin{aligned}
 \Delta_2(1, a) &= \widehat{\Delta}_1(\{1\}, a) = \cup \{C_{\epsilon}(\Delta_1(q, a)) \mid q \in \widehat{\Delta}_1(\{1\}, \epsilon)\} \\
 &= C_{\epsilon}(\Delta_1(1, a)) \cup C_{\epsilon}(\Delta_1(2, a)) \cup C_{\epsilon}(\Delta_1(3, a))
 \end{aligned}$$

Example (ϵ -elimination)


$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

- | Δ | a | b | c |
|----------|-----|-----|-----|
|----------|-----|-----|-----|



$$\begin{aligned}
 \Delta_2(1, a) &= \widehat{\Delta}_1(\{1\}, a) = \cup \{C_{\epsilon}(\Delta_1(q, a)) \mid q \in \widehat{\Delta}_1(\{1\}, \epsilon)\} \\
 &= C_{\epsilon}(\Delta_1(1, a)) \cup C_{\epsilon}(\Delta_1(2, a)) \cup C_{\epsilon}(\Delta_1(3, a)) \\
 &= C_{\epsilon}(\{1\}) \cup C_{\epsilon}(\emptyset) \cup C_{\epsilon}(\emptyset)
 \end{aligned}$$

Example (ϵ -elimination)


$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

Δ	a	b	c
1	{1, 2, 3}		



$$\begin{aligned}
 \Delta_2(1, a) &= \widehat{\Delta}_1(\{1\}, a) = \cup \{C_{\epsilon}(\Delta_1(q, a)) \mid q \in \widehat{\Delta}_1(\{1\}, \epsilon)\} \\
 &= C_{\epsilon}(\Delta_1(1, a)) \cup C_{\epsilon}(\Delta_1(2, a)) \cup C_{\epsilon}(\Delta_1(3, a)) \\
 &= C_{\epsilon}(\{1\}) \cup C_{\epsilon}(\emptyset) \cup C_{\epsilon}(\emptyset) \\
 &= \{1, 2, 3\}
 \end{aligned}$$

Example (ε -elimination)


$\text{NFA}_\varepsilon N_1 = (\{1, 2, 3\}, \{a, b, c\}, \varepsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ε
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$\text{NFA } N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_\varepsilon(\{q\}) \cap F_1 \neq \emptyset\}$

Δ	a	b	c
1	{1, 2, 3}		



Example (ϵ -elimination)


$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

Δ	a	b	c
1	{1, 2, 3}		



$$\Delta_2(1, b) = \widehat{\Delta}_1(\{1\}, b) = \cup \{C_{\epsilon}(\Delta_1(q, b)) \mid q \in \widehat{\Delta}_1(\{1\}, \epsilon)\}$$

Example (ϵ -elimination)


NFA $_{\epsilon}$ $N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

NFA $N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

Δ	a	b	c
1	{1, 2, 3}		



$$\begin{aligned}
 \Delta_2(1, b) &= \widehat{\Delta}_1(\{1\}, b) = \cup \{C_{\epsilon}(\Delta_1(q, b)) \mid q \in \widehat{\Delta}_1(\{1\}, \epsilon)\} \\
 &= C_{\epsilon}(\Delta_1(1, b)) \cup C_{\epsilon}(\Delta_1(2, b)) \cup C_{\epsilon}(\Delta_1(3, b))
 \end{aligned}$$

Example (ϵ -elimination)


$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

Δ	a	b	c
1	{1, 2, 3}		



$$\begin{aligned}
 \Delta_2(1, b) &= \widehat{\Delta}_1(\{1\}, b) = \cup \{C_{\epsilon}(\Delta_1(q, b)) \mid q \in \widehat{\Delta}_1(\{1\}, \epsilon)\} \\
 &= C_{\epsilon}(\Delta_1(1, b)) \cup C_{\epsilon}(\Delta_1(2, b)) \cup C_{\epsilon}(\Delta_1(3, b)) \\
 &= C_{\epsilon}(\emptyset) \cup C_{\epsilon}(\{2\}) \cup C_{\epsilon}(\emptyset)
 \end{aligned}$$

Example (ϵ -elimination)


$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

Δ	a	b	c
1	{1, 2, 3}	{2, 3}	



$$\begin{aligned}
 \Delta_2(1, b) &= \widehat{\Delta}_1(\{1\}, b) = \cup \{C_{\epsilon}(\Delta_1(q, b)) \mid q \in \widehat{\Delta}_1(\{1\}, \epsilon)\} \\
 &= C_{\epsilon}(\Delta_1(1, b)) \cup C_{\epsilon}(\Delta_1(2, b)) \cup C_{\epsilon}(\Delta_1(3, b)) \\
 &= C_{\epsilon}(\emptyset) \cup C_{\epsilon}(\{2\}) \cup C_{\epsilon}(\emptyset) \\
 &= \{2, 3\}
 \end{aligned}$$

Example (ϵ -elimination cont'd)



$NFA_{\epsilon} N_1 = (\{1, 2, 3\}, \{a, b, c\}, \epsilon, \Delta_1, \{1\}, \{3\})$ with

Δ_1	a	b	c	ϵ
1	{1}	\emptyset	\emptyset	{2}
2	\emptyset	{2}	\emptyset	{3}
3	\emptyset	\emptyset	{3}	\emptyset

$NFA N_2 = (\{1, 2, 3\}, \{a, b, c\}, \Delta_2, \{1\}, F_2)$ with

- $F_2 = \{q \mid C_{\epsilon}(\{q\}) \cap F_1 \neq \emptyset\}$

Δ	a	b	c
1	{1, 2, 3}	{2, 3}	{3}
2	\emptyset	{2, 3}	{3}
3	\emptyset	\emptyset	{3}



Effective Computation (subset construction)

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$

Effective Computation (subset construction)

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with

Effective Computation (subset construction)

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - 1 $Q_M \quad \quad \quad := 2^{Q_N}$

Effective Computation (subset construction)

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - ① $Q_M := 2^{Q_N}$
 - ② $\delta_M(A, a) := \widehat{\Delta}(A, a) \quad \forall A \subseteq Q_N \quad \forall a \in \Sigma$


Effective Computation (subset construction)

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - ① $Q_M \quad \quad \quad := 2^{Q_N}$
 - ② $\delta_M(A, a) \quad := \widehat{\Delta}(A, a) \quad \quad \quad \forall A \subseteq Q_N \quad \forall a \in \Sigma$
 - ③ $s_M \quad \quad \quad := S_N$

Effective Computation (subset construction)

- NFA $N = (Q_N, \Sigma, \Delta_N, S_N, F_N)$
- $L(N) = L(M)$ for some DFA $M = (Q_M, \Sigma, \delta_M, s_M, F_M)$ with
 - ① $Q_M \quad \quad \quad := 2^{Q_N}$
 - ② $\delta_M(A, a) \quad := \widehat{\Delta}(A, a) \quad \quad \quad \forall A \subseteq Q_N \quad \forall a \in \Sigma$
 - ③ $s_M \quad \quad \quad := S_N$
 - ④ $F_M \quad \quad \quad := \{A \subseteq Q_N \mid A \cap F_N \neq \emptyset\}$


Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	a	b

Δ	a	b


Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	<i>a</i>	<i>b</i>
<i>A</i>	<i>A</i>	<i>A</i>

Δ	<i>a</i>	<i>b</i>


Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	<i>a</i>	<i>b</i>
A	A	A
B	E	B

Δ	<i>a</i>	<i>b</i>


Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	a	b
A	A	A
B	E	B
C	D	F

Δ	a	b


Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b


Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A


Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

Example (subset construction)


 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbbaababbabbbaababba




Example (subset construction)


 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

ab**b**baababbabbbaababba


Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba




Example (subset construction)


 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbb**a**ababbabbbaababba


Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba




Example (subset construction)


 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$

Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababba**bb**baababba


Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba



Example (subset construction)




$A = \emptyset$
 $B = \{1\}$
 $C = \{2\}$
 $D = \{3\}$

$E = \{1, 2\}$
 $F = \{1, 3\}$
 $G = \{2, 3\}$
 $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba



Example (subset construction)




$A = \emptyset$
 $B = \{1\}$
 $C = \{2\}$
 $D = \{3\}$

$E = \{1, 2\}$
 $F = \{1, 3\}$
 $G = \{2, 3\}$
 $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba




Example (subset construction)

 $A = \emptyset$ $B = \{1\}$ $C = \{2\}$ $D = \{3\}$ $E = \{1, 2\}$ $F = \{1, 3\}$ $G = \{2, 3\}$ $H = \{1, 2, 3\}$


Δ	a	b
A	A	A
B	E	B
C	D	F
D	D	A

Δ	a	b
E	H	F
F	H	B
G	D	F
H	H	F

abbbaababbabbbaababba



Example (subset construction)



$$B = \{1\}$$

$$E = \{1, 2\}$$

$$F = \{1, 3\}$$


$$H = \{1, 2, 3\}$$

Δ	a	b
B	E	B

Δ	a	b
E	H	F
F	H	B
H	H	F

abbbaababbabbbaababba

remove inaccessible states



Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

① remove inaccessible states

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

- ① remove inaccessible states
- ② for every two different states, determine whether they are distinguishable (marking)

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

- ① remove inaccessible states
- ② for every two different states, determine whether they are distinguishable (marking)
- ③ **collapse** indistinguishable states

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

- ① remove inaccessible states
- ② for every two different states, determine whether they are distinguishable (marking)
- ③ collapse indistinguishable states

Marking Algorithm

given DFA $M = (Q, \Sigma, \delta, s, F)$ without inaccessible states

- ① tabulate all unordered pairs $\{p, q\}$ with $p, q \in Q$, initially unmarked

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

- ① remove inaccessible states
- ② for every two different states, determine whether they are distinguishable (marking)
- ③ collapse indistinguishable states

Marking Algorithm

given DFA $M = (Q, \Sigma, \delta, s, F)$ without inaccessible states

- ① tabulate all unordered pairs $\{p, q\}$ with $p, q \in Q$, initially unmarked
- ② mark $\{p, q\}$ if $p \in F$ and $q \notin F$ or vice versa

Minimization Algorithm

DFA $M = (Q, \Sigma, \delta, s, F)$

- 1 remove inaccessible states
- 2 for every two different states, determine whether they are distinguishable (marking)
- 3 collapse indistinguishable states


Marking Algorithm

given DFA $M = (Q, \Sigma, \delta, s, F)$ without inaccessible states


- 1 tabulate all unordered pairs $\{p, q\}$ with $p, q \in Q$, initially unmarked
- 2 mark $\{p, q\}$ if $p \in F$ and $q \notin F$ or vice versa
- 3 repeat until no change:

mark $\{p, q\}$ if $\{\delta(p, a), \delta(q, a)\}$ is marked for some $a \in \Sigma$

Example




Example



a
b
c
d
e
f
g
h
i
j
k
l


✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓

Example






Example




a	b	
✓	✓	c
✓	✓	✓ d
	✓	✓ e
✓	✓	✓ ✓ f
✓	✓	✓ ✓ ✓ g
✓	✓	✓ ✓ ✓ ✓ h
✓	✓	✓ ✓ ✓ ✓ i
✓	✓	✓ ✓ ✓ ✓ j
✓	✓	✓ ✓ ✓ ✓ ✓ k
✓	✓	✓ ✓ ✓ ✓ ✓ ✓ l

Example




a
✓ b
✓ ✓ c
✓ ✓ ✓ d
✓ ✓ ✓ e
✓ ✓ ✓ f
✓ ✓ ✓ g
✓ ✓ ✓ h
✓ ✓ ✓ i
✓ ✓ ✓ j
✓ ✓ ✓ k
✓ ✓ ✓ l

Example



a
 ✓ b
 ✓ ✓ c
 ✓ ✓ ✓ d
 ✓ ✓ ✓ ✓ e
 ✓ ✓ ✓ ✓ f
 ✓ ✓ ✓ ✓ g
 ✓ ✓ ✓ ✓ h
 ✓ ✓ ✓ ✓ i
 ✓ ✓ ✓ ✓ j
 ✓ ✓ ✓ ✓ k
 ✓ ✓ ✓ ✓ l

Example





a
✓ b
✓ ✓ c
✓ ✓ ✓ d
✓ ✓ ✓ ✓ e
✓ ✓ ✓ ✓ ✓ f
✓ ✓ ✓ ✓ ✓ ✓ g
✓ ✓ ✓ ✓ ✓ ✓ ✓ h
✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ i
✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ j
✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ k
✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ ✓ l


states d, g and h, k can be merged



Outline



- 1 Midterm
- 2 Finite Automata
- 3 Regular Expressions**











Topics Appear in the Midterm (Regular Expressions)

DFAs to regular expressions

Definition

regular expressions are restricted patterns which use only

$$\mathbf{a} \in \Sigma \quad \epsilon \quad \emptyset \quad \alpha + \beta \quad \alpha^* \quad \alpha\beta$$

Definition

regular expressions are restricted patterns which use only

$$\mathbf{a} \in \Sigma \quad \epsilon \quad \emptyset \quad \alpha + \beta \quad \alpha^* \quad \alpha\beta$$

Theorem

finite automata and regular expressions are **equivalent**:

Definition

regular expressions are restricted patterns which use only

$$\mathbf{a} \in \Sigma \quad \epsilon \quad \emptyset \quad \alpha + \beta \quad \alpha^* \quad \alpha\beta$$

Theorem

finite automata and regular expressions are **equivalent**:

for all $A \subseteq \Sigma^*$ ① A is regular
 \iff ② $A = L(\alpha)$ for some regular expression α

Proof. (1 \implies 2 – An idea)

given $\text{NFA}_\varepsilon N_\varepsilon = (Q, \Sigma, \varepsilon, \Delta, S, F)$

$\forall Y \subseteq Q \quad \forall u, v \in Q$ construct regular expression α_{uv}^Y such that

$x \in L(\alpha_{uv}^Y) \iff \exists \text{ a path from } u \text{ to } v \text{ labeled } x \text{ (} v \in \widehat{\Delta}(\{u\}, x) \text{)}$
such that all intermediate states belong to Y

Proof. (1 \implies 2 – An idea)given $\text{NFA}_\epsilon N_\epsilon = (Q, \Sigma, \epsilon, \Delta, S, F)$ $\forall Y \subseteq Q \quad \forall u, v \in Q$ construct regular expression α_{uv}^Y such that

$$x \in L(\alpha_{uv}^Y) \iff \boxed{\exists \text{ a path from } u \text{ to } v \text{ labeled } x \ (v \in \widehat{\Delta}(\{u\}, x)) \text{ such that all intermediate states belong to } Y}$$

Definitions

$$\bullet \alpha_{uv}^\emptyset := \begin{cases} a_1 + \dots + a_k & \text{if } u \neq v \text{ and } k > 0 \\ \emptyset & \text{if } u \neq v \text{ and } k = 0 \\ a_1 + \dots + a_k + \epsilon & \text{if } u = v \text{ and } k > 0 \\ \epsilon & \text{if } u = v \text{ and } k = 0 \end{cases} \quad \{a_1, \dots, a_k\} := \{a \in \Sigma \cup \{\epsilon\} \mid v \in \Delta(u, a)\}$$

Proof. (1 \implies 2 – An idea)given NFA $_{\epsilon}$ $N_{\epsilon} = (Q, \Sigma, \epsilon, \Delta, S, F)$ $\forall Y \subseteq Q \quad \forall u, v \in Q$ construct regular expression α_{uv}^Y such that

$$x \in L(\alpha_{uv}^Y) \iff \boxed{\exists \text{ a path from } u \text{ to } v \text{ labeled } x \text{ (} v \in \widehat{\Delta}(\{u\}, x) \text{) such that all intermediate states belong to } Y}$$

Definitions

- $$\alpha_{uv}^{\emptyset} := \begin{cases} a_1 + \dots + a_k & \text{if } u \neq v \text{ and } k > 0 \\ \emptyset & \text{if } u \neq v \text{ and } k = 0 \\ a_1 + \dots + a_k + \epsilon & \text{if } u = v \text{ and } k > 0 \\ \epsilon & \text{if } u = v \text{ and } k = 0 \end{cases} \quad \{a_1, \dots, a_k\} := \{a \in \Sigma \cup \{\epsilon\} \mid v \in \Delta(u, a)\}$$
- $$\alpha_{uv}^Y := \alpha_{uv}^{Y-\{q\}} + \alpha_{uq}^{Y-\{q\}} (\alpha_{qq}^{Y-\{q\}})^* \alpha_{qv}^{Y-\{q\}} \quad \text{for some fixed } q \in Y$$

Proof. (1 \implies 2 – An idea)

given $\text{NFA}_\varepsilon N_\varepsilon = (Q, \Sigma, \varepsilon, \Delta, S, F)$

$\forall Y \subseteq Q \quad \forall u, v \in Q$ construct regular expression α_{uv}^Y such that

$x \in L(\alpha_{uv}^Y) \iff \exists \text{ a path from } u \text{ to } v \text{ labeled } x \text{ (} v \in \widehat{\Delta}(\{u\}, x) \text{)}$
such that all intermediate states belong to Y

Theorem

$$L(N_\varepsilon) = L\left(\sum_{s \in S, t \in F} \alpha_{st}^Q\right)$$

Thanks! & Questions?