

CMPE 322/327 - Theory of Computation

Week 12: Halting Problem & Reduction

Burak Ekici

May 16-20, 2022

Outline

1 A Quick Recap

2 Halting Problem

3 Reduction

Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with

Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with

- 1 Q : finite set of states

Definition


Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with

- 1 Q : finite set of states
- 2 Σ : input alphabet

Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with


- 1 Q : finite set of states
- 2 Σ : input alphabet
- 3 $\Gamma \supseteq \Sigma$: **tape** alphabet



Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with


- 1 Q : finite set of states
- 2 Σ : input alphabet
- 3 $\Gamma \supseteq \Sigma$: tape alphabet
- 4 $\vdash \in \Gamma - \Sigma$: **left endmarker**



Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with


- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\Gamma \supseteq \Sigma$: tape alphabet
- ④ $\vdash \in \Gamma - \Sigma$: left endmarker
- ⑤ $\sqcup \in \Gamma - \Sigma$: **blank symbol**



Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with


- 1 Q : finite set of states
- 2 Σ : input alphabet
- 3 $\Gamma \supseteq \Sigma$: tape alphabet
- 4 $\vdash \in \Gamma - \Sigma$: left endmarker
- 5 $\sqcup \in \Gamma - \Sigma$: blank symbol
- 6 $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$: **(partial) transition function**



Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with


- 1 Q : finite set of states
- 2 Σ : input alphabet
- 3 $\Gamma \supseteq \Sigma$: tape alphabet
- 4 $\vdash \in \Gamma - \Sigma$: left endmarker
- 5 $\sqcup \in \Gamma - \Sigma$: blank symbol
- 6 $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$: (partial) transition function
- 7 $s \in Q$: **start** state



Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with


- 1 Q : finite set of states
- 2 Σ : input alphabet
- 3 $\Gamma \supseteq \Sigma$: tape alphabet
- 4 $\vdash \in \Gamma - \Sigma$: left endmarker
- 5 $\sqcup \in \Gamma - \Sigma$: blank symbol
- 6 $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$: (partial) transition function
- 7 $s \in Q$: start state
- 8 $t \in Q$: **accept** state



Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with


- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\Gamma \supseteq \Sigma$: tape alphabet
- ④ $\vdash \in \Gamma - \Sigma$: left endmarker
- ⑤ $\sqcup \in \Gamma - \Sigma$: blank symbol
- ⑥ $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$: (partial) transition function
- ⑦ $s \in Q$: start state
- ⑧ $t \in Q$: accept state
- ⑨ $r \in Q - \{t\}$: **reject** state



Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with

- ① Q : finite set of states
- ② Σ : input alphabet
- ③ $\Gamma \supseteq \Sigma$: tape alphabet
- ④ $\vdash \in \Gamma - \Sigma$: left endmarker
- ⑤ $\sqcup \in \Gamma - \Sigma$: blank symbol
- ⑥ $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$: (partial) transition function
- ⑦ $s \in Q$: start state
- ⑧ $t \in Q$: accept state
- ⑨ $r \in Q - \{t\}$: reject state




such that

$$\forall a \in \Gamma \ \exists b, c \in \Gamma \ \exists d, e \in \{L, R\}: \delta(\textcolor{red}{t}, a) = (\textcolor{red}{t}, b, d) \quad \text{and} \quad \delta(\textcolor{red}{r}, a) = (\textcolor{red}{r}, c, e)$$

Definition

Turing machine (TM) is 9-tuple $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$ with

- 1 Q : finite set of states
- 2 Σ : input alphabet
- 3 $\Gamma \supseteq \Sigma$: tape alphabet
- 4 $\vdash \in \Gamma - \Sigma$: left endmarker
- 5 $\sqcup \in \Gamma - \Sigma$: blank symbol
- 6 $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$: (partial) transition function
- 7 $s \in Q$: start state
- 8 $t \in Q$: accept state
- 9 $r \in Q - \{t\}$: reject state



such that

$$\forall a \in \Gamma \ \exists b, c \in \Gamma \ \exists d, e \in \{L, R\}: \delta(t, a) = (t, b, d) \quad \text{and} \quad \delta(r, a) = (r, c, e)$$

$$\forall p \in Q \ \exists q \in Q: \delta(p, \vdash) = (q, \vdash, R)$$

Decision Problems

- halting problem for TMs

instance: TM M , string x

question: does M halt on x ?

Decision Problems

- halting problem for TMs
 - instance: TM M , string x
 - question: does M halt on x ?
- uniform halting problem for TMs
 - instance: TM M
 - question: does M halt on all inputs?

Decision Problems

- halting problem for TMs

instance: TM M , string x
question: does M halt on x ?

- uniform halting problem for TMs

instance: TM M
question: does M halt on all inputs?

- membership problem for CFGs

instance: CFG G , string x
question: $x \in L(G)$?

Decision Problems

- halting problem for TMs

instance: TM M , string x

question: does M halt on x ?

- uniform halting problem for TMs

instance: TM M

question: does M halt on all inputs?

- membership problem for CFGs

instance: CFG G , string x

question: $x \in L(G)$?

- equivalence problem for regular expressions

instance: regular expressions α, β

question: $L(\alpha) = L(\beta)$?

Decision Problems

- | | |
|---|---|
| <ul style="list-style-type: none"> • halting problem for TMs <ul style="list-style-type: none"> instance: TM M, string x question: does M halt on x? • uniform halting problem for TMs <ul style="list-style-type: none"> instance: TM M question: does M halt on all inputs? | <ul style="list-style-type: none"> • membership problem for CFGs <ul style="list-style-type: none"> instance: CFG G, string x question: $x \in L(G)$? • equivalence problem for regular expressions <ul style="list-style-type: none"> instance: regular expressions α, β question: $L(\alpha) = L(\beta)$? |
|---|---|

Decision Problems as Membership Problems

- code instance of problem as string over some alphabet

Decision Problems

- halting problem for TMs
 - instance: TM M , string x
 - question: does M halt on x ?
- uniform halting problem for TMs
 - instance: TM M
 - question: does M halt on all inputs?
- membership problem for CFGs
 - instance: CFG G , string x
 - question: $x \in L(G)$?
- equivalence problem for regular expressions
 - instance: regular expressions α, β
 - question: $L(\alpha) = L(\beta)$?

Decision Problems as Membership Problems

- code instance of problem as string over some alphabet
- language is set of all strings that correspond to yes instances

Definition (Encoding of Membership Problem for Turing Machines)

$$\text{MP} = \{\text{enc}(M) \# \text{enc}(x) \mid x \in L(M)\}$$

Definition (Encoding of Membership Problem for Turing Machines)

$$\text{MP} = \{ \quad M \# \quad x \mid x \in L(M) \}$$

Definition (Encoding of Membership Problem for Turing Machines)

$$\text{MP} = \{ M \# x \mid x \in L(M) \}$$

Theorem (Universal Turing Machine)

\exists TM U such that $L(U) = \text{MP}$

Definition (Encoding of Membership Problem for Turing Machines)

$MP = \{ M \# x \mid x \in L(M) \}$

Theorem (Universal Turing Machine)

\exists TM U such that $L(U) = MP$

universal TM can simulate any TM

Definition (Encoding of Membership Problem for Turing Machines)

$MP = \{ M \# x \mid x \in L(M) \}$

Theorem (Universal Turing Machine)

\exists TM U such that $L(U) = MP$

universal TM can simulate any TM

Corollary

MP is r.e.

How to Prove Undecidability?

- diagonalization
- reduction

Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$ is uncountable

Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$ is uncountable

Proof. (Diagonalization)

- suppose $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$ is countable

	A_0	A_1	A_2	\dots
x_0				
x_1				
x_2				
\vdots				

Theorem

 $\Sigma \neq \emptyset \implies 2^{\Sigma^*} \text{ is uncountable}$

Proof. (Diagonalization)

- suppose $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$ is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$ is countable

	A_0	A_1	A_2	\dots
x_0				
x_1				
x_2				
\vdots				

Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$ is uncountable

Proof. (Diagonalization)

- suppose $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$ is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$ is countable
- create infinite table

	A_0	A_1	A_2	\dots
x_0				
x_1				
x_2				
\vdots				

Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$ is uncountable

Proof. (Diagonalization)

- suppose $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$ is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$ is countable
- create infinite table

	A_0	A_1	A_2	...
x_0	×			
x_1	✓			
x_2	×			
\vdots				

- $A_0 = \{x_1\}$

Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$ is uncountable

Proof. (Diagonalization)

- suppose $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$ is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$ is countable
- create infinite table

	A_0	A_1	A_2	...
x_0	×	✓		
x_1	✓	×		
x_2	×	✓		
\vdots				

- $A_0 = \{x_1\}$ $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$

Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$ is uncountable

Proof. (Diagonalization)

- suppose $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$ is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$ is countable
- create infinite table

	A_0	A_1	A_2	...
x_0	×	✓	×	
x_1	✓	×	×	
x_2	×	✓	✓	
\vdots				

- $A_0 = \{x_1\}$ $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$ $A_2 = \{x_2, x_5\}$

Theorem

 $\Sigma \neq \emptyset \implies 2^{\Sigma^*}$ is uncountable

Proof. (Diagonalization)

- suppose $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$ is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$ is countable
- create infinite table

	A_0	A_1	A_2	...
x_0	×	✓	×	
x_1	✓	×	×	
x_2	×	✓	✓	
\vdots				

- $A_0 = \{x_1\}$ $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$ $A_2 = \{x_2, x_5\}$
- define $B = \{x_i \mid x_i \notin A_i\} \in 2^{\Sigma^*}$

Theorem

 $\Sigma \neq \emptyset \implies 2^{\Sigma^*}$ is uncountable

Proof. (Diagonalization)

- suppose $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$ is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$ is countable
- create infinite table

	A_0	A_1	A_2	...
x_0	✓	✓	×	
x_1	✓	✓	×	
x_2	×	✓	×	
\vdots				

- $A_0 = \{x_1\}$ $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$ $A_2 = \{x_2, x_5\}$
- define $B = \{x_i \mid x_i \notin A_i\} \in 2^{\Sigma^*}$
- $B \neq A_i \forall i$

Theorem

 $\Sigma \neq \emptyset \implies 2^{\Sigma^*}$ is uncountable

Proof. (Diagonalization)

- suppose $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$ is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$ is countable
- create infinite table

	A_0	A_1	A_2	...
x_0	✓	✓	×	
x_1	✓	✓	×	
x_2	×	✓	×	
\vdots				

- $A_0 = \{x_1\}$ $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$ $A_2 = \{x_2, x_5\}$
- define $B = \{x_i \mid x_i \notin A_i\} \in 2^{\Sigma^*}$
- $B \neq A_i \forall i$ ⚡

Outline

- 1 A Quick Recap
- 2 Halting Problem
- 3 Reduction

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Proof. ()

- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Proof. ()

- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive
- $HP = L(K)$ for some total TM K

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Proof. ()

- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive
- $HP = L(K)$ for some total TM K
- construct TM N that on input x
 - constructs M_x from x

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Proof. ()


- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive
- $HP = L(K)$ for some total TM K
- construct TM N that on input x
 - constructs M_x from x
 - runs K on input $M_x \# x$

Theorem


halting problem (HP) for TMs is undecidable (not recursive)

Proof. ()


- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive
- $HP = L(K)$ for some total TM K
- construct TM N that on input x
 - constructs M_x from x
 - runs K on input $M_x \# x$
 - accepts if K rejects and loops if K accepts




design of TM N




design of TM N




design of TM N



design of TM N




design of TM N

design of TM *N*

	ϵ	0	1	00	01	10	...
M_ϵ	x	✓	x	x	✓	x	...
M_0	✓	✓	x	x	x	✓	
M_1	x	x	✓	✓	✓	✓	
M_{00}	x	✓	✓	x	✓	x	
M_{01}	x	x	✓	x	✓	x	
M_{10}	✓	x	✓	x	x	x	
\vdots	\vdots						

✓: halts


x: loops



design of TM *N*

	ϵ	0	1	00	01	10	...
M_ϵ	×	✓	×	×	✓	×	...
M_0	✓	✓	×	×	×	✓	
M_1	×	×	✓	✓	✓	✓	
M_{00}	×	✓	✓	×	✓	×	
M_{01}	×	×	✓	×	✓	×	
M_{10}	✓	×	✓	×	×	×	
\vdots	\vdots						

✓: accepts
×: rejects

design of TM N

	ϵ	0	1	00	01	10	...
M_ϵ	✓	✓	x	x	✓	x	...
M_0	✓	x	x	x	x	✓	
M_1	x	x	x	✓	✓	✓	
M_{00}	x	✓	✓	✓	✓	x	
M_{01}	x	x	✓	x	x	x	
M_{10}	✓	x	✓	x	x	✓	
\vdots	\vdots						

✓: halts

x: loops

$$L(N) = \{\epsilon, 00, 10, \dots\}$$

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Proof. ()

- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive
- $HP = L(K)$ for some total TM K
- construct TM N that on input x
 - constructs M_x from x
 - runs K on input $M_x \# x$
 - accepts if K rejects and loops if K accepts
- for all inputs x N halts on x \iff K rejects $M_x \# x$

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Proof. ()

- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive
- $HP = L(K)$ for some total TM K
- construct TM N that on input x
 - constructs M_x from x
 - runs K on input $M_x \# x$
 - accepts if K rejects and loops if K accepts
- for all inputs x N halts on $x \iff K$ rejects $M_x \# x \iff M_x$ does not halt on x

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Proof. ()

- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive
- $HP = L(K)$ for some total TM K
- construct TM N that on input x
 - constructs M_x from x
 - runs K on input $M_x \# x$
 - accepts if K rejects and loops if K accepts
- for all inputs x N halts on $x \iff K$ rejects $M_x \# x \iff M_x$ does not halt on x
- N is different from all M_x

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Proof. ()

- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive
- $HP = L(K)$ for some total TM K
- construct TM N that on input x
 - constructs M_x from x
 - runs K on input $M_x \# x$
 - accepts if K rejects and loops if K accepts
- for all inputs x N halts on $x \iff K$ rejects $M_x \# x \iff M_x$ does not halt on x
- N is different from all M_x ⚡

Theorem

halting problem (HP) for TMs is undecidable (not recursive)

Proof. (diagonalization)

- suppose $HP = \{M \# x \mid \text{TM } M \text{ halts on input } x\}$ is recursive
- $HP = L(K)$ for some total TM K
- construct TM N that on input x
 - constructs M_x from x
 - runs K on input $M_x \# x$
 - accepts if K rejects and loops if K accepts
- for all inputs x N halts on x \iff K rejects $M_x \# x$ \iff M_x does not halt on x
- N is different from all M_x ⚡

Membership problem for TMs

instance: TM M , string x
question: does M accept x ?

Membership problem for TMs

instance: TM M , string x
question: does M accept x ?

Theorem

- 1 membership problem for TMs is **undecidable**:
 $\text{MP} = \{M \# x \mid x \in L(M)\}$ is not recursive

Membership problem for TMs

instance: TM M , string x
question: does M accept x ?

Theorem

- 1 membership problem for TMs is undecidable:
 $MP = \{M \# x \mid x \in L(M)\}$ is not recursive
- 2 membership problem for TMs is **semi-decidable** (MP is r.e.):
 $MP = L(U)$

Membership problem for TMs

instance: TM M , string x
question: does M accept x ?

Theorem

- 1 membership problem for TMs is undecidable:
 $MP = \{M \# x \mid x \in L(M)\}$ is not recursive
- 2 membership problem for TMs is semi-decidable (MP is r.e.):
 $MP = L(U)$

Proof.

- 1 **reduction** from halting problem for TMs

Membership problem for TMs

instance: TM M , string x
question: does M accept x ?

Theorem

- 1 membership problem for TMs is undecidable:
 $MP = \{M \# x \mid x \in L(M)\}$ is not recursive
- 2 membership problem for TMs is semi-decidable (MP is r.e.):
 $MP = L(U)$

Proof.

- 1 **reduction** from halting problem for TMs (next section)

Outline

- 1 A Quick Recap
- 2 Halting Problem
- 3 Reduction

Definition


(many-one) **reduction** of set $A \subseteq \Sigma^*$ to set $B \subseteq \Delta^*$ is total computable function $\sigma: \Sigma^* \rightarrow \Delta^*$ such that

$$x \in A \iff \sigma(x) \in B$$

Definition

(many-one) **reduction** of set $A \subseteq \Sigma^*$ to set $B \subseteq \Delta^*$ is total computable function $\sigma: \Sigma^* \rightarrow \Delta^*$ such that


$$x \in A \iff \sigma(x) \in B$$



Definition

(many-one) reduction of set $A \subseteq \Sigma^*$ to set $B \subseteq \Delta^*$ is total computable function $\sigma: \Sigma^* \rightarrow \Delta^*$ such that


$$x \in A \iff \sigma(x) \in B$$



Definition

(many-one) reduction of set $A \subseteq \Sigma^*$ to set $B \subseteq \Delta^*$ is total computable function $\sigma: \Sigma^* \rightarrow \Delta^*$ such that

$$x \in A \iff \sigma(x) \in B$$




Notation

$$A \leq_m B$$

Definition

(many-one) reduction of set $A \subseteq \Sigma^*$ to set $B \subseteq \Delta^*$ is total computable function $\sigma: \Sigma^* \rightarrow \Delta^*$ such that

$$x \in A \iff \sigma(x) \in B$$



Notation

$$A \leq_m B$$

$$\sigma: A \rightarrow_m B$$

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x
 - ① computes $\sigma(x)$

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.


- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x
 - 1 computes $\sigma(x)$
 - 2 runs M on input $\sigma(x)$

Theorem


if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.


- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x
 - ① computes $\sigma(x)$
 - ② runs M on input $\sigma(x)$
 - ③ accepts if M accepts and rejects if M rejects




design of TM *N*




design of TM N



design of TM N



design of TM N



design of TM N

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x
 - ① computes $\sigma(x)$
 - ② runs M on input $\sigma(x)$
 - ③ accepts if M accepts and rejects if M rejects
- for all inputs x
 N accepts $x \iff M$ accepts $\sigma(x)$

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x
 - ① computes $\sigma(x)$
 - ② runs M on input $\sigma(x)$
 - ③ accepts if M accepts and rejects if M rejects
- for all inputs x

$$N \text{ accepts } x \iff M \text{ accepts } \sigma(x) \iff \sigma(x) \in B$$

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x
 - ① computes $\sigma(x)$
 - ② runs M on input $\sigma(x)$
 - ③ accepts if M accepts and rejects if M rejects
- for all inputs x

$$N \text{ accepts } x \iff M \text{ accepts } \sigma(x) \iff \sigma(x) \in B \iff x \in A$$

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x
 - ① computes $\sigma(x)$
 - ② runs M on input $\sigma(x)$
 - ③ accepts if M accepts and rejects if M rejects
- for all inputs x

$$N \text{ accepts } x \iff M \text{ accepts } \sigma(x) \iff \sigma(x) \in B \iff x \in A$$

- $L(N) = A$

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x
 - ① computes $\sigma(x)$
 - ② runs M on input $\sigma(x)$
 - ③ accepts if M accepts and rejects if M rejects
- for all inputs x

$$N \text{ accepts } x \iff M \text{ accepts } \sigma(x) \iff \sigma(x) \in B \iff x \in A$$

- $L(N) = A \implies A$ is r.e.

Theorem

if $A \leq_m B$ and B is r.e. then A is r.e.

Proof.

- $\sigma: A \rightarrow_m B$
- $B = L(M)$ for some TM M
- construct TM N that on input x
 - ① computes $\sigma(x)$
 - ② runs M on input $\sigma(x)$
 - ③ accepts if M accepts and rejects if M rejects
- for all inputs x

$$N \text{ accepts } x \iff M \text{ accepts } \sigma(x) \iff \sigma(x) \in B \iff x \in A$$

- $L(N) = A \implies A$ is r.e.

Theorem (Equivalent)

if $A \leq_m B$ and A is not r.e. then B is not r.e.

Theorem

if $A \leq_m B$ and B is recursive then A is recursive

Theorem

if $A \leq_m B$ and B is recursive then A is recursive

Proof.

- $\sim A \leq_m \sim B$

Theorem

if $A \leq_m B$ and B is recursive then A is recursive

Proof.

- $\sim A \leq_m \sim B$
- B and $\sim B$ are r.e.

Theorem

if $A \leq_m B$ and B is recursive then A is recursive

Proof.

- $\sim A \leq_m \sim B$
- B and $\sim B$ are r.e.
- A and $\sim A$ are r.e. (by previous theorem)

Theorem

if $A \leq_m B$ and B is recursive then A is recursive

Proof.

- $\sim A \leq_m \sim B$
- B and $\sim B$ are r.e.
- A and $\sim A$ are r.e. (by previous theorem)
- A is recursive

Theorem

if $A \leq_m B$ and B is recursive then A is recursive

Proof.

- $\sim A \leq_m \sim B$
- B and $\sim B$ are r.e.
- A and $\sim A$ are r.e. (by previous theorem)
- A is recursive

Theorem (Equivalent)

if $A \leq_m B$ and A is not recursive then B is not recursive

Theorem

$HP \leq_m MP$

Theorem

$HP \leq_m MP$

Proof.

reduction $\sigma: HP \rightarrow_m MP$

Theorem

$HP \leq_m MP$

Proof.

reduction $\sigma: HP \rightarrow_m MP$

- σ transforms $M \# x$ into $N \# x$


Theorem

$HP \leq_m MP$


Proof.

reduction $\sigma: HP \rightarrow_m MP$


- σ transforms $M \# x$ into $N \# x$
- TM N is exactly like M but it accepts if M accepts or rejects




design of TM N



design of TM N



design of TM N



design of TM N

Theorem

$HP \leq_m MP$

Proof.

reduction $\sigma: HP \rightarrow_m MP$

- σ transforms $M \# x$ into $N \# x$
- TM N is exactly like M but it accepts if M accepts or rejects
- $M \# x \in HP \iff M \text{ halts on } x$

Theorem

$HP \leq_m MP$

Proof.

reduction $\sigma: HP \rightarrow_m MP$

- σ transforms $M \# x$ into $N \# x$
- TM N is exactly like M but it accepts if M accepts or rejects
- $M \# x \in HP \iff M \text{ halts on } x$
 $\iff N \text{ accepts } x$

Theorem

$HP \leq_m MP$

Proof.

reduction $\sigma: HP \rightarrow_m MP$

- σ transforms $M \# x$ into $N \# x$
- TM N is exactly like M but it accepts if M accepts or rejects
- $M \# x \in HP \iff M \text{ halts on } x$
 $\iff N \text{ accepts } x$
 $\iff N \# x \in MP$

Theorem

$HP \leq_m MP$

Proof.

reduction $\sigma: HP \rightarrow_m MP$

- σ transforms $M \# x$ into $N \# x$
- TM N is exactly like M but it accepts if M accepts or rejects
- $M \# x \in HP \iff M \text{ halts on } x$
 $\iff N \text{ accepts } x$
 $\iff N \# x \in MP$
 $\iff \sigma(M \# x) \in MP$

All Together

1 MP is r.e.

(by corollary in w11.pdf on slide #29)

All Together

- 1 MP is r.e. (by corollary in w11.pdf on slide #29)
- 2 HP is **not recursive** (by theorem on slide #9)

All Together

- 1 MP is r.e. (by corollary in w11.pdf on slide #29)
- 2 HP is not recursive (by theorem on slide #9)
- 3 $A \leq_m B$ and B is r.e. $\Rightarrow A$ is r.e. (by the first theorem on slides #14 – 15)

All Together

- 1 MP is r.e. (by corollary in w11.pdf on slide #29)
- 2 HP is not recursive (by theorem on slide #9)
- 3 $A \leq_m B$ and B is r.e. \Rightarrow A is r.e. (by the first theorem on slides #14 – 15)
- 4 $A \leq_m B$ and B is recursive \Rightarrow A is recursive (by the first theorem on slide #16)

All Together

- 1 MP is r.e. (by corollary in w11.pdf on slide #29)
- 2 HP is not recursive (by theorem on slide #9)
- 3 $A \leq_m B$ and B is r.e. \Rightarrow A is r.e. (by the first theorem on slides #14 – 15)
- 4 $A \leq_m B$ and B is recursive \Rightarrow A is recursive (by the first theorem on slide #16)
- 5 **HP \leq_m MP** (by theorem on slides #17 – 18)

All Together

- 1 MP is r.e. (by corollary in w11.pdf on slide #29)
- 2 HP is not recursive (by theorem on slide #9)
- 3 $A \leq_m B$ and B is r.e. \Rightarrow A is r.e. (by the first theorem on slides #14 – 15)
- 4 $A \leq_m B$ and B is recursive \Rightarrow A is recursive (by the first theorem on slide #16)
- 5 $HP \leq_m MP$ (by theorem on slides #17 – 18)

Theorem

MP is not recursive

All Together

- ① MP is r.e. (by corollary in w11.pdf on slide #29)
- ② HP is not recursive (by theorem on slide #9)
- ③ $A \leq_m B$ and B is r.e. \Rightarrow A is r.e. (by the first theorem on slides #14 – 15)
- ④ $A \leq_m B$ and B is recursive \Rightarrow A is recursive (by the first theorem on slide #16)
- ⑤ $HP \leq_m MP$ (by theorem on slides #17 – 18)

Theorem

MP is not recursive

Proof.

by statements 5, contrapositive of 4 and 2

All Together

- ① MP is r.e. (by corollary in w11.pdf on slide #29)
- ② HP is not recursive (by theorem on slide #9)
- ③ $A \leq_m B$ and B is r.e. \Rightarrow A is r.e. (by the first theorem on slides #14 – 15)
- ④ $A \leq_m B$ and B is recursive \Rightarrow A is recursive (by the first theorem on slide #16)
- ⑤ $HP \leq_m MP$ (by theorem on slides #17 – 18)

Theorem

MP is not recursive

Proof.

by statements 5, contrapositive of 4 and 2

Theorem

HP is r.e.

All Together

- ① MP is r.e. (by corollary in w11.pdf on slide #29)
- ② HP is not recursive (by theorem on slide #9)
- ③ $A \leq_m B$ and B is r.e. \Rightarrow A is r.e. (by the first theorem on slides #14 – 15)
- ④ $A \leq_m B$ and B is recursive \Rightarrow A is recursive (by the first theorem on slide #16)
- ⑤ $HP \leq_m MP$ (by theorem on slides #17 – 18)

Theorem

MP is not recursive

Proof.

by statements 5, contrapositive of 4 and 2


Theorem

HP is r.e.


Proof.

by statements 5, 3 and 1


Summary




Summary




Summary



Summary



Summary



$TOTAL = \{M \mid \text{TM } M \text{ halts on all inputs}\}$

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\Rightarrow A$ is r.e.

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$
- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN}$ and $\sim \text{HP}$ is not r.e. $\implies \text{FIN}$ is not r.e.

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN}$ and $\sim \text{HP}$ is not r.e. $\implies \text{FIN}$ is not r.e.

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN}$ and **True** $\implies \text{FIN}$ is not r.e.

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

reduction $\sigma: \sim \text{HP} \rightarrow_m \text{FIN}$

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

reduction $\sigma: \sim \text{HP} \rightarrow_m \text{FIN}$

- σ transforms $M \# x$ into TM N

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

reduction $\sigma: \sim \text{HP} \rightarrow_m \text{FIN}$

- σ transforms $M \# x$ into TM N that

- 1 erases its input

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

reduction $\sigma: \sim \text{HP} \rightarrow_m \text{FIN}$

- σ transforms $M \# x$ into TM N that

- ① erases its input
- ② writes x on its tape

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

reduction $\sigma: \sim \text{HP} \rightarrow_m \text{FIN}$

- σ transforms $M \# x$ into TM N that

- ① erases its input
- ② writes x on its tape
- ③ runs M on input x

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

reduction $\sigma: \sim \text{HP} \rightarrow_m \text{FIN}$

- σ transforms $M \# x$ into TM N that

- ① erases its input
- ② writes x on its tape
- ③ runs M on input x
- ④ accepts if M halts on x

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

reduction $\sigma: \sim \text{HP} \rightarrow_m \text{FIN}$

- σ transforms $M \# x$ into TM N that

- ① erases its input
- ② writes x on its tape
- ③ runs M on input x
- ④ accepts if M halts on x

- $M \# x \in \sim \text{HP} \iff M$ does not halt on x

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

reduction $\sigma: \sim \text{HP} \rightarrow_m \text{FIN}$

- σ transforms $M \# x$ into TM N that

- 1 erases its input
- 2 writes x on its tape
- 3 runs M on input x
- 4 accepts if M halts on x

- $M \# x \in \sim \text{HP} \iff M \text{ does not halt on } x \iff L(N) = \emptyset$

Theorem

finiteness problem for TMs is not semi-decidable (is not r.e.):

instance: TM M

question: is $L(M)$ finite ?

Proof.

$A \leq_m B$ and B is r.e. $\implies A$ is r.e.

- $A \leq_m B$ and A is not r.e. $\implies B$ is not r.e. (by the second theorem on slides #14 – 15)

- $A := \sim \text{HP}$

- $B := \text{FIN} = \{M \mid L(M) \text{ is finite}\}$

- $\sim \text{HP} \leq_m \text{FIN} \implies \text{FIN}$ is not r.e.

reduction $\sigma: \sim \text{HP} \rightarrow_m \text{FIN}$

- σ transforms $M \# x$ into TM N that

- ① erases its input
- ② writes x on its tape
- ③ runs M on input x
- ④ accepts if M halts on x

- $M \# x \in \sim \text{HP} \iff M \text{ does not halt on } x \iff L(N) = \emptyset \iff N \in \text{FIN}$

Thanks! & Questions?