

# CMPE 322/327 - Theory of Computation

## Week 11: Turing Machines & Decision Problems

Burak Ekici

May 9-13, 2022

# Outline

- 1 A Quick Recap
- 2 Turing Machines
- 3 Decision Problems
- 4 Encoding
- 5 Diagonalization

## Definitions

- **NPDA** is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with

## Definitions

- NPDA is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - 1  $Q$ : finite set of **states**

## Definitions

- NPDA is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - 1  $Q$ : finite set of states
  - 2  $\Sigma$ : input alphabet

## Definitions

- NPDA is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - 1  $Q$ : finite set of states
  - 2  $\Sigma$ : input alphabet
  - 3  $\Gamma$ : stack alphabet

## Definitions

- NPDA is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - 1  $Q$ : finite set of states
  - 2  $\Sigma$ : input alphabet
  - 3  $\Gamma$ : stack alphabet
  - 4  $\delta$ : finite subset of  $(Q \times (\Sigma \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$

## Definitions

- **NPDA** is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - ①  $Q$ : finite set of states
  - ②  $\Sigma$ : input alphabet
  - ③  $\Gamma$ : stack alphabet
  - ④  $\delta$ : finite subset of  $(Q \times (\Sigma \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$
  - ⑤  $s \in Q$ : **start** state



## Definitions

- **NPDA** is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - ①  $Q$ : finite set of states
  - ②  $\Sigma$ : input alphabet
  - ③  $\Gamma$ : stack alphabet
  - ④  $\delta$ : finite subset of  $(Q \times (\Sigma \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$
  - ⑤  $s \in Q$ : start state
  - ⑥  $\perp \in \Gamma$ : **initial** stack symbol

## Definitions

- **NPDA** is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - ①  $Q$ : finite set of states
  - ②  $\Sigma$ : input alphabet
  - ③  $\Gamma$ : stack alphabet
  - ④  $\delta$ : finite subset of  $(Q \times (\Sigma \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$
  - ⑤  $s \in Q$ : start state
  - ⑥  $\perp \in \Gamma$ : initial stack symbol
  - ⑦  $F \subseteq Q$ : **final** states

## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], []), (1, \epsilon)), ((1, [, []), (1, [[]), ((1, \epsilon, \perp), (2, \epsilon))\}$

## Definitions

- NPDA is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - 1  $Q$ : finite set of states
  - 2  $\Sigma$ : input alphabet
  - 3  $\Gamma$ : stack alphabet
  - 4  $\delta$ : finite subset of  $(Q \times (\Sigma \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$
  - 5  $s \in Q$ : start state
  - 6  $\perp \in \Gamma$ : initial stack symbol
  - 7  $F \subseteq Q$ : final states
- configuration**: element of  $Q \times \Sigma^* \times \Gamma^*$

## Definitions

- NPDA is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - 1  $Q$ : finite set of states
  - 2  $\Sigma$ : input alphabet
  - 3  $\Gamma$ : stack alphabet
  - 4  $\delta$ : finite subset of  $(Q \times (\Sigma \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$
  - 5  $s \in Q$ : start state
  - 6  $\perp \in \Gamma$ : initial stack symbol
  - 7  $F \subseteq Q$ : final states
- configuration**: element of  $Q \times \Sigma^* \times \Gamma^*$  (current state, remaining input, stack content)

## Definitions

- NPDA is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - 1  $Q$ : finite set of states
  - 2  $\Sigma$ : input alphabet
  - 3  $\Gamma$ : stack alphabet
  - 4  $\delta$ : finite subset of  $(Q \times (\Sigma \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$
  - 5  $s \in Q$ : start state
  - 6  $\perp \in \Gamma$ : initial stack symbol
  - 7  $F \subseteq Q$ : final states
- configuration: element of  $Q \times \Sigma^* \times \Gamma^*$  (current state, remaining input, stack content)
- start configuration on input  $x$ :  $(s, x, \perp)$

## Definitions

- NPDA is septuple  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$  with
  - 1  $Q$ : finite set of states
  - 2  $\Sigma$ : input alphabet
  - 3  $\Gamma$ : stack alphabet
  - 4  $\delta$ : finite subset of  $(Q \times (\Sigma \cup \{\epsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$
  - 5  $s \in Q$ : start state
  - 6  $\perp \in \Gamma$ : initial stack symbol
  - 7  $F \subseteq Q$ : final states
- configuration: element of  $Q \times \Sigma^* \times \Gamma^*$  (current state, remaining input, stack content)
- start configuration on input  $x$ :  $(s, x, \perp)$
- next configuration relation** is binary relation  $\xrightarrow[M]{1}$  defined as:  $(p, ay, A\beta) \xrightarrow[M]{1} (q, y, \gamma\beta)$   
for all  $((p, a, A), (q, \gamma)) \in \delta$  with  $a \in \Sigma \cup \{\epsilon\}$  and  $y \in \Sigma^*, \beta \in \Gamma^*$

## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((1, [, [\perp), (1, [\perp)), ((1, ], \perp), (2, \epsilon))\}$

input:      [ [ ] [ [ ] ] ]



## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((1, [, [\perp), (1, [\perp)), ((1, ], [\perp), (1, \epsilon)), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:     [ [ ] [ [ ] ] ]  
state:     1  
stack:     ⊥

## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((\textcolor{red}{1}, [\textcolor{red}{\perp}), (1, [\perp)), ((1, \textcolor{blue}{]}, (1, \epsilon)), ((1, \textcolor{blue}{[}}, (1, [\textcolor{blue}{[})), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:      $\textcolor{blue}{[} \textcolor{blue}{[} \textcolor{blue}{]} \textcolor{blue}{[} \textcolor{blue}{[} \textcolor{blue}{]} \textcolor{blue}{]}$   
state:      $\textcolor{red}{1}$   
stack:      $\textcolor{red}{\perp}$

## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((1, [, []), (1, [\perp)), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:     [ [ ] [ [ ] ] ]  
state:     1 1  
stack:     ⊥ ⊥  
           [

## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((1, [, ], (1, \epsilon)), ((1, ], [), (1, [\perp)), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:     [ [ ] [ [ ] ] ]  
state:     1 1  
stack:     ⊥ ⊥  
           [



## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((1, [, ], (1, [\perp)), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:     [ [ ] [ [ ] ] ]  
state:     1 1 1  
stack:     ⊥ ⊥ ⊥  
           [ [  
              ]

## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((1, [, [\perp), (1, [\perp)), ((1, ], [\perp), (1, \epsilon)), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:     [ [ ] [ [ ] ] ]  
state:     1 1 1 1  
stack:     ⊥ ⊥ ⊥ ⊥  
           [ [ [  
           [

## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((1, [, [), (1, [\perp)), ((1, ], [), (1, \epsilon)), ((1, [\perp), (1, [\perp)), ((1, ], [), (1, \epsilon)), ((1, [\perp), (1, [\perp)), ((1, ], [), (1, \epsilon)), ((1, [\perp), (1, [\perp)), ((1, ], [), (1, \epsilon))\}$

input:     [ [ ] [ [ ] ] ]  
state:     1 1 1 1  
stack:     ⊥ ⊥ ⊥ ⊥  
           [ [ [



## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((1, [, [\perp), (1, [\perp)), ((1, ], [\perp), (1, \epsilon)), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:     [ [ ] [ [ ] ] ]  
state:     1 1 1 1 1  
stack:     ⊥ ⊥ ⊥ ⊥ ⊥  
           [ [ [ [

## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((\mathbf{1}, [, \mathbf{]), (1, \epsilon)), ((\mathbf{1}, [, \mathbf{]), (1, \mathbf{[ ]}), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:     [ [ ] [ [ ] ] ]  
state:     1 1 1 1 **1**  
stack:     ⊥ ⊥ ⊥ ⊥ ⊥  
           [ [ [ [  
              [     [







## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], [\]), (1, \epsilon)), ((1, [, [\]), (1, [[\]), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:    [ [ ] [ [ ] ] ]  
state:    1 1 1 1 1 1 1  
stack:     $\perp \perp \perp \perp \perp \perp \perp$   
          [ [ [ [ [ [ [

## Example

$A = \{x \in \{[, ]\}^* \mid x \text{ is balanced}\}$  is accepted by NPDA  $M = (Q, \Sigma, \Gamma, \delta, 1, \perp, F)$  with

- ①  $Q = \{1, 2\}$
- ②  $\Sigma = \{[, ]\}$
- ③  $\Gamma = \{\perp, [\}$
- ④  $F = \{2\}$
- ⑤  $s = 1$
- ⑥  $\delta = \{((1, [, \perp), (1, [\perp)), ((1, ], \perp), (1, \epsilon)), ((1, [, [\perp), (1, [\perp)), ((1, ], [\perp), (1, [\perp)), ((1, \epsilon, \perp), (2, \epsilon))\}$

input:     [ [ ] [ [ ] ] ]  
state:     1 1 1 1 1 1 1 1  
stack:     ⊥ ⊥ ⊥ ⊥ ⊥ ⊥ ⊥ ⊥  
            [ [ [ [ [ [ [ [























## Definitions

NPDA  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$

- $\xrightarrow[n]{M} = \left(\xrightarrow[1]{M}\right)^n \quad \forall n \geq 0$

## Definitions

NPDA  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$

- $\xrightarrow[M]{n} = \left(\xrightarrow[M]{1}\right)^n \quad \forall n \geq 0$
- $\xrightarrow[M]{*} = \bigcup_{n \geq 0} \xrightarrow[M]{n}$

## Definitions

NPDA  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$

- $\xrightarrow[M]{n} = \left(\xrightarrow[M]{1}\right)^n \quad \forall n \geq 0$
- $\xrightarrow[M]{*} = \bigcup_{n \geq 0} \xrightarrow[M]{n}$
- $x \in \Sigma^*$  is **accepted by final state** if  $(s, x, \perp) \xrightarrow[M]{*} (q, \varepsilon, \alpha)$  with  $q \in F$

## Definitions

NPDA  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$

- $\xrightarrow[M]{n} = (\xrightarrow[M]{1})^n \quad \forall n \geq 0$
- $\xrightarrow[M]{*} = \bigcup_{n \geq 0} \xrightarrow[M]{n}$
- $x \in \Sigma^*$  is accepted by final state if  $(s, x, \perp) \xrightarrow[M]{*} (q, \varepsilon, \alpha)$  with  $q \in F$
- $L_f(M) = \{x \in \Sigma^* \mid x \text{ is accepted by final state}\}$

## Definitions

NPDA  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$

- $\xrightarrow[M]{n} = \left(\xrightarrow[M]{1}\right)^n \quad \forall n \geq 0$
- $\xrightarrow[M]{*} = \bigcup_{n \geq 0} \xrightarrow[M]{n}$
- $x \in \Sigma^*$  is accepted by final state if  $(s, x, \perp) \xrightarrow[M]{*} (q, \varepsilon, \alpha)$  with  $q \in F$
- $L_f(M) = \{x \in \Sigma^* \mid x \text{ is accepted by final state}\}$
- $x \in \Sigma^*$  is **accepted by empty stack** if  $(s, x, \perp) \xrightarrow[M]{*} (q, \varepsilon, \varepsilon)$

## Definitions

NPDA  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$

- $\xrightarrow[M]{n} = \left(\xrightarrow[M]{1}\right)^n \quad \forall n \geq 0$
- $\xrightarrow[M]{*} = \bigcup_{n \geq 0} \xrightarrow[M]{n}$
- $x \in \Sigma^*$  is accepted by final state if  $(s, x, \perp) \xrightarrow[M]{*} (q, \varepsilon, \alpha)$  with  $q \in F$
- $L_f(M) = \{x \in \Sigma^* \mid x \text{ is accepted by final state}\}$
- $x \in \Sigma^*$  is accepted by empty stack if  $(s, x, \perp) \xrightarrow[M]{*} (q, \varepsilon, \varepsilon)$
- $L_e(M) = \{x \in \Sigma^* \mid x \text{ is accepted by empty stack}\}$

## Definitions

NPDA  $M = (Q, \Sigma, \Gamma, \delta, s, \perp, F)$

- $\xrightarrow[M]{n} = \left(\xrightarrow[M]{1}\right)^n \quad \forall n \geq 0$
- $\xrightarrow[M]{*} = \bigcup_{n \geq 0} \xrightarrow[M]{n}$
- $x \in \Sigma^*$  is accepted by final state if  $(s, x, \perp) \xrightarrow[M]{*} (q, \epsilon, \alpha)$  with  $q \in F$
- $L_f(M) = \{x \in \Sigma^* \mid x \text{ is accepted by final state}\}$
- $x \in \Sigma^*$  is accepted by empty stack if  $(s, x, \perp) \xrightarrow[M]{*} (q, \epsilon, \epsilon)$
- $L_e(M) = \{x \in \Sigma^* \mid x \text{ is accepted by empty stack}\}$

## Theorem

CFGs and NPDAs are **equivalent**:

- 1  $A = L(G)$  for some CFG  $G \iff$
- 2  $A = L_f(M)$  for some NPDA  $M$



## Theorem

CFGs and NPDAs are **equivalent**:

- ①  $A = L(G)$  for some CFG  $G \iff$
- ②  $A = L_f(M)$  for some NPDA  $M \iff$
- ③  $A = L_e(M)$  for some NPDA  $M$

## Theorem

CFGs and NPDA are **equivalent**:

- ①  $A = L(G)$  for some CFG  $G \iff$
- ②  $A = L_f(M)$  for some NPDA  $M \iff$
- ③  $A = L_e(M)$  for some NPDA  $M \iff$
- ④  $A = L_e(M) = L_f(M)$  for some NPDA  $M$

## Definition

A **deterministic pushdown automaton (DPDA)** is an octuple  $M = (Q, \Sigma, \Gamma, \delta, \perp, \dashv, s, F)$

- ①  $\dashv$  is a special symbol not in  $\Sigma$ , called the right endmarker
- ② for any  $p \in Q, a \in \Sigma \cup \{\varepsilon\}, A \in \Gamma$ , the set  $\delta \subseteq (Q \times (\Sigma \cup \{\dashv\} \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$  contains
  - at most one element of the form  $((p, a, A), (q, \beta))$
  - exactly one transition of the form  $((p, a, A), (q, \beta))$  or  $((p, \varepsilon, A), (q, \beta))$

## Definition

A deterministic pushdown automaton (DPDA) is an octuple  $M = (Q, \Sigma, \Gamma, \delta, \perp, \vdash, s, F)$

- ①  $\vdash$  is a special symbol not in  $\Sigma$ , called the right endmarker
- ② for any  $p \in Q$ ,  $a \in \Sigma \cup \{\varepsilon\}$ ,  $A \in \Gamma$ , the set  $\delta \subseteq (Q \times (\Sigma \cup \{\vdash\} \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$  contains
  - at most one element of the form  $((p, a, A), (q, \beta))$
  - exactly one transition of the form  $((p, a, A), (q, \beta))$  or  $((p, \varepsilon, A), (q, \beta))$

## Remark

- **DPDAs** are strictly less powerful than NPDAs

## Definition

A deterministic pushdown automaton (DPDA) is an octuple  $M = (Q, \Sigma, \Gamma, \delta, \perp, \vdash, s, F)$

- ①  $\vdash$  is a special symbol not in  $\Sigma$ , called the right endmarker
- ② for any  $p \in Q$ ,  $a \in \Sigma \cup \{\varepsilon\}$ ,  $A \in \Gamma$ , the set  $\delta \subseteq (Q \times (\Sigma \cup \{\vdash\} \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$  contains
  - at most one element of the form  $((p, a, A), (q, \beta))$
  - exactly one transition of the form  $((p, a, A), (q, \beta))$  or  $((p, \varepsilon, A), (q, \beta))$

## Remark

- DPDAs are strictly less powerful than NPDAs
- **deterministic context-free language** is set accepted by DPDA


## Definition

A deterministic pushdown automaton (DPDA) is an octuple  $M = (Q, \Sigma, \Gamma, \delta, \perp, \neg, s, F)$

- 1  $\neg$  is a special symbol not in  $\Sigma$ , called the right endmarker
- 2 for any  $p \in Q$ ,  $a \in \Sigma \cup \{\varepsilon\}$ ,  $A \in \Gamma$ , the set  $\delta \subseteq (Q \times (\Sigma \cup \{\neg\} \cup \{\varepsilon\}) \times \Gamma) \times (Q \times \Gamma^*)$  contains
  - at most one element of the form  $((p, a, A), (q, \beta))$
  - exactly one transition of the form  $((p, a, A), (q, \beta))$  or  $((p, \varepsilon, A), (q, \beta))$

## Remark

- DPDAs are strictly less powerful than NPDA
- **deterministic context-free language** is set accepted by DPDA



$$A = \{a^i b^j c^k \mid i = j \text{ or } j = k\}$$

## Theorem

context-free sets are effectively closed under

- union
- concatenation
- asterate
- homomorphic image
- homomorphic preimage

## Theorem

context-free sets are effectively closed under

- union
- concatenation
- asterate
- homomorphic image
- homomorphic preimage

context-free sets are **not** closed under

- intersection
- complement



## Theorem

deterministic context-free sets are effectively closed under

- complement
- homomorphic preimage

## Theorem

deterministic context-free sets are effectively closed under


- complement
- homomorphic preimage

deterministic context-free sets are **not** closed under

- union
- intersection
- concatenation
- asterate
- homomorphic image


# Outline

- 1 A Quick Recap
- 2 Turing Machines**
- 3 Decision Problems
- 4 Encoding
- 5 Diagonalization



recursively enumerable


Turing machines  
unrestricted grammars




$$\{A \mid A \subseteq \Sigma^*\} = 2^{\Sigma^*}$$

recursively enumerable

Turing machines  
unrestricted grammars



$$\{A \mid A \subseteq \Sigma^*\} = 2^{\Sigma^*}$$



## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with



## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with

- 1  $Q$ : finite set of states

## Definition


**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with

- 1  $Q$ : finite set of states
- 2  $\Sigma$ : input alphabet

## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with


- ①  $Q$ : finite set of states
- ②  $\Sigma$ : input alphabet
- ③  $\Gamma \supseteq \Sigma$ : **tape** alphabet



## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with


- ①  $Q$ : finite set of states
- ②  $\Sigma$ : input alphabet
- ③  $\Gamma \supseteq \Sigma$ : tape alphabet
- ④  $\vdash \in \Gamma - \Sigma$ : **left endmarker**



## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with


- 1  $Q$ : finite set of states
- 2  $\Sigma$ : input alphabet
- 3  $\Gamma \supseteq \Sigma$ : tape alphabet
- 4  $\vdash \in \Gamma - \Sigma$ : left endmarker
- 5  $\sqcup \in \Gamma - \Sigma$ : **blank symbol**



## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with


- 1  $Q$ : finite set of states
- 2  $\Sigma$ : input alphabet
- 3  $\Gamma \supseteq \Sigma$ : tape alphabet
- 4  $\vdash \in \Gamma - \Sigma$ : left endmarker
- 5  $\sqcup \in \Gamma - \Sigma$ : blank symbol
- 6  $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$ : **(partial) transition function**



## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with


- 1  $Q$ : finite set of states
- 2  $\Sigma$ : input alphabet
- 3  $\Gamma \supseteq \Sigma$ : tape alphabet
- 4  $\vdash \in \Gamma - \Sigma$ : left endmarker
- 5  $\sqcup \in \Gamma - \Sigma$ : blank symbol
- 6  $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$ : (partial) transition function
- 7  $s \in Q$ : **start** state



## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with

- 1  $Q$ : finite set of states
- 2  $\Sigma$ : input alphabet
- 3  $\Gamma \supseteq \Sigma$ : tape alphabet
- 4  $\vdash \in \Gamma - \Sigma$ : left endmarker
- 5  $\sqcup \in \Gamma - \Sigma$ : blank symbol
- 6  $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$ : (partial) transition function
- 7  $s \in Q$ : start state
- 8  $t \in Q$ : **accept** state






## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with

- ①  $Q$ : finite set of states
- ②  $\Sigma$ : input alphabet
- ③  $\Gamma \supseteq \Sigma$ : tape alphabet
- ④  $\vdash \in \Gamma - \Sigma$ : left endmarker
- ⑤  $\sqcup \in \Gamma - \Sigma$ : blank symbol
- ⑥  $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$ : (partial) transition function
- ⑦  $s \in Q$ : start state
- ⑧  $t \in Q$ : accept state
- ⑨  $r \in Q - \{t\}$ : **reject** state




## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with

- 1  $Q$ : finite set of states
- 2  $\Sigma$ : input alphabet
- 3  $\Gamma \supseteq \Sigma$ : tape alphabet
- 4  $\vdash \in \Gamma - \Sigma$ : left endmarker
- 5  $\sqcup \in \Gamma - \Sigma$ : blank symbol
- 6  $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$ : (partial) transition function
- 7  $s \in Q$ : start state
- 8  $t \in Q$ : accept state
- 9  $r \in Q - \{t\}$ : reject state

such that


$$\forall a \in \Gamma \ \exists b, c \in \Gamma \ \exists d, e \in \{L, R\}: \delta(t, a) = (t, b, d) \quad \text{and} \quad \delta(r, a) = (r, c, e)$$



## Definition

**Turing machine (TM)** is 9-tuple  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  with

- ①  $Q$ : finite set of states
- ②  $\Sigma$ : input alphabet
- ③  $\Gamma \supseteq \Sigma$ : tape alphabet
- ④  $\vdash \in \Gamma - \Sigma$ : left endmarker
- ⑤  $\sqcup \in \Gamma - \Sigma$ : blank symbol
- ⑥  $\delta: Q \times \Gamma \rightarrow Q \times \Gamma \times \{L, R\}$ : (partial) transition function
- ⑦  $s \in Q$ : start state
- ⑧  $t \in Q$ : accept state
- ⑨  $r \in Q - \{t\}$ : reject state



such that

$$\forall a \in \Gamma \ \exists b, c \in \Gamma \ \exists d, e \in \{L, R\}: \delta(t, a) = (t, b, d) \quad \text{and} \quad \delta(r, a) = (r, c, e)$$

$$\forall p \in Q \ \exists q \in Q: \delta(p, \vdash) = (q, \vdash, R)$$

## Example

$A = \{a^n b^n c^n \mid n \geq 0\} = L(M)$  for TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, 1, t, r)$  with

- ①  $Q = \{1, 2, 3, 4, 5, t, r\}$
- ②  $\Sigma = \{a, b, c\}$
- ③  $\Gamma = \{a, b, c, \vdash, \sqcup, X, x\}$

④ $\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(r, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$


## Example

$A = \{a^n b^n c^n \mid n \geq 0\} = L(M)$  for TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, 1, t, r)$  with

- ①  $Q = \{1, 2, 3, 4, 5, t, r\}$
- ②  $\Sigma = \{a, b, c\}$
- ③  $\Gamma = \{a, b, c, \vdash, \sqcup, X, x\}$

④ $\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, —, —)	(r, —, —)	(t, —, —)	(5, X, R)	(r, —, —)
2	(r, —, —)	(2, a, R)	(3, X, R)	(r, c, R)	(r, —, —)	(2, X, R)	(r, —, —)
3	(r, —, —)	(r, —, —)	(3, b, R)	(4, x, L)	(r, —, —)	(r, —, —)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, —, —)	(4, X, L)	(4, x, L)
5	(r, —, —)	(r, a, R)	(r, —, —)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	—	—	—	—	—	—	—
r	—	—	—	—	—	—	—

## Example



# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, X, R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, X, R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, X, R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, X, L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , X, R)	(5, X, R)	(5, $x$ , R)

 $a^2b^2c^2$ 
 $\vdash aabbcc \sqcup^\omega$   
1

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, X, R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, X, R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, X, R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, X, L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , X, R)	(5, X, R)	(5, $x$ , R)

$a^2b^2c^2$   
 $\vdash_{\text{1}} aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash_{\text{1}} \textcolor{red}{a}abbcc \sqcup^\omega$



# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, X, R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, X, R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, X, R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, X, L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , X, R)	(5, X, R)	(5, $x$ , R)

$a^2b^2c^2$   
 $\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash \textcolor{red}{a}bbcc \sqcup^\omega$   
 $\textcolor{blue}{1} \qquad \qquad \textcolor{blue}{1} \qquad \qquad \textcolor{red}{2}$

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, -, -)	(r, -, -)	(t, -, -)	(5, X, R)	(r, -, -)
2	(r, -, -)	(2, a, R)	(3, X, R)	(r, c, R)	(r, -, -)	(2, X, R)	(2, -, -)
3	(r, -, -)	(r, -, -)	(3, b, R)	(4, x, L)	(r, -, -)	(r, -, -)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, -, -)	(4, X, L)	(4, x, L)
5	(r, -, -)	(r, a, R)	(r, -, -)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)

$a^2b^2c^2$

$\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega$

1      1      2      2

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

$a^2b^2c^2$

$\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega$

1                      1                      2                      2                      3

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

$a^2b^2c^2$

$$\begin{array}{c}
\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \\
\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega
\end{array}$$

1 1 2 2 3 3

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

$a^2b^2c^2$

$\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega$

$\xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega$

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

  
 $a^2b^2c^2$ 
  

$$\begin{array}{ccccccc}
\vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega \\
1 & & 1 & & 2 & & 2 & & 3 \\
& \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & & \\
& & 3 & & 4 & & 4 & & 
\end{array}$$

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

  
 $a^2b^2c^2$ 
  

$$\begin{array}{ccccccc}
\underbrace{\vdash aabbcc \sqcup^\omega}_1 & \xrightarrow{M} & \underbrace{\vdash aabbcc \sqcup^\omega}_1 & \xrightarrow{M} & \underbrace{\vdash abbcc \sqcup^\omega}_2 & \xrightarrow{M} & \underbrace{\vdash abbcc \sqcup^\omega}_2 & \xrightarrow{M} & \underbrace{\vdash aXbcc \sqcup^\omega}_3 \\
& & \xrightarrow{M} & \underbrace{\vdash aXbcc \sqcup^\omega}_3 & \xrightarrow{M} & \underbrace{\vdash aXbxc \sqcup^\omega}_4 & \xrightarrow{M} & \underbrace{\vdash aXbxc \sqcup^\omega}_4 & \xrightarrow{M} & \underbrace{\vdash aXbxc \sqcup^\omega}_4
\end{array}$$

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

$a^2b^2c^2$

$$\begin{array}{l}
\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \\
\vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \\
\vdash aXbxc \sqcup^\omega
\end{array}$$



## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

$a^2b^2c^2$

$$\begin{array}{ccccccc}
\vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega \\
1 & & 1 & & 2 & & 2 & & 3 \\
& \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega \\
& & 3 & & 4 & & 4 & & 4 \\
& \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & & & & \\
& & 4 & & 1 & & & & 
\end{array}$$

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

$a^2b^2c^2$

$$\begin{array}{ccccccc}
\vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega \\
1 & & 1 & & 2 & & 2 & & 3 \\
& \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega \\
& & 3 & & 4 & & 4 & & 4 \\
& \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash Xbxc \sqcup^\omega \\
& & 4 & & 1 & & 2
\end{array}$$

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

  
 $a^2b^2c^2$ 
  

$$\begin{array}{ccccccc}
\underbrace{\vdash aabbcc \sqcup^\omega}_1 & \xrightarrow[M]{1} & \vdash \underbrace{aabbcc \sqcup^\omega}_1 & \xrightarrow[M]{1} & \vdash \underbrace{abbcc \sqcup^\omega}_2 & \xrightarrow[M]{1} & \vdash \underbrace{abbcc \sqcup^\omega}_2 & \xrightarrow[M]{1} & \vdash \underbrace{aXbcc \sqcup^\omega}_3 \\
& & \xrightarrow[M]{1} & \vdash \underbrace{aXbcc \sqcup^\omega}_3 & \xrightarrow[M]{1} & \vdash \underbrace{aXbxc \sqcup^\omega}_4 & \xrightarrow[M]{1} & \vdash \underbrace{aXbxc \sqcup^\omega}_4 & \xrightarrow[M]{1} & \vdash \underbrace{aXbxc \sqcup^\omega}_4 \\
& & \xrightarrow[M]{1} & \vdash \underbrace{aXbxc \sqcup^\omega}_4 & \xrightarrow[M]{1} & \vdash \underbrace{aXbxc \sqcup^\omega}_1 & \xrightarrow[M]{1} & \vdash \underbrace{Xbxc \sqcup^\omega}_2 & \xrightarrow[M]{1} & \vdash \underbrace{Xbxc \sqcup^\omega}_2
\end{array}$$

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

  
 $a^2b^2c^2$ 
  

$$\begin{array}{l}
\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \\
\vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \\
\vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \\
\vdash \vdash \vdash XXxc \sqcup^\omega
\end{array}$$

$$a^2 b^2 c^2$$

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

$\vdash aabbcc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aabbcc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \textcolor{blue}{abbcc} \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \textcolor{blue}{abbcc} \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aX\textcolor{blue}{bcc} \sqcup^\omega$
1		1		2		2		3
	$\xrightarrow[M]{1}$	$\vdash aX\textcolor{blue}{bcc} \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aX\textcolor{blue}{bxc} \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aX\textcolor{blue}{bxc} \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aX\textcolor{blue}{bxc} \sqcup^\omega$
		3		4		4		4
	$\xrightarrow[M]{1}$	$\vdash aX\textcolor{blue}{bxc} \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aX\textcolor{blue}{bxc} \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \textcolor{blue}{Xbxc} \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \textcolor{blue}{Xbxc} \sqcup^\omega$
		4		1		2		2
	$\xrightarrow[M]{1}$	$\vdash \textcolor{blue}{XXxc} \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \textcolor{blue}{XXxc} \sqcup^\omega$				
		3		3				

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

  
 $a^2b^2c^2$ 
  

$$\begin{array}{ccccccc}
\vdash aabbcc \sqcup^\omega & \xrightarrow{1/M} & \vdash aabbcc \sqcup^\omega & \xrightarrow{1/M} & \vdash abbcc \sqcup^\omega & \xrightarrow{1/M} & \vdash abbcc \sqcup^\omega & \xrightarrow{1/M} & \vdash aXbcc \sqcup^\omega \\
1 & & 1 & & 2 & & 2 & & 3 \\
& \xrightarrow{1/M} & \vdash aXbcc \sqcup^\omega & \xrightarrow{1/M} & \vdash aXbxc \sqcup^\omega & \xrightarrow{1/M} & \vdash aXbxc \sqcup^\omega & \xrightarrow{1/M} & \vdash aXbxc \sqcup^\omega \\
& & 3 & & 4 & & 4 & & 4 \\
& \xrightarrow{1/M} & \vdash aXbxc \sqcup^\omega & \xrightarrow{1/M} & \vdash aXbxc \sqcup^\omega & \xrightarrow{1/M} & \vdash \vdash Xbxc \sqcup^\omega & \xrightarrow{1/M} & \vdash \vdash Xbxc \sqcup^\omega \\
& & 4 & & 1 & & 2 & & 2 \\
& \xrightarrow{1/M} & \vdash \vdash \vdash XXxc \sqcup^\omega & \xrightarrow{1/M} & \vdash \vdash \vdash XXxc \sqcup^\omega & \xrightarrow{1/M} & \vdash \vdash \vdash XXxc \sqcup^\omega & & \\
& & 3 & & 3 & & 4 & & 
\end{array}$$

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

  
 $a^2b^2c^2$ 
  

$$\begin{array}{ccccccc}
\vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega \\
1 & & 1 & & 2 & & 2 & & 3 \\
& \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega \\
& & 3 & & 4 & & 4 & & 4 \\
& \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash Xbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash Xbxc \sqcup^\omega \\
& & 4 & & 1 & & 2 & & 2 \\
& \xrightarrow[M]{1} & \vdash \vdash \vdash XXxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash X\cancel{X}xx \sqcup^\omega \\
& & 3 & & 3 & & 4 & & 4
\end{array}$$

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

  
 $a^2b^2c^2$ 
  
 $\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash XXXc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash XXXx \sqcup^\omega$



$$a^2 b^2 c^2$$

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

$\vdash aabbcc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aabbcc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash abbcc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash abbcc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aXbcc \sqcup^\omega$
1		1		2		2		3
	$\xrightarrow[M]{1}$	$\vdash aXbcc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aXbxc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aXbxc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aXbxc \sqcup^\omega$
		3		4		4		4
	$\xrightarrow[M]{1}$	$\vdash aXbxc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash aXbxc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \vdash Xbxc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \vdash Xbxc \sqcup^\omega$
		4		1		2		2
	$\xrightarrow[M]{1}$	$\vdash \vdash \vdash XXxc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \vdash \vdash XXxc \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \vdash \vdash XXXx \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \vdash \vdash XXXx \sqcup^\omega$
		3		3		4		4
	$\xrightarrow[M]{1}$	$\vdash \vdash \vdash XXXx \sqcup^\omega$	$\xrightarrow[M]{1}$	$\vdash \vdash \vdash XXXx \sqcup^\omega$				
		4		4				

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

  
 $a^2b^2c^2$ 
  
 $\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash Xbxc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash \vdash \vdash XXxc \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \vdash XXxc \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \vdash XXxx \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \vdash XXxx \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash \vdash \vdash XXxx \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \vdash XXxx \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \vdash XXxx \sqcup^\omega$

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

  
 $a^2b^2c^2$ 
  
 $\vdash aabbcc \sqcup^\omega \xrightarrow{1/M} \vdash aabbcc \sqcup^\omega \xrightarrow{1/M} \vdash abbcc \sqcup^\omega \xrightarrow{1/M} \vdash abbcc \sqcup^\omega \xrightarrow{1/M} \vdash aXbcc \sqcup^\omega$ 
  
 $\xrightarrow{1/M} \vdash aXbcc \sqcup^\omega \xrightarrow{1/M} \vdash aXbxc \sqcup^\omega \xrightarrow{1/M} \vdash aXbxc \sqcup^\omega \xrightarrow{1/M} \vdash aXbxc \sqcup^\omega$ 
  
 $\xrightarrow{1/M} \vdash aXbxc \sqcup^\omega \xrightarrow{1/M} \vdash aXbxc \sqcup^\omega \xrightarrow{1/M} \vdash \vdash Xbxc \sqcup^\omega \xrightarrow{1/M} \vdash \vdash Xbxc \sqcup^\omega$ 
  
 $\xrightarrow{1/M} \vdash \vdash \vdash XXxc \sqcup^\omega \xrightarrow{1/M} \vdash \vdash \vdash XXxc \sqcup^\omega \xrightarrow{1/M} \vdash \vdash \vdash XXxx \sqcup^\omega \xrightarrow{1/M} \vdash \vdash \vdash XXxx \sqcup^\omega$ 
  
 $\xrightarrow{1/M} \vdash \vdash \vdash XXxx \sqcup^\omega \xrightarrow{1/M} \vdash \vdash \vdash XXxx \sqcup^\omega \xrightarrow{1/M} \vdash \vdash \vdash XXxx \sqcup^\omega \xrightarrow{1/M} \vdash \vdash \vdash X\color{red}{X}xx \sqcup^\omega$

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

  
 $a^2b^2c^2$ 
  
 $\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXc \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash XXXc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega$ 
  
 $\xrightarrow[M]{1} \vdash XXXx \sqcup^\omega$

$$a^2 b^2 c^2$$

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	$(1, \vdash, R)$	$(2, \vdash, R)$	$(r, -, -)$	$(r, -, -)$	$(t, -, -)$	$(5, X, R)$	$(r, -, -)$
2	$(r, -, -)$	$(2, a, R)$	$(3, X, R)$	$(r, c, R)$	$(r, -, -)$	$(2, X, R)$	$(2, -, -)$
3	$(r, -, -)$	$(r, -, -)$	$(3, b, R)$	$(4, x, L)$	$(r, -, -)$	$(r, -, -)$	$(3, x, R)$
4	$(1, \vdash, R)$	$(4, a, L)$	$(4, b, L)$	$(4, c, L)$	$(r, -, -)$	$(4, X, L)$	$(4, x, L)$
5	$(r, -, -)$	$(r, a, R)$	$(r, -, -)$	$(r, c, R)$	$(t, X, R)$	$(5, X, R)$	$(5, x, R)$

$\vdash aabbcc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash aabbcc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash abbcc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash abbcc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash aXbcc \sqcup^\omega$
1		1		2		2		3
$\xrightarrow{1/M}$		$\vdash aXbcc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash aXbxc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash aXbxc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash aXbxc \sqcup^\omega$
		3		4		4		4
$\xrightarrow{1/M}$		$\vdash aXbxc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash aXbxc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash \vdash Xbxc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash \vdash Xbxc \sqcup^\omega$
		4		1		2		2
$\xrightarrow{1/M}$		$\vdash \vdash XXxc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash \vdash XXxc \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash \vdash XXxx \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash \vdash XXxx \sqcup^\omega$
		3		3		4		4
$\xrightarrow{1/M}$		$\vdash \vdash XXxx \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash \vdash XXxx \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash \vdash XXxx \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash \vdash XXxx \sqcup^\omega$
		4		4		1		5
$\xrightarrow{1/M}$		$\vdash \vdash XXxx \sqcup^\omega$	$\xrightarrow{1/M}$	$\vdash \vdash XXxX \sqcup^\omega$				
		5		5				

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

$a^2b^2c^2$

$$\begin{array}{l}
\vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aabbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash abbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \\
\vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbcc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \\
\vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash aXbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \\
\vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash Xbxc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXc \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \\
\vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \\
\vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega \xrightarrow[M]{1} \vdash XXXx \sqcup^\omega
\end{array}$$

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	(2, $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

$a^2b^2c^2 \in L(M)$

$$\begin{array}{ccccccc}
\vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aabbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash abbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega \\
1 & & 1 & & 2 & & 2 & & 3 \\
& \xrightarrow[M]{1} & \vdash aXbcc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega \\
& & 3 & & 4 & & 4 & & 4 \\
& \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash aXbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash Xbxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash Xbxc \sqcup^\omega \\
& & 4 & & 1 & & 2 & & 2 \\
& \xrightarrow[M]{1} & \vdash \vdash \vdash XXxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxc \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega \\
& & 3 & & 3 & & 4 & & 4 \\
& \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega \\
& & 4 & & 4 & & 1 & & 5 \\
& \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXxx \sqcup^\omega & \xrightarrow[M]{1} & \vdash \vdash \vdash XXXxx \sqcup^\omega \\
& & 5 & & 5 & & 5 & & t
\end{array}$$

## Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	( $r$ , $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)

acb

$\vdash acb \sqcup^\omega$   
1



# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	( $r$ , $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)
$t$	$-$	$-$	$-$	$-$	$-$	$-$	$-$
$r$	$-$	$-$	$-$	$-$	$-$	$-$	$-$

acb

$$\underset{1}{\vdash} \text{acb} \sqcup^\omega \xrightarrow[M]{1} \vdash \underset{1}{\text{acb}} \sqcup^\omega$$

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	( $r$ , $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)
$t$	$-$	$-$	$-$	$-$	$-$	$-$	$-$
$r$	$-$	$-$	$-$	$-$	$-$	$-$	$-$

acb

$$\underset{1}{\vdash} \text{acb} \sqcup^\omega \xrightarrow[M]{1} \vdash \underset{1}{\text{acb}} \sqcup^\omega \xrightarrow[M]{1} \vdash \underset{2}{\text{cb}} \sqcup^\omega$$

# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, -, -)	(r, -, -)	(t, -, -)	(5, X, R)	(r, -, -)
2	(r, -, -)	(2, a, R)	(3, X, R)	(r, c, R)	(r, -, -)	(2, X, R)	(r, -, -)
3	(r, -, -)	(r, -, -)	(3, b, R)	(4, x, L)	(r, -, -)	(r, -, -)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, -, -)	(4, X, L)	(4, x, L)
5	(r, -, -)	(r, a, R)	(r, -, -)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	—	—	—	—	—	—	—
r	—	—	—	—	—	—	—

acb  $\notin L(M)$

$$\underset{1}{\vdash} \text{acb} \sqcup^\omega \xrightarrow[M]{1} \vdash \underset{1}{\text{acb}} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{2}{\text{cb}} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{r}{\text{cb}} \sqcup^\omega$$

# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, —, —)	(r, —, —)	(t, —, —)	(5, X, R)	(r, —, —)
2	(r, —, —)	(2, a, R)	(3, X, R)	(r, c, R)	(r, —, —)	(2, X, R)	(r, —, —)
3	(r, —, —)	(r, —, —)	(3, b, R)	(4, x, L)	(r, —, —)	(r, —, —)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, —, —)	(4, X, L)	(4, x, L)
5	(r, —, —)	(r, a, R)	(r, —, —)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	—	—	—	—	—	—	—
r	—	—	—	—	—	—	—

$acb \notin L(M)$

$$\underset{1}{\vdash} \underset{1}{a} \underset{2}{c} b \sqcup^{\omega} \xrightarrow[M]{1} \vdash \underset{1}{a} \underset{2}{c} b \sqcup^{\omega} \xrightarrow[M]{1} \vdash \vdash \underset{2}{c} b \sqcup^{\omega} \xrightarrow[M]{1} \vdash \vdash \underset{r}{c} b \sqcup^{\omega}$$

abca

$$\underset{1}{\vdash} a b c a \sqcup^{\omega}$$

# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, —, —)	(r, —, —)	(t, —, —)	(5, X, R)	(r, —, —)
2	(r, —, —)	(2, a, R)	(3, X, R)	(r, c, R)	(r, —, —)	(2, X, R)	(r, —, —)
3	(r, —, —)	(r, —, —)	(3, b, R)	(4, x, L)	(r, —, —)	(r, —, —)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, —, —)	(4, X, L)	(4, x, L)
5	(r, —, —)	(r, a, R)	(r, —, —)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	—	—	—	—	—	—	—
r	—	—	—	—	—	—	—

$acb \notin L(M)$

$$\vdash_{1} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{1} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{2} cb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{r} cb \sqcup^{\omega}$$

abca

$$\vdash_{1} abca \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{1} abca \sqcup^{\omega}$$

# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, -, -)	(r, -, -)	(t, -, -)	(5, X, R)	(r, -, -)
2	(r, -, -)	(2, a, R)	(3, X, R)	(r, c, R)	(r, -, -)	(2, X, R)	(r, -, -)
3	(r, -, -)	(r, -, -)	(3, b, R)	(4, x, L)	(r, -, -)	(r, -, -)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, -, -)	(4, X, L)	(4, x, L)
5	(r, -, -)	(r, a, R)	(r, -, -)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	—	—	—	—	—	—	—
r	—	—	—	—	—	—	—

$acb \notin L(M)$

$$\vdash_{\text{1}} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{\text{1}} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{\text{2}} cb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{\text{r}} cb \sqcup^{\omega}$$

abca

$$\vdash_{\text{1}} abca \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{\text{1}} abca \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{\text{2}} bca \sqcup^{\omega}$$

# Example

$\delta$	$\vdash$	$a$	$b$	$c$	$\sqcup$	$X$	$x$
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	( $t$ , $-$ , $-$ )	(5, $X$ , R)	( $r$ , $-$ , $-$ )
2	( $r$ , $-$ , $-$ )	(2, $a$ , R)	(3, $X$ , R)	( $r$ , $c$ , R)	( $r$ , $-$ , $-$ )	(2, $X$ , R)	( $r$ , $-$ , $-$ )
3	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $b$ , R)	(4, $x$ , L)	( $r$ , $-$ , $-$ )	( $r$ , $-$ , $-$ )	(3, $x$ , R)
4	(1, $\vdash$ , R)	(4, $a$ , L)	(4, $b$ , L)	(4, $c$ , L)	( $r$ , $-$ , $-$ )	(4, $X$ , L)	(4, $x$ , L)
5	( $r$ , $-$ , $-$ )	( $r$ , $a$ , R)	( $r$ , $-$ , $-$ )	( $r$ , $c$ , R)	( $t$ , $X$ , R)	(5, $X$ , R)	(5, $x$ , R)
$t$	$-$	$-$	$-$	$-$	$-$	$-$	$-$
$r$	$-$	$-$	$-$	$-$	$-$	$-$	$-$

$acb \notin L(M)$

$$\vdash_{1} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{1} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{2} cb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{r} cb \sqcup^{\omega}$$

$abca$

$$\vdash_{1} abca \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{1} abca \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{2} bca \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{3} Xca \sqcup^{\omega}$$

# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, -, -)	(r, -, -)	(t, -, -)	(5, X, R)	(r, -, -)
2	(r, -, -)	(2, a, R)	(3, X, R)	(r, c, R)	(r, -, -)	(2, X, R)	(r, -, -)
3	(r, -, -)	(r, -, -)	(3, b, R)	(4, x, L)	(r, -, -)	(r, -, -)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, -, -)	(4, X, L)	(4, x, L)
5	(r, -, -)	(r, a, R)	(r, -, -)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	-	-	-	-	-	-	-
r	-	-	-	-	-	-	-

$acb \notin L(M)$

$$\vdash_{1} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{1} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{2} cb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{r} cb \sqcup^{\omega}$$

abca

$$\vdash_{1} abca \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{1} abca \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{2} bca \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{3} Xca \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{4} Xxa \sqcup^{\omega}$$



# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, -, -)	(r, -, -)	(t, -, -)	(5, X, R)	(r, -, -)
2	(r, -, -)	(2, a, R)	(3, X, R)	(r, c, R)	(r, -, -)	(2, X, R)	(r, -, -)
3	(r, -, -)	(r, -, -)	(3, b, R)	(4, x, L)	(r, -, -)	(r, -, -)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, -, -)	(4, X, L)	(4, x, L)
5	(r, -, -)	(r, a, R)	(r, -, -)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	-	-	-	-	-	-	-
r	-	-	-	-	-	-	-

$acb \notin L(M)$

$$\vdash_{1} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{1} acb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{2} cb \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{r} cb \sqcup^{\omega}$$

abca

$$\vdash_{1} abca \sqcup^{\omega} \xrightarrow[M]{1} \vdash_{1} abca \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{2} bca \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{3} Xca \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{4} Xxa \sqcup^{\omega} \xrightarrow[M]{1} \vdash\vdash_{4} Xxa \sqcup^{\omega}$$

# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, -, -)	(r, -, -)	(t, -, -)	(5, X, R)	(r, -, -)
2	(r, -, -)	(2, a, R)	(3, X, R)	(r, c, R)	(r, -, -)	(2, X, R)	(r, -, -)
3	(r, -, -)	(r, -, -)	(3, b, R)	(4, x, L)	(r, -, -)	(r, -, -)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, -, -)	(4, X, L)	(4, x, L)
5	(r, -, -)	(r, a, R)	(r, -, -)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	-	-	-	-	-	-	-
r	-	-	-	-	-	-	-

$acb \notin L(M)$

$$\underset{1}{\vdash} acb \sqcup^\omega \xrightarrow[M]{1} \vdash \underset{1}{acb} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{2}{cb} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{r}{cb} \sqcup^\omega$$

abca

$$\begin{aligned} \underset{1}{\vdash} abca \sqcup^\omega &\xrightarrow[M]{1} \vdash \underset{1}{abca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{2}{bca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{3}{Xca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{4}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{4}{Xxa} \sqcup^\omega \\ &\xrightarrow[M]{1} \vdash \vdash \underset{1}{Xxa} \sqcup^\omega \end{aligned}$$

# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, -, -)	(r, -, -)	(t, -, -)	(5, X, R)	(r, -, -)
2	(r, -, -)	(2, a, R)	(3, X, R)	(r, c, R)	(r, -, -)	(2, X, R)	(r, -, -)
3	(r, -, -)	(r, -, -)	(3, b, R)	(4, x, L)	(r, -, -)	(r, -, -)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, -, -)	(4, X, L)	(4, x, L)
5	(r, -, -)	(r, a, R)	(r, -, -)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	-	-	-	-	-	-	-
r	-	-	-	-	-	-	-

$acb \notin L(M)$

$$\underset{1}{\vdash} \underset{1}{ac} b \sqcup^\omega \xrightarrow[M]{1} \vdash \underset{1}{ac} b \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{2}{cb} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{r}{cb} \sqcup^\omega$$

abca

$$\begin{aligned} \underset{1}{\vdash} \underset{1}{ab} ca \sqcup^\omega &\xrightarrow[M]{1} \vdash \underset{1}{ab} ca \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{2}{bca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{3}{Xca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{4}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{4}{Xxa} \sqcup^\omega \\ &\xrightarrow[M]{1} \vdash \vdash \underset{1}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{5}{Xxa} \sqcup^\omega \end{aligned}$$

# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, -, -)	(r, -, -)	(t, -, -)	(5, X, R)	(r, -, -)
2	(r, -, -)	(2, a, R)	(3, X, R)	(r, c, R)	(r, -, -)	(2, X, R)	(r, -, -)
3	(r, -, -)	(r, -, -)	(3, b, R)	(4, x, L)	(r, -, -)	(r, -, -)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, -, -)	(4, X, L)	(4, x, L)
5	(r, -, -)	(r, a, R)	(r, -, -)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	-	-	-	-	-	-	-
r	-	-	-	-	-	-	-

$acb \notin L(M)$

$$\underset{1}{\vdash} acb \sqcup^\omega \xrightarrow[M]{1} \vdash \underset{1}{acb} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{2}{cb} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{r}{cb} \sqcup^\omega$$

abca

$$\begin{aligned} \underset{1}{\vdash} abca \sqcup^\omega &\xrightarrow[M]{1} \vdash \underset{1}{abca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{2}{bca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{3}{Xca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{4}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{4}{Xxa} \sqcup^\omega \\ &\xrightarrow[M]{1} \vdash \vdash \underset{1}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{5}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{5}{Xxa} \sqcup^\omega \end{aligned}$$

# Example

$\delta$	$\vdash$	a	b	c	$\sqcup$	X	x
1	(1, $\vdash$ , R)	(2, $\vdash$ , R)	(r, -, -)	(r, -, -)	(t, -, -)	(5, X, R)	(r, -, -)
2	(r, -, -)	(2, a, R)	(3, X, R)	(r, c, R)	(r, -, -)	(2, X, R)	(r, -, -)
3	(r, -, -)	(r, -, -)	(3, b, R)	(4, x, L)	(r, -, -)	(r, -, -)	(3, x, R)
4	(1, $\vdash$ , R)	(4, a, L)	(4, b, L)	(4, c, L)	(r, -, -)	(4, X, L)	(4, x, L)
5	(r, -, -)	(r, a, R)	(r, -, -)	(r, c, R)	(t, X, R)	(5, X, R)	(5, x, R)
t	-	-	-	-	-	-	-
r	-	-	-	-	-	-	-

$acb \notin L(M)$

$$\underset{1}{\vdash} \underset{1}{ac} \sqcup^\omega \xrightarrow[M]{1} \vdash \underset{1}{acb} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{2}{cb} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{r}{cb} \sqcup^\omega$$

$abca \notin L(M)$

$$\begin{aligned} \underset{1}{\vdash} \underset{1}{ab} \underset{1}{ca} \sqcup^\omega &\xrightarrow[M]{1} \vdash \underset{1}{ab} \underset{1}{ca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{2}{bca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{3}{Xca} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{4}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{4}{Xxa} \sqcup^\omega \\ &\xrightarrow[M]{1} \vdash \vdash \underset{1}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{5}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{5}{Xxa} \sqcup^\omega \xrightarrow[M]{1} \vdash \vdash \underset{r}{Xxa} \sqcup^\omega \end{aligned}$$

## Definitions

- **configuration:** element of  $Q \times \{y \sqcup^{\omega} \mid y \in \Gamma^*\} \times \mathbb{N}$

## Definitions

- configuration: element of  $Q \times \{y \sqcup^\omega \mid y \in \Gamma^*\} \times \mathbb{N}$
- **start configuration** on input  $x \in \Sigma^*$ :  $(s, \vdash x \sqcup^\omega, 0)$

## Definitions

- configuration: element of  $Q \times \{y \sqcup^\omega \mid y \in \Gamma^*\} \times \mathbb{N}$
- start configuration on input  $x \in \Sigma^*$ :  $(s, \vdash x \sqcup^\omega, 0)$
- **next configuration relation** is binary relation  $\xrightarrow[M]{1}$  defined as:

$$(p, z, n) \xrightarrow[M]{1} \begin{cases} (q, z', n-1) & \text{if } \delta(p, z_n) = (q, b, L) \\ (q, z', n+1) & \text{if } \delta(p, z_n) = (q, b, R) \end{cases}$$

with

- $z_n$ :  $n$ -th symbol of  $z$



## Definitions

- configuration: element of  $Q \times \{y \sqcup^\omega \mid y \in \Gamma^*\} \times \mathbb{N}$
- start configuration on input  $x \in \Sigma^*$ :  $(s, \vdash x \sqcup^\omega, 0)$
- **next configuration relation** is binary relation  $\xrightarrow[M]{1}$  defined as:

$$(p, z, n) \xrightarrow[M]{1} \begin{cases} (q, z', n-1) & \text{if } \delta(p, z_n) = (q, b, L) \\ (q, z', n+1) & \text{if } \delta(p, z_n) = (q, b, R) \end{cases}$$

with

- $z_n$ :  $n$ -th symbol of  $z$

## Definitions

- configuration: element of  $Q \times \{y \sqcup^\omega \mid y \in \Gamma^*\} \times \mathbb{N}$
- start configuration on input  $x \in \Sigma^*$ :  $(s, \vdash x \sqcup^\omega, 0)$
- **next configuration relation** is binary relation  $\xrightarrow[M]{1}$  defined as:

$$(p, z, n) \xrightarrow[M]{1} \begin{cases} (q, z', n-1) & \text{if } \delta(p, z_n) = (q, b, L) \\ (q, z', n+1) & \text{if } \delta(p, z_n) = (q, b, R) \end{cases}$$

with

- $z_n$ :  $n$ -th symbol of  $z$
- $z'$ : string obtained from  $z$  by substituting  $b$  for  $z_n$  (at position  $n$ )

## Definitions

- configuration: element of  $Q \times \{y \sqcup^\omega \mid y \in \Gamma^*\} \times \mathbb{N}$
- start configuration on input  $x \in \Sigma^*$ :  $(s, \vdash x \sqcup^\omega, 0)$
- next configuration relation is binary relation  $\xrightarrow[M]{1}$  defined as:

$$(p, z, n) \xrightarrow[M]{1} \begin{cases} (q, z', n-1) & \text{if } \delta(p, z_n) = (q, b, L) \\ (q, z', n+1) & \text{if } \delta(p, z_n) = (q, b, R) \end{cases}$$

with

- $z_n$ :  $n$ -th symbol of  $z$
- $z'$ : string obtained from  $z$  by substituting  $b$  for  $z_n$  (at position  $n$ )
- $\xrightarrow[M]{n} = (\xrightarrow[M]{1})^n \quad \forall n \geq 0 \quad \xrightarrow[M]{*} = \bigcup_{n \geq 0} \xrightarrow[M]{n}$

## Definitions

- configuration: element of  $Q \times \{y \sqcup^\omega \mid y \in \Gamma^*\} \times \mathbb{N}$
- start configuration on input  $x \in \Sigma^*$ :  $(s, \vdash x \sqcup^\omega, 0)$
- next configuration relation is binary relation  $\xrightarrow[M]{1}$  defined as:

$$(p, z, n) \xrightarrow[M]{1} \begin{cases} (q, z', n-1) & \text{if } \delta(p, z_n) = (q, b, L) \\ (q, z', n+1) & \text{if } \delta(p, z_n) = (q, b, R) \end{cases}$$

with

- $z_n$ :  $n$ -th symbol of  $z$
- $z'$ : string obtained from  $z$  by substituting  $b$  for  $z_n$  (at position  $n$ )
- $\xrightarrow[M]{n} = (\xrightarrow[M]{1})^n \quad \forall n \geq 0$        $\xrightarrow[M]{*} = \bigcup_{n \geq 0} \xrightarrow[M]{n}$

## Definitions

- $x \in \Sigma^*$  is **accepted** by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (t, y, n)$

## Definitions

- $x \in \Sigma^*$  is accepted by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (t, y, n)$
- $L(M)$  is set of strings accepted by  $M$

## Definitions

- $x \in \Sigma^*$  is accepted by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (t, y, n)$
- $L(M)$  is set of strings accepted by  $M$
- $x \in \Sigma^*$  is **rejected** by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (r, y, n)$

## Definitions

- $x \in \Sigma^*$  is accepted by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (t, y, n)$
- $L(M)$  is set of strings accepted by  $M$
- $x \in \Sigma^*$  is rejected by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (r, y, n)$
- $M$  **halts** on  $x$  if it accepts or rejects  $x$ , otherwise  $M$  **loops** on  $x$



## Definitions

- $x \in \Sigma^*$  is accepted by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (t, y, n)$
- $L(M)$  is set of strings accepted by  $M$
- $x \in \Sigma^*$  is rejected by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (r, y, n)$
- $M$  halts on  $x$  if it accepts or rejects  $x$ , otherwise  $M$  loops on  $x$
- $M$  is **total** if it halts on all inputs

## Definitions

- $x \in \Sigma^*$  is accepted by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (t, y, n)$
- $L(M)$  is set of strings accepted by  $M$
- $x \in \Sigma^*$  is rejected by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (r, y, n)$
- $M$  halts on  $x$  if it accepts or rejects  $x$ , otherwise  $M$  loops on  $x$
- $M$  is total if it halts on all inputs
- set  $A$  is
  - **recursively enumerable (r.e.)** if  $A = L(M)$  for some TM  $M$

## Definitions

- $x \in \Sigma^*$  is accepted by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (t, y, n)$
- $L(M)$  is set of strings accepted by  $M$
- $x \in \Sigma^*$  is rejected by  $M$  if  $\exists y \exists n$  such that  $(s, \vdash x \sqcup^\omega, 0) \xrightarrow[M]{*} (r, y, n)$
- $M$  halts on  $x$  if it accepts or rejects  $x$ , otherwise  $M$  loops on  $x$
- $M$  is total if it halts on all inputs
- set  $A$  is
  - recursively enumerable (r.e.) if  $A = L(M)$  for some TM  $M$
  - **recursive** if  $A = L(M)$  for some total TM  $M$

## Theorem

- every recursive set is r.e.

## Theorem

- every recursive set is r.e.
- not every r.e. set is recursive

## Theorem


- every recursive set is r.e.
- not every r.e. set is recursive
- every context-free set is recursive

## Theorem

- every recursive set is r.e.
- not every r.e. set is recursive
- every context-free set is recursive
- not every recursive set is context-free

## Theorem

- every recursive set is r.e.
- not every r.e. set is recursive
- every context-free set is recursive
- not every recursive set is context-free (A)



$$A = \{a^n b^n c^n \mid n \geq 0\}$$



## Theorem

if  $A$  and  $\sim A$  are r.e. then  $A$  is recursive

## Theorem

if  $A$  and  $\sim A$  are r.e. then  $A$  is recursive

## Proof.

- $A = L(M_1)$  for TM  $M_1 = TM(Q_1, \Sigma, \Gamma_1, \vdash, \sqcup, \delta_1, s_1, t_1, r_1)$
- $\sim A = L(M_2)$  for TM  $M_2 = TM(Q_2, \Sigma, \Gamma_2, \vdash, \sqcup, \delta_2, s_2, t_2, r_2)$

## Theorem

if  $A$  and  $\sim A$  are r.e. then  $A$  is recursive

## Proof.


- $A = L(M_1)$  for TM  $M_1 = TM(Q_1, \Sigma, \Gamma_1, \vdash, \sqcup, \delta_1, s_1, t_1, r_1)$
- $\sim A = L(M_2)$  for TM  $M_2 = TM(Q_2, \Sigma, \Gamma_2, \vdash, \sqcup, \delta_2, s_2, t_2, r_2)$
- define TM  $M$  that simulates  $M_1$  and  $M_2$  on separate tracks of its single tape

## Theorem

if  $A$  and  $\sim A$  are r.e. then  $A$  is recursive

## Proof.

- $A = L(M_1)$  for TM  $M_1 = TM(Q_1, \Sigma, \Gamma_1, \vdash, \sqcup, \delta_1, s_1, t_1, r_1)$
- $\sim A = L(M_2)$  for TM  $M_2 = TM(Q_2, \Sigma, \Gamma_2, \vdash, \sqcup, \delta_2, s_2, t_2, r_2)$
- define TM  $M$  that simulates  $M_1$  and  $M_2$  on separate tracks of its single tape





## Theorem

if  $A$  and  $\sim A$  are r.e. then  $A$  is recursive

## Proof.

- $A = L(M_1)$  for TM  $M_1 = TM(Q_1, \Sigma, \Gamma_1, \vdash, \sqcup, \delta_1, s_1, t_1, r_1)$
- $\sim A = L(M_2)$  for TM  $M_2 = TM(Q_2, \Sigma, \Gamma_2, \vdash, \sqcup, \delta_2, s_2, t_2, r_2)$
- define TM  $M$  that simulates  $M_1$  and  $M_2$  on separate tracks of its single tape



## Theorem

recursive sets are effectively closed under complement

## Theorem

recursive sets are effectively closed under complement

## Theorem

extending TMs with

- nondeterminism

does not change the class of recursive (enumerable) languages

## Theorem

recursive sets are effectively closed under complement

## Theorem

extending TMs with

- nondeterminism
- two-way infinite tape

does not change the class of recursive (enumerable) languages



## Theorem

recursive sets are effectively closed under complement

## Theorem

extending TMs with

- nondeterminism
- two-way infinite tape
- multiple tapes

does not change the class of recursive (enumerable) languages

## Theorem

recursive sets are effectively closed under complement

## Theorem

extending TMs with

- nondeterminism
- two-way infinite tape
- multiple tapes
- multidimensional tape

does not change the class of recursive (enumerable) languages

## Theorem

recursive sets are effectively closed under complement

## Theorem

extending TMs with

- nondeterminism
- two-way infinite tape
- multiple tapes
- multidimensional tape
- multiple heads

does not change the class of recursive (enumerable) languages

# Outline

- 1 A Quick Recap
- 2 Turing Machines
- 3 Decision Problems**
- 4 Encoding
- 5 Diagonalization

## Decision Problems

- halting problem for TMs

instance: TM  $M$ , string  $x$

question: does  $M$  halt on  $x$ ?

## Decision Problems

- halting problem for TMs
  - instance: TM  $M$ , string  $x$
  - question: does  $M$  halt on  $x$ ?
- uniform halting problem for TMs
  - instance: TM  $M$
  - question: does  $M$  halt on all inputs?

## Decision Problems

- halting problem for TMs

instance: TM  $M$ , string  $x$   
question: does  $M$  halt on  $x$ ?

- uniform halting problem for TMs

instance: TM  $M$   
question: does  $M$  halt on all inputs?

- membership problem for CFGs

instance: CFG  $G$ , string  $x$   
question:  $x \in L(G)$ ?

## Decision Problems

- halting problem for TMs

instance: TM  $M$ , string  $x$

question: does  $M$  halt on  $x$ ?

- uniform halting problem for TMs

instance: TM  $M$

question: does  $M$  halt on all inputs?

- membership problem for CFGs

instance: CFG  $G$ , string  $x$

question:  $x \in L(G)$ ?

- equivalence problem for regular expressions

instance: regular expressions  $\alpha, \beta$

question:  $L(\alpha) = L(\beta)$ ?



## Decision Problems

- halting problem for TMs
  - instance: TM  $M$ , string  $x$
  - question: does  $M$  halt on  $x$ ?
- uniform halting problem for TMs
  - instance: TM  $M$
  - question: does  $M$  halt on all inputs?
- membership problem for CFGs
  - instance: CFG  $G$ , string  $x$
  - question:  $x \in L(G)$ ?
- equivalence problem for regular expressions
  - instance: regular expressions  $\alpha, \beta$
  - question:  $L(\alpha) = L(\beta)$ ?

## Decision Problems as Membership Problems

- code instance of problem as string over some alphabet

## Decision Problems

- |  |   |
|--|---|
| <ul style="list-style-type: none"><li>• halting problem for TMs<br/>instance: TM <math>M</math>, string <math>x</math><br/>question: does <math>M</math> halt on <math>x</math>?</li></ul> | <ul style="list-style-type: none"><li>• membership problem for CFGs<br/>instance: CFG <math>G</math>, string <math>x</math><br/>question: <math>x \in L(G)</math>?</li></ul>                                |
| <ul style="list-style-type: none"><li>• uniform halting problem for TMs<br/>instance: TM <math>M</math><br/>question: does <math>M</math> halt on all inputs?</li></ul>                    | <ul style="list-style-type: none"><li>• equivalence problem for regular expressions<br/>instance: regular expressions <math>\alpha, \beta</math><br/>question: <math>L(\alpha) = L(\beta)</math>?</li></ul> |


## Decision Problems as Membership Problems

- code instance of problem as string over some alphabet
- language is set of all strings that correspond to yes instances

## Theorem

for every set  $A$  exactly one of following alternatives holds


- 1  $A$  and  $\sim A$  are recursive



## Theorem

for every set  $A$  exactly one of following alternatives holds


- 1  $A$  and  $\sim A$  are recursive
- 2  $A$  and  $\sim A$  are not r.e.



## Theorem

for every set  $A$  exactly one of following alternatives holds


- ①  $A$  and  $\sim A$  are recursive
- ②  $A$  and  $\sim A$  are not r.e.
- ③ one of  $A$  and  $\sim A$  is r.e. and other is not r.e.



## Theorem

for every set  $A$  exactly one of following alternatives holds


- ①  $A$  and  $\sim A$  are recursive
- ②  $A$  and  $\sim A$  are not r.e.
- ③ one of  $A$  and  $\sim A$  is r.e. and other is not r.e.



## Theorem

for every set  $A$  exactly one of following alternatives holds


- ①  $A$  and  $\sim A$  are recursive
- ②  $A$  and  $\sim A$  are not r.e.
- ③ one of  $A$  and  $\sim A$  is r.e. and other is not r.e.



## Theorem

for every set  $A$  exactly one of following alternatives holds

- ①  $A$  and  $\sim A$  are recursive
- ②  $A$  and  $\sim A$  are not r.e.
- ③ one of  $A$  and  $\sim A$  is r.e. and other is not r.e.






## Theorem

for every set  $A$  exactly one of following alternatives holds


- ①  $A$  and  $\sim A$  are recursive
- ②  $A$  and  $\sim A$  are not r.e.
- ③ one of  $A$  and  $\sim A$  is r.e. and other is not r.e.



## Theorem

for every set  $A$  exactly one of following alternatives holds

- ①  $A$  and  $\sim A$  are recursive
- ②  $A$  and  $\sim A$  are not r.e.
- ③ one of  $A$  and  $\sim A$  is r.e. and other is not r.e.



$A \setminus \sim A$	A	B	C
A	①	x	x
B	x	x	③
C	x	③	②

## Definitions

decision problem  $P$  is

- **decidable** if there is algorithm ( TM that always halts ) for  $P$ :
  - 1 **accepts** for every yes instance
  - 2 **rejects** for every no instance

## Definitions

decision problem  $P$  is

- decidable if there is algorithm ( TM that always halts ) for  $P$ :
  - ① accepts for every yes instance
  - ② rejects for every no instance
- **semi-decidable** if there is procedure ( TM ) for  $P$ :
  - ① **accepts** for every yes instance
  - ② **rejects** or **loops** for every no instance

## Definitions

decision problem  $P$  is

- decidable if there is algorithm ( TM that always halts ) for  $P$ :
  - ① accepts for every yes instance
  - ② rejects for every no instance
- semi-decidable if there is procedure ( TM ) for  $P$ :
  - ① accepts for every yes instance
  - ② rejects or loops for every no instance

## Remark

decision problem  $P$  is ( semi- )decidable  $\iff$  encoding of  $P$  is recursive(ly enumerable )

## Definitions

decision problem  $P$  is

- decidable if there is algorithm ( TM that always halts ) for  $P$ :
  - ① accepts for every yes instance
  - ② rejects for every no instance
- semi-decidable if there is procedure ( TM ) for  $P$ :
  - ① accepts for every yes instance
  - ② rejects or loops for every no instance

## Remark

decision problem  $P$  is ( semi- )decidable  $\iff$  encoding of  $P$  is recursive(ly enumerable )

# Outline

- 1 A Quick Recap
- 2 Turing Machines
- 3 Decision Problems
- 4 Encoding**
- 5 Diagonalization

## Theorem

- $\exists$  computable mappings  
 $N \mapsto \text{enc}(N)$  from TMs to bit strings



## Theorem

- $\exists$  computable mappings

$N \mapsto \text{enc}(N)$  from TMs to bit strings

$x \mapsto \text{dec}(x)$  from bit strings to TMs

## Theorem

- $\exists$  computable mappings

$N \mapsto \text{enc}(N)$  from TMs to bit strings

$x \mapsto \text{dec}(x)$  from bit strings to TMs

such that

$$\text{dec}(\text{enc}(N)) = N$$

$$\forall \text{ TM } N$$

## Theorem

- $\exists$  computable mappings

$N \mapsto \text{enc}(N)$  from TMs to bit strings

$x \mapsto \text{dec}(x)$  from bit strings to TMs

such that

$$\text{dec}(\text{enc}(N)) = N$$

$$\forall \text{ TM } N$$

$$\text{enc}(\text{dec}(x)) = \begin{cases} x \end{cases}$$

if  $x = \text{enc}(N)$  for some TM  $N$

$$\forall x \in \{0, 1\}^*$$

## Theorem

- $\exists$  computable mappings

$N \mapsto \text{enc}(N)$  from TMs to bit strings

$x \mapsto \text{dec}(x)$  from bit strings to TMs

such that

$$\text{dec}(\text{enc}(N)) = N$$

$$\forall \text{ TM } N$$

$$\text{enc}(\text{dec}(x)) = \begin{cases} x & \text{if } x = \text{enc}(N) \text{ for some TM } N \\ \text{enc}(N_{\text{trivial}}) & \text{otherwise} \end{cases}$$

$$\forall x \in \{0, 1\}^*$$

where  $N_{\text{trivial}}$  is some fixed TM with  $L(N_{\text{trivial}}) = \emptyset$

## Example (Encoding)

- TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$

## Example (Encoding)

- TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  without loss of generality
  - $Q = \{0, 1, \dots, n-1\}$

$$\text{enc}(M) = 0^n$$

## Example (Encoding)

- TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  without loss of generality
  - $Q = \{0, 1, \dots, n-1\}$
  - $\Gamma = \{0, 1, \dots, m-1\}$

$$\text{enc}(M) = 0^n 10^m$$

## Example (Encoding)

- TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  without loss of generality
  - $Q = \{0, 1, \dots, n-1\}$
  - $\Gamma = \{0, 1, \dots, m-1\}$
  - $\Sigma = \{0, 1, \dots, k-1\}$

$$\text{enc}(M) = 0^n 10^m 10^k$$



## Example (Encoding)

- TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  without loss of generality
  - $Q = \{0, 1, \dots, n-1\}$
  - $\Gamma = \{0, 1, \dots, m-1\}$
  - $\Sigma = \{0, 1, \dots, k-1\}$

$$\text{enc}(M) = 0^n 10^m 10^k 10^s 10^t 10^r$$

## Example (Encoding)

- TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  without loss of generality
  - $Q = \{0, 1, \dots, n-1\}$
  - $\Gamma = \{0, 1, \dots, m-1\}$
  - $\Sigma = \{0, 1, \dots, k-1\}$
  - $\vdash = u$

$$\text{enc}(M) = 0^n 10^m 10^k 10^s 10^t 10^r 10^u$$

## Example (Encoding)

- TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  without loss of generality
  - $Q = \{0, 1, \dots, n-1\}$
  - $\Gamma = \{0, 1, \dots, m-1\}$
  - $\Sigma = \{0, 1, \dots, k-1\}$
  - $\vdash = u$
  - $\sqcup = v$

$$\text{enc}(M) = 0^n 10^m 10^k 10^s 10^t 10^r 10^u 10^v 1$$

## Example (Encoding)

- TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  without loss of generality

- $Q = \{0, 1, \dots, n-1\}$
- $\Gamma = \{0, 1, \dots, m-1\}$
- $\Sigma = \{0, 1, \dots, k-1\}$
- $\vdash = u$
- $\sqcup = v$

$$\text{enc}(M) = 0^n 10^m 10^k 10^s 10^t 10^r 10^u 10^v 1 \dots 0^p 10^a 10^q 10^b 10^c 1 \dots$$

$$\text{for all } ((p, a), (q, b, d)) \in \delta \text{ with } c = \begin{cases} 0 & \text{if } d = L \\ 1 & \text{if } d = R \end{cases}$$

## Example (Encoding)

- TM  $M = (Q, \Sigma, \Gamma, \vdash, \sqcup, \delta, s, t, r)$  without loss of generality

- $Q = \{0, 1, \dots, n-1\}$
- $\Gamma = \{0, 1, \dots, m-1\}$
- $\Sigma = \{0, 1, \dots, k-1\}$
- $\vdash = u$
- $\sqcup = v$

$$\text{enc}(M) = 0^n 10^m 10^k 10^s 10^t 10^r 10^u 10^v 1 \dots 0^p 10^a 10^q 10^b 10^c 1 \dots \quad (\text{order is not important})$$

$$\text{for all } ((p, a), (q, b, d)) \in \delta \text{ with } c = \begin{cases} 0 & \text{if } d = L \\ 1 & \text{if } d = R \end{cases}$$

- string  $x = a_1 a_2 \dots a_n \in \Sigma^*$   
 $\text{enc}(x) = 0^{a_1} 10^{a_2} 1 \dots 0^{a_n} 1$

## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, 3, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	$(0, 0, R)$	$(0, 1, R)$	$(0, 2, R)$	$(0, 3, R)$
1	$(1, 0, R)$	$(1, 1, R)$	$(1, 2, R)$	$(1, 3, R)$

## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, 3, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	$(0, 0, R)$	$(0, 1, R)$	$(0, 2, R)$	$(0, 3, R)$
1	$(1, 0, R)$	$(1, 1, R)$	$(1, 2, R)$	$(1, 3, R)$

- $\text{enc}(N_{\text{trivial}}) = 001$

## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, 3, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	$(0, 0, R)$	$(0, 1, R)$	$(0, 2, R)$	$(0, 3, R)$
1	$(1, 0, R)$	$(1, 1, R)$	$(1, 2, R)$	$(1, 3, R)$

- $\text{enc}(N_{\text{trivial}}) = 00100001$



## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, 3, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	$(0, 0, R)$	$(0, 1, R)$	$(0, 2, R)$	$(0, 3, R)$
1	$(1, 0, R)$	$(1, 1, R)$	$(1, 2, R)$	$(1, 3, R)$

- $\text{enc}(N_{\text{trivial}}) = 00100001001$

## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, 3, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	$(0, 0, R)$	$(0, 1, R)$	$(0, 2, R)$	$(0, 3, R)$
1	$(1, 0, R)$	$(1, 1, R)$	$(1, 2, R)$	$(1, 3, R)$

- $\text{enc}(N_{\text{trivial}}) = 001000010011011$

## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, 3, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	$(0, 0, R)$	$(0, 1, R)$	$(0, 2, R)$	$(0, 3, R)$
1	$(1, 0, R)$	$(1, 1, R)$	$(1, 2, R)$	$(1, 3, R)$

- $\text{enc}(N_{\text{trivial}}) = 001000010011011\mathbf{001}$

## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, \textcolor{red}{3}, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	$(0, 0, R)$	$(0, 1, R)$	$(0, 2, R)$	$(0, 3, R)$
1	$(1, 0, R)$	$(1, 1, R)$	$(1, 2, R)$	$(1, 3, R)$

- $\text{enc}(N_{\text{trivial}}) = 001000010011011001\textcolor{red}{0001}$

## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, 3, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	(0, 0, R)	(0, 1, R)	(0, 2, R)	(0, 3, R)
1	(1, 0, R)	(1, 1, R)	(1, 2, R)	(1, 3, R)

- $\text{enc}(N_{\text{trivial}}) = 0010000100110110010001$   
111101

## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, 3, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	(0, 0, R)	(0, 1, R)	(0, 2, R)	(0, 3, R)
1	(1, 0, R)	(1, 1, R)	(1, 2, R)	(1, 3, R)

- $\text{enc}(N_{\text{trivial}}) = 0010000100110110010001$   
111101101101011001100101100011000101

## Example

- $N_{\text{trivial}} = (\{0, 1\}, \{0, 1\}, \{0, 1, 2, 3\}, 2, 3, \delta, 0, 1, 0)$  with

$\delta$	0	1	2	3
0	$(0, 0, R)$	$(0, 1, R)$	$(0, 2, R)$	$(0, 3, R)$
1	$(1, 0, R)$	$(1, 1, R)$	$(1, 2, R)$	$(1, 3, R)$

- $\text{enc}(N_{\text{trivial}}) = 0010000100110110010001$   
111101101101011001100101100011000101  
011011010101010101001010010101000101000101

## Theorem

- $\exists$  computable mappings

$N \mapsto \text{enc}(N)$  from TMs to bit strings

$x \mapsto \text{dec}(x)$  from bit strings to TMs

such that

$$\text{dec}(\text{enc}(N)) = N$$

$$\forall \text{ TM } N$$

$$\text{enc}(\text{dec}(x)) = \begin{cases} x & \text{if } x = \text{enc}(N) \text{ for some TM } N \\ \text{enc}(N_{\text{trivial}}) & \text{otherwise} \end{cases}$$

$$\forall x \in \{0, 1\}^*$$

where  $N_{\text{trivial}}$  is some fixed TM with  $L(N_{\text{trivial}}) = \emptyset$

- $\exists$  **computable enumeration** of all TMs with input alphabet  $\{0, 1\}$

$M_{\epsilon}, M_0, M_1, M_{00}, M_{01}, M_{10}, \dots$



## Definition (Encoding of Membership Problem for Turing Machines)

$$\text{MP} = \{\text{enc}(M) \# \text{enc}(x) \mid x \in L(M)\}$$

## Definition (Encoding of Membership Problem for Turing Machines)

$$MP = \{ M \# x \mid x \in L(M) \}$$

### Definition (Encoding of Membership Problem for Turing Machines)

$MP = \{ M \# x \mid x \in L(M) \}$

### Theorem ( Universal Turing Machine )

$\exists$  TM  $U$  such that  $L(U) = MP$

### Definition (Encoding of Membership Problem for Turing Machines)

$MP = \{ M \# x \mid x \in L(M) \}$

### Theorem ( Universal Turing Machine )

$\exists$  TM  $U$  such that  $L(U) = MP$

universal TM can simulate any TM

### Definition (Encoding of Membership Problem for Turing Machines)

$MP = \{ M \# x \mid x \in L(M) \}$

### Theorem ( Universal Turing Machine )

$\exists$  TM  $U$  such that  $L(U) = MP$

universal TM can simulate any TM

### Corollary

MP is r.e.

## How to Prove Undecidability?

- diagonalization
- reduction

## How to Prove Undecidability?

- diagonalization
- reduction

# Outline

- 1 A Quick Recap
- 2 Turing Machines
- 3 Decision Problems
- 4 Encoding
- 5 Diagonalization**



## Definitions

- set  $A$  is **countably infinite** if there exists bijection  $f: \mathbb{N} \rightarrow A$

## Definitions

- set  $A$  is countably infinite if there exists bijection  $f: \mathbb{N} \rightarrow A$
- set is **countable** if it is finite or countably infinite

## Definitions

- set  $A$  is countably infinite if there exists bijection  $f: \mathbb{N} \rightarrow A$
- set is countable if it is finite or countably infinite

## Example

- $\mathbb{N} \quad f(n) = n$

## Definitions

- set  $A$  is countably infinite if there exists bijection  $f: \mathbb{N} \rightarrow A$
- set is countable if it is finite or countably infinite

## Example

- $\mathbb{N} \quad f(n) = n$
- $\mathbb{Z} \quad f(n) = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ -\frac{n+1}{2} & \text{if } n \text{ is odd} \end{cases}$

## Definitions

- set  $A$  is countably infinite if there exists bijection  $f: \mathbb{N} \rightarrow A$
- set is countable if it is finite or countably infinite

## Example

- $\mathbb{N} \quad f(n) = n$
- $\mathbb{Z} \quad f(n) = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ -\frac{n+1}{2} & \text{if } n \text{ is odd} \end{cases}$
- $\mathbb{Q} \quad f(n) = \dots$

## Definitions

- set  $A$  is countably infinite if there exists bijection  $f: \mathbb{N} \rightarrow A$
- set is countable if it is finite or countably infinite

## Example

- $\mathbb{N} \quad f(n) = n$
- $\mathbb{Z} \quad f(n) = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ -\frac{n+1}{2} & \text{if } n \text{ is odd} \end{cases}$
- $\mathbb{Q} \quad f(n) = \dots$

## Theorem (Schröder–Bernstein)

$\exists$  bijection  $A \rightarrow B \iff \exists$  injective functions  $f: A \rightarrow B$  and  $g: B \rightarrow A$

## Definitions

- set  $A$  is countably infinite if there exists bijection  $f: \mathbb{N} \rightarrow A$
- set is countable if it is finite or countably infinite

## Example

- $\mathbb{N} \quad f(n) = n$
- $\mathbb{Z} \quad f(n) = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ -\frac{n+1}{2} & \text{if } n \text{ is odd} \end{cases}$
- $\mathbb{Q} \quad f(n) = n \quad g(0) = 0 \quad g(\frac{p}{q}) = 2^p 3^q \quad g(-\frac{p}{q}) = 2^p 3^q 5$

## Theorem (Schröder-Bernstein)

$\exists$  bijection  $A \rightarrow B \iff \exists$  injective functions  $f: A \rightarrow B$  and  $g: B \rightarrow A$

## Definitions

- set  $A$  is countably infinite if there exists bijection  $f: \mathbb{N} \rightarrow A$
- set is countable if it is finite or countably infinite

## Example

- $\mathbb{N} \quad f(n) = n$
- $\mathbb{Z} \quad f(n) = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ -\frac{n+1}{2} & \text{if } n \text{ is odd} \end{cases}$
- $\mathbb{Q} \quad f(n) = n \quad g(0) = 0 \quad g(\frac{p}{q}) = 2^p 3^q \quad g(-\frac{p}{q}) = 2^p 3^q 5$
- $\Sigma^*$  for finite alphabet  $\Sigma$  is countable

## Theorem (Schröder–Bernstein)

$\exists$  bijection  $A \rightarrow B \iff \exists$  injective functions  $f: A \rightarrow B$  and  $g: B \rightarrow A$



## Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Proof. (Diagonalization)

- suppose  $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$  is countable

	$A_0$	$A_1$	$A_2$	$\dots$
$x_0$				
$x_1$				
$x_2$				
$\vdots$				
$\vdots$				

## Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Proof. (Diagonalization)

- suppose  $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$  is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$  is countable

	$A_0$	$A_1$	$A_2$	$\dots$
$x_0$				
$x_1$				
$x_2$				
$\vdots$				
$\vdots$				

## Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Proof. (Diagonalization)

- suppose  $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$  is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$  is countable
- create infinite table

	$A_0$	$A_1$	$A_2$	$\dots$
$x_0$				
$x_1$				
$x_2$				
$\vdots$				
$\vdots$				

## Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Proof. (Diagonalization)

- suppose  $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$  is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$  is countable
- create infinite table

	$A_0$	$A_1$	$A_2$	$\dots$
$x_0$	×			
$x_1$	✓			
$x_2$	×			
$\vdots$				
$\vdots$				

- $A_0 = \{x_1\}$

## Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Proof. (Diagonalization)

- suppose  $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$  is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$  is countable
- create infinite table

	$A_0$	$A_1$	$A_2$	...
$x_0$	×	✓		
$x_1$	✓	×		
$x_2$	×	✓		
⋮				
⋮				

- $A_0 = \{x_1\}$      $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$

## Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Proof. (Diagonalization)

- suppose  $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$  is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$  is countable
- create infinite table

	$A_0$	$A_1$	$A_2$	$\dots$
$x_0$	x	✓	x	
$x_1$	✓	x	x	
$x_2$	x	✓	✓	
$\vdots$				

- $A_0 = \{x_1\}$     $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$     $A_2 = \{x_2, x_5\}$

## Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Proof. (Diagonalization)

- suppose  $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$  is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$  is countable
- create infinite table

	$A_0$	$A_1$	$A_2$	$\dots$
$x_0$	x	✓	x	
$x_1$	✓	x	x	
$x_2$	x	✓	✓	
$\vdots$				

- $A_0 = \{x_1\}$     $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$     $A_2 = \{x_2, x_5\}$
- define  $B = \{x_i \mid x_i \notin A_i\} \in 2^{\Sigma^*}$



## Theorem

$\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Proof. (Diagonalization)

- suppose  $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$  is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$  is countable
- create infinite table

	$A_0$	$A_1$	$A_2$	...
$x_0$	✓	✓	×	
$x_1$	✓	✓	×	
$x_2$	×	✓	×	
⋮				
⋮				

- $A_0 = \{x_1\}$     $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$     $A_2 = \{x_2, x_5\}$
- define  $B = \{x_i \mid x_i \notin A_i\} \in 2^{\Sigma^*}$
- $B \neq A_i \forall i$

## Theorem

 $\Sigma \neq \emptyset \implies 2^{\Sigma^*}$  is uncountable

## Proof. (Diagonalization)

- suppose  $2^{\Sigma^*} = \{A_0, A_1, A_2, \dots\}$  is countable
- $\Sigma^* = \{x_0, x_1, x_2, \dots\}$  is countable
- create infinite table

	$A_0$	$A_1$	$A_2$	...
$x_0$	✓	✓	×	
$x_1$	✓	✓	×	
$x_2$	×	✓	×	
⋮				
⋮				

- $A_0 = \{x_1\}$     $A_1 = \{x_0, x_2, x_7, x_8, \dots\}$     $A_2 = \{x_2, x_5\}$
- define  $B = \{x_i \mid x_i \notin A_i\} \in 2^{\Sigma^*}$
- $B \neq A_i \forall i$  ⚡

Thanks! & Questions?