

$$A = \begin{pmatrix} -1 & 0 \\ 1 & 2 \end{pmatrix}$$

$$B = \begin{pmatrix} 1 & 0 & 3 \\ 0 & 1 & 0 \end{pmatrix}$$

$$C = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} \quad D = \begin{pmatrix} 1 & 2 \end{pmatrix}$$

$B \quad C = \begin{pmatrix} 1 & 0 & 3 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix}$
 $\begin{matrix} 2 \times 3 & 3 \times 1 \\ \text{product } 2 \times 1 \end{matrix}$
 $= 1 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 1 \end{pmatrix} + 3 \begin{pmatrix} 3 \\ 0 \end{pmatrix} = \begin{pmatrix} 10 \\ 0 \end{pmatrix}$

$$A \quad B = \begin{pmatrix} -1 & 0 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 & 3 \\ 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} -1 \\ 0 \end{pmatrix} \perp \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$= \begin{pmatrix} 1 \begin{pmatrix} -1 \\ 1 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 2 \end{pmatrix} & 0 \begin{pmatrix} -1 \\ 1 \end{pmatrix} + 1 \begin{pmatrix} 0 \\ 2 \end{pmatrix} & 3 \begin{pmatrix} -1 \\ 1 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 2 \end{pmatrix} \end{pmatrix}$$

$$= \begin{pmatrix} -1 & 0 & -3 \\ 1 & 2 & 3 \end{pmatrix}$$

$$A = \begin{pmatrix} -1 & 0 \\ 1 & 2 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 0 & 3 \\ 0 & 1 & 0 \end{pmatrix} \quad C = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} \quad D = \begin{pmatrix} 1 & 2 \end{pmatrix}$$

Can $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ be written as a linear combination of $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 2 \end{pmatrix}$?

Does $\begin{pmatrix} -1 & 0 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ have a solution?

Does the system of linear eqns $\begin{cases} -x_1 + 0x_2 = 1 \\ x_1 + 2x_2 = 2 \end{cases}$ have a solution?

$$A = \begin{pmatrix} -1 & 0 \\ 1 & 2 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 0 & 3 \\ 0 & 1 & 0 \end{pmatrix} \quad C = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} \quad D = \begin{pmatrix} 1 & 2 \end{pmatrix}$$

$$DA = \begin{pmatrix} 1 & 2 \end{pmatrix} \begin{pmatrix} \boxed{-1} & \boxed{0} \\ \boxed{1} & \boxed{2} \end{pmatrix} = \begin{pmatrix} \boxed{1} & \boxed{4} \end{pmatrix}$$

$1 \times 2 \qquad \qquad 2 \times 2 \qquad \qquad 1 \times 2$

$$= 1 \begin{pmatrix} -1 & 0 \end{pmatrix} + 2 \begin{pmatrix} 1 & 2 \end{pmatrix}$$

$$= \begin{pmatrix} -1 & 0 \end{pmatrix} + \begin{pmatrix} 2 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 4 \end{pmatrix}.$$

Solve

$$-x_1 + 0x_2 = 1$$

$$x_1 + 2x_2 = 2$$

$$\begin{pmatrix} -1 & 0 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}.$$

$$F = \begin{pmatrix} 2 & -1 \\ 1 & 1 \\ 0 & 2 \end{pmatrix} \quad \underline{z} = \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}.$$

1. Can $\begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}$ be written as a linear combination of $\begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$?

$\hat{=}$ Can I choose numbers x_1 and x_2 so that

$$x_1 \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}.$$

$$\begin{pmatrix} x_1 & x_2 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ 1 & 1 \\ 0 & 2 \end{pmatrix}$$

$1 \times 2 \qquad \qquad 3 \times 2$

$$\begin{pmatrix} 2 & -1 \\ 1 & 1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \stackrel{?}{=} \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}.$$

$3 \times 2 \qquad 2 \times 1$

$$\begin{pmatrix} 2 & -1 \\ 1 & 1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}.$$

$$x_1 \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}.$$

$$\begin{pmatrix} 2x_1 - x_2 \\ x_1 + x_2 \\ 0 + 2x_2 \end{pmatrix} = \begin{pmatrix} 2x_1 \\ x_1 \\ 0 \end{pmatrix} + \begin{pmatrix} -x_2 \\ x_2 \\ 2x_2 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}.$$

$$\begin{cases} 2x_1 - x_2 = 3 \\ x_1 + x_2 = 0 \\ 0 + 2x_2 = -2 \end{cases}$$

$$x_1 = 1 \quad x_2 = -1.$$

$$\begin{pmatrix} 2 & -1 & 3 & 3 \\ 1 & 1 & 0 & 1 \\ 0 & 2 & -2 & -2 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -1 \\ 1 & 1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_1 \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}.$$

1. Is $\begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}$ a linear combination of $\begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$?

2. Is there a vector $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$ s.t. $\begin{pmatrix} 2 & -1 \\ 1 & 1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}$?

3. Solve the system

$$2x_1 - x_2 = 3$$

$$x_1 + x_2 = 0.$$

$$0x_1 + 2x_2 = -2$$

$$\begin{pmatrix} 2 & -1 & 3 & 1 & 3 \\ 1 & 1 & 0 & 2 & -1 \\ 0 & 2 & -2 & 2 & 2 \end{pmatrix}$$

$$\begin{pmatrix} \textcircled{2} & -1 & \boxed{3} & 1 & 3 \\ 1 & 1 & 0 & 2 & -1 \\ 0 & 2 & -2 & 2 & 2 \end{pmatrix} \xrightarrow{R_2 \leftarrow R_2 - \frac{1}{2} R_1} \begin{pmatrix} \textcircled{2} & -1 & 3 & 1 & 3 \\ 0 & \textcircled{3/2} & -3/2 & 3/2 & -5/2 \\ 0 & 2 & -2 & 2 & 2 \end{pmatrix}$$

pivots.

$$(1 \ 1 \ 0 \ 2 \ -1) - \frac{1}{2} (2 \ -1 \ 3 \ 1 \ 3) \\ = (1 \ 1 \ 0 \ 2 \ -1) + (-1 \ 1/2 \ -3/2 \ -1/2 \ -3/2)$$

$$\begin{pmatrix} \textcircled{2} & -1 & 3 & 1 & 3 \\ 0 & \textcircled{3/2} & -3/2 & 3/2 & -5/2 \\ 0 & \underline{2} & -2 & 2 & 2 \end{pmatrix} \xrightarrow{R_3 \leftarrow R_3 - \frac{4}{3} R_2} \begin{pmatrix} \textcircled{2} & -1 & 3 & 1 & 3 \\ 0 & \textcircled{3/2} & -3/2 & 3/2 & -5/2 \\ 0 & 0 & \boxed{0} \rightarrow \boxed{0} \rightarrow \textcircled{16/3} \end{pmatrix}$$

↓ normalize leading entries.

$$\begin{pmatrix} \textcircled{1} & -1/2 & 3/2 & 1/2 & 0 \\ 0 & \textcircled{1} & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & \textcircled{1} \end{pmatrix} \xleftarrow{\begin{matrix} R_2 \leftarrow R_2 + \frac{5}{3} R_3 \\ R_1 \leftarrow R_1 - \frac{2}{3} R_3 \end{matrix}} \begin{pmatrix} \textcircled{1} & -1/2 & 3/2 & 1/2 & 3/2 \\ 0 & \textcircled{1} & -1 & 1 & \boxed{-5/3} \\ 0 & 0 & 0 & 0 & \textcircled{1} \end{pmatrix}$$

$$\downarrow R_1 \leftarrow R_1 + \frac{1}{2} R_2$$

RREF: $\begin{pmatrix} \textcircled{1} & 0 & 1 & 1 & 0 \\ 0 & \textcircled{1} & -1 & 1 & 0 \\ 0 & 0 & 0 & \underline{0} & \textcircled{1} \end{pmatrix}$

Original: $\begin{pmatrix} \textcircled{2} & -1 & \boxed{3} & 1 & 3 \\ 1 & 1 & 0 & 2 & -1 \\ 0 & 2 & -2 & \underline{2} & \underline{2} \end{pmatrix}$

$$\begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} - \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ -2 \end{pmatrix}$$

$$\begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} + \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$$