

A .
(Square $n \times n$)

$$\underline{A} \underline{x} = \lambda \underline{x} \quad (\text{non-zero } \underline{x}, \text{ numbers } \lambda).$$

$$\underline{A} \underline{x} - \lambda \underline{I}_n \underline{x} = \underline{0}$$
$$(\underline{A} - \lambda \underline{I}_n) \underline{x} = \underline{0} \quad \checkmark$$

A has free cols iff $\det(A) = 0$.

If A has free cols; say $\underline{c}_j = \sum_{i < j} \alpha_i \underline{c}_i$

$$\det \begin{bmatrix} | & & | & | & & | \\ \underline{c}_1 & \dots & \underline{c}_{j-1} & \underline{c}_j & \dots & \underline{c}_n \\ | & & | & | & & | \end{bmatrix} = \det \begin{bmatrix} | & & | & & & | \\ \underline{c}_1 & \dots & \underline{c}_{j-1} & \alpha_1 \underline{c}_1 + \alpha_2 \underline{c}_2 + \dots + \alpha_{j-1} \underline{c}_{j-1} & & \underline{c}_n \\ | & & | & & & | \end{bmatrix}$$
$$= \sum_{i < j} \alpha_i \det \begin{bmatrix} | & & | & | & & | \\ \underline{c}_1 & \dots & \underline{c}_{j-1} & \underline{c}_i & \underline{c}_{j+1} & \dots & \underline{c}_n \\ | & & | & | & & | \end{bmatrix} = 0.$$

If $\det(A) = 0$ then it has free cols.

OR If A has no free cols, then $\det(A) \neq 0$.

$\det(A) = \pm$ product of pivots $\neq 0$.

depending on how many row exchanges in elimination.

Check $\det(A - \lambda I) = 0$.

$$A = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$$

$$A - \lambda I = \begin{pmatrix} 1-\lambda & 0 \\ -1 & 1-\lambda \end{pmatrix}$$

$$\lambda I = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}$$

$\det(A - \lambda I) = ?$ use linearity properties.

First row $(1-\lambda) \begin{pmatrix} 1 & 0 \end{pmatrix} + 0 \begin{pmatrix} 0 & 1 \end{pmatrix}$

$$\det(A - \lambda I) = \det \begin{pmatrix} 1-\lambda & [1 \ 0] \\ -1 & (1-\lambda) \end{pmatrix} + \det \begin{pmatrix} 0 & [0 \ 1] \\ -1 & 1-\lambda \end{pmatrix}$$

$$= (1-\lambda) \det \begin{pmatrix} 1 & 0 \\ -1 & 1-\lambda \end{pmatrix} + 0 \cancel{\det \begin{pmatrix} 0 & 1 \\ -1 & 1-\lambda \end{pmatrix}}$$

Second row: $-1 [1 \ 0] + (1-\lambda)[0 \ 1]$

$$(1-\lambda) \det \begin{bmatrix} 1 & 0 \\ -1 [1 \ 0] + (1-\lambda)[0 \ 1] \end{bmatrix} = (1-\lambda) \det \begin{bmatrix} 1 & 0 \\ -1 [1 \ 0] \end{bmatrix} + (1-\lambda) \det \begin{bmatrix} 1 & 0 \\ (1-\lambda)[0 \ 1] \end{bmatrix}$$

$$= (1-\lambda)(-1) \det \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} + (1-\lambda)^2 \det \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \underline{(1-\lambda)^2}$$

$$A = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \quad \det(A - \lambda I) = \underline{(1-\lambda)^2}$$

roots of $\det(A - \lambda I) = 0$ are 1, 1.

$$A - 1I = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} - 1 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ -1 & 0 \end{bmatrix}$$

$$\text{null}(A - I) = \left\{ \alpha \begin{pmatrix} 0 \\ 1 \end{pmatrix} : \alpha \in \mathbb{R} \right\}$$

$$\text{rref}(A - I) = \text{rref} \begin{bmatrix} 0 & 0 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \begin{array}{l} \underline{C2} = \underline{0C1} \\ \underline{C2} - \underline{0C1} = \underline{0} \end{array}$$

characteristic polynomial.

Roots of $\det(A - \lambda I) = 0$: Eigenvalues.

If a root is repeated k times in $\det(A - \lambda I) = 0$, we say that eigenvalue has algebraic multiplicity k .

Square Symmetric real A :

$$A^T = A \quad \text{eg: } A = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}.$$

$$A - \lambda I = \begin{pmatrix} 1-\lambda & -1 \\ -1 & 1-\lambda \end{pmatrix}.$$

$$\det(A - \lambda I) = (1-\lambda)^2 - 1 = \lambda^2 - 2\lambda + 1 - 1 = \lambda^2 - 2\lambda.$$

Roots: 0, 2.

$$\text{null}(A - 0I) = \text{null}(A) = \left\{ \alpha \begin{pmatrix} 1 \\ 1 \end{pmatrix} : \alpha \in \mathbb{R} \right\}.$$

$$\text{rref}(A) = \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}, \quad \begin{array}{l} \underline{c2} = -\underline{c1} \\ \underline{c2} + \underline{c1} = 0. \end{array}$$

$$\text{null}(A - 2I) = \text{null}\left[\begin{pmatrix} -1 & -1 \\ -1 & -1 \end{pmatrix}\right] = \left\{ \alpha \begin{pmatrix} 1 \\ -1 \end{pmatrix} : \alpha \in \mathbb{R} \right\}.$$

Dot product of $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ -1 \end{pmatrix} = 0$.

The eigenvectors (one from each of $\text{null}(A)$ and $\text{null}(A - 2I)$)
 $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ form a max linearly indep set in $\mathbb{R}^2$.

$$\text{For all } \underline{x} \in \mathbb{R}^2, \underline{x} = \alpha \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \beta \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

$$A \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$A \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

$$A \underline{x} = A \left[\alpha \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \beta \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right].$$

$$= A \left[\alpha \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right] + A \left[\beta \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right].$$

$$= \alpha A \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \beta A \begin{pmatrix} 1 \\ -1 \end{pmatrix} =$$

$$= \alpha \cdot 0 \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix} + 2\beta \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

$$A(A \underline{x}) = 4\beta \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

$$B = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} + \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix}$$

$$= \begin{pmatrix} 3/2 & 1/2 \\ 1/2 & 3/2 \end{pmatrix}.$$

$$\lambda I = \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}$$

Eigenvalues of B.

$$\det(B - \lambda I) = \det \begin{pmatrix} 3/2 - \lambda & 1/2 \\ 1/2 & 3/2 - \lambda \end{pmatrix}.$$

$$\begin{bmatrix} \frac{3}{2} - \lambda & 1/2 \\ 1/2 & 3/2 - \lambda \end{bmatrix} \xrightarrow{r_2 \leftarrow r_2 - \frac{1}{2} r_1} \begin{bmatrix} 3/2 - \lambda & 1/2 \\ 0 & \left(\frac{3}{2} - \lambda\right) - \frac{1}{2} \cdot \frac{1}{2} \end{bmatrix} \quad \lambda \neq 3/2.$$

$$\det(B) = \left(\frac{3}{2} - \lambda\right) \left(\frac{3}{2} - \lambda\right) - 1/4 = \lambda^2 - 3\lambda + 9/4 - 1/4 = \lambda^2 - 3\lambda + 2.$$

$$\text{Roots: } 2, 1$$

$$= (\lambda - 2)(\lambda - 1)$$

$$\lambda = 2 \quad B - \lambda I = \begin{pmatrix} \frac{3}{2} - \lambda & 1/2 \\ 1/2 & 3/2 - \lambda \end{pmatrix} = \begin{pmatrix} -1/2 & 1/2 \\ 1/2 & -1/2 \end{pmatrix} \quad \begin{array}{l} \underline{c2} = -\underline{c1}. \\ \underline{c1} + \underline{c2} = 0. \end{array}$$

$\therefore$ Eigenvector for eigenvalue $\lambda = 2$ $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$.

$$\lambda = 1 \quad B - \lambda I = \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix} \quad \underline{c2} = \underline{c1}.$$

Eigenvector for eigenvalue $\lambda = 1$ $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$.

$$\underline{x} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

$$B \underline{x} = \frac{1}{2} B \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \frac{1}{2} B \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \frac{1}{2} \cdot 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \frac{1}{2} \cdot 1 \cdot \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

$$\begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{ is in null } (B - 2I) \quad (B - 2I) \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 0. \quad B \begin{pmatrix} 1 \\ 1 \end{pmatrix} - 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 0.$$

$$B \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$B \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 1 \cdot \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

$$B(Bx) = B\left[\frac{1}{2} \cdot 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \frac{1}{2} \cdot 1 \begin{pmatrix} 1 \\ -1 \end{pmatrix}\right] = \frac{1}{2} \cdot 2^2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \frac{1}{2} \cdot 1^2 \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

$$B^n(x) = \frac{1}{2} \cdot 2^n \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \frac{1}{2} \cdot (1)^n \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

Spectral Decomposition Theorem.

Suppose A is square, symmetric. $n \times n$ matrix.

(i) A has n real eigenvalues (possibly repeated).

$\det(A - \lambda I)$ has n real roots (possibly repeated).

(ii) Eigenvectors corresponding to distinct eigenvalues are orthogonal.

(iii) Suppose λ^* is an eigenvalue $(A - \lambda^* I)$ has free columns.

$$\dim[\text{null}(A - \lambda^* I)] = \# \text{ free cols in } A - \lambda^* I$$

$$= \# \lambda^* \text{ is repeated among roots of } \det(A - \lambda I)$$

$\Rightarrow$ we get n linearly indpt eigenvectors.

$\therefore$ eigenvectors form a basis for $\mathbb{R}^n$.

we can also get n orthogonal vectors as a basis for $\mathbb{R}^n$.

Normalize all so that they have length 1.

$$\underline{v}_1 \dots \underline{v}_n.$$

$$A \underline{v}_1 = \lambda_1 \underline{v}_1,$$

$$A \underline{v}_2 = \lambda_2 \underline{v}_2$$

$$\vdots$$

$$A \underline{v}_n = \lambda_n \underline{v}_n.$$

$$\underline{v}_i^T \underline{v}_j = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j. \end{cases}$$

$$V = \begin{bmatrix} | & & | \\ \underline{v}_1 & \dots & \underline{v}_n \\ | & & | \end{bmatrix}.$$

$$\underline{V}^{-1} = \underline{V}^T.$$

$$\begin{aligned} \underline{V}^T \underline{V} &= \begin{bmatrix} i,j^{\text{th}} \text{ entry} \\ = \underline{v}_i^T \underline{v}_j \end{bmatrix} \\ &= \begin{bmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{bmatrix} \end{aligned}$$

$$AV = A \begin{bmatrix} v_1 & \dots & v_n \end{bmatrix} = \begin{bmatrix} \lambda_1 v_1 & \lambda_2 v_2 & \dots & \lambda_n v_n \end{bmatrix}$$

$$= V \underbrace{\begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix}}_{\Lambda}$$

$$= V \Lambda$$

$$A \underbrace{V V^T}_I = V \Lambda V^T$$

$$\therefore A = V \Lambda V^T$$

$$\det(A - \lambda I) = 0.$$

Roots

$$\lambda_1 \geq \lambda_2 \geq \lambda_3 \dots \geq \lambda_n.$$

Distinct roots

$$\left\{ \begin{array}{l} \lambda_{i_1} \text{ repeats } n_1 \text{ times.} \\ \lambda_{i_2} \text{ repeats } n_2 \text{ times.} \\ \vdots \\ \lambda_{i_k} \text{ ——— } n_k \text{ times.} \end{array} \right.$$

$$i_1=1 \quad i_2=2 \quad i_3=4 \quad i_4=5$$

$$\lambda_1 \geq \lambda_2 = \lambda_3 > \lambda_4 > \lambda_5 = \lambda_6$$

$$1, 0.5, 0.5, 0.2, -0.5, -0.5$$

$$\lambda_1 \text{ has multiplicity } 1.$$

$$\lambda_2 \text{ has ——— } 2.$$

$$\lambda_4 \text{ has ——— } 1.$$

$$\lambda_5 \text{ has ——— } 2.$$

$$\text{Clearly } \underline{n_1} + n_2 + \dots + n_k = n.$$

$$1 + 2 + 1 + 2 = 6.$$

$\text{null}(A - \lambda_{i_1} I)$ contributes n_1 linearly independent vectors.

$\text{null}(A - \lambda_{i_2} I)$ ——— n_2 ———

$\vdots$

$\text{null}(A - \lambda_{i_k} I)$ ——— n_k ———

Concatenate these lists (because vectors in diff eigenspaces above are going to be orthogonal if A is symmetric) to form a list of $n_1 + n_2 + \dots + n_k = n$ linearly indep vectors in $\mathbb{R}^n$.

Any vector in $\mathbb{R}^n$ is a linear combination of the above.

Let $v_1, \dots, v_n$ be n mutually orthogonal (unit-length) vectors.

$$A = V \Lambda V^T$$

$$V = \begin{bmatrix} v_1 & \dots & v_n \end{bmatrix}$$

$$\Lambda = \begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix}$$

$$AV = V\Lambda.$$

$$V^T AV = \underbrace{(V^T V)}_{=I} \Lambda = \Lambda = \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix}$$

* Interpret $A\underline{x}$ for any $\underline{x} \in \mathbb{R}^n$.

$$\underline{x} = \alpha_1 \underline{v}_1 + \dots + \alpha_n \underline{v}_n \quad (\text{because } \underline{v}_1, \dots, \underline{v}_n \text{ is a basis}).$$

$$A\underline{x} = \alpha_1 A\underline{v}_1 + \alpha_2 A\underline{v}_2 + \dots + \alpha_n A\underline{v}_n.$$

$$= \alpha_1 \lambda_1 \underline{v}_1 + \alpha_2 \lambda_2 \underline{v}_2 + \dots + \alpha_n \lambda_n \underline{v}_n$$

(for symmetric matrices).

* Suppose $X: m \times n$ matrix.

$$(AB)^T = B^T A^T.$$

$$(X^T X)^T = X^T (X^T)^T = X^T X. \quad (n \times n)$$

$$(X X^T)^T = (X^T)^T X^T = X X^T. \quad (m \times m)$$

$m \geq n$. (X is tall).

Eigenvalues of $X^T X$ are $\lambda_1, \dots, \lambda_n \geq 0$.

If $\lambda_n < 0$.

$$(X^T X) \underline{v}_n = \lambda_n \underline{v}_n.$$

$$\underline{v}_n^T (X^T X) \underline{v}_n = \lambda_n \underline{v}_n^T \underline{v}_n = \lambda_n \quad (\text{we chose } \underline{v}_n \text{ that way}).$$

$$\|\underline{w}_n\|^2 = \underline{w}_n^T \underline{w}_n = \underline{w}_n^T \underbrace{X \underline{v}_n}_{\underline{w}_n} = \underbrace{(\underline{v}_n^T X^T)}_{\underline{w}_n^T} \underbrace{(X \underline{v}_n)}_{\underline{w}_n}.$$