




# Forecasting and time variability analysis of Ozone concentrations using nitrogen oxide and meteorological variables as predictors

This manuscript ([permalink](#)) was automatically generated from [uiceds/cee-492-term-project-fall-2022-hydrograds@6d24888](#) on October 30, 2022.

## Authors

---

- **Jiewen Luo**  
·  [Noomi-Luo](#)  
Department of CEE, University of Illinois at Urbana&Champaign
- **Rourou Ji**  
·  [Jadeji](#)  
Department of CEE, University of Illinois at Urbana&Champaign
- **Bernardo Burbano**  
·  [BernieJBA](#)  
Department of CEE, University of Illinois at Urbana&Champaign

## CEE 492 Final Project Selection

---

### 1. Dataset description:

The dataset used in this project is a CSV file about the air quality in northern Taiwan collected in 2015 [<https://www.kaggle.com/datasets/nelsonchu/air-quality-in-northern-taiwan>], which include air quality data and meteorological monitoring data for research and analysis, originally from Environmental Protection Administration, Executive Yuan, R.O.C. (Taiwan). There are 25 observation stations in total. Columns in this CSV file are the following:

1. Time - The first column is the observation time of 2015
2. Station - The second column is the station name, there are 25 observation stations, those stations are showing at the table [1](#).

**Table 1:** A table contain all stations in Taiwan.

		station		
Banqiao	Cailiao	Datong	Dayuan	Guanyin
Guting	Keelung	Longtan	Pingzhen	Sanchong
Shilin	Songshan	Tamsui	Taoyuan	Tucheng
Wanhua	Wanli	Xindian	Xinzhuang	Xizhi
Yangming	Yonghe	Zhongli	Zhongshan	Linkou

3. Items - From the third column to the last one

#### 4. item - unit - description

- SO<sub>2</sub> - ppb - Sulfur dioxide
- CO - ppm - Carbon monoxide
- O<sub>3</sub> - ppb - ozone
- PM<sub>10</sub> - µg/m<sup>3</sup> - Particulate matter
- PM<sub>2.5</sub> - µg/m<sup>3</sup> - Particulate matter
- NO<sub>x</sub> - ppb - Nitrogen oxides
- NO - ppb - Nitric oxide
- NO<sub>2</sub> - ppb - Nitrogen dioxide
- THC - ppm - Total Hydrocarbons
- NMHC - ppm - Non-Methane Hydrocarbon
- CH<sub>4</sub> - ppm - Methane
- UVB - UVI - Ultraviolet index
- AMB\_TEMP - Celsius - Ambient air temperature
- RAINFALL - mm
- RH - % - Relative humidity
- WIND\_SPEED - m/sec - The average of the last ten minutes per hour
- WIND\_DIREC - degrees - The average of the last ten minutes per hour
- WS\_HR - m/sec - The average of an hour
- WD\_HR - degrees - The average of an hour
- PH\_RAIN - PH - Acid rain
- RAIN\_COND - µS/cm - Conductivity of acid rain

## Proposal:

---


The purpose of this project is to predict O<sub>3</sub> concentrations using measurements of concentration of other pollutants and available meteorological measurements. Ozone might be formed when heat and sunlight cause chemical reactions between oxides of nitrogen (NO<sub>x</sub>) and Volatile Organic Compounds (VOC), which are also known as Hydrocarbons. Therefore it could be hypothesized that using measurements of NO<sub>x</sub> as an independent variable a model could be developed to predict O<sub>3</sub> concentrations. Additionally, meteorological variables such as air temperature, relative humidity(RH) and ultraviolet index (UVB - UVI) could be included as independent variables to assess their influence on temporal variability of ozone. As an additional step wind-related variables such as mean wind velocity and direction will be included to study their effect on temporal variability of ozone.

After the air quality data has been processed the strongest O<sub>3</sub> predictors will be determined using PCA. PCA could be used to identify the main axes of variance within the dataset and explore underlying correlations that exist in a set of variables. Variables that are highly correlated cluster together. Using PCA 2D figures per each pair of variables are not needed, instead all the variables could be visualized simultaneously. Differences on PC1 are more important than differences on PC2. After plotting PCA plots, a heatmap could also be plotted to check the results. As additional criteria to identify the strongest predictors a LSTM network (long short-term memory network) can be used since the data used is time dependent. The network should contain several LSTM layers and fully-connected layers. The output should contain the pollution concentration and will point out the weights assigned to each correlated criterion, the values of such weights should also indicate what the strongest predictors are. Once the strongest predictors have been identified, genetic programming will be used to develop the models to predict O<sub>3</sub> concentrations.

## Exploratory Data Analysis:

---

In order to explore the relation between the dependent variable and independent variables several scatter plots were created between meteorological variables, pollutant concentrations and ozone concentrations. Additionally, a heatmap was generated to investigate the correlation values between ozone concentration and independent variables. The most correlated variables are RH(relative humidity) and UVB(Ultraviolet index). RH is negatively correlated with ozone,value of -0.51, while UVB is positively correlated, value of 0.51. Another relevant observation was that concentrations of nitrogen-related chemicals are highly correlated between each other, which is close to the truth.



**Figure 1:** Heatmap for the variables we considered in this analysis

Furthermore, the fraction of available measurements, meaning the number of data points available divided by the number of hours in a year, was computed for all stations and all measured variables. This computation helped visualize the stations that missed the least data points as well as the variables whose values are recorded the most consistently through different stations. The station with the highest fraction of available measurements was Banquiao, as seen in the table ?? For this reason the remaining portion of this EDA was devoted to this station.


	Dependent Variable	Independent variable: Nitrogen based				Independent variable: Metereological variables							
Station	O3	NMHC	NO	NO2	NOx	Temperature	RH	UVB	W Direction 10 min	W Speed 10min	W Speed hourly	Wind Direction hourly	Average Fraction
Banqiao	0.991	0.984	0.966	0.966	0.966	0.991	0.991	0.991	0.991	0.991	0.991	0.991	0.984
Cailliao	0.978	0.000	0.958	0.958	0.958	0.977	0.978	0.000	0.977	0.977	0.977	0.977	0.794
Dayuan	0.988	0.000	0.975	0.975	0.975	0.980	0.955	0.000	0.852	0.910	0.922	0.922	0.770
Guanyin	0.980	0.000	0.969	0.969	0.969	0.973	0.980	0.000	0.977	0.980	0.979	0.977	0.798
Guting	0.988	0.966	0.983	0.983	0.983	0.985	0.983	0.000	0.985	0.985	0.985	0.985	0.901
Keelung	0.987	0.970	0.977	0.977	0.977	0.977	0.980	0.000	0.986	0.986	0.986	0.986	0.891
Linkou	0.973	0.000	0.966	0.966	0.966	0.973	0.973	0.000	0.972	0.972	0.964	0.972	0.793
Longtan	0.982	0.000	0.961	0.961	0.961	0.982	0.982	0.000	0.982	0.982	0.981	0.981	0.798
Pingzhen	0.988	0.000	0.968	0.968	0.968	0.988	0.988	0.000	0.983	0.988	0.988	0.988	0.802
Shilin	0.983	0.000	0.957	0.957	0.957	0.983	0.983	0.000	0.983	0.983	0.980	0.983	0.797
Songshan	0.985	0.967	0.976	0.976	0.976	0.985	0.985	0.000	0.984	0.984	0.984	0.984	0.891
Tamsui	0.961	0.000	0.956	0.956	0.956	0.000	0.000	0.961	0.000	0.000	0.000	0.000	0.348
Taoyuan	0.983	0.958	0.976	0.976	0.976	0.982	0.982	0.976	0.982	0.982	0.982	0.982	0.978
Tucheng	0.981	0.974	0.965	0.965	0.965	0.980	0.981	0.000	0.980	0.980	0.980	0.980	0.886
Wanhua	0.986	0.000	0.979	0.979	0.979	0.986	0.986	0.000	0.986	0.986	0.986	0.986	0.805
Wanli	0.984	0.000	0.976	0.976	0.976	0.983	0.976	0.000	0.976	0.982	0.983	0.975	0.800
Xindian	0.985	0.000	0.979	0.979	0.979	0.985	0.982	0.000	0.985	0.985	0.985	0.985	0.804
Xinzhuang	0.978	0.000	0.970	0.970	0.970	0.978	0.978	0.000	0.978	0.978	0.978	0.978	0.798
Xizhi	0.963	0.000	0.954	0.954	0.954	0.962	0.962	0.000	0.962	0.962	0.958	0.958	0.784
Yangming	0.979	0.000	0.964	0.964	0.964	0.979	0.979	0.000	0.000	0.000	0.000	0.000	0.441
Yonghe	0.987	0.972	0.978	0.978	0.978	0.984	0.987	0.000	0.987	0.987	0.987	0.987	0.893
Zhongli	0.988	0.973	0.967	0.967	0.967	0.986	0.986	0.000	0.986	0.986	0.985	0.985	0.890
Zhongshan	0.986	0.976	0.976	0.976	0.976	0.986	0.986	0.000	0.986	0.986	0.986	0.986	0.893

**Figure 2:** Data availability of air quality stations

Other relevant statistics from the Banquo air quality station are shown as Table [2](#)

**Table 2:** Descriptive statistics of features that compose in Banqiao Station

	count	mean	std	min	25%	50%	75%	max
AMB_TEMP	8682	1.18E-15	1.00	-2.47	-0.89	0.15	0.68	2.25
NMHC	8619	-4.74E-15	1.00	-1.32	-0.62	-0.30	0.29	16.26
NO	8462	3.20E-15	1.00	-0.75	-0.50	-0.32	0.05	23.55
NO2	8462	1.56E-15	1.00	-1.94	-0.78	-0.10	0.57	5.48
NOx	8462	1.62E-15	1.00	-1.54	-0.69	-0.20	0.42	14.67
O3	8685	-8.78E-16	1.00	-1.35	-0.80	-0.07	0.54	6.06
RH	8684	-1.56E-15	1.00	-4.85	-0.75	0.17	0.76	2.35
UVB	8684	3.53E-15	1.00	-0.56	-0.56	-0.56	0.12	4.81
WD_Hour	8680	2.64E-16	1.00	-1.58	-0.78	-0.59	1.01	2.32
WIND_DIREC_10min	8682	6.96E-16	1.00	-1.56	-0.78	-0.58	1.01	2.31
WIND_SPEED_10min	8682	-6.21E-15	1.00	-1.43	-0.72	-0.18	0.53	7.92
WS_HR	8680	-2.83E-15	1.00	-1.67	-0.80	-0.12	0.56	7.86




**Figure 3:** Scatter plots of hourly measurements of variables


As described in previous sections, the dataset consists of hourly observations of ozone (dependent variable) and several pollutant concentrations and meteorological measurements (independent variables). The first step was to plot ozone against all of the independent variables to visualize if the data collapsed into any identifiable pattern, thus to later on use such a pattern to identify potential models. The measurements in an hourly time scale did not show any discernible pattern between the dependent and independent variables. As seen in figure 3.

Plotting the raw data, i.e. the available measurements without any processing or transformation, did not yield any insights that could help elucidate the relation between the variables. Therefore, the data was normalized. Notwithstanding, normalization did not translate into plots where patterns could be identified. Thus, the data was processed again following two consecutive steps. First the values were averaged over a day and over a month producing a dataset of daily and monthly measurement. Second, such values were standardized by dividing them by the corresponding daily and monthly averages.


The resulting daily and monthly standardized averages were plotted against time. Plotting the daily averaged variables shown plots where the fluctuations of the values happened in a relatively short time and thus such fluctuations obscured any pattern that could be observed in the data, as seen in the next figure. Conversely, when the monthly standardized averages were plotted against time it was visible that the pollutants concentrations shown similar time patterns as seen in figures 4.



**Figure 4:** Daily concentrations of ozone



**Figure 5:** Standardized pollutants and ozone monthly concentration changes



**Figure 6:** Standardized meteorological measurements and standardized ozone monthly concentration changes



In regards to pollutant concentrations, O<sub>3</sub> peaked in the months when concentration of the nitrogen based pollutants and non-methane hydrocarbons dropped. This is especially the case for NO concentrations (green line). This pattern of corresponding decreasing pollutant concentrations and increasing ozone could suggest that the pollutant concentrations are negatively correlated with ozone concentrations. This is also consistent with figure 5 (correlation plot)

In regards to the meteorological variables, UVB (ultraviolet index) and air temperature peak in the same months. Both temperature and UVB experience an increase in their values from the beginning of the year peaking in June. After June, both values experience a steady decrease. No matching pattern can be observed in terms of the relation of the latter two variables and ozone concentrations.

The increase or decrease of the values of wind direction shown in figure... correspond to a relative shift in direction of the wind compared to the yearly wind direction. These shifts in the direction of the wind can be used later on the forecasting of O<sub>3</sub> concentration. Wind direction could help elucidate if O<sub>3</sub> concentration from upwind neighboring locations could affect O<sub>3</sub> values in the location of interest, Banquiao.

## Predictive Modeling

---

As a first iteration, a linear model with multiple independent variables will be optimized using available standardized measurements of ozone. A first model will be produced only using standardized pollutant values, and a second model will include as additional variables wind direction and upwind station standardized ozone concentrations.



**Figure 7:** Potential upwind station with respect to Banquiao

If after training, the linear model mean square error (computed using predictions of ozone and observation) is above a certain threshold, a more involved model will be used. Two candidates for the second iteration of the predictive model will be considered. A fully connected neural network and a model produced with genetic programming packages in python for model discovery.

In order to train the neural network, measurements from Banquiao of the first 8 months of the year will be used. The variables that will be used in the training are: pollutant and ozone measurements, wind direction, ozone measurements from the 6 potential upwind stations shown in figure. In order to validate the model, data from the remaining 4 months will be used in order to assess the model ability to forecast ozone concentrations.



# References

---