Determination of the Refractive Indexes of Air and Acrylic

Yi Hu

March 6, 2016

Date Performed: March 2, 2016 Partners: Yi Hu

Trina Le

Instructor: Kyle Slinker

1 Objective

To determine the atomic weight of magnesium via its reaction with oxygen and to study the stoichiometry of the reaction (as defined in 1.1):

1.1 Definitions

Stoichiometry The relationship between the relative quantities of substances taking part in a reaction or forming a compound, typically a ratio of whole integers.

Atomic mass The mass of an atom of a chemical element expressed in atomic mass units. It is approximately equivalent to the number of protons and neutrons in the atom (the mass number) or to the average number allowing for the relative abundances of different isotopes.

2 Experimental Data

Mass of empty crucible	$7.28\mathrm{g}$
Mass of crucible and magnesium before heating	$8.59\mathrm{g}$
Mass of crucible and magnesium oxide after heating	$9.46\mathrm{g}$
Balance used	#4
Magnesium from sample bottle	#1

3 Sample Calculation

 $\begin{array}{lll} \text{Mass of magnesium metal} &= 8.59\,\mathrm{g} - 7.28\,\mathrm{g} \\ &= 1.31\,\mathrm{g} \\ \\ \text{Mass of magnesium oxide} &= 9.46\,\mathrm{g} - 7.28\,\mathrm{g} \\ &= 2.18\,\mathrm{g} \\ \\ \text{Mass of oxygen} &= 2.18\,\mathrm{g} - 1.31\,\mathrm{g} \\ &= 0.87\,\mathrm{g} \end{array}$

Because of this reaction, the required ratiotwo significant figures).

4 Results and Conclusions

The atomic weight of magnesium is concluded to rmined by the stoichiometry of its chemical combination with oxygen. This result is in agreement with the accepted value.



Figure 1: Figure caption.

5 Discussion of Experimental Uncertainty

6 Answers to Definitions

- a. The atomic weight of an element is the relative weight of one of its atoms compared to C-12 with a weight of 12.0000000..., hydrogen with a weight of 1.008, to oxygen with a weight of 16.00. Atomic weight is also the average weight of all the atoms of that element as they occur in nature.
- b. The units of atomic weight are two-fold, with an identical numerical value. They are g/mole of atoms (or just g/mol) or amu/atom.
- c. Percentage discrepancy between an accepted (literature) value and an experimental value is

 $\frac{\text{experimental result} - \text{accepted result}}{\text{accepted result}}$

References

[1] Smith, J. M. and Jones, A. B. (2012). Chemistry. Publisher, 7th edition.