

# Computational Data Mining

Fatemeh Mansoori

# Course material

- <https://github.com/uisf-course/CDM>

# Multiplication $Ax$ Using rows of $A$

- Multiply  $A$  times  $x$  using the three rows of  $A$  :

$$\text{By rows} \quad \begin{bmatrix} 2 & 3 \\ 2 & 4 \\ 3 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2x_1 + 3x_2 \\ 2x_1 + 4x_2 \\ 3x_1 + 7x_2 \end{bmatrix} = \begin{matrix} \text{inner products} \\ \text{of the rows} \\ \text{with } x = (x_1, x_2) \end{matrix}$$

- a row at a time
- are also known as "dot products" because of the dot notation

$$\text{row} \cdot \text{column} = (2, 3) \cdot (x_1, x_2) = 2x_1 + 3x_2$$

- use this for computing but not for understanding.

# Multiplication $Ax$ Using Columns of $A$

- For Understanding  $Ax$  using vector approach.

**By columns** 
$$\begin{bmatrix} 2 & 3 \\ 2 & 4 \\ 3 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix} + x_2 \begin{bmatrix} 3 \\ 4 \\ 7 \end{bmatrix} = \begin{matrix} \text{combination} \\ \text{of the columns} \\ \mathbf{a}_1 \text{ and } \mathbf{a}_2 \end{matrix}$$

- vector approach sees  $Ax$  as a "linear combination" of  $\mathbf{a}_1$  and  $\mathbf{a}_2$
- linear combination of  $\mathbf{a}_1$  and  $\mathbf{a}_2$  includes two steps :
  - Multiply the columns  $\mathbf{a}_1$  and  $\mathbf{a}_2$  by "scalars"  $x_1$  and  $x_2$
  - Add vectors  $x_1 \mathbf{a}_1 + x_2 \mathbf{a}_2 = Ax$ .
- $Ax$  is a linear combination of the columns of  $A$
- leads us to the column space of  $A$ .
  - all combinations of the columns
  - Space created by  $Ax$  for all vectors  $x$

# Column space of A $C(A)$

- All combinations  $x_1 a_1 + x_2 a_2 = Ax$  produce what part of the full 3D space?
- produce a plane
  - contains the complete line in the direction of  $a_1 = (2, 2, 3)$
  - contains the complete line in the direction of  $a_2 = (3, 4, 7)$
  - *sum* of any vector on one line plus any vector on the other line
  - This addition fills out an infinite plane containing the two lines.
  - does not fill out the whole 3-dimensional space
- If  $b = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$  is in the column space of A ?

## Column space of A $C(A)$

- What are the column spaces of

$$A_2 = \begin{bmatrix} 2 & 3 & 5 \\ 2 & 4 & 6 \\ 3 & 7 & 10 \end{bmatrix} \text{ and } A_3 = \begin{bmatrix} 2 & 3 & 1 \\ 2 & 4 & 1 \\ 3 & 7 & 1 \end{bmatrix} ?$$

# What is all possible column spaces inside $\mathbb{R}^3$

- The zero vector  $(0, 0, 0)$
- A line of all vectors  $x_1 a_1$
- A plane of all vectors  $x_1 a_1 + x_2 a_2$
- The whole  $\mathbb{R}^3$  with all vectors  $x_1 a_1 + x_2 a_2 + x_3 a_3$

# Matrix C

- create a matrix  $C$  whose columns independence columns of  $A$
- construction of  $C$  from the  $n$  columns of  $A$  :
  - If column 1 of  $A$  is not all zero, put it into the matrix  $C$ .
  - If column 2 of  $A$  is not a multiple of column 1, put it into  $C$ .
  - If column 3 of  $A$  is not a combination of columns 1 and 2, put it into  $C$ .  
*Continue.*
- At the end  $C$  will have columns ( $r \leq n$ ).
- They will be a "basis" for the column space of  $A$ .
- A basis for a subspace is a full set of independent vectors
- All vectors in the space are combinations of the basis vectors



# Example

- What is matrix C and R for the matrices below :

$$A = \begin{bmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \\ 0 & 1 & 2 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 & 5 \\ 1 & 2 & 5 \\ 1 & 2 & 5 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}$$

# Rank of the matrix

**The rank of a matrix is the dimension of its column space.**

- Different basis, but always the same number of vectors
- That number  $r$  is the "dimension" of the column space of  $A$  and  $C$

# Matrix R

- The matrix  $C$  connects to  $A$  by a third matrix  $R$ :  $A = CR$
- Their shapes are :
  - $(m \text{ by } n) = (m \text{ by } r) (r \text{ by } n)$ .

$$A = \begin{bmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \\ 0 & 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \end{bmatrix} = CR$$

$C$  multiplies the first column  $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$  of  $R$ , this produces column 1 of  $C$  and  $A$ .

$C$  multiplies the second column  $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$  of  $R$ , we get column 2 of  $C$  and  $A$ .

•  $C$  multiplies the third column  $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$  of  $R$ , we get  $2(\text{column 1}) + 2(\text{column 2})$ .

## Matrix R

$$A = \begin{bmatrix} 1 & 2 & 5 \\ 1 & 2 & 5 \\ 1 & 2 & 5 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 5 \end{bmatrix} = CR$$

All three matrices have rank  $r = 1$   
**Column Rank = Row Rank**

The number of *independent columns* equals the number of *independent rows*

# Matrix-Matrix Multiplication $AB$

- **Inner products** (*rows times columns*) produce each of the numbers in  $AB = C$ :

**row 2 of  $A$**   
**column 3 of  $B$**   
**give  $c_{23}$  in  $C$**

$$\begin{bmatrix} \cdot & \cdot & \cdot \\ a_{21} & a_{22} & a_{23} \\ \cdot & \cdot & \cdot \end{bmatrix} \begin{bmatrix} \cdot & \cdot & b_{13} \\ \cdot & \cdot & b_{23} \\ \cdot & \cdot & b_{33} \end{bmatrix} = \begin{bmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & c_{23} \\ \cdot & \cdot & \cdot \end{bmatrix}$$

dot product  $c_{23} = (\text{row 2 of } A) \cdot (\text{column 3 of } B)$  is a sum of  $a$ 's times  $b$ 's:

$$c_{23} = a_{21} b_{13} + a_{22} b_{23} + a_{23} b_{33} = \sum_{k=1}^3 a_{2k} b_{k3}$$

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}.$$

# Matrix-Matrix Multiplication $AB$

- columns of  $A$  times rows of  $B$ .
- Outer product :
  - *one column  $u$  times one row  $v^T$*

<b>“Outer product”</b>	$uv^T = \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix} \begin{bmatrix} 3 & 4 & 6 \end{bmatrix} = \begin{bmatrix} 6 & 8 & 12 \\ 6 & 8 & 12 \\ 3 & 4 & 6 \end{bmatrix} =$	<b>“rank one matrix”</b>
------------------------	--	--------------------------

- $m$  by  $1$  matrix (a column  $u$ ) times a  $1$  by  $p$  matrix (a row  $v^T$ ) gives an  $m$  by  $p$  matrix

what is special about the rank one matrix  $uv^T$ :

# AB multiplication

- All column of  $uv^T$  are multiply of  $u = \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$
- All rows are multiply of  $v^T = [3 \ 4 \ 6]$
- column space of  $uv^T$  is one-dimensional: *the line in the direction of  $u$*
- All nonzero matrices  $uv^T$  have rank one

AB multiplication

**$AB = \text{Sum of Rank One Matrices}$**

$$AB = \left[ \begin{array}{c|ccc|c} & & & & \\ & & & & \\ a_1 & & \dots & & a_n \\ & & & & \\ & & & & \end{array} \right] \left[ \begin{array}{c} \text{---} b_1^* \text{---} \\ \vdots \\ \text{---} b_n^* \text{---} \end{array} \right] = a_1 b_1^* + a_2 b_2^* + \dots + a_n b_n^*.$$

**sum of rank 1 matrices**

$$\begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 \\ 0 & 5 \end{bmatrix} =$$



# Question

- Suppose the column space of an  $m$  by  $n$  matrix is all of  $\mathbf{R}^3$ . What can you say about  $m$ ? What can you say about  $n$ ? What can you say about the rank  $r$ ?
- Suppose  $A$  is the 3 by 3 matrix ones(3, 3) of all ones. Find two independent vectors  $x$  and  $y$  that solve  $Ax = 0$  and  $Ay = 0$ . Write that first equation  $Ax = \mathbf{0}$  (with numbers) as a combination of the columns of  $A$ . Why don't I ask for a third independent vector with  $Az = 0$ ?