





lickcalc: Easy analysis of lick microstructure in experiments of rodent ingestive behaviour

K. Linnea Volcko¹ and James E. McCutcheon¹

¹ Dept. of Psychology, UiT The Arctic University of Norway, Tromsø, Norway  Corresponding author

DOI: 10.xxxxxx/draft

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Lick microstructure is a term used to describe the information that can be obtained from a detailed study of when individual licks occur, when a rodent is drinking. Rather than simply recording total intake (volume consumed), lick microstructure examines how licks are grouped, and the spacing of these groups of licks. This type of analysis can provide important insight into why an animal is drinking, such as if it is driven by taste or as a response to the consequences of consumption (e.g. feeling “full”). The simplicity of using lickcalc, requiring only a drag-and-drop of files, will make microstructural analysis accessible to any who wish to use it while providing sophisticated analyses with high scientific value.

Statement of need

lickcalc is a software suite that performs microstructural lick analysis on data files containing timestamps of lick onset and offset, in MedAssociates or csv/txt format. In addition to providing an overview of individual files (corresponding to a particular session for a single subject), results of the analysis can be added to a table that is exportable to Excel.

Microstructural analysis was first described in John D. Davis & Smith (1992) and has since then been used understand diverse phenomena. In-depth reviews on many of these, and microstructural parameters used to study them, are available (Alexander W. Johnson, 2018; Naneix et al., 2020; Smith, 2001). Briefly, although much of the foundational work on drinking microstructure was on licking for nutritive solutions (e.g., sucrose solutions), microstructural analysis can also be used to study intake of water (McKay & Daniels, 2013; Santollo et al., 2021), ethanol (Patwell et al., 2021), and other tastants such as non-caloric artificial sweeteners, sodium and quinine (Lin et al., 2012; Spector & St. John, 1998; Verharen et al., 2019). Lick microstructure has been used to shed light on, for example, how licking is affected by neuropeptides (McKay & Daniels, 2013), enzymes in the mouth (Chometton et al., 2022), ovarian hormones (Santollo et al., 2021), nutrient restriction (Naneix et al., 2020), response to alcohol (Patwell et al., 2021), and diet (Alexander W. Johnson, 2012). The number of lick bouts over a session are thought to reflect postingestive feedback from the consumed fluid, whereas the number of licks in a bout are thought to reflect palatability of the solution.

Lick microstructure can provide a great deal of interesting information about why an animal is drinking. Often, changes in microstructure are accompanied by changes in total intake, but this is not always the case: sometimes, equal intake will be achieved by quite different licking patterns that indicate changes in orosensory and postingestive feedback (A. W. Johnson et al., 2010; Volcko et al., 2020). Analyzing lick microstructure is therefore highly valuable in understanding how a manipulation affects appetite; if X causes an animal to feel more satiated after drinking, that may lead to a different interpretation than if X were to reduce the palatability of the solution. Because of the value of microstructural data, many labs habitually

record and analyze it. There are many others, however, that have not yet begun collecting these data. Investing in lickometers can be costly, but there are an increasing number of alternatives to commercial products such as those produced by Med Associates. Several open-source lickometer designs are now available (e.g. Frie & Khokhar (2024); Monfared et al. (2024); Petersen et al. (2024); Silva et al. (2024); Raymond et al. (2018)).

Recording individual licks with high temporal resolution is necessary for microstructural analysis of drinking behavior, but another barrier to reporting microstructure is its analysis. This problem can now easily be solved by lickcalc. lickcalc does not require any special software or coding knowledge: all the user has to do is drag a file with timestamps of lick onset (and, ideally, offset) into the program and lickcalc will generate a detailed microstructural analysis, with a high degree of user control over key parameters. After loading a file and setting parameters, lickcalc displays results, the detail of which are an additional advantage of using the program. Tables show values for number of licks, number of bursts, and burst size (among others) – the values that are often reported and used to draw inferences about post-ingestive and orosensory feedback of the solution. But importantly, there are several charts displayed that show information that helps with quality control of the data and challenges the user to think critically about which parameters they have chosen. In short, lickcalc makes microstructural analysis accessible to any who have timestamps of lick onsets, while providing a lot of information on the data underlying the analysis which helps in choosing the appropriate parameters.

Key features

lickcalc has several features that make it exceptionally user-friendly at the same time as providing a sophisticated and detailed microstructural analysis. Some of these features include:

- **ease of use:** a file in Med Associates or CSV/TXT format simply needs to be dragged into the lickcalc software in order to perform the analyses. Parameters can be set manually using sliders, and results exported to Excel with the push of a button.
- **flexibility:** the user sets key parameters appropriate for their experimental setup and data. Data can be examined by whole-session or different epochs.
- **customization:** by using the configuration file, users can change default settings to match their preferences and avoid manually changing settings for each file loaded.
- **results compilation:** data from multiple sessions and/or individuals can be exported into a single Excel file, which streamlines the analysis process.
- **detail of analysis:** one of the benefits of using lickcalc is the level of detail it provides. In addition to the properties often reported (e.g. burst number, burst size), lickcalc computes and displays attributes of the data that are important in establishing the quality of the data and determining appropriate parameters for its microstructural analysis. Four charts are:

- 1) *intra-burst lick frequency*, or how often certain interlick intervals within a burst of licking occur. While a rodent is licking, its tongue makes rhythmic protrusions that are under the control of a central pattern generator (Travers et al., 1997). Rats typically lick 6-7 times per second (John D. Davis & Smith, 1992), while mice lick at a slightly higher rate (A. W. Johnson et al., 2010). In addition to these species differences, there are also strain differences (A. W. Johnson et al., 2010; ?). Because intra-burst lick rate is under the control of the central pattern generator, it should remain relatively stable across mice and conditions (unless a manipulation is expected to cause changes in the central pattern generator). A typical chart for a mouse might show a sharp peak around an intra-burst ILI of around 129, which corresponds to a lick rate of 7.75 Hz. Much smaller peaks are often present at the harmonics of the intra-burst ILI (e.g., a primary peak at 129 will have smaller

peaks at 258, 387, and so on), often because of “missed licks” in which the mouse attempts to lick but its tongue misses the spout. A large number of these, or other differences from the expected pattern of results, may indicate problems with the experimental setup (e.g. if the animal fails to reach the spout frequently, then perhaps the spout is too far away).

2) *lick length* is only available when lick offset is included in the data file. As with intraburst lick frequency, lick length should show little variability and the graph will have a sharp peak. Occasionally a lickometer will register longer licks than normal. This may be because a “lick” is registered by something other than a tongue, such as if a rodent grabs the spout with its paws, or if a fluid droplet hangs between the spout and the cage and is thus able to complete the electrical circuit. Concerns about data quality may be warranted with increasing number and duration of long licks. `lickcalc` displays both the number and maximum duration (s) of licks above the threshold that the user has set. There is also an option to remove these problematic licks from the dataset.

3) *burst frequency*, or how often certain burst sizes occurred. This is informative because burst size, by virtue of being a mean (mean licks per burst), does not take into account potentially relevant information about the distribution of how many licks are in a burst. For example, a burst size of 80 could result from bursts all containing between 70 and 90 licks, or from many single licks and one or two burst with a lot of licks. The latter case might raise some questions about how reliable the burst size value is. Although single licks occur, they can also be caused by non-tongue contact with the lickometer. Changing the minimum licks/burst parameter can perhaps filter out some of these suspect “licks.”

4) *Weibull probability*. The Weibull analysis, as described in J. D. Davis (1996), uses a mathematical equation to fit the data to a survival function. Although used by some (Aja et al., 2001; Moran et al., 1998; Spector & St. John, 1998), it is still relatively rare to find Weibull probabilities in microstructural analyses. The Weibull function can be used on several aspects of data, such as lick rate across a session, but in the `lickcalc` program the Weibull probability is calculated for burst size. It plots the probability that, given n licks, the mouse will continue to lick. This makes it sensitive to the number of licks per burst parameter that is set by the user. The Weibull alpha means.... And the Weibull beta means...

Design and usage

`lickcalc` is hosted by UiT The Arctic University of Norway and can be accessed at <https://lickcalc.uit.no>. Alternatively, it can be installed locally following instructions in the repository. To use `lickcalc`, the user drags in a file in Med Associates or CSV/TXT format and indicates which column contains the lick onsets and, if applicable, the lick offsets. A plot is automatically generated that displays a histogram of licks across the session. Session length defaults to the time of the last licks. This can be manually changed, or set in the optional “config file”. Session length can be set in seconds, minutes, or hours. The bin size (licks per unit of time) can be changed manually or in the config file. The user can toggle between the default histogram and a plot showing cumulative licks.

A microstructural analysis is, in essence, a division of individual licks into groups of licks. To perform this grouping, the user must set several parameters. One of these is the inter-lick interval (ILI), which is the minimum amount of time licks must be separated by in order to be considered separate groups. Early studies identified ILI of 251-500 ms as separating “bursts” of licking, and pauses of more than 500 ms as separating “clusters” of licking (i.e. a cluster of licks is made up of several bursts of licking). Others have argued that ILI of 1 s better reflects separation of lick bursts (Spector & St. John, 1998). In `lickcalc`, the user may set

the ILI between 250 ms and 2 s. Another parameter that needs to be decided prior to the lick analysis is the minimum number of licks per burst. `lickcalc` allows between 1 and 5 licks. The appropriate number of minimum licks/burst may vary depending on experimental set up, and the likelihood that a single lick represents a lick rather than, for example, a tail touching the spout. Finally, in `lickcalc`, the user must set a “long-lick threshold” between 0.1 and 1 s. This parameter is only available when lick offset is included. Licks that are longer than the set threshold are counted as “long” and may indicate a problem (e.g. the mouse holding the spout with its paws) rather than a true lick. The user can decide whether to remove “long licks” or not. All of these parameters can be set manually or through the config file. Four plots are generated (see Key Features section above), and tables are displayed showing values of several properties: total licks, intraburst frequency, number of long licks, maximum duration of long licks, number of bursts, mean licks per burst, weibull alpha, weibull beta, and weibull r-squared.

To save these data, the user has two options. The first is to export an excel file for the given data file. The user sets the animal ID and chooses which data should be exported in addition to the summary of the microstructural analysis. Data for “session histogram,” “intraburst frequency,” “lick length,” “burst histogram,” “burst probability,” and “burst details” are available. These allow the user to recreate the plots displayed in `lickcalc`. The second option for saving the data is to add the microstructural data to “results summary” table in the app. The results in this table remain even as new data files are loaded, so the data from many sessions (and/or individual animals) can be exported into a single Excel file. In addition to the data from the whole session, the user can choose to divide the session into epochs, or to examine only the first *n* bursts, or perform a trial-based analysis (e.g., for Davis rig experiments). Each of these analysis epochs can be added to the results summary table. The table contains the data as well as the parameters (e.g., minimum burst size) used to generate them. Finally, a batch process feature is available allowing multiple files to be analysed using the same parameters.

Acknowledgements

We acknowledge contributions from colleagues in the field of ingestive behaviour who have thought deeply about the meaning of patterns of licking. In particular, the following have either contributed data for us to test or have advised on the design of the program and analysis: (in alphabetical order) Derek Daniels, Samantha Fortin, Kevin Myers, Jess Santollo, Lindsey Schier, Alan Spector.

References

- Aja, S., Schwartz, G. J., Kuhar, M. J., & Moran, T. H. (2001). Intracerebroventricular CART peptide reduces rat ingestive behavior and alters licking microstructure. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 280(6), R1613–R1619. <https://doi.org/10.1152/ajpregu.2001.280.6.R1613>
- Chometton, S., Jung, A.-H., Mai, L., Dal Bon, T., Ramirez, A. O., Pittman, D. W., & Schier, L. A. (2022). A glucokinase-linked sensor in the taste system contributes to glucose appetite. *Molecular Metabolism*, 64, 101554. <https://doi.org/10.1016/j.molmet.2022.101554>
- Davis, J. D. (1996). Deterministic and probabilistic control of the behavior of rats ingesting liquid diets. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 270(4), R793–R800. <https://doi.org/10.1152/ajpregu.1996.270.4.R793>
- Davis, John D., & Smith, G. P. (1992). Analysis of the microstructure of the rhythmic tongue movements of rats ingesting maltose and sucrose solutions. *Behavioral Neuroscience*, 106(1), 217–228. <https://doi.org/10.1037/0735-7044.106.1.217>

- 188 Frie, J. A., & Khokhar, J. Y. (2024). FARESHARE: An open-source apparatus for assessing
189 drinking microstructure in socially housed rats. *NPP—Digital Psychiatry and Neuroscience*,
190 2(1), 1. <https://doi.org/10.1038/s44277-024-00002-z>
- 191 Johnson, Alexander W. (2012). Dietary manipulations influence sucrose acceptance in diet
192 induced obese mice. *Appetite*, 58(1), 215–221. <https://doi.org/10.1016/j.appet.2011.09.015>
- 193
- 194 Johnson, Alexander W. (2018). Characterizing ingestive behavior through licking microstructure:
195 Underlying neurobiology and its use in the study of obesity in animal models. *International*
196 *Journal of Developmental Neuroscience: The Official Journal of the International Society for*
197 *Developmental Neuroscience*, 64, 38–47. <https://doi.org/10.1016/j.ijdevneu.2017.06.012>
- 198 Johnson, A. W., Sherwood, A., Smith, D. R., Wosiski-Kuhn, M., Gallagher, M., & Holland, P.
199 C. (2010). An analysis of licking microstructure in three strains of mice. *Appetite*, 54(2),
200 320–330. <https://doi.org/10.1016/j.appet.2009.12.007>
- 201 Lin, J.-Y., Amodeo, L. R., Arthurs, J., & Reilly, S. (2012). Taste neophobia and palatability:
202 The pleasure of drinking. *Physiology & Behavior*, 106(4), 515–519. <https://doi.org/10.1016/j.physbeh.2012.03.029>
- 203
- 204 McKay, N. J., & Daniels, D. (2013). Glucagon-like peptide-1 receptor agonist administration
205 suppresses both water and saline intake in rats. *Journal of Neuroendocrinology*, 25(10),
206 929–938. <https://doi.org/10.1111/jne.12086>
- 207 Monfared, M., Mascret, Q., Marroquin-Rivera, A., Blanc-Árabe, L., Lebouleux, Q., Lévesque,
208 J., Gosselin, B., & Labonté, B. (2024). High-throughput low-cost digital lickometer system
209 for the assessment of licking behaviours in mice. *Journal of Neuroscience Methods*, 410,
210 110221. <https://doi.org/10.1016/j.jneumeth.2024.110221>
- 211 Moran, T. H., Katz, L. F., Plata-Salaman, C. R., & Schwartz, G. J. (1998). Disordered
212 food intake and obesity in rats lacking cholecystokinin A receptors. *American Journal*
213 *of Physiology-Regulatory, Integrative and Comparative Physiology*, 274(3), R618–R625.
214 <https://doi.org/10.1152/ajpregu.1998.274.3.R618>
- 215 Naneix, F., Peters, K. Z., & McCutcheon, J. E. (2020). Investigating the Effect of Physiological
216 Need States on Palatability and Motivation Using Microstructural Analysis of Licking.
217 *Neuroscience*, 447, 155–166. <https://doi.org/10.1016/j.neuroscience.2019.10.036>
- 218 Patwell, R., Yang, H., Pandey, S. C., & Glover, E. J. (2021). An operant ethanol self-
219 administration paradigm that discriminates between appetitive and consummatory behaviors
220 reveals distinct behavioral phenotypes in commonly used rat strains. *Neuropharmacology*,
221 201, 108836. <https://doi.org/10.1016/j.neuropharm.2021.108836>
- 222 Petersen, N., Adank, D. N., Quan, Y., Edwards, C. M., Hallal, S. D., Taylor, A., Winder, D. G.,
223 & Doyle, M. A. (2024). A Novel Mouse Home Cage Lickometer System Reveals Sex- and
224 Housing-Based Influences on Alcohol Drinking. *eNeuro*, 11(10), ENEURO.0234–24.2024.
225 <https://doi.org/10.1523/ENEURO.0234-24.2024>
- 226 Raymond, M. A., Mast, T. G., & Breza, J. M. (2018). An open-source lickometer and
227 microstructure analysis program. *HardwareX*, 4, e00035. <https://doi.org/10.1016/j.ohx.2018.e00035>
- 228
- 229 Santollo, J., Edwards, A. A., Howell, J. A., & Myers, K. E. (2021). Bidirectional effects
230 of estradiol on the control of water intake in female rats. *Hormones and Behavior*, 133,
231 104996. <https://doi.org/10.1016/j.yhbeh.2021.104996>
- 232 Silva, A., Carriço, P., Fernandes, A. B., Saraiva, T., Oliveira-Maia, A. J., & Da
233 Silva, J. A. (2024). High-Precision Optical Fiber-Based Lickometer. *Eneuro*, 11(7),
234 ENEURO.0189–24.2024. <https://doi.org/10.1523/ENEURO.0189-24.2024>
- 235 Smith, G. P. (2001). John Davis and the meanings of licking. *Appetite*, 36(1), 84–92.

- 236 <https://doi.org/10.1006/appe.2000.0371>
- 237 Spector, A. C., & St. John, S. J. (1998). Role of taste in the microstructure of quinine
238 ingestion by rats. *American Journal of Physiology-Regulatory, Integrative and Comparative*
239 *Physiology*, 274(6), R1687–R1703. <https://doi.org/10.1152/ajpregu.1998.274.6.R1687>
- 240 Travers, J. B., Dinardo, L. A., & Karimnamazi, H. (1997). Motor and Premotor Mechanisms
241 of Licking. *Neuroscience & Biobehavioral Reviews*, 21(5), 631–647. [https://doi.org/10.1016/S0149-7634\(96\)00045-0](https://doi.org/10.1016/S0149-7634(96)00045-0)
- 242
- 243 Verharen, J. P. H., Roelofs, T. J. M., Menting-Henry, S., Luijendijk, M. C. M., Vanderschuren,
244 L. J. M. J., & Adan, R. A. H. (2019). Limbic control over the homeostatic need for sodium.
245 *Scientific Reports*, 9(1), 1050. <https://doi.org/10.1038/s41598-018-37405-w>
- 246 Volcko, K. L., Brakey, D. J., Przybysz, J. T., & Daniels, D. (2020). Exclusively drinking
247 sucrose or saline early in life alters adult drinking behavior by laboratory rats. *Appetite*,
248 149, 104616. <https://doi.org/10.1016/j.appet.2020.104616>

DRAFT