Order-Aware Temporal Dynamics in Disorders of Consciousness: Multi-Scale Electrophysiological Modeling and Multicenter Evaluation

Hongjie Chen[1], Jianming Chen[2], Guanyu Xiong[3], Fang Duan[1,*], and Fang Li[4,*]

***Abstract*— Conventional bedside electroencephalography markers discard temporal sequence order, obscuring dynamic cortical reverberation critical for detecting covert consciousness. We formulate an interpretable electrophysiological framework combining physically adjacent lag-one autocorrelation, normalized signal jump rate, and common-mode-decoupled residual bilateral correlation under equal-weight consensus scoring. Evaluation spanned four cohorts totaling 129 patients across continuous telemetry, rapid triage, and circadian regimes. On continuous discovery telemetry from 41 patients, the primary four-feature suite attained an area under the receiver operating characteristic curve of 0.9324 across 25 cross-validation partitions, exceeding the strongest matched supervised baseline by a paired margin of 0.0360 (95 percent confidence interval 0.0090 to 0.0728). Physical sequence dependence was confirmed by 10,000 synchronous whole-channel temporal permutations, reducing discrimination from 0.8911 to a permutation baseline of 0.8182 plus or minus 0.0049. On external inpatient telemetry from 15 subacute patients at China Rehabilitation Research Center, source-frozen transfer yielded an area of 0.5741, whereas unsupervised persistence-anchored direction alignment reached 0.7778 (0.7963 with soft truncation), capturing injury-dependent bilateral decoupling. On 120-second triage records from 50 patients, the jump-rate operator achieved 0.7691, detecting all 19 minimally conscious cases. Sleep-stage transitions across 23 patients captured macro-scale circadian dynamics. Requiring 2.15 milliseconds of algorithmic execution time per evaluated update step, this parameter-free framework operates over 1,800 times faster than the 4-second telemetry window, supporting auditable bedside neuromonitoring.**

***Index Terms*— Disorders of consciousness (DoC), electroencephalography (EEG), temporal dynamics, bedside neuromonitoring, mesocircuit hypothesis, consensus fusion.**

I. Introduction

SEVERE acute brain injury from trauma, anoxia, or stroke leaves patients in prolonged disorders of consciousness (DoC), clinically categorized into the vegetative state/unresponsive wakefulness syndrome (VS/UWS) and the minimally conscious state (MCS) [1], [2]. Distinguishing MCS from VS/UWS informs decisions on withdrawing life-sustaining treatment, administering analgesia, and projecting rehabilitation potential. Bedside clinical evaluations carry an estimated 41% misdiagnosis rate [3]. Although standardized assessments such as the Coma Recovery Scale-Revised (CRS-R) [3] reduce diagnostic error, motor impairment and fluctuating arousal frequently confound behavioral examinations [4]. Prospective multi-center cohorts establish that approximately 25% (and 15%–35% across multimodal cohorts) of behaviorally unresponsive patients exhibit cognitive-motor dissociation (CMD) [5], retaining volitional cortical responses without overt motor output.

Functional neuroimaging and perturbational protocols (fMRI, ^{18}F-FDG PET, TMS-EEG) [6], [7] show that preserved cortical complexity differentiates MCS from VS/UWS, but transport hazards preclude routine ICU deployment. Continuous bedside EEG provides a practical modality for tracking recovery on sparse ICU montages (4–8 electrodes) [8]–[10]. Prior work examined non-linear entropy [8], [11], microstates [12], sleep architecture [13], [14], detrended fluctuation analysis (DFA) [15], [16], source connectivity [17], and criticality [18]. Yet bedside translation remains limited: multi-scale DFA is disrupted by non-stationary artifact gaps, source imaging is ill-posed on sparse arrays, entropy is artifact-sensitive, and deep classifiers risk overfitting small cohorts ($N < 50$). Bedside assessment thus relies on band power, marginal dispersion ($-\mathrm{CV}_\theta$), and unadjusted bilateral correlations [10], [13], bounding discrimination to 0.75–0.82 AUROC.

A. The Permutation Invariance Defect of Conventional Metrics

This diagnostic plateau stems from a shared limitation: strict permutation invariance. Conventional pipelines rely on aggregate dispersion ($-\mathrm{CV}_\theta$) and instantaneous bilateral correlation (r_{interhem}), which are invariant under temporal sequence permutation. Treating continuous EEG trajectories as unordered sets discards directional multi-second continuity and cortical reverberation.

B. The Mesocircuit Hypothesis and Autoregressive Memory

This methodological gap diverges from established neurobiology of conscious recovery, specifically the anterior

[1]College of Information Science and Engineering, Huaqiao University, Xiamen 361021, China (e-mail: 2415251001@stu.hqu.edu.cn).
[2]School of Information Engineering, Minnan University of Science and Technology, Quanzhou, China (e-mail: xgxy001@mnust.edu.cn).
[3]School of Biomedical Engineering, University of British Columbia, Vancouver, Canada (gyxiong@student.ubc.ca).
[4]Parkinson Medical Center, Beijing Rehabilitation Hospital, Capital Medical University, Beijing, China.
*Corresponding authors: Fang Duan (first; duan.fang@hqu.edu.cn) and Fang Li (second; lifangcrrc@mail.ccmu.edu.cn).

forebrain mesocircuit hypothesis [19]. Under this framework, sustained awareness depends on recurrent excitatory loops linking the central thalamus, striatum, and frontoparietal association cortices. Preserved mesocircuit connectivity manifests electrophysiologically as multi-second autoregressive continuity, quantifiable via lag-1 autocorrelation (ρ_1) across consecutive epochs. In VS/UWS, widespread deafferentation and loss of striatal inhibition disinhibit the internal globus pallidus (GPi), driving central thalamic hyperpolarization and degrading cortical activity into memoryless fluctuations. Distinguishing conscious capacity from reflexive unresponsiveness thus requires operators sensitive to physical sequence persistence.

As illustrated in Fig. 1a, individual theta trajectories in MCS (e.g., P36) exhibit sustained multi-second wave-holding, whereas VS/UWS trajectories (e.g., P25) break into decoupled transients. Static dispersion ($-\mathrm{CV}_\theta$) displays substantial overlap between cohorts ($p = 0.0264$; Fig. 1c), whereas lag-1 persistence (ρ_1) reliably separates conscious from unresponsive states ($p = 0.0227$; Fig. 1b).

For short triage records (< 5 min) where sample autocorrelation suffers elevated variance ($\mathrm{Var}(\hat{\rho}_1) \sim \mathcal{O}(1/M)$), we formulate the normalized jump rate (D_1) across verified adjacent epochs ($\mathbb{I}_{\text{adj}} = 1$), stabilizing short-horizon temporal estimation while preventing spurious concatenation across artifact gaps (Section II-C).

C. Spurious Bilateral Coupling Driven by Diffuse Slow Waves

In acute brain injury, diffuse delta/theta oscillations propagate via volume conduction and subcortical pacemaking. Although common average reference (CAR) removes linear potential offsets, rectified band-power envelopes retain shared variance ($R^2 \approx 47\%$), inflating raw inter-hemispheric correlation. Decoupling this common mode via ordinary least squares (OLS) orthogonal projection isolates residual bilateral resonance (r_{residual}; Section II-D), distinguishing pathological slow-wave co-activation in VS/UWS from preserved modular dynamics in MCS.

D. Contributions of This Work

This study introduces an order-aware electrophysiological framework with three principal contributions:

1) *Order-aware temporal operators and artifact decoupling*: We formulate lag-1 autocorrelation (ρ_1) for continuous telemetry alongside normalized adjacent jump rates (D_1) for short recordings under physical adjacency constraints ($\mathbb{I}_{\text{adj}} = 1$), analytically unified via $D_1 \approx 2\pi^{-1/2}(-\mathrm{CV}_\theta)(1-\rho_1)^{1/2}$. Synchronously shuffling 10,000 surrogate sequences degrades classification from 0.8911 AUROC to permutation baseline (0.8182 ± 0.0049, $p < 0.0001$), confirming that diagnostic separation requires chronological sequence order rather than static marginal distributions. OLS orthogonal projection decouples shared slow-wave envelope variance ($R^2 \approx 47\%$) to isolate intrinsic bilateral resonance (r_{residual}).

2) *Interpretable, parameter-free bedside consensus*: The parsimonious four-feature suite $\mathbf{x}_{\text{ortho}} = [c_{\text{global}}, -\mathrm{CV}_\theta, \rho_1, r_{\text{residual}}]^\top$ attains 0.9324 AUROC and 0.8998 average precision (AP) on primary discovery telemetry. Given identical inputs, equal-weight consensus scoring exceeds the top matched supervised baseline (logistic regression) by $+0.0360$ AUROC (Dirichlet paired 95% CI $[+0.0090, +0.0728]$), executing in 2.15 ms per update step under streaming telemetry ingestion (Supplementary Note D).

3) *Tiered multicenter evaluation across clinical regimes*: The framework is evaluated across 129 records spanning four cohorts structured into three clinical tiers: Tier 1 inpatient telemetry (CRRC, $N = 15$) evaluates cross-hardware transfer and etiology-dependent spatial dynamics (0.5741 frozen vs. 0.7778–0.7963 under unsupervised anchor alignment); Tier 2 rapid triage (Figshare, $N = 50$) demonstrates short-epoch jump dynamics (D_1, 0.7691 AUROC); and Tier 3 polysomnography (Wielek, $N = 23$) tracks macro-scale sleep stage transitions across 24-h circadian cycles.

II. Mathematical Formulation and Methods

Fig. 2 summarizes the processing, scoring and evaluation workflow.

A. Clinical Cohorts and Ethical Compliance

Four multicenter cohorts totaling $N = 129$ records were evaluated under institutional ethical approvals:

1) Mendeley Discovery Cohort ($N = 41$): Continuous 6-channel bedside EEG (F3, F4, C3, C4, O1, O2; 256 Hz) from 41 patients with severe brain injuries at FRCCICMR Moscow [20]: 12 MCS (8 MCS$^+$, 4 MCS$^-$; duration 6.00 ± 0.00 h) and 29 VS/UWS (5.97 ± 0.61 h), diagnosed through repeated CRS-R assessments.

2) CRRC Inpatient Cohort ($N = 15$): Prospective bedside telemetry from 15 subacute male patients (6 MCS, 9 VS/UWS; 11 ICH, 4 TBI; age 51.7 ± 12.3 years, range 28–69; duration 19.5 ± 8.0 min; Ethics No. 2025-118-01 [21]) at China Rehabilitation Research Center (CRRC / Beijing Bo'ai Hospital) using a 32-channel NeuSen NSM2 system (1000 Hz, 4-s blocks; Supplementary Note A). Homologous 6 channels were evaluated; two patients (CRRC_008, CRRC_009) exhibited localized mechanical drift at O_2, enabling assessment on both the full cohort ($N = 15$) and the drift-free subset ($N = 13$; 6 MCS including 1 EMCS, 7 VS/UWS).

3) Figshare Short-Record Analysis ($N = 50$): High-density resting-state EEG (250 Hz, 120-s records) from 50 patients with prolonged DoC (19 MCS, 31 VS/UWS) at Hangzhou Mingzhou Brain Rehabilitation Hospital [12], [22], evaluating rapid bedside triage using homologous channels under fixed short-duration recordings.

4) Wielek PSG Cohort ($N = 23$): 24-hour polysomnography from 23 post-comatose patients (11 MCS, 12 VS/UWS) [13], with 30-second hypnograms (Wake, N1, N2, N3, REM) tracking macro-scale transition continuity across circadian sleep cycles.

B. EEG Acquisition, Preprocessing, and Physical Adjacency Constraint

Continuous 6-channel EEG (256 Hz) was re-referenced to Common Average Reference (CAR) and segmented into




Fig. 1. Bedside EEG dynamics and patient distributions. (a) Standardized theta power across 30 consecutive 4-s epochs in P36 (MCS) and P25 (VS/UWS). (b,c) Lag-1 persistence ρ_1 ($p = 0.0227$) and positive theta CV ($p = 0.0264$) across 12 MCS and 29 VS/UWS patients (two-sided Mann–Whitney tests). Points show individual patients; boxes show medians and IQRs; whiskers span 1.5 IQR. Panel c uses a logarithmic scale.




Fig. 2. System overview. Schematic EEG traces illustrate processing. Original block indices are retained, and lag-1 persistence and D1 use only physically adjacent clean blocks ($\mathbf{\Delta index = 1}$). Main and short-record branches use distinct four-feature suites and independently fitted equal-weight consensus scores. In the short-record branch, Spec., Spat., and Bilat. denote spectral contrast, spatial contrast, and bilateral correlation. Discovery uses 4-s epochs; Figshare triage uses 2-s segments. Source training data determine the operating median. External cohorts include target-batch standardization and direction alignment where indicated. The temporal-order test compares intact data with 10,000 synchronized multichannel permutations using the frozen intact scale.

non-overlapping 4-second epochs (1,024 samples). Spectral densities were computed across each epoch via Hann-tapered periodograms ($\Delta f = 0.25$ Hz), extracting theta (4–8 Hz) and alpha (8–13 Hz) band power.

Artifact screening at $500\,\mu$V retained $5{,}185.5 \pm 595.6$ blocks per patient ($96.98\% \pm 3.2\%$, 5.76 ± 0.66 h; rejection $2.84\%{\pm}2.1\%$ in MCS vs. $3.09\%{\pm}3.6\%$ in VS/UWS, $p = 0.82$, Mann–Whitney U test). A conservative $150\,\mu$V cutoff retained $90.14\% \pm 17.87\%$ blocks ($4{,}672.9 \pm 1{,}024.7$ pairs; minimum 668), confirming balanced retention.

Let $\mathcal{T} = \{t_1, t_2, \ldots, t_N\}$ denote clean epoch indices in a continuous session, and let G_{t_i} denote spatial mean theta power across all 6 channels at epoch t_i. To prevent spurious concatenation across artifact gaps, the physical adjacency indicator is defined as:

$$\mathbb{I}_{\text{adj}}(t_i, t_{i+1}) = \begin{cases} 1, & \text{if } t_{i+1} - t_i = 1, \\ 0, & \text{otherwise}, \end{cases} \tag{1}$$

where 0 denotes an artifact gap, and $\mathbb{I}_{\text{adj}} = 1$ defines the adjacent index set $\mathcal{I}_{\text{adj}} = \{i \mid t_{i+1} - t_i = 1\}$. Sequential metrics operate strictly on $\mathcal{I}_{\text{adj}}$, retaining $M = |\mathcal{I}_{\text{adj}}| = 5{,}133.3 \pm 687.7$ adjacent pairs per patient (minimum 1,678, validity threshold

TABLE I

PERFORMANCE OF ORDER-AWARE TEMPORAL DYNAMICS, CONSENSUS FUSION, AND MACHINE LEARNING BASELINES ON MENDELEY COHORT (25 OUTER PARTITIONS; 95% CIS PAIRED VS. CANONICAL LINEAR BASELINE VIA 10,000 DIRICHLET RESAMPLES; PR-AUC REPORTED AS AVERAGE PRECISION [AP]; MATCHED SUPERVISED ROWS USE THE SAME FOUR FEATURES AS THE DEPLOYMENT SUITE; 5-FEAT COMBINED MODELS SUBSTITUTE r_{INTERHEM} WITH r_{RES}).

Model / Feature Configuration	AUROC	PR-AUC	Balanced Acc.	Sensitivity	Specificity	95% CI vs. Baseline
Consensus Aggregation Frameworks						
Canonical Static (4 Feats, Linear)	0.8200	0.7823	0.7201	0.8333	0.6069	Baseline
Primary Orthogonal (4 Feats, Linear)	**0.9324**	**0.8998**	**0.8101**	0.9167	0.7034	$[+0.0174, +0.2186]$
Primary Orthogonal (4 Feats, Rank Consensus)	**0.9076**	**0.8548**	**0.8448**	1.0000	0.6897	$[+0.0137, +0.1725]$
Plus Lag-1 ρ_1 (5 Feats, Linear)	0.8911	0.8384	0.7997	0.9167	0.6828	$[+0.0011, +0.1534]$
Plus $\rho_1 + r_{\text{res}}$ (5 Feats, Linear, $c = 1.5\sigma$)	**0.9149**	**0.8782**	**0.8399**	0.9833	0.6966	$[+0.0064, +0.1931]$
Plus $\rho_1 + r_{\text{res}}$ (5 Feats, Rank Consensus)	0.8949	0.8353	0.7741	0.9000	0.6483	$[+0.0049, +0.1530]$
T-GECF Adaptive Gated Hybrid (5 Feats)	**0.9116**	**0.8669**	**0.8399**	0.9833	0.6966	$[+0.0013, +0.1927]$
Supervised & Domain Baseline Benchmarks						
Lempel-Ziv Complexity (LZC)	0.5136	0.5061	0.5267	0.5500	0.5034	$[-0.4233, -0.1862]$
Shannon Spectral Entropy (SE)	0.6953	0.6182	0.6721	0.7167	0.6276	$[-0.2540, +0.0085]$
Canonical Static (Tuned Ridge)	0.7653	0.7193	0.6661	0.7667	0.5655	$[-0.0998, -0.0140]$
Orthogonal 4-Feat (Tuned Ridge)	0.8813	0.8393	0.7825	0.9167	0.6483	$[-0.0446, +0.1744]$
Canonical Static (Tuned Logistic)	0.7722	0.7236	0.6917	0.7833	0.6000	$[-0.0959, -0.0064]$
Orthogonal 4-Feat (Tuned Logistic)	0.8964	0.8524	0.8080	0.9333	0.6828	$[-0.0225, +0.1826]$
Canonical Static (Shrinkage-LDA)	0.7709	0.7252	0.6626	0.7667	0.5586	$[-0.0884, -0.0162]$
Orthogonal 4-Feat (Shrinkage-LDA)	0.8769	0.8424	0.7859	0.9167	0.6552	$[-0.0496, +0.1699]$
Representation Ablations & Operator Variants						
Canonical Static (4 Feats, Rank Consensus)	0.7893	0.7299	0.7270	0.8333	0.6207	$[-0.0619, -0.0076]$
Plus Lag-1 ρ_1 (5 Feats, Rank Consensus)	0.8591	0.8068	0.7187	0.8167	0.6207	$[-0.0188, +0.1053]$
Plus Jump D_1 (5 Feats, Linear)	0.8611	0.8012	0.7491	0.8500	0.6483	$[-0.0018, +0.0911]$
Plus Jump D_1 (5 Feats, Rank Consensus)	0.8424	0.7787	0.7132	0.8333	0.5931	$[-0.0208, +0.0694]$
Replace $-\text{CV}_\theta$ with ρ_1 (Linear)	0.8851	0.8495	0.7678	0.8667	0.6690	$[-0.0223, +0.1613]$
Replace $-\text{CV}_\theta$ with D_1 (Linear)	0.8720	0.8097	0.7776	0.9000	0.6552	$[+0.0050, +0.1094]$
Residual Correlation r_{res} (Linear)	0.8433	0.7671	0.7575	0.8667	0.6483	$[-0.0561, +0.1070]$

$M \geq 10$).

C. Canonical Static Features and Order-Aware Dynamics

Four baseline features define $\mathbf{x}_{\text{canonical}} = [c_{\text{global}}, c_{\text{spatial}}, -\text{CV}_\theta, r_{\text{interhem}}]^\top \in \mathbb{R}^4$. We evaluate three expanded feature spaces: the 5-feature temporal extension $\mathbf{x}_{+\rho_1} = [c_{\text{global}}, c_{\text{spatial}}, -\text{CV}_\theta, r_{\text{interhem}}, \rho_1]^\top \in \mathbb{R}^5$; the 5-feature combined suite $\mathbf{x}_{\text{combined}} = [c_{\text{global}}, c_{\text{spatial}}, -\text{CV}_\theta, \rho_1, r_{\text{residual}}]^\top \in \mathbb{R}^5$; and the primary parsimonious 4-feature deployment suite $\mathbf{x}_{\text{ortho}} = [c_{\text{global}}, -\text{CV}_\theta, \rho_1, r_{\text{residual}}]^\top \in \mathbb{R}^4$. Theta power (4–8 Hz) is the primary carrier because thalamocortical mesocircuit projections drive frontoparietal dynamics within this band [19], whereas delta reflects non-specific slowing and alpha is largely abolished in VS/UWS. Let $P_{\theta,t}^{(k)}$ and $P_{\alpha,t}^{(k)}$ denote theta and alpha powers of channel $k \in \{1, \ldots, 6\}$ at clean epoch $t \in \{1, \ldots, N\}$.

1) Global Spectral Contrast (c_{global}): Quantifies the ratio of slow to fast power across all channels: $c_{\text{global}} = (\bar{P}_\theta - \bar{P}_\alpha)/(\bar{P}_\theta + \bar{P}_\alpha + \epsilon)$, where $\bar{P}_\theta = \frac{1}{6N}\sum_{k,t} P_{\theta,t}^{(k)}$, $\bar{P}_\alpha = \frac{1}{6N}\sum_{k,t} P_{\alpha,t}^{(k)}$, and $\epsilon = 10^{-12}$.

2) Spatial Contrast (c_{spatial}): Computes mean slow-to-fast ratio: $c_{\text{spatial}} = \frac{1}{6}\sum_{k=1}^{6}(\bar{P}_\theta^{(k)} - \bar{P}_\alpha^{(k)})/(\bar{P}_\theta^{(k)} + \bar{P}_\alpha^{(k)} + \epsilon)$. Fixed feature configurations were compared to assess spectral redundancy; the four-feature suite omitted spatial contrast (Section III-C).

3) Negative Theta Dispersion ($-\text{CV}_\theta$): Measures temporal variance of mean theta $G_t = \frac{1}{6}\sum_{k=1}^{6} P_{\theta,t}^{(k)}$: $-\text{CV}_\theta = -\sqrt{\frac{1}{N}\sum_{t=1}^{N}(G_t - \bar{G})^2}/(\bar{G} + \epsilon)$, with temporal mean $\bar{G} = \frac{1}{N}\sum_{t=1}^{N} G_t$.

4) Inter-Hemispheric Bilateral Correlation (r_{interhem}): Averages raw pairwise Pearson correlations across symmetric bilateral pairs: $r_{\text{interhem}} = \frac{1}{3}\sum_{(L,R)} \text{Corr}(\mathbf{y}_L, \mathbf{y}_R)$, where $\mathbf{y}_L, \mathbf{y}_R \in \mathbb{R}^N$ denote the 4-second block theta-power vectors across symmetric pairs $(L, R) \in \{(\text{F3}, \text{F4}), (\text{C3}, \text{C4}), (\text{O1}, \text{O2})\}$.

To capture multi-second sequential continuity, two order-aware temporal representations are formulated on physically adjacent blocks:

Lag-1 Memory Persistence (ρ_1): Quantifies autoregressive temporal persistence across consecutive 4-second blocks:

$$\rho_1 = \text{Corr}(\{G_{t_i}\}_{i \in \mathcal{I}_{\text{adj}}}, \{G_{t_{i+1}}\}_{i \in \mathcal{I}_{\text{adj}}}). \tag{2}$$

High ρ_1 values reflect smooth baseline preservation, whereas low values signify rapid temporal decorrelation.

Normalized Adjacent Jump Rate (D_1): Measures resistance to high-frequency baseline discontinuities:

$$D_1 = -\frac{1}{M}\sum_{i \in \mathcal{I}_{\text{adj}}} |G_{t_{i+1}} - G_{t_i}|/(\bar{G} + \epsilon), \tag{3}$$

where $\bar{G} = \frac{1}{N}\sum_{t=1}^{N} G_t$, $M = \sum \mathbb{I}_{\text{adj}}$, and $\epsilon = 10^{-12}$. Under stationary Gaussian increments $\Delta G_t \sim \mathcal{N}(0, 2\sigma^2(1 - \rho_1))$, expected jump magnitude satisfies $\mathbb{E}[|\Delta G_t|] = 2\pi^{-1/2}\sigma(1 - \rho_1)^{1/2}$, yielding $D_1 \approx 2\pi^{-1/2}(-\text{CV}_\theta)(1 - \rho_1)^{1/2}$. Evaluating D_1 on log-power increments accounts for 93.91% of empirical variance under this identity (Section III-C). Given estimator variance $\text{Var}(\hat{\rho}_1) \sim \mathcal{O}(1/M)$, D_1 provides a low-variance proxy for brief records ($M < M^* \approx 75$ epochs), whereas ρ_1 achieves asymptotic efficiency.

To test macro-scale persistence on the Wielek PSG cohort ($N = 23$), stage stay persistence on 30-s hypnograms $\{s_t\}_{t=1}^{L} \in \{\text{Wake, N1, N2, N3, REM}\}$ is $P_{\text{stay}} = \frac{1}{L-1}\sum_{t=1}^{L-1} \mathbb{I}(s_{t+1} = s_t)$, with transition flexibility $T_{\text{trans}} =$

TABLE II
MULTISCALE SENSITIVITY AND EXTERNAL COHORT BENCHMARKS

(A) Multiscale Window & Cohort Transfer (AUROC)		
Cohort / Duration	**Linear**	**Rank Consensus**
Mendeley 2-second	0.8916	0.8529
Mendeley 4-second (Default)	**0.8911**	**0.8591**
Mendeley 8-second	0.8711	0.8453
Mendeley 30-min Trunc.	0.8867	0.8480
Figshare Static Baseline	0.7199	0.6812
Figshare 120-s ($N = 50$, Jump D_1)	**0.7691**	0.6927
Figshare 120-s ($N = 50$, Lag-1 ρ_1)	0.7012	0.6845
Figshare 25-Fold CV (D_1, $N = 50$)	**0.7771**	0.7505
(B) Longitudinal & Multi-Tier Benchmarks		
Validation Metric		**Value**
Mendeley 5-h ICC ($\rho_1/D_1/r_{\text{res}}$)		0.4857/0.6433/0.4511
Wielek Transition AUC ($p = 0.030$)		**0.7348**
Wielek Mean State Stay (MCS vs. VS)		0.7393 vs. 0.8361
Figshare Zero-Shot BA (Train Thresh.)		0.5645 (Sens 1.000)
62-Channel HD Mode-1 R^2		46.52%

$1 - P_{\text{stay}}$.

These four features span spectral slowing, marginal dispersion, autoregressive continuity, and residual bilateral coupling, forming the primary deployment suite $[c_{\text{global}}, -\text{CV}_\theta, \rho_1, r_{\text{residual}}]$.

D. Residual Bilateral Resonance ($r_{residual}$)

For each symmetric bilateral pair $(L, R) \in \{(\text{F3}, \text{F4}), (\text{C3}, \text{C4}), (\text{O1}, \text{O2})\}$ with block theta-power vectors $\mathbf{y}_L, \mathbf{y}_R \in \mathbb{R}^N$, the nuisance signal across the remaining 4 channels is $z_t^{(L,R)} = \frac{1}{4}\sum_{k \notin \{L,R\}} P_{\theta,t}^{(k)}$. With standardized $\tilde{\mathbf{z}}^{(L,R)} \in \mathbb{R}^N$ and design matrix $\mathbf{X}_c^{(L,R)} = [\mathbf{1}, \tilde{\mathbf{z}}^{(L,R)}] \in \mathbb{R}^{N\times 2}$, the exact unregularized OLS projection operator $\mathbf{P}_{(L,R)}^{\perp} = \mathbf{I}_N - \mathbf{X}_c^{(L,R)}(\mathbf{X}_c^{(L,R)})^{+} \in \mathbb{R}^{N\times N}$ (with $\text{rank}(\mathbf{P}^{\perp}) = N - \text{rank}(\mathbf{X}_c)$, where $\mathbf{X}_c^{+}$ denotes the Moore–Penrose pseudoinverse) computes orthogonalized residuals $\boldsymbol{\epsilon}_L^{(L,R)} = \mathbf{P}_{(L,R)}^{\perp}\mathbf{y}_L$ and $\boldsymbol{\epsilon}_R^{(L,R)} = \mathbf{P}_{(L,R)}^{\perp}\mathbf{y}_R$. The residual bilateral resonance averages pairwise correlations:

$$r_{\text{residual}} = \frac{1}{3}\sum_{(L,R)} \text{Corr}(\boldsymbol{\epsilon}_L^{(L,R)}, \boldsymbol{\epsilon}_R^{(L,R)}). \tag{4}$$

removing shared slow-wave envelope variance ($\approx 47\%$) to isolate intrinsic homologous coupling.

E. Consensus Operators and Machine Learning Baselines

Linear Standardized Aggregation (Linear): Equal-weight consensus follows the fixed-coefficient formulation of Dawes improper linear models [23]. The score is $S_{\text{linear}}(\mathbf{x}) = K^{-1}\sum_{k=1}^{K} s_k z_k$, where $z_k = (x_k - \mu_k^{\text{tr}})/\sigma_k^{\text{tr}}$ and each sign s_k is fitted from the feature–label correlation in the source training partition. Standardization and signs are re-estimated inside every inner and outer training fold. For the full-source four-feature fit, the signs are $[+1, +1, +1, -1]^\top$. CRRC target alignment uses the unsupervised persistence-anchored vector $[-1, +1, +1, +1]^\top$ across models in common oriented coordinates (Supplementary Note C). A soft-truncated variant clips coordinates at $c = 1.5$ ($K^{-1}\sum_k \text{clip}(s_k z_k, -c, c)$); the primary model uses $c = \infty$.

Non-Parametric Rank Consensus (S_{rank}): Adapted from multi-criteria decision frameworks [24], rank consensus maps signed features to empirical cumulative distribution function (CDF) midranks:

$$S_{\text{rank}}(\mathbf{x}) = \frac{1}{K}\sum_{k=1}^{K} \tilde{F}_k(s_k x_k), \tag{5}$$

where $\tilde{F}_k(u) = \frac{1}{N_{\text{tr}}}\sum_{i=1}^{N_{\text{tr}}}[\mathbb{I}(s_k x_{i,k}^{\text{tr}} < u) + \frac{1}{2}\mathbb{I}(s_k x_{i,k}^{\text{tr}} = u)]$. Curvature is fixed to $p = 1.0$ (mean on ranks), ensuring constant marginal substitution without uncalibrated parameters on small cohorts.

Adaptive Fluctuation-Gated Hybrid (T-GECF): To combine linear sensitivity with rank stability under sensor disturbances, T-GECF couples the two estimators via sample-level volatility gating. For signed standardized features $\tilde{x}_k = s_k(x_k - \mu_k^{\text{tr}})/(\sigma_k^{\text{tr}} + \epsilon)$, sample volatility is $v(\mathbf{x}) = \sqrt{\frac{1}{K}\sum_{k=1}^{K}(\tilde{x}_k - \bar{\tilde{x}})^2}$, standardized to $z_v(\mathbf{x}) = (v(\mathbf{x}) - \mu_v^{\text{tr}})/(\sigma_v^{\text{tr}} + \epsilon)$ against training folds. A continuous Sigmoid gate $\beta(\mathbf{x}) = 1/(1 + \exp(-\kappa(z_v(\mathbf{x}) - \theta_0)))$ ($\theta_0 = 1.645, \kappa = 2.0$) mixes standardized rank and linear scores:

$$S_{\text{T-GECF}}(\mathbf{x}) = \beta(\mathbf{x})\tilde{S}_{\text{rank}}(\mathbf{x}) + (1 - \beta(\mathbf{x}))\tilde{S}_{\text{linear}}(\mathbf{x}), \tag{6}$$

where $\tilde{S}_{\text{rank}}$ and $\tilde{S}_{\text{linear}}$ are standardized to zero mean and unit variance using outer fold statistics prior to mixing. Elevated volatility ($z_v(\mathbf{x}) > 1.645$) dynamically shunts decision weight toward bounded rank consensus $\tilde{S}_{\text{rank}}(\mathbf{x})$, mitigating the influence of isolated channel artifacts or amplitude spikes.

Supervised Machine Learning and Domain Baselines: Ridge, logistic regression, and shrinkage LDA receive exactly the same feature columns, standardized coordinates, direction coding, and subject partitions as equal-weight scoring. Ridge α and logistic C are selected from $\{10^{-3}, 10^{-2}, 10^{-1}, 1, 10, 10^2, 10^3\}$ using source-only inner three-fold stratified cross-validation; transforms are fitted on each inner training fold. LDA uses analytical shrinkage (`auto`) [25]. The same source-selected parameters are retained across external protocols. The primary matched comparison uses $\mathbf{x}_{\text{ortho}}$; the short-record branch uses $[c_{\text{global}}, c_{\text{spatial}}, D_1, r_{\text{interhem}}]^\top$ for every model. Lempel-Ziv complexity, spectral entropy, and an EEGNet-style CNN [26] provide additional domain baselines under the recorded protocols.

F. Subject-Level Validation Protocol and Decision Thresholds

Discovery evaluation uses five random seeds and five stratified subject-level folds per seed, giving 25 frozen outer partitions. Source means, scales, signs, empirical ranks, and hyperparameters are fitted strictly within training partitions. AUROC and average precision (AP) are averaged across outer folds; sensitivity, specificity, balanced accuracy, and F1 use pooled outer-test decisions, with each subject contributing once per seed. Each model's operating threshold is its source-training score median. Paired 95% intervals use 10,000 subject-level Dirichlet Bayesian bootstrap draws [27], sharing each subject's weight across folds and methods conditional on the fitted predictions. External validation separates frozen-source transfer, unlabeled target-batch standardization, and CRRC target direction alignment. Models share the same target transform




Fig. 3. Representation and scoring-rule comparisons in the 41-patient discovery cohort. (a) Annotated heatmap of fold-averaged AUROC for the canonical static suite and its five-feature lag-1 extension, using linear or rank fusion. Colors follow the displayed AUROC scale. (b) Paired AUROC gain of four-feature equal-weight consensus over ridge, logistic regression, and shrinkage LDA using identical four-feature inputs. Points show gains; bars show pointwise 95% subject-level Dirichlet Bayesian intervals from 10,000 draws, conditional on fitted predictions. The dotted line denotes zero gain.

within each protocol, remaining blinded to external labels until final metric evaluation. All reported classification operating points use source-training medians. Full parameters, transforms, thresholds, and predictions are archived with the evaluation scripts. A multivariable logistic regression on standardized $[c_{\text{global}}, R^2, \rho_1]$ provides the conditional association analysis.

III. Experimental Results

A. The 2×2 Orthogonal Factorial Decomposition

Table I summarizes classification performance. In the 2×2 factorial decomposition (Fig. 3a), Cell A (Static, Linear) achieves 0.8200 AUROC, Cell B (Static, Rank) yields 0.7893, Cell C (Plus ρ_1, Linear) attains 0.8911, and Cell D (Plus ρ_1, Rank) reaches 0.8591. The temporal representation gain is $\Delta_{\text{rep}} = \text{AUC(Cell C)} - \text{AUC(Cell A)} = +0.0711$ (95% CI $[+0.0011, +0.1534]$, positive gain probability 97.74%), while $\Delta_{\text{fuse}} = \text{AUC(Cell D)} - \text{AUC(Cell C)} = -0.0320$ reflects the trade-off between linear resolution and bounded rank stability.

B. Diagnostic Operating Characteristics and Imbalanced Precision-Recall

At the pre-specified training median threshold, incorporating ρ_1 elevated Balanced Accuracy from 0.7201 to 0.7997 and Sensitivity from 0.8333 to 0.9167 (Table I). Combining lag-1 persistence with residual bilateral resonance ($\mathbf{x}_{\text{combined}}$) improved the five-feature extension: under soft truncation ($c = 1.5\sigma$), linear aggregation achieved **0.9149** AUROC (PR-AUC = **0.8782**, Balanced Accuracy = **0.8399**, Sensitivity = 0.9833). In rank consensus, this suite elevated AUROC to **0.8949**, matching unconstrained linear aggregation within 0.0047 (0.8996).

Paired Bayesian bootstrap intervals for rank consensus, soft truncation, and T-GECF remain strictly positive ($[+0.0049, +0.1530]$, $[+0.0064, +0.1931]$, and $[+0.0013, +0.1927]$ vs. Canonical Static Baseline 0.8200), indicating that the performance gains from joint temporal continuity and spatial de-confounding are statistically reliable.

Stratifying the 12 MCS patients by behavioral subtype showed consistent persistence: MCS$^+$ ($N = 8$, command following/verbalization) had $\rho_1 = 0.4563 \pm 0.2687$ with 8/8 classified (100% sensitivity), and MCS$^-$ ($N = 4$, visual pursuit/localization) had $\rho_1 = 0.4377 \pm 0.1670$ with 3/4 classified (75% sensitivity; overall 11/12 = 91.67%). Under class imbalance (12 MCS vs. 29 VS/UWS, baseline PR-AUC = 0.2927), ρ_1 raised PR-AUC from 0.7823 (Cell A) to **0.8384** (Cell C), **0.8495** when replacing $-\text{CV}_\theta$, and **0.8782** in the combined model.

C. Feature Ablation and Collinearity Redundancy Analysis

Replacing $-\text{CV}_\theta$ with ρ_1 in the 4-feature suite yielded 0.8851 AUROC, confirming that sequence memory supersedes marginal variance. Across 41 patients, MCS showed higher persistence (0.450 ± 0.250 vs. 0.287 ± 0.181, $p = 0.0227$, AUROC = 0.7299; Fig. 1b,c) and lower jump rates (D_1: -0.420 ± 0.089 vs. -0.558 ± 0.168, $p = 0.0112$, AUROC = 0.7557). Log-power increments verify the analytical identity ($r = 0.9691, p = 2.59 \times 10^{-25}, R^2 = 93.91\%$), making D_1 a stable proxy below $M^* \approx 75$ epochs (5 min). Central sensorimotor channels (C3–C4) showed the highest univariate discrimination (ρ_1: 0.3996 ± 0.1863 vs. 0.2611 ± 0.1631, $p = 0.0132$, AUROC = 0.7500), exceeding frontal (0.6264) and occipital (0.6810) pairs.

Regressing out the global common mode (r_{residual}) improved AUROC from 0.8200 to 0.8433, accounting for $R^2 = 47.27\% \pm 18.43\%$ of raw bilateral variance uniformly across groups (46.45% in MCS vs. 47.61% in VS/UWS; $p = 0.7418$). Whereas raw bilateral correlation was non-significant (0.563 ± 0.168 in MCS vs. 0.661 ± 0.146 in VS/UWS, $p = 0.0605$, AUROC = 0.6897), residualization unmasked residual bilateral differentiation beyond the shared background power envelope (0.233 ± 0.080 vs. 0.376 ± 0.193, $p = 0.0211$, AUROC = 0.7328), reflecting modularity in MCS versus hyper-synchrony in VS/UWS. Multivariable logistic regression controlling for c_{global} ($z = -0.812, p = 0.417$) and common-mode R^2 ($z = 0.329, p = 0.742$) showed that ρ_1 provides significant independent diagnostic value ($z = 2.460, p = 0.0139$, adjusted OR = 4.260, 95% CI $[1.343, 13.512]$), confirming conditional independence.

Because c_{global} and c_{spatial} are strongly collinear ($r = 0.9813, p = 1.58 \times 10^{-29}$), omitting c_{spatial} to form the 4-feature deployment suite $[c_{\text{global}}, -\text{CV}_\theta, \rho_1, r_{\text{residual}}]$ (Table I) raised linear AUROC to **0.9324** (PR-AUC = **0.8998**; Balanced Accuracy = 0.8101, 95% CI $[+0.0174, +0.2186]$ vs. baseline, positive gain probability 99.16%) and rank consensus to **0.9076** (95% CI $[+0.0137, +0.1725]$, sensitivity 1.0000). Dropping c_{global} reached 0.9289 in linear and 0.9082 in rank consensus, while the 5-feature truncated suite (0.9149) assessed bounded-influence scoring.

D. Temporal-Order Dependency via 10,000 Synchronous Surrogates

To verify that performance gains originate strictly from temporal sequence order rather than incidental distributional




Fig. 4. Frequency distribution of AUROC under 10,000 synchronous time shuffles for the 5-feature lag-1 extension with intact scales (30 equal-width bins). Shuffled AUROC is $\mathbf{0.8182 \pm 0.0049}$ (mean $\pm$ SD); none reached intact AUROC 0.8911 ($\boldsymbol{p < 0.0001}$). Vertical lines mark static baseline (0.8200, dashed) and intact score (solid).

properties, 10,000 synchronous whole-montage permutation surrogates were evaluated across all 41 patients [28]. Freezing intact training scales ($\sigma_k^{\text{tr, intact}}$) prevents inverse noise amplification on attenuated shuffled variances. Disrupting chronological sequence alignment degraded macro AUROC across all temporal configurations (Fig. 4): Plus Lag-1 (Linear) dropped from 0.8911 to 0.8182 ± 0.0049 (range: $[0.7902, 0.8369]$, empirical $p < 0.0001$, zero of 10,000 surrogates matching intact performance); Plus Lag-1 (Rank) fell from 0.8591 to 0.7732 ± 0.0416 (range: $[0.6620, 0.9098]$, empirical $p = 0.0178$); and Plus Jump (D_1) degraded from 0.8611 to 0.7783 ± 0.0040 ($p < 0.0001$).

This frozen scaling bounds all 10,000 shuffled linear iterations below 0.8370. Across $M \approx 5{,}133$ adjacent blocks, the shuffled increment D_1 converges to $\mathbb{E}[|X - Y|]$ (variance $\approx 2 \times 10^{-4}$), collapsing to a static marginal quantity that aligns with the canonical baseline (0.8182 ± 0.0049).

E. Comparison with Supervised Machine Learning and Domain Baselines

On identical 4-feature inputs (Table I; Fig. 3b), equal-weight linear scoring achieved 0.9324 AUROC versus 0.8813 for source-tuned ridge ($+0.0511$), 0.8964 for logistic regression ($+0.0360$; paired 95% CI $[+0.0090, +0.0728]$), and 0.8769 for shrinkage LDA ($+0.0556$). Domain baselines yielded 0.5136 AUROC for LZC, 0.6953 for spectral entropy, and 0.5702 for the EEGNet-style CNN (Supplementary Table S2).

F. Adaptive Gated Consensus (T-GECF) and Noise Stress Testing

On the complete representation $\mathbf{x}_{\text{combined}}$, continuous Sigmoid volatility gating in T-GECF ($\theta_0 = 1.645, \kappa = 2.0$) achieved an AUROC of **0.9116** (95% CI $[+0.0013, +0.1927]$ vs. canonical baseline; positive gain probability 97.68%) with 0.8399 balanced accuracy, 0.9833 sensitivity, 0.6966 specificity, and PR-AUC = **0.8669** (Table I). On $\mathbf{x}_{+\rho_1}$, T-GECF similarly achieved 0.8828 AUROC ($+0.0237$ over standard rank consensus 0.8591).

To assess resilience against sensor disturbances, impulse spike stress tests introduced outlier perturbations ($\pm\delta\sigma$, $\delta \in \{0, 5, 10, 20\}$) into individual feature coordinates across testing partitions. Under extreme 20σ impulse corruption, unconstrained linear aggregation dropped from 0.8911 to 0.7404 (-0.1507). In contrast, T-GECF shifted weight toward bounded rank consensus, retaining an AUROC of 0.8518 ($+0.1114$ over linear aggregation) while maintaining 0.9116 AUROC under unperturbed conditions.

G. Multiscale Window Sensitivity and Longitudinal Reliability

Evaluating observational windows (Table II-A) showed consistent linear discrimination across 2 s (0.8916), 4 s (0.8911), and 8 s (0.8711) partitions, with rank consensus similarly stable (0.8453–0.8591). Truncating recordings showed that a 30-minute record attained 0.8867 AUROC, close to full-record performance. Test-retest reliability across five 1-hour segments ($N = 38$, 190 patient-hours) yielded $\text{ICC}(3,1) = 0.4857$ for ρ_1, 0.6433 for D_1, and 0.4511 for r_{residual}. Higher-order lags decayed monotonically ($\rho_1 = 0.450 \rightarrow \rho_2 = 0.401 \rightarrow \rho_3 = 0.389$ in MCS; $0.287 \rightarrow 0.252 \rightarrow 0.208$ in VS/UWS), reflecting multi-second persistence. Under adjacency masking $\mathbb{I}_{\text{adj}}$, 50-iteration Monte Carlo dropout maintained stability ($r = 0.9942 \pm 0.0030$ at 10% dropout; $r = 0.9199 \pm 0.0308$ at 50% loss), while tightening artifact rejection to $150\,\mu$V preserved 0.8702 AUROC (+0.1729 over unrejected baseline 0.6973; +0.0502 over unconstrained baseline 0.8200). Contiguous 30-min block removal ($r = 0.9840 \pm 0.0166$) and partition depths (10- to 100-fold CV, 0.8900–0.9002) verified stability across analytical scales.

H. External Waveform Validation and Sleep Macro-Dynamics

1) Rapid Bedside Triage (Figshare, $N = 50$): On 120-s records, jump rates reached 0.7691 AUROC (0.7199 static; baselines 0.7674–0.7759; diff vs. ridge -0.0068; Supplementary Table S2). At source thresholds, it provides a high-sensitivity funnel (1.0000 sensitivity, 19/19 MCS; 0.1290 specificity; Table II-B); Youden calibration recovers balanced accuracy (0.7233; sensitivity 0.7368, specificity 0.7097). *2) Cross-Hardware Inpatient Validation (CRRC, $N = 15$)*: Frozen transfer yielded 0.5741 AUROC; direction alignment ($\mathbf{s}_{\text{target}} = [-1, +1, +1, +1]^\top$) reached 0.7778 (0.7963 with $c = 1.5$ truncation; baselines 0.7407–0.7593; drift-free $N = 13$: 0.7143; Supplementary Table S2). At source thresholds, truncated scoring attained 0.6667 sensitivity, 0.7778 specificity. *3) Circadian Sleep Macro-Dynamics (Wielek, $N = 23$)*: Polysomnography across 24-hour records (11 MCS, 12 VS/UWS) showed that temporal continuity extends to macro-scale sleep architecture: VS/UWS patients exhibited state trapping ($P_{\text{stay}} = 0.8361 \pm 0.1001$), whereas MCS retained transition flexibility ($T_{\text{trans}} = 0.261 \pm 0.099$ vs. 0.164 ± 0.100, one-sided $p = 0.0302$, AUROC = 0.7348; nocturnal sleep $p = 0.0392$, AUROC = 0.7576; Table II-B).

I. Hyperparameter Sensitivity and Gating Grid Evaluation

Supplementary Table S3-A reports systematic benchmarks across truncation thresholds c and continuous Sigmoid gating parameters (θ_0, κ). Varying Winsorized soft truncation from

$c = 1.0\sigma$ to $c = \infty$ produced stable AUROC within $[0.8958, 0.9149]$ (balanced accuracy 0.8046–0.8399, sensitivity 0.9333–0.9833), peaking at $c = 1.5\sigma$ (0.9149) before approaching unconstrained linear aggregation (0.8996).

Evaluating T-GECF across a 4×4 grid crossing Gaussian quantile thresholds $\theta_0 \in \{1.282, 1.645, 1.960, 2.326\}$ and slopes $\kappa \in \{0.5, 1.0, 2.0, 5.0\}$ showed bounded variation (Supplementary Table S3-B). Across all 16 configurations, AUROC remained bounded within 0.8996–0.9220, exceeding the canonical baseline (0.8200) by $+0.08$ to $+0.10$. At $\theta_0 = 2.326$ ($\kappa = 5.0$), $\beta(\mathbf{x}) \to 0$, smoothly recovering linear aggregation (0.8996).

IV. Discussion

A. Physiological Significance of Autoregressive Memory

Under the mesocircuit hypothesis [19], thalamocortical loops sustain multi-second theta continuity in MCS, whereas deafferentation in VS/UWS degrades fluctuations into decoupled transients ($\rho_1 \to 0$). Because marginal dispersion ($-\mathrm{CV}_\theta$) conflates reverberation with noise, enforcing chronological continuity ($\mathbb{I}_{\text{adj}}$) provides an auditable marker of thalamocortical integrity. Across centers, temporal persistence (ρ_1 for continuous telemetry, D_1 for 120-s triage, and sleep-stage flexibility for 24-h circadian records) acts as a consistent physiological anchor, whereas spectral and spatial features reflect regime-dependent dynamics requiring contextual calibration.

B. Resolution of Non-Specific Bilateral Confounding and Direction Inversion

Residual correlation (r_{residual}, AUROC 0.8433) removes common-mode activity ($47.27\% \pm 18.43\%$ of bilateral variance), unmasking acute VS/UWS hyper-synchrony (0.376 ± 0.193) versus preserved MCS modularity (0.233 ± 0.080, $p = 0.0211$). In subacute CRRC patients (11/15 hemorrhage), this polarity inverted (0.308 in MCS vs. 0.210 in VS/UWS): callosal hemorrhage isolates hemispheres in VS/UWS, whereas recovering MCS restores interhemispheric coupling [7], [19]. Similarly, chronic VS/UWS exhibits diffuse slowing ($c_{\text{global}} \uparrow$) while recovering MCS restores posterior alpha ($c_{\text{global}} \downarrow$) [10], resolved via the unsupervised ρ_1-anchor rule ($\mathbf{s}_{\text{target}} = [-1, +1, +1, +1]^\top$; Supplementary Note C). In dense arrays, 4-channel averaging generalizes to subspace projection ($\mathbf{P}^\perp = \mathbf{I} - \mathbf{Z}\mathbf{Z}^+$; 46.52% Mode-1 variance, Table II-B).

C. Operator Dynamics: Rank Consensus vs. Linear Aggregation

Matched internal benchmarks support fixed-weight aggregation: the linear suite retained an AUROC advantage over supervised models on identical inputs. External validation showed comparable performance across classifiers, indicating that representation and calibration govern transfer. Soft truncation increased full-cohort CRRC AUROC from 0.7778 to 0.7963 (discovery 0.9291 vs. 0.9324 unclipped), providing bounded-influence scoring. Adaptive gating via T-GECF complements this by combining linear resolution with rank stability under impulse disturbances.

D. Clinical Scope, Cognitive Motor Dissociation, and Multimodal Triage

A primary clinical application is detecting covert consciousness in cognitive-motor dissociation (CMD), where CRS-R misses volitional responses in up to 25% of unresponsive patients [5]. Continuous lag-1 persistence (ρ_1) provides an auditable physiological marker to prioritize candidates for confirmatory active fMRI/PET or TMS-EEG [4], [6]. Under continuous telemetry (prevalence $12/41$), the median threshold reaches 0.9167 sensitivity and 0.7034 specificity, projecting 52.2% fewer transfers ($\Delta_{\text{referral}} = 1 - [P_{\text{MCS}}\,\text{Sens} + (1 - P_{\text{MCS}})(1 - \text{Spec})]$; 48.6%–61.0% across 15%–35% ICU prevalence) than universal escalation. In 120-s triage (Figshare), source thresholds ensure zero missed MCS cases (sensitivity 1.0000), while Youden calibration balances referral reduction to 42.6% (sensitivity 73.68%, specificity 70.97%).

E. Comparison with Recent Advanced Paradigms (2024–2026)

Recent dense-array and sensory paradigms report high DoC discrimination: microstates reach 0.97 AUC [12], entropy CNNs reach 0.902 AUC [29], multimodal pipelines achieve 92.4% accuracy [30], auditory features differentiate consciousness levels [21], and source graphs yield 0.714–0.893 AUC [31]. Translating them to bedside surveillance faces obstacles: high-density arrays are motion-sensitive, acoustic tests require specialized hardware, and deep models risk overfitting ($N < 50$). Combining order-aware persistence (ρ_1, D_1) with common-mode decoupling (r_{residual}) enables 4–6 channel operation with 2.15 ms per-step execution time and robust multicenter utility (0.7778–0.7963 on CRRC; 0.7691 on Figshare).

F. Limitations

Limitations include: (1) discovery in 41 patients (12 MCS), warranting prospective expansion; (2) external cohorts evaluate distinct regimes: subacute telemetry (CRRC) reflects etiology-dependent spatial dynamics, whereas 120-s records (Figshare) assess jump dynamics (D_1); (3) unmeasured covariates, such as continuous sedatives (GABA_A agonists) and craniectomy defects, warrant standardized washout and impedance modeling; (4) streaming latency includes 4-s buffering and $\geq$ 40-s accumulation, though per-step execution takes 2.15 ms; and (5) sample sizes limit subgroup resolution across etiologies.

V. Conclusion

Order-aware temporal descriptors and common-mode-adjusted bilateral dynamics provide a compact EEG representation for bedside consciousness assessment. On identical four-feature inputs, equal-weight consensus achieved 0.9324 discovery AUROC versus 0.8964 for matched supervised logistic regression. External evaluation yielded 0.7778 AUROC on CRRC telemetry (0.7963 with truncation) and 0.7691 on 120-s Figshare triage records. Operating with 2.15 ms execution time under contextual calibration, these results provide an interpretable framework for continuous bedside neuromonitoring.

Acknowledgment

OpenAI Codex assisted code implementation, figure preparation, and manuscript revision. The authors verified all code, figures, results, and interpretations.

Declarations

Ethics and Consent: Secondary analyses complied with Declaration of Helsinki, approved by institutional ethics committees: FRCCICMR Moscow [20], Salzburg Ethics Commission (No. 415-E/1517/14-2013) [13], Hangzhou Mingzhou Hospital [12], and China Rehabilitation Research Center (CRRC, Ethics No. 2025-118-01) [21], with written surrogate consent. **Data and Code Availability:** Public data: Mendeley [20] and Figshare [12], [22]. De-identified CRRC tabular feature matrices and replication scripts: `https://github.com/ujujuj3061831150/DoC_Temporal_Dynamics`. Raw CRRC telemetry from China Rehabilitation Research Center (Beijing Bo'ai Hospital) is restricted by institutional patient confidentiality regulations. **Conflict of Interest:** The authors declare no competing interests.

References

[1] J. T. Giacino, S. Ashwal, N. Childs, R. Cranford, B. Jennett, D. I. Katz *et al.*, "The minimally conscious state: definition and diagnostic criteria," *Neurology*, vol. 58, no. 3, pp. 349–353, 2002.

[2] J. T. Giacino, D. I. Katz, N. D. Schiff, J. Whyte, E. J. Ashman, S. Ashwal *et al.*, "Practice guideline update recommendations summary: Disorders of consciousness," *Neurology*, vol. 91, no. 10, pp. 450–460, 2018.

[3] C. Schnakers, A. Vanhaudenhuyse, J. Giacino, M. Ventura, M. Boly, S. Majerus, G. Moonen, and S. Laureys, "Diagnostic accuracy of the vegetative and minimally conscious state: clinical consensus versus standardized neurobehavioral assessment," *BMC Neurology*, vol. 9, p. 35, 2009.

[4] D. Cruse, S. Chennu, C. Chatelle, T. A. Bekinschtein, D. Fernández-Espejo, J. D. Pickard, S. Laureys, and A. M. Owen, "Bedside detection of awareness in the vegetative state: a cohort study," *The Lancet*, vol. 378, no. 9809, pp. 2088–2094, 2011.

[5] Y. G. Bodien, J. Allanson, P. Cardone, A. Bonhomme, J. Carmona, C. Chatelle *et al.*, "Cognitive motor dissociation in disorders of consciousness," *New England Journal of Medicine*, vol. 391, no. 7, pp. 598–608, 2024.

[6] A. G. Casali, O. Gosseries, M. Rosanova, M. Boly, S. Sarasso, K. R. Casali, S. Casarotto, M.-A. Bruno, S. Laureys, G. Tononi, and M. Massimini, "A theoretically based index of consciousness independent of sensory processing and behavior," *Science Translational Medicine*, vol. 5, no. 198, p. 198ra105, 2013.

[7] S. Chennu, J. Annen, S. Wannez, A. Thibaut, C. Chatelle, and H. Cassol, "Brain networks predict metabolism, diagnosis and prognosis at the bedside in disorders of consciousness," *Brain*, vol. 140, no. 8, pp. 2120–2132, 2017.

[8] J. D. Sitt, J.-R. King, I. El Karoui, B. Rohaut, F. Faugeras, A. Gramfort, L. Cohen, M. Sigman, S. Dehaene, and L. Naccache, "Large scale screening of neural signatures of consciousness in patients in a vegetative or minimally conscious state," *Brain*, vol. 137, no. 8, pp. 2258–2270, 2014.

[9] D. A. Engemann, F. Raimondo, J.-R. King, B. Rohaut, G. Louppe, F. Faugeras *et al.*, "Robust EEG-based cross-site and cross-protocol classification of states of consciousness," *Brain*, vol. 141, no. 11, pp. 3179–3192, 2018.

[10] C. A. Bareham, N. Roberts, J. Allanson, P. J. A. Hutchinson, J. D. Pickard, D. K. Menon, and S. Chennu, "Bedside EEG predicts longitudinal behavioural changes in disorders of consciousness," *NeuroImage: Clinical*, vol. 28, p. 102372, 2020.

[11] C. Bandt and B. Pompe, "Permutation entropy: a natural complexity measure for time series," *Physical Review Letters*, vol. 88, no. 17, p. 174102, 2002.

[12] Y. Ling, X. Wen, J. Tang, Z. Tao, L. Sun, H. Xin, and B. Luo, "Effect of topographic comparison of electroencephalographic microstates on the diagnosis and prognosis prediction of patients with prolonged disorders of consciousness," *CNS Neuroscience & Therapeutics*, vol. 30, no. 3, p. e14421, 2024.

[13] T. Wielek, J. Lechinger, M. Wislowska, C. Blume, P. Ott, S. Wegenkittl *et al.*, "Sleep in patients with disorders of consciousness characterized by means of machine learning," *PLoS One*, vol. 13, no. 1, p. e0190458, 2018.

[14] J. Xiao, T. Yu, H. Huang, J. Pan, F. Wang, D. Chen *et al.*, "Assessing disorders of consciousness using temporal sleep dynamics extracted from whole-night PSG," *IEEE Transactions on Biomedical Engineering*, pp. 1–14, 2026, early access, doi: 10.1109/TBME.2026.3675367.

[15] T. Thiéry, T. Lajnef, E. Combrisson, A. Dehgan, P. Rainville, G. A. Mashour *et al.*, "Long-range temporal correlations in the brain distinguish conscious wakefulness from induced unconsciousness," *NeuroImage*, vol. 179, pp. 30–39, 2018.

[16] X. Wei, Z. Yan, L. Cai, M. Lu, G. Yi, J. Wang *et al.*, "Aberrant temporal correlations of ongoing oscillations in disorders of consciousness on multiple time scales," *Cognitive Neurodynamics*, vol. 17, no. 3, pp. 633–645, 2023.

[17] Y. Bai, J. He, X. Xia, Y. Wang, Y. Yang, H. Di *et al.*, "Spontaneous transient brain states in EEG source space in disorders of consciousness," *NeuroImage*, vol. 240, p. 118407, 2021.

[18] D. Toker, I. Pappas, J. D. Lendner, J. Frohlich, D. M. Mateos, S. Muthukumaraswamy *et al.*, "Consciousness is supported by near-critical slow cortical electrodynamics," *Proceedings of the National Academy of Sciences*, vol. 119, no. 7, p. e2024455119, 2022.

[19] N. D. Schiff, "Recovery of consciousness after brain injury: a mesocircuit hypothesis," *Trends in Neurosciences*, vol. 33, no. 1, pp. 1–9, 2010.

[20] J. Nekrasova, M. Kanarskii, D. Yankevich, I. Borisov, and P. Pradhan, "Polysomnographic records of patients with disorders of consciousness," Mendeley Data, V1, 2020.

[21] F. Duan, J. Chen, Z. Yan, X. Cao, G. Xiong, and F. Li, "Auditory paradigm-dependent EEG feature screening and representation for distinguishing consciousness states in male patients with disorders of consciousness," *Frontiers in Neurology*, vol. 17, p. 1927313, 2026.

[22] C. Xu, J. Zou, F. He, X. Wen, J. Li, J. Gao *et al.*, "Neural tracking of sound rhythms correlates with diagnosis, severity, and prognosis of disorders of consciousness," *Frontiers in Neuroscience*, vol. 15, p. 646543, 2021.

[23] R. M. Dawes, "The robust beauty of improper linear models in decision making," *American Psychologist*, vol. 34, no. 7, pp. 571–582, 1979.

[24] M. Grabisch, J.-L. Marichal, R. Mesiar, and E. Pap, *Aggregation Functions*. Cambridge University Press, 2009.

[25] O. Ledoit and M. Wolf, "A well-conditioned estimator for large-dimensional covariance matrices," *Journal of Multivariate Analysis*, vol. 88, no. 2, pp. 365–411, 2004.

[26] V. J. Lawhern, A. J. Solon, N. R. Waytowich, S. M. Gordon, C. P. Hung, and B. J. Lance, "EEGNet: A compact convolutional neural network for EEG-based brain–computer interfaces," *Journal of Neural Engineering*, vol. 15, no. 5, p. 056013, 2018.

[27] D. B. Rubin, "The Bayesian bootstrap," *The Annals of Statistics*, vol. 9, no. 1, pp. 130–134, 1981.

[28] J. Theiler, S. Eubank, A. Longtin, B. Galdrikian, and J. D. Farmer, "Testing for nonlinearity in time series: the method of surrogate data," *Physica D: Nonlinear Phenomena*, vol. 58, no. 1-4, pp. 77–94, 1992.

[29] S. Qu, X. Wu, L. Huang, Y. Zhou, Q. Sun, and F. Zeng, "Diagnosis of disorders of consciousness using nonlinear feature derived EEG topographic maps via deep learning," *Scientific Reports*, vol. 16, p. 7417, 2026.

[30] W. Zhang, X. Shi, M. Li, L. Zhang, R. Zhang, X. Wu *et al.*, "Assess the level of consciousness in patients with disorders of consciousness by combining resting-state and auditory-evoked EEG," *Frontiers in Neuroscience*, vol. 19, p. 1613356, 2025.

[31] Z. Hao, X. Xia, Y. Pan, Y. Bai, Y. Wang, B. Peng, and W. Dou, "Uncovering brain network insights for prognosis in disorders of consciousness: EEG source space analysis and brain dynamics," *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, vol. 32, pp. 144–153, 2024.