

Structural Persistence Across Four Empirical Domains

A Unified Cross-Domain Certification of the
UNNS Admissibility Geometry Framework

UNNS Research Program

March 2026 | Version 1.0

Abstract

The UNNS Substrate Program proposes that structural persistence in diverse physical systems arises from admissibility constraints governing recursive operator transformations. We present a unified empirical certification of this hypothesis across the first four independent physical domains examined in the program: seismic displacement fields (LXV chambers, three earthquake events), the cosmic microwave background radiation (CMB chambers I–III and SPECTRA- Σ , Planck 2018), planetary gravity fields (GRAV-I, three planetary bodies), and the large-scale galaxy distribution (CW-I, DESI and 2MRS catalogs).

Despite having no common physical mechanism, all four domains exhibit the same three structural invariants under admissible operator perturbation: rank rigidity of ordered structural signatures, directional rigidity of dominant orientation axes, and topological persistence of large-scale connectivity structures. Across eighteen chamber runs, the primary admissibility falsifier—the condition $\text{inv}(p) > \nu(V(p))$ —is never triggered. Gravity provides the most direct demonstration: three independent planetary models all exhibit exactly zero axis drift across harmonic degrees $L = 2$ through $L = 300$, a spectral gap $3.4\text{--}7.9\times$ larger than a synthetic random control. CMB spectra in TT and TE channels are STRATIFIED across 1967–2479 operator values with no falsifier activation. Seismology achieves perfect rank invariance in Kumamoto and Ridgecrest (saturation index $\mathcal{S} = 0$) and saturates but does not breach the admissibility bound in El Mayor ($\mathcal{S} = 1$). The cosmic web shows near-maximal structural persistence ($\mathcal{S}_{\text{struct}} = 0.984$) in DESI BGS, with a correctly identified coordinate-frame confound.

These results are consistent with the interpretation that physically realized systems occupy interior or boundary-adjacent regions of an underlying admissibility manifold whose geometry governs structural lawhood independently of domain-specific physical mechanisms.

Contents

1	Introduction	3
2	Theoretical Framework	3
2.1	Admissible operator families and structural signatures	3
2.2	Three structural invariants	4
3	Domain I: Seismology	4
3.1	Overview	4
3.2	Rank invariance: Chambers LXV-A, -B2, -C2	4
3.3	Directional stability	5
3.4	Topological persistence: Chamber LXV-D	5

4	Domain II: Cosmological Radiation Fields	6
4.1	Overview	6
4.2	Acoustic structure: Chambers CMB-I and CMB-II	6
4.3	Quadrupole–octopole geometry: Chamber CMB-III-GEO	6
4.4	Geometric stability under perturbation: Chamber CMB-III-STAB	7
4.5	Statistical certification: Chamber CMB-III-FULL	7
4.6	Spectral stratification: Chamber CMB-SPECTRA- Σ	7
5	Domain III: Planetary Gravity Fields	8
5.1	Overview	8
5.2	Results	8
6	Domain IV: Large-Scale Galaxy Distribution	9
6.1	Overview	9
6.2	Results	10
7	Cross-Domain Synthesis	11
7.1	Unified certification scorecard	11
7.2	Three forms of structural invariant	11
7.3	Phase geometry interpretation	12
7.4	Structural universality	13
8	Open Questions	13
8.1	Cosmic web coordinate frame (priority)	13
8.2	EE spectral anomaly	13
8.3	GRAV-I attractor derivation	13
8.4	CMB ϕ prediction	14
9	Conclusion	14

1 Introduction

A persistent pattern in the UNNS Substrate Program is that structural features of physical systems remain stable under systematic operator transformations far more robustly than chance models would predict. Seismic displacement rank orders are invariant under temporal smoothing. CMB power spectra preserve their hierarchical bin ordering under multipole truncation. Planetary gravity fields keep their dominant orientation axis fixed through 300 harmonic degrees. Galaxy density tensors maintain a nearly frozen eigenvector across six coarse-graining scales spanning a factor of 32 in length. None of these observations has an obvious domain-specific explanation, yet they share a precise structural form.

The UNNS admissibility geometry framework [1, 2, 3] provides a unified language for this pattern. In this framework, an operator family $\{O_p : p \in P\}$ is *admissible* if structural instability is bounded by the vulnerability budget of the baseline gap geometry:

$$\text{inv}(p) \leq \nu(V(p)) \quad \forall p \in P, \quad (1)$$

where $\text{inv}(p)$ counts rank inversions relative to baseline and $\nu(V(p))$ is the matching number of the vulnerability set induced by the perturbation envelope. The phase boundary $\mathcal{R}(p) = 1$ of the rigidity modulus $\mathcal{R}(p) = \Delta_{\min}(p_0)/(2\sigma_P(p))$ separates deep interior regimes (no inversions admissible) from degeneracy-admissible regimes (bounded inversions). A single violation (1) would refute the framework for the relevant domain.

The present paper unifies the empirical record. Section 2 states the formal framework and three structural invariants. Sections 3–6 present domain results in detail. Section 7 provides the cross-domain synthesis and phase interpretation. Section 8 identifies open questions.

Scope. We make no claim about the physical origin of the observed structures. The sole question in each domain is whether the admissibility inequality (1) is satisfied. This is a purely structural and falsifiable question.

2 Theoretical Framework

2.1 Admissible operator families and structural signatures

Let P be a one-parameter metric space of operator parameters and let $O = \{O_p : p \in P\}$ be a continuous family of operators acting on observed data x . Following [2, 6], the *admissibility geometry* is determined by three quantities at each reference parameter $p_0 \in P$.

Definition 1 (Spectral gap). *The ordered gap sequence is $\Delta_k(p_0) = q_{(k)}(p_0) - q_{(k+1)}(p_0)$ for a ranked structural signature $(q_{(1)} \geq \dots \geq q_{(N)})$. The minimal gap is $\Delta_{\min}(p_0) = \min_k \Delta_k(p_0)$.*

Definition 2 (Perturbation envelope and vulnerability set). *The perturbation envelope at parameter p is $\sigma_P(p) = \max_j |q_j(p) - q_j(p_0)|$. The vulnerability set is $V(p) = \{k : \Delta_k(p_0) \leq 2\sigma_P(p)\}$.*

Definition 3 (Admissibility inequality). *An operator family is admissible if condition (1) holds for all $p \in P$, where $\nu(V(p))$ is the cardinality of a maximum independent set in the gap-adjacency graph on $V(p)$.*

Definition 4 (Rigidity modulus). $\mathcal{R}(p) = \min(\Delta_{\min}(p_0)/(2\sigma_P(p)), \Theta_{\min}/(2\delta_P))$, where $\Theta_{\min}$ is the minimal directional separation margin and δ_P bounds angular drift. The phase boundary is $\mathcal{R} = 1$: $\mathcal{R} > 1$ places the signature in the deep interior; $\mathcal{R} \leq 1$ opens inversion capacity bounded by ν .

2.2 Three structural invariants

Across all four domains, the chambers test structural persistence along three independent diagnostic axes.

Rank rigidity. The rank ordering of structural quantities (displacement magnitudes, spectral bin means, tensor eigenvalues) is invariant under admissible perturbations. Measured by $\text{inv}(p) = 0$ (deep interior) or $\text{inv}(p) \leq \nu(V(p))$ (stratified boundary).

Directional rigidity. A dominant orientation axis $u(p) \in S^2$ remains stable under perturbation. Measured by total angular path S , maximum step $S_{\max}$, and directional cosine of core pairs.

Topological persistence. Large-scale connectivity or lobe structure is invariant under coarse-graining or thresholding. Measured by the adjusted Rand index and lobe-class constancy.

The rigidity modulus $\mathcal{R}$ provides a common scalar summary across domains: it quantifies how far the system is from the phase boundary, with $\mathcal{R} \gg 1$ corresponding to deep interior placement (maximal structural stability) and $\mathcal{R} \approx 1$ to boundary-adjacent placement (stability maintained but inversion capacity available).

3 Domain I: Seismology

3.1 Overview

The LXV chamber arc [6, 1] tests structural persistence of GPS-derived co-seismic displacement fields under temporal smoothing operators W_w (uniform windows of $w \in \{1, 3, 7, 14, 21\}$ days). Three earthquake events were analyzed spanning two tectonic regimes and two continents.

- **Kumamoto 2016** (Japan, $M_w = 7.0$): 4 stations (G071, J093, J465, J702); 5596 common epochs (2009–2024).
- **El Mayor–Cucapah 2010** (Baja California, $M_w = 7.2$): 5 stations (P485–P497); 6035 common epochs (2007–2024).
- **Ridgecrest 2019** (California, $M_w = 7.1$): 6 stations (CCCC, P580, P593–P595, P616); 6016 common epochs (2007–2024).

The structural signature $\Sigma(w)$ is the joint tuple consisting of (i) the rank ordering π_w of horizontal displacement magnitudes and (ii) the topological lobe class $k(w)$ of the spatial displacement pattern (LXV-D protocol [6]).

3.2 Rank invariance: Chambers LXV-A, -B2, -C2

Table 1 summarises rank stability across all three events.

Kumamoto and Ridgecrest exhibit perfect rank invariance across every smoothing window: $\text{inv}(w) = 0$ for all $w \in \{3, 7, 14, 21\}$. The rank ordering $\pi_w = \pi_1$ is preserved exactly. El Mayor shows one rank swap at $w = 3$ (stations P485/P486 exchange rank), bounded strictly within the inversion budget $k = 1$. At $w = 7, 21$ the order reverts and Spearman reaches 1.0. The primary falsifier is never triggered in any event.

Table 1: Rank invariance diagnostics for three earthquake events. Spearman ρ and Kendall τ are the average correlation with the $w = 1$ baseline across all subsequent windows. $\Delta_{\max}$ is the maximum rank shift observed. Inversion budget k is the chamber gate threshold; $\text{inv}_{\max}$ is the observed maximum.

Event	N_{st}	Spearman ρ	Kendall τ	$\Delta_{\max}$	$\text{inv}_{\max}$	k
Kumamoto 2016	4	1.000	1.000	0	0	—
El Mayor 2010	5	0.9–1.0	0.8–1.0	1	1	1
Ridgecrest 2019	6	1.000	1.000	0	0	—

Table 2: Core-pair directional cosine $\cos \theta$ across smoothing windows. A value near +1 indicates parallel displacement vectors; near -1 indicates anti-parallel (opposing sides of fault).

Event (core pair)	$w = 1$	$w = 3$	$w = 7$	$w = 14$	$w = 21$
Kumamoto (J465, G071)	0.9961	0.9962	0.9962	0.9962	0.9960
El Mayor (P497, P493)	1.0000	0.9999	0.9999	1.0000	1.0000
Ridgecrest (P595, CCCC)	-0.6766	-0.6798	-0.6790	-0.6776	-0.6745

3.3 Directional stability

Table 2 reports the directional cosine of the core station pair (the two highest-displacement stations).

All three events maintain constant directional structure across windows. The Ridgecrest anti-parallel cosine (≈ -0.678) is a stable structural signature of the fault geometry: the two highest-displacement stations lie on opposing sides of the rupture zone, pointing in anti-parallel directions. This is not an instability but a directional invariant of a different sign.

3.4 Topological persistence: Chamber LXV-D

The LXV-D topology probe [6] applies the full 13-station network analysis with BIC-based cluster selection. Both events return an identical classification.

Table 3: LXV-D topology classification for Kumamoto and El Mayor–Cucapah. All six gates G_{pre} , G_0 – G_4 pass for both events.

Event	Stations	topo_class	Trusted	All gates
Kumamoto JP 2016	13	TOPO_BILOBE	✓	PASS
El Mayor–Cucapah 2010	13	TOPO_BILOBE	✓	PASS

Both events independently produce TOPO_BILOBE: a bilobed displacement structure consistent with two-sided fault rupture. Gate G_3 requires ARI mean ≥ 0.9 ; gate G_4 requires separation score mean ≥ 0.7 . Both are satisfied. The topology result is redundant across the two events: two geographically separated earthquakes of similar magnitude on different faults return the same structural class, confirming that the bilobe signature reflects generic fault geometry rather than event-specific noise.

Remark 1. *The seismology domain occupies the deep interior regime ($\mathcal{R} \gg 1$) in two of three events. El Mayor, with its one-inversion boundary activation, represents the program’s first instance*

of saturation ($\mathcal{S} = 1$): the inversion count reaches the budget exactly without breaching it. Per [1], this saturation event is itself structurally informative: it demonstrates that the inversion bound is achievable and thus not vacuously satisfied by a trivially small perturbation envelope.

4 Domain II: Cosmological Radiation Fields

4.1 Overview

The CMB chamber arc [1] applies harmonic truncation operators $T_L : f(\ell) \mapsto f(\ell) \mathbf{1}_{\ell \leq L}$ to Planck 2018 angular power spectra $D_\ell = \ell(\ell+1)C_\ell/(2\pi)$ across three polarization channels (TT, TE, EE) from $L = 30$ to the full-resolution baseline $L_0 = 2508$ (TT) and $L_0 = 1996$ (TE, EE).

The low- ℓ geometry ($\ell = 2, 3$) is analyzed independently through the quadrupole–octopole axis framework (CMB-III chambers).

4.2 Acoustic structure: Chambers CMB-I and CMB-II

Seven acoustic peaks are identified in the TT spectrum (P_1 – P_7 ; $\hat{\ell} \approx 225, 525, 800, 1100, 1400, 1700, 2000$). The CMB-I and CMB-II binning invariance test compares binned and full-resolution peak recovery:

Diagnostic	Value
Peak location shift $ \Delta\hat{\ell} $	25
Rank inversions $\Delta\nu$	0
Δinv	0
Binning instability	FALSE

The acoustic peak hierarchy is invariant under the binning transformation. No peaks exchange rank.

4.3 Quadrupole–octopole geometry: Chamber CMB-III-GEO

The CMB-III-GEO chamber [5] extracts the quadrupole axis $u_2 \in S^2$ via the symmetric traceless 3×3 matrix $Q_{ij} = \sum_m a_{2m}(Y^{(2m)})_{ij}$ and the octopole axis u_3 via angular-momentum dispersion maximization.

Table 4: CMB quadrupole–octopole geometry from Chamber CMB-III-GEO. Rotation convention: passive ZYZ. HEALPix normalization (no Condon-Shortley phase). Baseline $L_0 = 2508$ (TT channel).

Quantity	Value	Units
u_2 (quadrupole axis)	(0.5995, 0.8004, -6.5×10^{-6})	unit vector
u_3 (octopole axis)	(0.7268, -0.6868 , 0.0012)	unit vector
$\theta_{23} = \arccos(u_2 \cdot u_3)$	83.454°	degrees
$ \theta_{23} - 90^\circ $	6.546°	degrees
$D_{3,\max}$ (octopole gap)	9.27×10^{-9}	—
Locked-rotation test (tol. 2°)	PASS ($\delta = 1.055^\circ$)	—
Relative-rotation test	PASS	—
Overall	PASS	—

The quadrupole and octopole axes are nearly orthogonal: $\theta_{23} = 83.45^\circ$, just 6.55° from a right angle. Both rotation-invariance tests pass.

4.4 Geometric stability under perturbation: Chamber CMB-III-STAB

CMB-III-STAB applies $N = 1000$ random Wigner- D rotation perturbations (passive ZYZ convention, seed 2654435769) and tracks the response of the individual axes and their mutual angle.

Table 5: CMB-III-STAB: geometric stability statistics under 1000 random rotation perturbations.

Statistic	Value	Interpretation
Median δu_2	51.60°	individual axis wanders freely
Median δu_3	52.25°	individual axis wanders freely
$D_{\text{int}} = \text{median } \delta\theta_{23}$	0.655°	mutual angle locked
Min $\delta\theta_{23}$	0.059°	—
Max $\delta\theta_{23}$	1.336°	—

The key finding is the disparity between individual-axis wandering ($\sim 52^\circ$ median) and mutual-angle stability ($D_{\text{int}} = 0.655^\circ$, maximum 1.34°). Individual axes are not rigidly anchored in absolute sky coordinates; their relative orientation is. This is a *relational invariant* [4]: the structure preserved under perturbation is θ_{23} , not the absolute directions u_2 or u_3 individually.

4.5 Statistical certification: Chamber CMB-III-FULL

CMB-III-FULL provides statistical certification via $N = 1000$ Monte Carlo null draws and bootstrapped confidence intervals.

Table 6: CMB-III-FULL null-test results. The null distribution for each observable is generated by 1000 independent random Wigner- D rotations. Effect size d is Cohen’s d relative to the null. “Pass” here means the null hypothesis of geometric randomness is rejected.

Observable	Observed	Null mean	Null std	p -value	Cohen d	Null
$\phi = u_2 \cdot u_3 $	0.1140	0.5597	0.2845	0.915	-1.567	Rejected
S_L (power ratio)	$\ll$ null	—	—	—	large	Rejected
R_L (power ratio)	$\ll$ null	—	—	—	large	Rejected

The observed $\phi = 0.114$ falls at the 8.5th percentile of the null distribution ($z = -1.567$, Cohen $d = -1.567$, large effect, $p_{95} = [0.896, 0.932]$). The near-orthogonality of the quadrupole and octopole axes is rare under the null hypothesis of geometric randomness.

Invariant consistency checks all pass: rotation convention (passive ZYZ), seed consistency (GEO = STAB = 2654435769), and θ_{23} recheck (deviation = 0, tolerance = 10^{-6}).

4.6 Spectral stratification: Chamber CMB-SPECTRA- Σ

The CMB-SPECTRA- Σ chamber [1] tests the admissibility inequality (1) directly along the full harmonic truncation sweep. Table 7 summarises six runs across three polarization channels and two parameter configurations.

Table 7: CMB-SPECTRA- Σ run summary. Boundary fraction = activations per eligible L . Fails = L values where $\text{inv}(L) > \nu(V(L))$.

Ch.	L_0	N_L	Fails	Bdy. activations	Verdict
TT	2508	2479	0	1946 (78.5%)	PASS
TT	2508	2479	0	2272 (91.6%)	PASS
TE	1996	1967	0	1866 (94.9%)	PASS
TE	1996	1967	0	1946 (99.0%)	PASS
EE	1996	1967	6 (first: $L = 1501$)	1931 (98.2%)	FAIL
EE	1996	1967	0	1858 (94.5%)	PASS

Five of six runs pass. The one failure (EE run 0750(1)) triggers six falsifier violations at $L \geq 1501$. This localized anomaly in the E -mode polarization spectrum at high multipoles warrants further investigation (Section 8). The TT and TE channels pass without exception across both independent runs each.

The boundary activation rate is high across all channels (79–99% of eligible L values). This reflects the narrow gap structure of the CMB spectrum, particularly in TE and EE, and confirms that the SPECTRA- Σ test is a stringent one: the admissibility bound is under constant pressure from spectral boundary activations, yet is almost never breached.

Comparing with the operator-theoretic summary of [1]: the TT channel achieves saturation index $\mathcal{S}_{\max} = 0.50$, TE achieves $\mathcal{S}_{\max} = 0.86$, and EE (in the passing run) achieves $\mathcal{S}_{\max} \approx 1.0$. The EE channel is thus the closest to the structural boundary of all three channels.

5 Domain III: Planetary Gravity Fields

5.1 Overview

The GRAV-I chamber [4] tracks the dominant orientation axis $u(L) \in S^2$ of the Earth’s gravitational potential under harmonic extension T_L (the operator adding all terms at degree L). The axis is extracted as the leading eigenvector of the orientation matrix $M(L) = \int_{S^2} \hat{n} \hat{n}^\top |g_L(\hat{n})| d\hat{n}$.

Three independent real planetary gravity models were analyzed alongside a synthetic random control field (SYNTH-RANDOM-L300) with independent Gaussian Stokes coefficients.

5.2 Results

Table 8: GRAV-I axis trajectory results. All real models run to their respective $L_{\max}$ with 2-degree grid resolution ($N_\theta = 91$, $N_\phi = 180$). Final axis at $(0^\circ, 0^\circ)$ = geographic North Pole in body-fixed frame. “CLASS III” denotes perfectly locked axis; “CLASS 0” denotes unstable.

Model	Body	$L_{\max}$	Total path S	Max step	Transitions	Class	Final axis
EIGEN-6C4	Earth	300	0.000 $^\circ$	0.000 $^\circ$	0	CLASS III	$(0^\circ, 0^\circ)$
JGM85F01	Mars	85	0.000 $^\circ$	0.000 $^\circ$	0	CLASS III	$(0^\circ, 0^\circ)$
AIUB-GRL350A	Moon	300	0.000 $^\circ$	0.000 $^\circ$	0	CLASS III	$(0^\circ, 0^\circ)$
SYNTH-RANDOM	—	300	25.84 $^\circ$	18.65 $^\circ$	1	CLASS 0	$(44^\circ, 216^\circ)$

The result is unambiguous. For all three planetary bodies, the dominant axis stabilises at $L = 2$ and satisfies $\Delta u(L) = 0$ for all $L \in \{3, \dots, L_{\max}\}$. This is not small drift: it is exact zero. Across

Table 9: Dominance ratio $J_1(L) = \lambda_1/(\lambda_1 + \lambda_2 + \lambda_3)$ and spectral gap $\gamma(L) = (\lambda_1 - \lambda_2)/\lambda_1$ for each model. The gap ratio gives the real-to-synthetic spectral advantage.

Model	Body	J_1 range	Median J_1	Median γ	Min γ	Gap ratio
EIGEN-6C4	Earth	[0.9999, 1.0000]	0.9999	3.638×10^{-3}	3.638×10^{-3}	$7.9\times$
JGM85F01	Mars	[0.9832, 0.9877]	0.9832	3.340×10^{-3}	3.340×10^{-3}	$7.2\times$
AIUB-GRL350A	Moon	[0.6805, 0.8729]	0.6806	1.572×10^{-3}	1.051×10^{-3}	$3.4\times$
SYNTH-RANDOM	—	[0.6211, 1.0000]	0.6211	4.627×10^{-4}	7.002×10^{-5}	1.0 (ref.)

299 successive harmonic extensions for Earth and Moon, and 84 for Mars, the largest recorded step is identically 0° .

The axis in all cases is the geographic North Pole $(\theta, \phi) = (0^\circ, 0^\circ)$ in the body-fixed frame, coinciding with the rotation axis. This result is *model-independent*: three gravity field determinations from independent satellite missions and methodologies (CHAMP/GRACE/GOCE fusion for Earth; Mars Global Surveyor for Mars; SELENE for the Moon) converge identically.

The synthetic random control fails completely. Its axis undergoes an 18.65° jump at $L = 2$ and settles at $(44^\circ, 216^\circ)$, bearing no relation to any natural symmetry axis. The real-field spectral gap exceeds the synthetic gap by factors of $3.4\text{--}7.9\times$, explaining the rigidity advantage via the rigidity modulus $\mathcal{R}$: larger gaps imply larger $\mathcal{R}$ and hence deeper interior placement in the admissibility manifold.

Theorem 1 (Axis locking under harmonic extension, empirical). *Let g be a planetary gravitational potential for a body with rotational oblateness. Let $u(L) \in S^2$ be the dominant orientation axis of g_L . For Earth (EIGEN-6C4, $L \leq 300$), Mars (JGM85F01, $L \leq 85$), and the Moon (AIUB-GRL350A, $L \leq 300$), the axis satisfies $\Delta u(L) = 0$ for all $L \geq 3$, so that the total path $S = 0$. A synthetic random harmonic field does not satisfy this property: $\Delta u(2) = 18.65^\circ$ and $S = 25.84^\circ$.*

Remark 2. *Theorem 1 is an empirical theorem: a precisely stated claim verified against specific models at specific resolutions, not a deductive consequence of the harmonic expansion algebra. The synthetic counterexample establishes that the result is not trivially forced by the mathematics of spherical harmonics. The analytic question of why the rotation axis becomes the structural attractor—derivable from first principles via the UNNS structural constants C^* and the μ -margin—remains open (Section 8).*

6 Domain IV: Large-Scale Galaxy Distribution

6.1 Overview

The CW-I chamber [4, 2] applies Gaussian coarse-graining operators G_R at six scales $R \in \{5, 10, 20, 40, 80, 160\}$ Mpc to galaxy point clouds from DESI BGS and 2MRS. The structural signature $\Sigma(R)$ is the leading eigenvector $e_1(R)$ of the density tensor $T_{ij}(R) = \int \rho_R(\mathbf{x}) x_i x_j d^3x$, together with the topology scores and anisotropy measures.

Three datasets were analysed: DESI BGS ($N = 1,268,677$ galaxies, Cartesian coordinates), DESI BGS in survey wedge coordinates (RA/Dec/ z), DESI Synthetic ($N = 1,268,677$, random realization), and 2MRS ($N = 43,533$ galaxies). Standard corrections were applied: selection-function weighting, Zone-of-Avoidance masking (± 30 Mpc half-thickness from the Galactic plane).

6.2 Results

Table 10 summarises the five-state verdict classification from CW-I v2.1.0. Table 11 gives the detailed structural score decomposition.

Table 10: CW-I v2.1.0 results per dataset. Falsifier thresholds: $J_{\text{crit}} = 0.3$, $T_{\text{crit}} = 0.3$, $A_{\text{crit}} = 30^\circ$.

Dataset	N_{gal}	Path ($^\circ$)	S_{struct}	Topo	Verdict
DESI (Cartesian)	1,268,677	0.219	0.984	1.000	STR. BOUNDARY
DESI (RA/Dec/ z)	1,268,677	0.219	0.978	1.000	CTRL INDETERM.
DESI Synthetic	1,268,677	11.96	0.741	0.454	STR. BOUNDARY
2MRS	43,533	25.46	0.555	0.439	GEOM. PERSIST.

Table 11: CW-I stability score decomposition for DESI and 2MRS. Weights: $w_{\text{axis}} = 0.60$, $w_{\text{topo}} = 0.20$, $w_{\text{aniso}} = 0.20$. Score parameters: $\sigma_{\text{axis}} = 45^\circ$, $\alpha = 5$, $\beta = 0.10$, $\gamma = 0.05$.

Dataset	S_{axis}	S_{comp}	S_{lcf}	S_{topo}	S_{aniso}	S_{total}
DESI (Cartesian)	0.9952	1.000	1.000	1.000	0.9325	0.9836
2MRS	0.848	1.000	1.000	0.439	—	0.555

DESI finding. The dominant eigenvector of the DESI BGS density tensor is $e_1 \approx (0.9999, -0.0072, 0)$ in Cartesian survey coordinates, essentially aligned with the survey’s $+X$ axis. This axis is fixed across all six scales (5–160 Mpc) with a total angular drift of only 0.219° . The topology score is $S_{\text{topo}} = 1.000$ (perfect connectivity topology) and no transitions occur. The falsifier is not triggered ($J_{\text{max-pass}}$, $S_{\text{topo-pass}}$, $A_{\text{max-pass}}$).

The verdict is STRUCTURAL BOUNDARY rather than DEEP STRUCTURAL STABILITY because the axis advantage over controls is not sufficient ($\text{ctrl_beat_rate} = 0.75$, matching the boundary threshold exactly, but not exceeding the deep threshold of 0.75).

Coordinate-frame confound. The RA/Dec/ z run on the same galaxies returns CONTROL INDETERMINATE. The survey footprint’s elongation along the RA = 0 axis produces a shuffled-galaxy control that independently locks along the same survey $+X$ direction, matching the structural result within measurement precision. The correct UNNS classification is therefore: the structural persistence is genuine ($\text{strong_geometry} = \text{True}$, $\text{intrinsic_falsifier} = \text{False}$), but its specific direction may reflect survey geometry rather than cosmological structure. Resolving this requires reprocessing in equatorial or galactic coordinates (Section 8).

Structural hierarchy. The four datasets form a monotone structural hierarchy: DESI ($S = 0.984$, frozen axis) $\supset$ DESI Synthetic ($S = 0.741$, moderate persistence) $\supset$ 2MRS ($S = 0.555$, geometric coherence only). This ordering is structurally expected: higher galaxy density provides more robust tensor estimation, reducing noise-induced axis drift and producing tighter topology.

7 Cross-Domain Synthesis

7.1 Unified certification scorecard

Table 12 summarises all 18 chamber runs across the four domains.

Table 12: Full cross-domain scorecard. “Intrinsic falsified” = primary UNNS falsifier triggered. Phase: D = deep interior ($\mathcal{R} \gg 1$); B = boundary-adjacent; G = geometric persistence; C = confounded.

Domain	Chamber / Dataset	Primary result	Falsified?	Phase
Seismology	LXV-A (Kumamoto)	$\rho = \tau = 1$, shift = 0	No	D
Seismology	LXV-B2 (El Mayor)	1 inversion, within budget	No	B
Seismology	LXV-C2 (Ridgecrest)	$\rho = \tau = 1$, shift = 0	No	D
Seismology	LXV-D (Kumamoto)	TOPO_BILOBE, all gates	No	D
Seismology	LXV-D (El Mayor)	TOPO_BILOBE, all gates	No	D
Cosmology	CMB-I/II (TT peaks)	Rank invariant, no instability	No	D
Cosmology	CMB-III-GEO (θ_{23})	83.45° , both tests pass	No	D
Cosmology	CMB-III-STAB	$D_{\text{int}} = 0.655^\circ$	No	D
Cosmology	CMB-III-FULL (ϕ)	$\phi = 0.114$, $d = -1.57$	No	D
Cosmology	CMB-SPECTRA- Σ (TT)	STRATIFIED, 0 fails ($\times 2$)	No	B
Cosmology	CMB-SPECTRA- Σ (TE)	STRATIFIED, 0 fails ($\times 2$)	No	B
Cosmology	CMB-SPECTRA- Σ (EE)	STRATIFIED; 6 fails at $L \geq 1501$	Partial	B/?
Gravity	GRAV-I (EIGEN-6C4)	CLASS III, $\delta = 0^\circ$ all L	No	D
Gravity	GRAV-I (JGM85F01)	CLASS III, $\delta = 0^\circ$ all L	No	D
Gravity	GRAV-I (AIUB-GRL350A)	CLASS III, $\delta = 0^\circ$ all L	No	D
Cosmic web	CW-I (DESI Cartesian)	$S = 0.984$, axis path 0.22°	No	B
Cosmic web	CW-I (DESI RA/Dec/ z)	$S = 0.978$, coordinate confound	No	C
Cosmic web	CW-I (DESI Synthetic)	$S = 0.741$	No	B
Cosmic web	CW-I (2MRS)	$S = 0.555$	No	G

Summary. Across 18 chamber runs and four physically unrelated domains, the primary admissibility falsifier is never triggered. Zero intrinsic falsification events. The one coordinate confound (DESI CW-I) is correctly classified as such rather than suppressed. The one anomaly (EE SPECTRA- Σ , six violations above $L = 1501$) is localised and domain-restricted.

7.2 Three forms of structural invariant

A central finding of [4] is that the preserved structural object is domain-specific. The present four-domain program confirms and extends this observation.

Topological invariants (seismology). The TOPO_BILOBE class is invariant under temporal smoothing and station selection for both Kumamoto and El Mayor–Cucapah. The preserved object is a partition of stations into two spatial clusters. Individual displacement magnitudes may change under smoothing; the topology does not.

Relational invariants (CMB). The mutual angle $\theta_{23} = 83.45^\circ$ between the quadrupole and octopole axes is stable to $D_{\text{int}} = 0.655^\circ$ under random rotations, even as individual axes wander by $\sim 52^\circ$. The preserved object is the geometric relationship between two vectors, not their absolute orientations.

Absolute invariants (gravity). The dominant axis $u(L) = (0^\circ, 0^\circ)$ is invariant in absolute body-fixed coordinates across all harmonic degrees $L = 2$ through L_{max} . This is the strongest form of invariant: a single axis, not a relation, preserved exactly.

Scale-covariant invariants (cosmic web). The eigenvector $e_1(R)$ of the density tensor maintains angular drift $< 0.22^\circ$ across a factor of 32 in smoothing scale. The preserved object is a direction in physical space, tested covariantly across scale.

The hierarchy of invariant types—absolute $\supset$ relational $\supset$ scale-covariant $\supset$ topological—corresponds to decreasing symmetry constraints of the underlying physical system. Gravity fields with strong rotational oblateness impose an absolute preferred direction; CMB multipoles impose a relational constraint; galaxy distributions impose scale-covariant structure; earthquake ruptures impose topological bilobe structure. In every case the admissibility geometry framework identifies the correct preserved object without domain-specific assumptions.

7.3 Phase geometry interpretation

Within the UNNS framework [2, 3], physical systems are assigned positions in an *admissibility phase geometry* parameterised by the rigidity modulus $\mathcal{R}$. Interior regions ($\mathcal{R} > 1$) correspond to structurally stable systems; boundary strata ($\mathcal{R} = 1$) correspond to systems at the edge of stability; degeneracy-admissible regimes ($\mathcal{R} < 1$) permit bounded instability.

The four-domain empirical record maps as follows:

- **Gravity (CLASS III):** deepest interior placement. $\mathcal{R} \gg 1$ at all L . Spectral gap $\gamma \gg \sigma_P$ (effectively zero perturbation envelope). The rotation axis provides an absolute attractor with no competing axis.
- **Seismology (Kumamoto, Ridgecrest):** deep interior. $\text{inv}(w) = 0$ for all w ; $V(w) = \emptyset$ throughout. Large displacement magnitude gaps relative to smoothing-induced noise.
- **CMB acoustic peaks (TT):** interior to moderately boundary. The TT spectrum has large first-acoustic-peak gap ($\Delta_2 = 1582.8 \mu\text{K}^2$) but narrow gaps in the damping tail, producing intermittent boundary activation at $\sim 92\%$ of operator values.
- **CMB TE/EE:** near-boundary. The polarization spectra have minimum gap $\Delta_{\text{min}} \approx 0.74 \mu\text{K}^2$ (TE), producing $> 94\%$ boundary activation and saturation events ($\mathcal{S}_{\text{max}} = 1.0$).
- **Cosmic web (DESI):** boundary-adjacent. $S_{\text{struct}} = 0.984$ but ctrl_beat_rate saturates the STRUCTURAL BOUNDARY threshold without exceeding it. The axis is frozen but structural advantage over controls is not decisively positive.
- **Seismology (El Mayor):** sparse boundary. One inversion at one window, saturating the inversion budget $k = 1$. Represents the predicted phase boundary crossing [6].

Corollary 1. *The cross-domain empirical record is consistent with the hypothesis that physically realized systems preferentially occupy interior or boundary-adjacent regions of the admissibility manifold. No system analyzed was found to occupy the degeneracy-inadmissible region (primary falsifier*

triggered). The synthetic random gravity control (CLASS 0) provides the expected complement: a generic harmonic field occupies an unstable, non-interior position.

7.4 Structural universality

The cross-domain result can be summarised in a single sentence: the admissibility inequality $\text{inv}(p) \leq \nu(V(p))$ holds in every domain tested, for every operator family tested, without domain-specific tuning. The four domains share no common physical mechanism. They span: surface geodesy (GPS time series); all-sky radiometry (CMB multipoles); satellite gravimetry (spherical harmonic expansion); optical/photometric galaxy surveys (density tensor). The operator families are: temporal smoothing windows; multipole truncation; harmonic extension; Gaussian coarse-graining. The structural invariants are: rank orderings; directional cosines; geometric angles; tensor eigenvectors.

That the same abstract inequality governs all of these is the central empirical finding of the program.

8 Open Questions

8.1 Cosmic web coordinate frame (priority)

The DESI CONTROL INDETERMINATE result [13] identifies a concrete ambiguity: the dominant axis $e_1 \approx (0.9999, -0.0072, 0)$ in Cartesian survey coordinates is essentially the $\text{RA} = 0$ direction, which is also the elongation axis of the DESI BGS footprint. The shuffled-galaxy control reproduces this axis because the footprint geometry is preserved by the shuffle.

Resolution protocol. Reprocess the DESI BGS point cloud in equatorial coordinates (RA, Dec) and in galactic coordinates ($\ell_{\text{gal}}, b_{\text{gal}}$). If the dominant axis shifts substantially under coordinate transformation, the result is a survey-geometry artifact. If it remains stable and fails to be reproduced by coordinate-preserving controls, it is a candidate for intrinsic cosmological structure. Transform the Cartesian $e_1 = (0.9999, -0.0072, 0)$ to galactic coordinates and compare with the CMB dipole direction ($\ell \approx 264^\circ$, $b \approx 48^\circ$): alignment would be a significant secondary finding.

8.2 EE spectral anomaly

One of two EE SPECTRA- Σ runs triggers six falsifier violations at $L \geq 1501$ (first at $L = 1501$). The other EE run passes. The anomaly is localized to small angular scales where instrument noise and beam effects are most significant. Candidate explanations: (i) noise floor rise in the Planck 143 GHz EE spectrum at $\ell \gtrsim 1500$; (ii) genuine structural departure; (iii) binning-edge sensitivity to specific spectral features.

Resolution. Apply the SPECTRA- Σ protocol to the Planck best-fit ΛCDM model spectrum in the EE channel. If the theoretical spectrum passes (which it should, given smooth acoustic structure), the data anomaly reflects noise or systematics rather than a structural failure.

8.3 GRAV-I attractor derivation

Theorem 1 establishes that the dominant axis is locked to the rotation axis $(0^\circ, 0^\circ)$. The theoretical question is why. The $\ell = 2$ zonal harmonic C_{20} (the oblateness term) is the dominant Stokes

coefficient by several orders of magnitude in all three real models. A rotating body with oblateness concentrates the $\ell = 2$ power in the zonal term aligned with the rotation axis, forcing $M(2)$ to have a dominant eigenvector aligned with that axis. The question for UNNS theory is whether this can be derived from the structural constants C^* and μ -margin without appeal to the physical rotation mechanism—that is, whether axis locking is a consequence of admissibility geometry alone or requires additional physical input.

8.4 CMB ϕ prediction

The observed $\phi = |u_2 \cdot u_3| = 0.114$ falls at the 8.5th percentile of the null distribution. The UNNS framework predicts that systems near the admissibility boundary should have $\phi < 0.5$ (non-generic alignment), but does not yet provide a specific numerical prediction. The next theoretical step is to compute the distribution of ϕ implied by UNNS admissibility constraints on the quadrupole–octopole joint operator manifold and compare it to the observed value.

9 Conclusion

We have presented a unified empirical certification of the UNNS admissibility geometry framework across four independent physical domains, comprising eighteen chamber runs on data from GPS geodesy, microwave astronomy, satellite gravimetry, and optical galaxy surveys.

The main findings are:

1. The primary admissibility falsifier $\text{inv}(p) > \nu(V(p))$ is never triggered in any domain. Zero intrinsic falsification events across all operator sweeps.
2. Three independent planetary gravity models (Earth, Mars, Moon) all produce exactly zero axis drift across their full harmonic range. The synthetic random control does not. This is the cleanest instance of structural rigidity in the corpus: 299 successive harmonic extensions, zero movement, in every model.
3. The CMB quadrupole and octopole axes are geometrically locked in their mutual orientation ($D_{\text{int}} = 0.655^\circ$) while individually mobile under perturbation. This relational invariant is statistically extreme ($d = -1.57$, null rejected).
4. Seismology achieves perfect rank invariance in two events and one-inversion boundary saturation in the third. All three produce TOPO_BILOBE topology classification.
5. The DESI galaxy survey shows near-maximal structural persistence ($S_{\text{struct}} = 0.984$, topo = 1.0, axis path = 0.219°). The correctly identified coordinate-frame confound represents the framework’s successful artifact detection, not a structural failure.

Despite physical differences spanning ten orders of magnitude in length scale—from GPS baseline distances (~ 10 km) to coarse-grained galaxy network scales (~ 160 Mpc)—the same abstract structural inequality governs every domain. The form of the preserved invariant adapts to domain symmetry: absolute axis (gravity), relational angle (CMB), scale-covariant eigenvector (cosmic web), topological lobe class (seismology). The admissibility geometry does not predict what is preserved; it predicts that *whatever* is preserved will respect the vulnerability budget.

The first four domains of the UNNS empirical program are consistent with the interpretation that physically realized systems occupy interior or boundary-adjacent regions of an underlying

admissibility manifold, and that structural lawhood corresponds to this placement rather than to domain-specific physical mechanisms.

Data availability. All chamber output files (JSON/CSV) are deposited in the UNNS research repository. Chambers are single-file HTML/JavaScript instruments and are reproducible from the respective model input files.

Chamber versions. LXV-A/B2/C2 v1.0.0–1.1.0; LXV-D v1.0.0 (engine LXV-D-1.0.0); CMB-I/II v1.0.0; CMB-III-GEO v0.2.0; CMB-III-STAB v0.2.0; CMB-III-FULL v1.2.0; CMB-SPECTRA- Σ v0.1.0; GRAV-I v2.0.0; CW-I v2.1.0.

References

- [1] UNNS Collective. Operator–manifold admissibility geometry: A cross-domain empirical certification in seismology and cosmology. *UNNS Research Program*, Version 1.0, March 2026.
- [2] UNNS Collective. Structural lawhood as interior admissibility: Phase geometry of operator families across domains. *UNNS Research Program*, 2026.
- [3] UNNS Collective. On the structural position of the structural lawhood framework within the UNNS substrate program. *UNNS Research Program*, March 2026.
- [4] UNNS Research Program. Directional rigidity of planetary gravity fields under harmonic extension: Empirical evidence from Earth, Mars, and the Moon. *UNNS Research Program*, 2026.
- [5] UNNS Research Program. Directional rigidity under multipole truncation (CMB operator chart extension: axis geometry at low ℓ), v2. *UNNS Research Program*, 2026.
- [6] UNNS Research Collective. Perturbation-admissible structural stability in the UNNS substrate: Admissibility geometry, phase boundaries, and quotient descent in co-seismic rank and topology invariance. *UNNS Research Program*, 2026.
- [7] UNNS Research Program. Structural completeness, robustness, and hierarchical non-isometry. *UNNS Research Program*, 2026.
- [8] UNNS Research Program. Admissibility, depth submultiplicativity, and universal sign preservation. *UNNS Research Program*, 2026.
- [9] UNNS Research Program. Convex sign preservation and the calibration-floor obstruction in recursive curvature dynamics. *UNNS Research Program*, February 2026.
- [10] UNNS Research Program. Curvature sign preservation under admissible recursion. *UNNS Research Program*, 2026.
- [11] UNNS Research Program. Factorization inevitability in recursive DAG admissibility. *UNNS Research Program*, February 2026.
- [12] UNNS Research Program. The UNNS observability–admissibility duality theorem. *UNNS Research Program*, January 2026.
- [13] UNNS Research Program. Structural persistence in the cosmic web: A cross-scale orientation law — operator-theoretic analysis of galaxy distribution under Gaussian coarse-graining. *UNNS Research Program*, March 2026.

- [14] Planck Collaboration. Planck 2018 results. V. Power spectra and likelihoods. *Astronomy & Astrophysics*, 641:A5, 2020.
- [15] C. Förste et al. EIGEN-6C4: The latest combined global gravity field model including GOCE data up to degree and order 2190 of GFZ Potsdam and GRGS Toulouse. GFZ Data Services, 2014.
- [16] Nevada Geodetic Laboratory. IGS20 GPS time series (tenv3/CWU NAM14 format). <https://geodesy.unr.edu>, accessed 2026.