

XCS229i Problem Set 3

Due Sunday, January 17 at 11:59pm PT.

Guidelines

1. These questions require thought, but do not require long answers. Please be as concise as possible.
2. If you have a question about this homework, we encourage you to post your question on our Slack channel, at <http://xcs229i-scpd.slack.com/>
3. Familiarize yourself with the collaboration and honor code policy before starting work.
4. For the coding problems, you may not use any libraries except those defined in the provided started code. In particular, ML-specific libraries such as `scikit-learn` are not permitted.

Submission Instructions

Written Submission: Some questions in this assignment require a written response. For these questions, you should submit a PDF with your solutions online in the online student portal. As long as the PDF is legible and organized, the course staff has no preference between a handwritten and a typeset \LaTeX submission. If you wish to typeset your submission and are new to \LaTeX , you can get started with the following:

- Type responses only in `submission.tex`.
- Submit the compiled PDF, **not** `submission.tex`.
- Use the commented instructions within the `Makefile` and `README.md` to get started.

Honor code

We strongly encourage students to form study groups. Students may discuss and work on homework problems in groups. However, each student must write down the solutions independently, and without referring to written notes from the joint session. In other words, each student must understand the solution well enough in order to reconstruct it by him/herself. In addition, each student should write on the problem set the set of people with whom s/he collaborated. Further, because we occasionally reuse problem set questions from previous years, we expect students not to copy, refer to, or look at the solutions in preparing their answers. It is an honor code violation to intentionally refer to a previous year's solutions.

1. Constructing kernels

In class, we saw that by choosing a kernel $K(x, z) = \phi(x)^T \phi(z)$, we can implicitly map data to a high dimensional space, and have a learning algorithm (e.g SVM or logistic regression) work in that space. One way to generate kernels is to explicitly define the mapping ϕ to a higher dimensional space, and then work out the corresponding K .

However in this question we are interested in direct construction of kernels. I.e., suppose we have a function $K(x, z)$ that we think gives an appropriate similarity measure for our learning problem, and we are considering plugging K into the SVM as the kernel function. However for $K(x, z)$ to be a valid kernel, it must correspond to an inner product in some higher dimensional space resulting from some feature mapping ϕ . Mercer's theorem tells us that $K(x, z)$ is a (Mercer) kernel if and only if for any finite set $\{x^{(1)}, \dots, x^{(n)}\}$, the square matrix $K \in \mathbb{R}^{n \times n}$ whose entries are given by $K_{ij} = K(x^{(i)}, x^{(j)})$ is symmetric and positive semidefinite. You can find more details about Mercer's theorem in the notes, though the description above is sufficient for this problem.

Now here comes the question: Let K_1, K_2 be kernels over $\mathbb{R}^d \times \mathbb{R}^d$, let $a \in \mathbb{R}^+$ be a positive real number, let $f : \mathbb{R}^d \mapsto \mathbb{R}$ be a real-valued function, let $\phi : \mathbb{R}^d \rightarrow \mathbb{R}^p$ be a function mapping from \mathbb{R}^d to \mathbb{R}^p , let K_3 be a kernel over $\mathbb{R}^p \times \mathbb{R}^p$, and let $p(x)$ a polynomial over x with *positive* coefficients.

For each of the functions K below, state whether it is necessarily a kernel. If you think it is, prove it; if you think it isn't, give a counter-example.

[**Hint:** For part (e), the answer is that K is indeed a kernel. You still have to prove it, though. (This one may be harder than the rest.) This result may also be useful for another part of the problem.]

- (a) [6 points (Written)] $K(x, z) = K_1(x, z) + K_2(x, z)$
- (b) [6 points (Written)] $K(x, z) = K_1(x, z) - K_2(x, z)$
- (c) [6 points (Written)] $K(x, z) = aK_1(x, z)$
- (d) [6 points (Written)] $K(x, z) = -aK_1(x, z)$
- (e) [6 points (Written)] $K(x, z) = K_1(x, z)K_2(x, z)$
- (f) [6 points (Written)] $K(x, z) = f(x)f(z)$
- (g) [6 points (Written)] $K(x, z) = K_3(\phi(x), \phi(z))$
- (h) [6 points (Written)] $K(x, z) = p(K_1(x, z))$

This handout includes space for every question that requires a written response. Please feel free to use it to handwrite your solutions (legibly, please). If you choose to typeset your solutions, the `README.md` for this assignment includes instructions to regenerate this handout with your typeset L^AT_EX solutions.

1.a

1.b

1.c

1.d

1.e

1.f

1.g

1.h