Dirac operators in coarse geometry

Ulrich Bunke*

October 23, 2026

Abstract

Contents

1 Overview **2**
1.1 A geometric operator and its analytic realization . . . 2
1.2 Which part of the spectrum is stable? . . . 4
1.3 The spectrum near zero and index obstructions . . . 5
1.4 From functional calculus to coarse geometry . . . 6
1.5 Positivity, support, and scalar curvature . . . 7
1.6 Decomposition and escape to infinity . . . 8
1.7 The symbol, the index map, and assembly . . . 8
1.8 Plan of the course . . . 9
1.9 References and further reading . . . 9

2 Dirac operators **12**
2.1 Aim and conventions . . . 12
2.2 Principal symbols . . . 12
2.3 Motivation: a first-order square root . . . 13
2.4 Symbols and the characterization of Dirac-type operators . . . 14
2.5 Clifford modules and Dirac bundles . . . 15
2.6 Examples from geometry . . . 17
2.6.1 The de Rham operator . . . 17
2.6.2 The spin Dirac operator and twisting . . . 18
2.6.3 A one-dimensional model . . . 20

*Fakultät für Mathematik, Universität Regensburg, 93040 Regensburg, ulrich.bunke@mathematik.uni-regensburg.de

2.7 The Weitzenbock formula and positivity 20
2.8 Gradings and Callias type operators 22
2.8.1 A worked Callias example on the line 23
2.9 Transition to the analytic theory 24
2.10 Exercises and suggested lecture structure 24

1 Overview

Dirac operators occur in differential geometry, global analysis, and mathematical physics. They connect local geometric data with global spectral invariants. The main purpose of this course is to explain one part of this connection: how a Dirac operator on a complete Riemannian manifold determines an index class which remembers the large-scale geometry of the manifold.

The main construction has the form

$$D \longmapsto \operatorname{ind} \mathcal{X}(D) \in K\mathcal{X}_*(M).$$

Here $K\mathcal{X}(M)$ denotes the complex coarse K-homology spectrum of M (Definition 8.18). The index relates the large-scale geometry of M to the spectral and localization properties of D near zero. Its degree is determined by the grading and Clifford symmetries of D. The analytic construction is developed in Section 9, after the construction of its target in Section 8.

1.1 A geometric operator and its analytic realization

Let M be a Riemannian manifold without boundary and let $E \to M$ be a finite-rank Hermitian vector bundle of positive rank. A Dirac-type operator is a formally symmetric first-order differential operator

$$D : C_c^\infty(M, E) \longrightarrow C_c^\infty(M, E)$$

whose square is of Laplace type. Thus its leading second-order part is determined by the Riemannian metric. For a geometric Dirac operator, this takes the more precise form of a Weitzenbock formula

$$D^2 = (\nabla^E)^* \nabla^E + \mathcal{R}, \tag{1.1}$$

where $\mathcal{R}$ is the curvature endomorphism described in Proposition 2.20. Examples include $d + d^*$ on differential forms and the spin Dirac operator. Adding a potential preserves the principal symbol (Proposition 2.16); Callias potentials impose additional conditions which make their contribution to positivity explicit (Definition 2.22 and Proposition 2.23).

The principal symbol recovers the Riemannian metric. By Proposition 2.11 and (2.4),

$$[D, f] = c(df), \qquad c(df)^* c(df) = |df|^2 \mathrm{id}_E$$

for every real-valued smooth function f. Thus the pointwise operator norm is $\|c(df_x)\| = |df_x|$. If df is bounded, the commutator, initially defined on compactly supported smooth sections, extends to a bounded multiplication operator with

$$\|[D, f]\|_{B(L^2(M,E))} = \|df\|_\infty.$$

The pointwise identity determines the norm of every cotangent vector, since each such vector is df_x for a suitable f.

Example 1.1. Justify the equality between the operator norm of multiplication by $c(df)$ and $\|df\|_\infty$. Further explain how one can recover the Riemannian metric from these norms. □

To discuss the spectrum, we must regard D as an unbounded operator on $L^2(M, E)$ and specify its domain. Write D_0 for the initial operator with domain $C_c^\infty(M, E)$. Formal symmetry of D_0 does not by itself imply selfadjointness.

Example 1.2. The operator $D_0 = i\partial_x$ on $C_c^\infty((0, 1)) \subseteq L^2((0, 1))$ is symmetric. We consider symmetric extensions $D_\pm$ with domains

$$\mathrm{dom}(D_\pm) = \{f \in C^\infty([0, 1]) : f(0) = \pm f(1)\}.$$

Indeed, integration by parts gives the boundary terms $i(f(1)\overline{g(1)} - f(0)\overline{g(0)})$ which vanish if both f and g are in $\mathtt{dom}(D_\pm)$. The operators $D_\pm$ happen to be essentially selfadjoint, see Proposition 3.18.

The functions $e^{-2\pi i n x}$ and $e^{-2\pi i(n+1/2)x}$, $n \in \mathbb{Z}$, form orthonormal eigenbases for the selfadjoint closures $\bar{D}_+$ and $\bar{D}_-$, respectively. Hence

$$\sigma(\bar{D}_+) = 2\pi\mathbb{Z}, \qquad \sigma(\bar{D}_-) = 2\pi(\mathbb{Z} + 1/2).$$

For λ outside the respective set, $(\lambda - \bar{D}_\pm)^{-1}$ exists as a bounded operator.

This example shows that the choice of a selfadjoint extension matters if we are interested in spectral properties. □

Note that the interval $(0, 1)$ is not complete as a Riemannian manifold. The following says that there is no choice on complete Riemannian manifolds.

Theorem 1.3 (Analytic starting point)**.** *On a complete Riemannian manifold without boundary, a formally symmetric Dirac-type operator D_0 is essentially selfadjoint.*

Its closure $\overline{D_0}$ is the unique selfadjoint extension of D_0. This is Theorem 3.13.

From now on, D denotes this selfadjoint realization unless an initial domain is specified. On an incomplete manifold, essential selfadjointness can fail, see Example 1.2. The domain is part of the spectral problem.

1.2 Which part of the spectrum is stable?

For the selfadjoint realization $(D, \mathtt{dom}(D))$ (which we abbreviate just by D), the spectrum $\sigma(D)$ is a closed subset of $\mathbb{R}$. Its different parts respond differently to geometric changes. The *essential spectrum* is what remains after removing isolated eigenvalues of finite multiplicity. It is invariant under relatively compact perturbations, and suitable changes confined to a compact region leave it unchanged. This makes geometry at infinity central to its study. The precise perturbation and decomposition statements are Theorem 4.5 and the decomposition principle, Theorem 4.9.

Individual eigenvalues can move under small perturbations. Changes in a compact region may create eigenvalues in a gap of the essential spectrum. Their positions and multiplicities are often difficult to compute, although they may carry substantial geometric or arithmetic information. Spectral problems involving automorphic forms provide one source of such questions.

The continuous spectral subspace further splits into absolutely continuous and singular continuous parts. Scattering theory studies the absolutely continuous part and its stability under suitable decay or trace-class hypotheses. Stability of the essential spectrum as a set does not by itself imply stability of this finer spectral type. We will mainly need the distinction between discrete and essential spectrum, together with estimates near zero.

Example 1.4. Start with $D_0 = -i\,d/dx$ on $C_c^\infty(\mathbb{R}) \subseteq L^2(\mathbb{R})$. Its selfadjoint closure has domain $H^1(\mathbb{R})$. Under the unitary Fourier transform it becomes multiplication by ξ with domain

$$\{v \in L^2(\mathbb{R}) : \xi v(\xi) \in L^2(\mathbb{R})\}.$$

Its spectrum is $\mathbb{R}$ and is purely absolutely continuous, but its L^2-kernel is zero. Thus absence of harmonic sections does not imply invertibility. In particular, counting kernel dimensions does not detect continuous spectrum at zero. See the line calculation in Section 4.8 and the approximate-eigenvector criterion in Theorem 4.3.

Example 1.5. We will see later that $i\,d/dx$ on $\mathbb{R}$ has a non-trivial coarse index. Since this index is unchanged by bounded selfadjoint potentials and vanishes for invertible operators, such a potential cannot create a spectral gap at zero. Applying the same argument after subtracting a real constant excludes a gap anywhere else.

In contrast, the operator

$$D := \begin{pmatrix} i\,d/dx & 0 \\ 0 & -i\,d/dx \end{pmatrix}$$

on $L^2(\mathbb{R}, \mathbb{C}^2)$ has zero coarse index: its two summands have opposite index classes. For $\epsilon > 0$, consider the constant selfadjoint potential

$$V := \begin{pmatrix} 0 & \epsilon \\ \epsilon & 0 \end{pmatrix}, \qquad \|V\| = \epsilon.$$

Since $DV + VD = 0$ and $V^2 = \epsilon^2 \mathrm{id}$, we obtain

$$(D+V)^2 = \left(-\frac{d^2}{dx^2} + \epsilon^2\right) \mathrm{id} \geq \epsilon^2 \mathrm{id}.$$

Thus $D+V$ is invertible and has a spectral gap $(-\epsilon, \epsilon)$. In fact, Fourier transformation gives

$$\sigma(D+V) = (-\infty, -\epsilon] \cup [\epsilon, \infty).$$

This illustrates how in the case of a trivial coarse index adding an arbitrarily small potential can open a gap. □

1.3 The spectrum near zero and index obstructions

For a selfadjoint operator, invertibility means that zero is outside the spectrum. Equivalently, there is $\varepsilon > 0$ such that

$$\|Du\| \geq \varepsilon \|u\| \qquad (u \in \mathrm{dom}(D)). \tag{1.2}$$

The estimate implies closed range and zero kernel. Selfadjointness then implies that the range is also dense, so D^{-1} exists and is bounded, with $\|D^{-1}\| \leq \varepsilon^{-1}$. Conversely, a bounded inverse gives the estimate. For Dirac operators it suffices to establish this bound on the initial core; see Corollary 3.17. An index extracts stable information about the obstruction to such invertibility, usually with prescribed symmetries.

Consider first a compact manifold and an odd Dirac operator on a $\mathbb{Z}/2$-graded bundle $E = E^+ \oplus E^-$. Its closure has the form

$$D = \begin{pmatrix} 0 & D^- \\ D^+ & 0 \end{pmatrix}, \qquad D^- = (D^+)^*.$$

The chiral operator D^+, viewed as a bounded map $H^1(M, E^+) \to L^2(M, E^-)$, is Fredholm, and

$$\mathrm{index}(D^+) = \dim \ker D^+ - \dim \ker D^- \in \mathbb{Z}.$$

This integer is unchanged by compact perturbations and is zero if D is invertible.

Let $F : H^+ \to H^-$ be a bounded Fredholm operator. There is a finite-rank operator $T : H^+ \to H^-$ such that $F + T$ is invertible if and only if $\mathrm{index}(F) = 0$. This can be applied to the chiral part of the bounded transform $D(1+D^2)^{-1/2}$.

Keeping the grading matters. An ungraded selfadjoint Fredholm operator has ordinary Fredholm index zero, and its finite-dimensional kernel can always be shifted away from zero by a finite-rank selfadjoint perturbation. The index of D^+ instead obstructs an invertible compact perturbation respecting the odd graded form.

The fact that the obstruction to invertible perturbations is given by an integer is related to the K-theory fact $K_0(\mathbb{K}) \cong \mathbb{Z}$.

On a noncompact manifold, a Dirac operator need not be Fredholm. We must replace $K(\mathbb{K})$ by a K-theory of an algebra which also records where an operator acts and how far it moves sections. The coarse index in coarse K-homology provides such a replacement. Its nonvanishing obstructs invertibility, while its vanishing need not make the original operator invertible. The analytic index construction and its vanishing property are discussed in Section 9.

1.4 From functional calculus to coarse geometry

The analytic construction leading to the coarse index class rests on three facts: essential selfadjointness (Theorem 3.13), finite propagation of e^{itD} (Theorem 5.7), and local compactness of $h(D)$ for $h \in C_0(\mathbb{R})$ (Lemma 4.6). The last property means that multiplication by a compactly supported continuous function on either side makes $h(D)$ compact; compare Definition 6.7.

For h with integrable, compactly supported Fourier transform, Proposition 5.13 expresses $h(D)$ using wave operators for bounded times. It therefore has finite propagation. Approximation extends the construction to every $h \in C_0(\mathbb{R})$, giving

$$h(D) \in C^*(M, E).$$

Here $C^*(M, E)$ is the norm closure of the locally compact finite propagation operators on $L^2(M, E)$ (Definition 6.10). This functional calculus statement is Theorem 6.20. Taking account of gradings and Clifford actions turns the functional calculus homomorphism into the index class.

The geometric Roe algebra uses the controlled Hilbert space $L^2(M, E)$, but the target of the index should depend on M rather than on the particular bundle. The Roe category (Definition 6.13) supplies such a target: its K-theory is, by definition, $K\mathcal{X}(M)$. For proper metric spaces, an ample Roe algebra gives an equivalent K-theory model by Theorem 8.21. Stabilization of the geometric module, when necessary, allows us to use this comparison.

Coarse geometry abstracts the data entering this construction. A bornological coarse space is a set with a coarse structure, which controls propagation, and a compatible bornology, which specifies the bounded regions used for formulating local compactness (Definition 7.1). Its morphisms are proper controlled maps (Definition 7.6). The resulting functor

$$K\mathcal{X} : \mathbf{BC} \longrightarrow \mathbf{Sp}$$

is a coarse homology theory by Theorem 8.19. The axioms are given in Definition 8.4. In particular, it inverts coarse equivalences, such as $\mathbb{Z}^n \hookrightarrow \mathbb{R}^n$ (Lemma 8.5 and Example 7.9).

Excision and vanishing on flasques yield

$$K\mathcal{X}(\mathbb{R}^n) \simeq \Sigma^n KU, \tag{1.3}$$

by Proposition 8.9 and Proposition 8.20.

1.5 Positivity, support, and scalar curvature

The local formula (1.1) yields a global estimate when its endomorphism term has a uniform positive lower bound. If $\mathcal{R} \geq \varepsilon^2 \mathrm{id}_E$ everywhere, then integration by parts gives

$$\|Du\|^2 = \|\nabla^E u\|^2 + \langle \mathcal{R}u, u\rangle \geq \varepsilon^2 \|u\|^2$$

for compactly supported smooth u. The estimate extends to the selfadjoint domain and implies invertibility; see Corollaries 2.21 and 3.17.

For the untwisted spin Dirac operator on a complete spin manifold, the Lichnerowicz formula (2.6) reads

$$D^2 = \nabla^*\nabla + \frac{\mathrm{scal}_g}{4}.$$

Consequently a uniform bound $\mathrm{scal}_g \geq c > 0$ implies a spectral gap and

$$\operatorname{ind} \mathcal{X}(D) = 0. \tag{1.4}$$

On a compact manifold, strict positivity supplies a uniform lower bound automatically. On a noncompact manifold, the condition $\mathrm{scal}_g(x) > 0$ at each point does not by itself supply the estimate needed here.

Uniform positivity outside a subset A has a more local consequence. It gives an index with support near A, whose image is the global coarse index. The support is expressed by the big family $\langle A\rangle$ of (7.2); its operator model is the support ideal of Definition 6.17. The supported index is discussed in Section 9.12. If A is compact, Corollary 4.10 gives a gap in the essential spectrum and hence Fredholmness, but discrete eigenvalues near zero may remain. Positivity outside A therefore need not imply invertibility on the whole manifold.

The contrapositive of (1.4) produces obstructions to uniformly positive scalar curvature. In comparing different metrics, one must also compare their coarse structures and the index classes in a common target. The course will formulate this compatibility rather than treating the target group as independent of every change of metric. When the primary index vanishes because of positivity, the corresponding trivialization can carry additional information. This leads to secondary, or ρ-, invariants of positive scalar curvature metrics.

1.6 Decomposition and escape to infinity

A useful index theory must permit calculations. Coarse homology theories satisfy excision, expressed by a Mayer–Vietoris sequence for suitable decompositions. If $M = M_- \cup M_+$ is coarsely excisive in the sense of Definition 8.6, Proposition 8.7 provides a boundary map

$$\partial : K\mathcal{X}_j(M) \longrightarrow K\mathcal{X}_{j-1}(M_- \cap M_+).$$

For a manifold partitioned by a hypersurface, with compatible Dirac data and product structure near the hypersurface, the partitioned index theorem identifies the boundary of the index with the hypersurface index, with compatible orientation conventions. This reduces a geometric calculation by one dimension.

Another basic mechanism is the Eilenberg swindle. On a ray, a controlled shift pushes the entire space away from every bounded subset. An infinite direct sum of successive shifts absorbs one additional copy. Additivity then forces the coarse K-homology of the ray to vanish. Spaces admitting the requisite uniformly controlled shift are called flasque (Definition 8.1). Example 8.2 checks the ray, and the proof of Theorem 8.19 implements the swindle in the Roe category.

Example 1.6. Both closed half-lines are flasque. Excision for their union $\mathbb{R}$ therefore gives

$$K\mathcal{X}(\mathbb{R}) \simeq \Sigma K\mathcal{X}(*).$$

Iteration and $K\mathcal{X}(*) \simeq KU$ give (1.3); the corresponding groups are computed in (8.6). Thus the vanishing on rays is compatible with nonvanishing on the whole line. Translating the line does not push its entire image away from a bounded set. We will prove these statements directly from the axioms in Section 8.

1.7 The symbol, the index map, and assembly

The passage from a Dirac operator to its index has a homotopy-theoretic interpretation. The cone construction of Section 10 gives a functor $\mathcal{O}^\infty : \mathbf{UBC} \to \mathbf{BC}$, where $\mathbf{UBC}$ denotes uniform bornological coarse spaces. We define local K-homology by

$$K^{\mathcal{X}} := \Sigma^{-1} K\mathcal{X} \circ \mathcal{O}^\infty : \mathbf{UBC} \longrightarrow \mathbf{Sp}.$$

The cone boundary and coarse suspension give the index map

$$a : K^{\mathcal{X}} \longrightarrow K\mathcal{X} \circ \iota,$$

where $\iota : \mathbf{UBC} \to \mathbf{BC}$ forgets the uniform structure. Section 11 will construct the symbol class $\sigma(D)$ in local K-homology $K^{\mathcal{X}}(M)$ and establish

$$a_M(\sigma(D)) = \operatorname{ind} \mathcal{X}(D).$$

The index map a_M makes the transition from the symbol to the analytic index accessible to methods of homotopy theory. For example, naturality shows that the index map is compatible with Mayer–Vietoris boundaries. Support conditions refine both its source and its target.

Coarse assembly applies this construction at increasing scales. For a symmetric entourage U containing the diagonal, the Rips complex $P_U(Z)$ has as simplices the finite subsets $F \subseteq Z$ with $F \times F \subseteq U$. With the standard uniform and bornological coarse structures on these complexes, the spectrum-level assembly map is the map

$$\mu_Z : \operatorname*{colim}_{U \in \mathcal{C}} K^{\mathcal{X}}(P_U(Z)) \longrightarrow K\mathcal{X}(Z).$$

induced by the index maps of the uniform bornological coarse spaces $P_U(Z)$ and the identifications $\mathtt{colim}_{U \in \mathcal{C}} K\mathcal{X}(P_U(Z)) \simeq K\mathcal{X}(Z)$. Here $\mathcal{C}$ is the poset of coarse entourages of Z. The Rips colimit is a coarsification, as explained in Section 8. The coarse Baum–Connes problem asks under which coarse geometric hypotheses assembly μ_Z is an equivalence.

1.8 Plan of the course

The first part develops the analytic foundations. Section 2 introduces Dirac bundles, their operators, and positivity formulas. Section 3 constructs selfadjoint realizations, and Section 4 studies spectral stability and examples. Section 5 provides the propagation and functional calculus estimates used in the index construction.

The second part introduces the geometric target. Section 6 constructs controlled Hilbert spaces and Roe algebras. Section 7 formulates boundedness and control independently of a metric. Section 8 develops coarse homology theories, Rips coarsification, and coarse K-homology, together with their first calculations.

Section 9 constructs the coarse index class. Sections 10 and 11 then develop the cone construction and the index map. Section 12 combines analytic vanishing with topological calculations to obtain applications to positive scalar curvature. The aim is to understand both the construction of the invariant and the geometric arguments which make it useful.

1.9 References and further reading

The following references complement the exposition of this course. Roe's books provide an entry into coarse geometry and its connection with index theory; the analytic references supply background for Dirac operators and their spectra. The final group develops the operator-algebraic and homotopy-theoretic index constructions.

Coarse geometry and coarse homology.

1. J. Roe, *Lectures on Coarse Geometry*, University Lecture Series **31**, American Mathematical Society, 2003 (publisher). A systematic introduction to coarse spaces, translation algebras, asymptotic dimension, and coarse embeddings. This is background for Sections 6 and 7.

2. U. Bunke and A. Engel, *Homotopy Theory with Bornological Coarse Spaces*, Lecture Notes in Mathematics **2269**, Springer, 2020 (publisher; arXiv:1607.03657). The reference for the axiomatic framework used in Sections 7 and 8, including coarse homology theories, coarsification, and coarse K-homology.

Dirac operators, spectral theory, and analytic foundations.

1. H. B. Lawson, Jr., and M.-L. Michelsohn, *Spin Geometry*, Princeton Mathematical Series **38**, Princeton University Press, 1989 (book). Background on Clifford algebras, spinor bundles, Dirac operators, and index theory, especially relevant to Section 2 and the scalar-curvature applications.

2. T. Friedrich, *Dirac Operators in Riemannian Geometry*, Graduate Studies in Mathematics **25**, American Mathematical Society, 2000 (publisher). A geometric introduction emphasizing spinors, curvature identities, and eigenvalue estimates for the Dirac operator.

3. J. Roe, *Elliptic Operators, Topology and Asymptotic Methods*, second edition, Pitman Research Notes in Mathematics **395**, Longman, 1998 (publisher). An introduction to elliptic analysis and the heat-kernel proof of the index theorem, complementing the functional calculus and heat operator discussion in Section 5.

4. T. Kato, *Perturbation Theory for Linear Operators*, Classics in Mathematics, Springer, 1995, reprint of the 1980 edition (publisher). A general reference for closed operators, selfadjointness, and spectral perturbation theory, providing analytic background for Sections 3 and 4.

5. J. Eichhorn, *Elliptic differential operators on noncompact manifolds*, in *Seminar Analysis of the Karl-Weierstrass-Institute of Mathematics, 1986/87*, Teubner-Texte zur Mathematik **106**, Teubner, Leipzig, 1988, pp. 4–169. An early systematic treatment of elliptic operators on noncompact manifolds, providing foundations for the analysis of Dirac operators on complete manifolds. This is background for the selfadjointness, spectral theory, and analytic estimates in Sections 3–5.

6. P. R. Chernoff, *Essential self-adjointness of powers of generators of hyperbolic equations*, Journal of Functional Analysis **12** (1973), 401–414 (article). A foundational source on essential selfadjointness through hyperbolic propagation, including powers of Dirac operators on complete manifolds. Compare Theorems 3.13 and 5.7, and Proposition 5.9.

7. C. Bär, *The Dirac operator on hyperbolic manifolds of finite volume*, Journal of Differential Geometry **54** (2000), 439–488 (arXiv:math/0010233). A detailed noncompact spectral example: the spin structure affects whether the spectrum is discrete or fills the real line. This complements the discussion of geometry at infinity in Section 4.

Coarse index theory, assembly, and positive scalar curvature.

1. J. Roe, *Index Theory, Coarse Geometry, and Topology of Manifolds*, CBMS Regional Conference Series in Mathematics **90**, American Mathematical Society, 1996 (publisher). A concise introduction to the interaction between index theory, coarse geometry, operator algebras, and assembly, including methods for calculating coarse indices.

2. N. Higson and J. Roe, *Analytic K-Homology*, Oxford University Press, 2000 (publisher). Analytic foundations for K-homology, operator-theoretic index classes, and assembly maps. This provides background for the passage from a Dirac operator to its symbol and index classes.

3. R. Willett and G. Yu, *Higher Index Theory*, Cambridge University Press, 2020 (publisher). A modern treatment of Roe and localization algebras, differential operators, and higher assembly maps; a complementary approach to Sections 6–11.

4. J. Roe, *Positive curvature, partial vanishing theorems, and coarse indices*, preprint, 2012 (arXiv:1210.6100). Explains localization of the coarse index when the Weitzenbock curvature term is uniformly positive away from a subset. This is directly relevant to Section 9.12.

5. U. Bunke and A. Engel, *The coarse index class with support* (arXiv:1706.06959). The main reference for the categorical construction followed in Section 9, including the comparison with Roe algebras and index classes with support.

6. U. Bunke and A. Engel, *Coarse assembly maps* (arXiv:1706.02164). Develops assembly for coarse homology theories and identifies its source using locally finite homology. This is background for the cone and index maps in Sections 10 and 11.

7. U. Bunke and M. Ludewig, *Coronas and Callias type operators in coarse geometry* (arXiv:2411.01646). Relates local symbols, support conditions, corona pairings, and coarse indices of Callias type operators. It also treats compatibility with partitions and Mayer–Vietoris boundaries, extending the perspective of this course.

2 Dirac operators

2.1 Aim and conventions

The starting point of this course is a geometric differential operator. We want to understand how its symbol determines a Clifford module, how connections produce such operators, and how curvature and potentials control their squares. The analytic realization on an L^2-space will be the subject of the next lecture.

Throughout this lecture M is a smooth Riemannian manifold without boundary and $E \to M$ is a finite-rank complex Hermitian vector bundle. Hermitian inner products are linear in the second variable.

2.2 Principal symbols

Let $\pi : T^*M \to M$ be the projection, let E, F be smooth complex vector bundles, and let $P : C^\infty(M, E) \to C^\infty(M, F)$ be a differential operator of order at most n.

Definition 2.1. *The principal symbol of P in order n is the smooth bundle homomorphism $\sigma_P : \pi^*E \to \pi^*F$ defined by*

$$\sigma_P(\xi)e = \frac{1}{n!}\big(\underbrace{[\cdots[[P, f], f], \ldots, f]}_{n \text{ commutators}} s\big)(x), \qquad \xi \in T^*_xM,$$

where f is a real-valued smooth function with $df_x = \xi$, and s is a section of E with $s(x) = e$. For $n = 0$ the formula means $\sigma_P(\xi) = P_x$.

For justification see Exercise 2.2. In local coordinates and bundle frames,

$$P = \sum_{|\alpha| \le n} a_\alpha(x)\partial_x^\alpha \quad \Longrightarrow \quad \sigma_P(\xi) = \sum_{|\alpha| = n} a_\alpha(x)\xi^\alpha.$$

In particular, the symbol is homogeneous of degree n in ξ.

Exercise 2.2. Verify the coordinate formula and deduce independence of f and s and smoothness of σ_P. □

Exercise 2.3. Let $Q : C^\infty(M, G) \to C^\infty(M, E)$ have order at most m. Show that the symbol of PQ in order $n + m$ satisfies $\sigma_{PQ} = \sigma_P \circ \sigma_Q$. Using Hermitian metrics and the Riemannian volume density, show that

$$\sigma_{P^*}(\xi) = (-1)^n \sigma_P(\xi)^* \ .$$

□

Exercise 2.4. Check that $\sigma_{\partial_{x_j}}(\xi) = \xi_j$ and that

$$\Delta = -\sum_{i,j} g^{ij}\nabla_i\nabla_j + \text{terms of order at most one} \quad \Longrightarrow \quad \sigma_\Delta(\xi) = -|\xi|^2 \mathtt{id}_E.$$

□

Definition 2.5. *We say that P is elliptic if $\sigma_P(\xi)$ is invertible for all $\xi \neq 0$.*

Exercise 2.6. If P has order less than n, its principal symbol in order n vanishes. □

2.3 Motivation: a first-order square root

Definition 2.7. *A second-order differential operator L on E is of* Laplace type *if its principal symbol satisfies $\sigma_L(\xi) = -|\xi|^2\mathtt{id}_E$. A first-order differential operator P is of* Dirac type *if P^2 is of Laplace type.*

Example 2.8. On differential forms $C^\infty(M, \Lambda^* T^*_{\mathbb{C}} M)$ the exterior derivative is a first order differential operator with $d^2 = 0$. Let d^* denote its formal adjoint with respect to the Riemannian volume density and the induced Hermitian metric. Then

$$D_{\mathrm{dR}} := d + d^*, \qquad D^2_{\mathrm{dR}} = dd^* + d^*d =: \Delta.$$

Here Δ is called the Hodge Laplacian.

Exercise 2.9. Check that the Hodge Laplacian is of Laplace type.

Thus the Hodge Laplacian is the square of a formally selfadjoint first-order operator of Dirac type. □

We discuss the problem of representing a Laplace-type operator as a square of a Dirac type operator.

Example 2.10. In Euclidean space, look for a constant-coefficient operator

$$D = \sum_{j=1}^m c_j \partial_{x_j} \quad \text{such that} \quad D^2 = -\sum_{j=1}^m \partial^2_{x_j}.$$

acting on $\mathbb{C}^n$-valued functions. Multiplication gives

$$D^2 = \sum_j c_j^2 \partial^2_{x_j} + \sum_{j<k} (c_j c_k + c_k c_j)\partial_{x_j}\partial_{x_k}.$$

Consequently the required relations are

$$c_j c_k + c_k c_j = -2\delta_{jk}\mathtt{id}.$$

These are precisely the Clifford relations. Moreover D is formally selfadjoint if $c_j^* = -c_j$, since ∂_{x_j} is formally skew-adjoint. This calculation motivates the consideration of Clifford algebras. □

For an euclidean vector space V let $\mathrm{Cliff}(V)$ be the unital complex C^*-algebra generated by the subspace V of antiselfadjoint elements with $v^2 = -\|v\|^2$. If $\dim(V) = 2n$, then $\mathrm{Cliff}(V) \cong \mathtt{Mat}_{2^n}(\mathbb{C})$ is simple and has a unique (up to isomorphism) irreducible 2^n-dimensional representation Δ called the spinor module. If E is a $\mathrm{Cliff}(V)$-module, then we have a canonical isomorphism

$$E \cong \Delta \otimes \mathtt{Hom}_{\mathrm{Cliff}(V)}(\Delta, V) \ .$$

The Clifford algebra has a grading induced by the action of -1 on V. For any oriented orthonormal basis of V, the element

$$\gamma := i^n v_1 \cdots v_{2n}$$

is an even selfadjoint involution and anticommutes with the generators. It induces a grading on any module.

If $\dim(V) = 2n+1$, then $\mathrm{Cliff}(V) \cong \mathtt{Mat}_{2^n}(\mathbb{C}) \oplus \mathtt{Mat}_{2^n}(\mathbb{C})$ and it has two irreducible 2^n-dimensional representations $\Delta^{\pm}$ corresponding to the projections onto the two summands. After choosing an orientation of V, the summands are determined by the eigenvalue ± 1 of the central selfadjoint involution

$$z := i^{n+1} v_1 \cdots v_{2n+1}$$

for any oriented orthonormal basis of V. If E is a $\mathrm{Cliff}(V)$-module, then we have a canonical isomorphism

$$E \cong \Delta^+ \otimes \mathtt{Hom}_{\mathrm{Cliff}(V)}(\Delta^+, V) \oplus E \cong \Delta^- \otimes \mathtt{Hom}_{\mathrm{Cliff}(V)}(\Delta^-, V) \ .$$

2.4 Symbols and the characterization of Dirac-type operators

Assume that P is first order. In coordinates, if $P = \sum_j a_j(x)\partial_{x_j} + b(x)$, then

$$\sigma_P(\xi) = \sum_j a_j(x)\xi_j.$$

P be a first-order operator and put $c(\xi) = \sigma_P(\xi)$.

Proposition 2.11. *Then P is of Dirac type if and only if*

$$c(\xi)c(\eta) + c(\eta)c(\xi) = -2\langle \xi, \eta \rangle \mathtt{id}_E \ . \tag{2.1}$$

If P is formally selfadjoint, then $c(\xi)^* = -c(\xi)$. Every Dirac-type operator is elliptic.

Proof. The second-order symbol of P^2 is $c(\xi)^2$. Hence the Dirac-type condition is $c(\xi)^2 = -|\xi|^2\mathtt{id}$. Polarizing gives (2.1); the converse follows by setting $\eta = \xi$. For formally selfadjoint P, the adjoint symbol formula gives

$$c(\xi) = \sigma_{P^*}(\xi) = -\sigma_P(\xi)^* = -c(\xi)^* \ .$$

Finally, for $\xi \neq 0$ the inverse of $c(\xi)$ is $-|\xi|^{-2}c(\xi)$, proving ellipticity. □

Thus a Dirac-type operator induces a Clifford multiplication $c : T^*M \to \mathtt{End}(E)$. If the operator is formally selfadjoint, this multiplication takes covectors to skew-adjoint endomorphisms.

2.5 Clifford modules and Dirac bundles

Definition 2.12. *A* Hermitian Clifford module *is a Hermitian bundle E with a smooth real-linear map $c : T^*M \to \mathtt{End}(E)$ satisfying (2.1) and $c(\xi)^* = -c(\xi)$. A* Clifford connection *is a Hermitian connection ∇^E such that*

$$\nabla^E_X(c(\xi)s) = c(\nabla^{\mathrm{LC}}_X\xi)s + c(\xi)\nabla^E_X s. \tag{2.2}$$

A Dirac bundle *is the data (E, c, ∇^E).*

Here ∇^{LC} is the Levi-Civita connection, also acting on covectors.

Lemma 2.13. *Every Hermitian Clifford module admits a Clifford connection.*

The proof is given after the construction of the spin Dirac bundle below; see the proof below.

Definition 2.14. *Given a Dirac bundle, define its* geometric Dirac operator *by*

$$D : C^\infty(M, E) \xrightarrow{\nabla} C^\infty(M, T^*_{\mathbb{C}}M \otimes E) \xrightarrow{c} C^\infty(M, E) \ . \tag{2.3}$$

Locally it is given by

$$D = \sum_{j=1}^m c(e^j)\nabla^E_{e_j} \ ,$$

where (e_j) is a local orthonormal frame and (e^j) its dual. This expression is independent of the frame.

The Leibniz rule gives

$$D(fs) = fDs + c(df)s. \tag{2.4}$$

Thus $\sigma_D(\xi) = c(\xi)$, and D is of Dirac type.

Proposition 2.15. *The geometric Dirac operator is formally selfadjoint:*

$$\int_M \langle Ds, t\rangle \, d\mathrm{vol}_g = \int_M \langle s, Dt\rangle \, d\mathrm{vol}_g, \qquad s, t \in C_c^\infty(M, E).$$

Proof. For a smooth real vector field V, define its divergence by

$$\int_M V(f) \, d\mathrm{vol}_g = -\int_M f \, \operatorname{div} V \, d\mathrm{vol}_g, \qquad f \in C_c^\infty(M).$$

If $d\mathrm{vol}_g = \rho \, |dx^1 \cdots dx^m|$, integration by parts gives $\operatorname{div} V = \rho^{-1} \sum_j \partial_{x_j}(\rho V^j)$, which also equals $\sum_j g(\nabla^{\mathrm{LC}}_{e_j} V, e_j)$. Extend this definition complex-linearly to complex vector fields.

Define the complex vector field V by $g(V, X) = \langle c(X^\flat)s, t\rangle$ for real vector fields X, where g is extended complex-bilinearly. Fix a point p and choose Riemannian normal coordinates centred at p. Then $g_{ij}(p) = \delta_{ij}$ and $\partial_k g_{ij}(p) = 0$, so the volume density $\rho = \sqrt{\det(g_{ij})}$ satisfies $\rho(p) = 1$ and $d\rho(p) = 0$. Orthonormalizing the coordinate frame gives a frame (e_j) which agrees with (∂_{x_j}) to first order at p; in particular, $\nabla^{\mathrm{LC}} e_j(p) = 0$. Writing $V = \sum_j V^j \partial_{x_j}$, we therefore have $e_j \langle c(e^j)s, t\rangle(p) = \partial_{x_j} V^j(p)$. Using skew-adjointness, Clifford compatibility, and metric compatibility, the following calculation is evaluated at p:

$$\begin{aligned}
\langle Ds, t\rangle - \langle s, Dt\rangle &= \sum_j \big(\langle c(e^j)\nabla^E_{e_j} s, t\rangle - \langle s, c(e^j)\nabla^E_{e_j} t\rangle \big) \\
&= \sum_j \big(\langle \nabla^E_{e_j}(c(e^j)s), t\rangle + \langle c(e^j)s, \nabla^E_{e_j} t\rangle \big) \\
&= \sum_j e_j\big(\langle c(e^j)s, t\rangle \big) = \sum_j \partial_{x_j} V^j \\
&= \rho^{-1} \sum_j \partial_{x_j}(\rho V^j) = \operatorname{div} V.
\end{aligned}$$

The penultimate equality uses $d\rho(p) = 0$. Since p was arbitrary, the resulting identity holds on all of M.

The vector field V is compactly supported. Choose $f \in C_c^\infty(M)$ equal to one on a neighbourhood of its support. Then $V(f) = 0$ and $f \operatorname{div} V = \operatorname{div} V$. The defining integration-by-parts identity therefore gives $\int_M \operatorname{div} V \, d\mathrm{vol}_g = 0$, proving the assertion. No orientation is required because we use the volume density. □

Proposition 2.16. *Let P be a formally selfadjoint Dirac-type operator, and let c be its symbol. Choose a Clifford connection on (E, c) and let D be the associated geometric Dirac operator. Then*

$$P = D + \Phi, \qquad \Phi \in C^\infty(M, \mathtt{End}(E)), \quad \Phi^* = \Phi.$$

Conversely, every such zero-order perturbation is a formally selfadjoint Dirac-type operator with the same symbol.

Proof. The first-order symbols of P and D agree, so their difference has order zero. Formal selfadjointness of both operators implies fibrewise selfadjointness of their difference. Conversely, adding a zero-order term changes neither the first-order symbol nor the second-order symbol of the square. □

The choice of connection in this statement is not unique. The important distinction is that a Dirac-type operator is characterized by its principal symbol, whereas a geometric Dirac operator comes with specified connection data. In particular, an arbitrary selfadjoint Φ need not be a Callias potential: its anticommutator with D may have order one.

Remark 2.17. The special feature of geometric Dirac operators in local index theory is the pointwise cancellation of negative powers of t in the supertrace of the heat kernel. Let M be closed of even dimension m, let E be a graded Hermitian Clifford module, and let its Clifford connection preserve the grading. The associated geometric Dirac operator is odd. Its heat kernel has the expansion

$$k_t(x,x) \sim (4\pi t)^{-m/2} \sum_{j\geq 0} t^j a_j(x), \qquad a_0(x) = \mathtt{id}_{E_x}.$$

Getzler rescaling shows that

$$\operatorname{str}(a_j(x)) = 0 \quad (j < m/2), \qquad \operatorname{str}(k_t(x,x)) = (4\pi)^{-m/2} \operatorname{str}(a_{m/2}(x)) + O(t).$$

Thus the local supertrace has a finite limit as $t \downarrow 0$; the cancellation occurs before integration over M.

See E. Getzler, *A short proof of the local Atiyah–Singer index theorem*, Topology **25** (1986), 111–117, and N. Berline, E. Getzler, M. Vergne, *Heat Kernels and Dirac Operators*, Springer, corrected reprint, 2004. □

2.6 Examples from geometry

2.6.1 The de Rham operator

Take $E = \Lambda^* T^*M \otimes \mathbb{C}$ with its induced metric and Levi-Civita connection. If $\varepsilon(\xi)$ denotes exterior multiplication and $\iota_{\xi^\sharp}$ denotes contraction, put

$$c(\xi) = \varepsilon(\xi) - \iota_{\xi^\sharp}.$$

The identities

$$\varepsilon(\xi)\iota_{\eta^\sharp} + \iota_{\eta^\sharp}\varepsilon(\xi) = \langle \xi, \eta \rangle \mathtt{id}, \quad \varepsilon(\xi)^* = \iota_{\xi^\sharp}$$

together with the vanishing anticommutators between two exterior multiplications or two contractions prove the Clifford and adjoint relations. Torsion-freeness of the Levi-Civita connection gives

$$d = \sum_j \varepsilon(e^j)\nabla_{e_j}, \qquad d^* = -\sum_j \iota_{e_j}\nabla_{e_j}.$$

Hence (2.3) is exactly $d + d^*$. This example exists on every Riemannian manifold.

The grading into even and odd forms makes D_{dR} odd. On a closed manifold Hodge theory identifies its kernel with de Rham cohomology, and therefore

$$\operatorname{index}(D^+_{\mathrm{dR}}) = \sum_k (-1)^k \dim H^k_{\mathrm{dR}}(M;\mathbb{C}) = \chi(M).$$

Here D^+ maps even forms to odd forms.

2.6.2 The spin Dirac operator and twisting

Let $m \geq 1$. It follows from the structure of real Cliffford algebras $\mathrm{Cliff}^{\mathbb{R}}(\mathbb{R}^m)$ that we can fix an irreducible complex Clifford module Δ_m of dimension $2^{\lfloor m/2 \rfloor}$ and an antilinear isometry J_m such that

$$J_m^2 = \varepsilon_m \mathtt{id}, \qquad J_m c(v) = \eta_m c(v) J_m \quad (v \in \mathbb{R}^m).$$

The two signs have different meanings: ε_m specifies the square of J_m, whereas η_m specifies its commutation with Clifford multiplication. The signs are given by

$m \bmod 8$	0	1	2	3	4	5	6	7
ε_m	1	1	-1	-1	-1	-1	1	1
η_m	1	-1	1	1	1	-1	1	1

In particular, for $m \equiv 0, 2, 4, 6 \pmod 8$ the commutation sign η_m is always $+1$, while ε_m is respectively $+1, -1, -1, +1$. Thus J_m is a real structure when $\varepsilon_m = 1$ and a quaternionic structure when $\varepsilon_m = -1$. In odd dimension we fix one of the two irreducible Clifford-module types.

Let M be oriented of dimension $m \geq 1$.

Definition 2.18. *A* spin structure*, in spinor-bundle language, is a Hermitian Clifford module $S \to M$ of complex rank $2^{\lfloor m/2 \rfloor}$ equipped with an antiunitary bundle map $J : S \to S$ such that (S, c, J) is locally isomorphic to (Δ_m, c, J_m) over oriented orthonormal frames. In particular,*

$$J^2 = \varepsilon_m \mathtt{id}_S, \qquad Jc(\xi) = \eta_m c(\xi) J$$

for real covectors ξ. Isomorphisms preserve the Hermitian metric, Clifford action, and J.

Remark 2.19. This recovers the usual spin structure. Indeed, for a fixed oriented orthonormal frame, two unitary Clifford-module identifications differ by a scalar $z \in U(1)$, by irreducibility. Preservation of the antilinear map J forces $z = \bar{z}$, hence $z = \pm 1$. Since J_m commutes with products of two Clifford vectors, the spin representation preserves J_m. Consequently the compatible spinor frames form precisely a $\mathrm{Spin}(m)$-lift of the oriented orthonormal frame bundle. The datum J therefore removes the scalar ambiguity of the complex Clifford module alone. If we forget J, then we obtain the definition of a Spin^c-structure. □

The Levi-Civita connection lifts uniquely to this spin frame bundle, since the differential of $\mathrm{Spin}(m) \to \mathrm{SO}(m)$ is an isomorphism of Lie algebras. It therefore induces a Hermitian connection on S. More explicitly, in a local spinor trivialization covering an orthonormal frame (e_j), put

$$\omega_{jk}(X) := g(\nabla^{\mathrm{LC}}_X e_j, e_k), \qquad A_X := \frac{1}{4}\sum_{j,k} \omega_{jk}(X) c(e^j)c(e^k).$$

The induced connection is $\nabla^S_X = X + A_X$. The matrices $c(e^j)$ are constant in this trivialization, and $\omega_{jk} = -\omega_{kj}$ implies $A^*_X = -A_X$, so the connection preserves the metric. The Clifford relations give

$$[c(e^j)c(e^k), c(e^\ell)] = 2\delta_{j\ell}c(e^k) - 2\delta_{k\ell}c(e^j),$$

and hence

$$[A_X, c(e^\ell)] = \sum_k \omega_{\ell k}(X) c(e^k) = c(\nabla^{\mathrm{LC}}_X e^\ell).$$

Together with the Leibniz rule, this proves

$$\nabla^S_X(c(\xi)s) = c(\nabla^{\mathrm{LC}}_X \xi)s + c(\xi)\nabla^S_X s.$$

Thus the Levi-Civita connection defines a Clifford connection. Moreover, J is constant in a compatible spinor trivialization and commutes with each product $c(e^j)c(e^k)$, so $\nabla^S J = 0$.

This is the unique Hermitian Clifford connection preserving J. Indeed, the difference of two Hermitian Clifford connections commutes with all Clifford multiplications, so irreducibility makes it $i\alpha\mathtt{id}_S$ for a real one-form α. Compatibility with J, which is antilinear, forces $\alpha = 0$. No auxiliary connection is needed.

The spin Dirac operator is

$$D_S = \sum_j c(e^j)\nabla^S_{e_j}.$$

In even dimension $S = S^+ \oplus S^-$ has its parallel chirality grading and D_S interchanges the summands. In odd dimension one chooses one of the two irreducible complex Clifford modules; the resulting spinor bundle has no such chirality grading.

For any Hermitian bundle W with Hermitian connection, the tensor product $S \otimes W$ is a Dirac bundle with Clifford action $c \otimes \mathtt{id}_W$ and connection

$$\nabla^{S\otimes W} = \nabla^S \otimes \mathtt{id}_W + \mathtt{id}_S \otimes \nabla^W.$$

Its operator is the twisted spin Dirac operator D_W. More generally, one may twist any Dirac bundle in the same way. Twisting changes curvature terms but preserves the Clifford symbol.

Proof of Lemma 2.13. Choose a contractible coordinate neighbourhood U with an oriented orthonormal frame. The local spin structure gives spinor bundles with Clifford connections induced by the Levi-Civita connection, as above. If $\dim M = 2n$, the classification of complex Clifford modules gives a unitary evaluation isomorphism

$$S \otimes F \xrightarrow{\cong} E|_U, \qquad F := \operatorname{Hom}_{\mathtt{Cl}(T^*U)}(S, E|_U),$$

where S has rank 2^n and F is the smooth multiplicity bundle. Choose any Hermitian connection on F. The tensor product connection $\nabla^S \otimes 1 + 1 \otimes \nabla^F$ is then a Clifford connection on $E|_U$.

If $\dim M = 2n+1$, use the two irreducible Clifford modules to write

$$E|_U \cong (S_+ \otimes F_+) \oplus (S_- \otimes F_-).$$

Both local spinor bundles have rank 2^n and carry the connections induced by the spin connection. Apply the tensor product construction to each summand. Only local spin structures are used, so no global orientation or spin structure on M is required.

For a locally finite cover by such neighbourhoods, choose a subordinate smooth partition of unity (ρ_α) and set $\nabla^E = \sum_\alpha \rho_\alpha \nabla^\alpha$. This is a Hermitian connection and satisfies (2.2): both conditions are affine in the connection, and $\sum_\alpha \rho_\alpha = 1$. □

2.6.3 A one-dimensional model

On $\mathbb{R}$ with its usual metric, take the trivial line bundle, $c(dx) = i$, and the trivial connection. Then

$$D = i\frac{d}{dx}, \qquad D^2 = -\frac{d^2}{dx^2}.$$

On the circle $\mathbb{R}/2\pi\mathbb{Z}$ with periodic sections the modes e^{ikx} satisfy $De^{ikx} = -ke^{ikx}$. The same local formula on $\mathbb{R}$ has a very different global spectral behavior. This contrast motivates keeping track of the global geometry and the operator domain in the next lectures.

2.7 The Weitzenbock formula and positivity

Use the curvature convention

$$R^E(X,Y) = [\nabla^E_X, \nabla^E_Y] - \nabla^E_{[X,Y]},$$

and define the connection Laplacian by

$$(\nabla^E)^*\nabla^E = -\sum_j \left(\nabla^E_{e_j}\nabla^E_{e_j} - \nabla^E_{\nabla^{\mathrm{LC}}_{e_j} e_j} \right).$$

It is non-negative on compactly supported sections after integration.

Proposition 2.20 (Weitzenbock formula)**.** *For the geometric Dirac operator of a Dirac bundle,*

$$D^2 = (\nabla^E)^*\nabla^E + \mathcal{R}_E, \qquad \mathcal{R}_E = \sum_{j<k} c(e^j)c(e^k)R^E(e_j, e_k). \tag{2.5}$$

The expression $\mathcal{R}_E$ is a globally defined selfadjoint bundle endomorphism.

Proof. At a chosen point use a normal orthonormal frame. Compatibility of the connection with c eliminates derivatives of the Clifford coefficients there. The diagonal terms in D^2 are $-\sum_j \nabla^E_{e_j}\nabla^E_{e_j}$. Pairing the off-diagonal terms with the Clifford relations gives

$$\sum_{j<k} c(e^j)c(e^k)\left(\nabla^E_{e_j}\nabla^E_{e_k} - \nabla^E_{e_k}\nabla^E_{e_j}\right).$$

The frame brackets vanish at that point, so these are precisely the curvature terms. This proves the formula pointwise, hence globally. The difference of the two globally defined operators is zero order. Both are formally selfadjoint, so that difference is fibrewise selfadjoint. □

For the untwisted spin Dirac operator the spin curvature identity

$$R^S(X,Y) = \frac{1}{4}\sum_{a,b}\langle R^{\mathrm{LC}}(X,Y)e_a, e_b\rangle c(e^a)c(e^b)$$

and the curvature symmetries and first Bianchi identity yield the *Lichnerowicz formula*

$$D_S^2 = (\nabla^S)^*\nabla^S + \frac{\mathrm{scal}_g}{4}\mathtt{id}_S. \tag{2.6}$$

The contraction of spin curvature to scalar curvature is recorded here as a standard identity; the preceding proof explains the general Weitzenbock formula. For the twisted spin operator,

$$D_W^2 = (\nabla^{S\otimes W})^*\nabla^{S\otimes W} + \frac{\mathrm{scal}_g}{4}\mathtt{id} + \sum_{j<k} c(e^j)c(e^k)\otimes R^W(e_j, e_k).$$

Thus positive scalar curvature alone does not give the same positivity statement for an arbitrary twist.

Corollary 2.21. *If $\mathcal{R}_E \geq a^2\mathtt{id}$ on an open subset $U \subset M$, where $a > 0$, then*

$$\|Ds\|^2_{L^2} = \|\nabla^E s\|^2_{L^2} + \int_M \langle \mathcal{R}_E s, s\rangle \,\mathrm{dvol}_g \geq a^2\|s\|^2_{L^2}, \qquad s \in C^\infty_c(U, E).$$

Proof. Integrate (2.5) against s and use formal selfadjointness and the definition of the connection Laplacian. □

In particular $\mathrm{scal}_g \geq \kappa > 0$ gives this estimate for D_S with $a = \sqrt{\kappa}/2$. On a complete manifold, after essential selfadjointness has been proved, the estimate on all of M implies a spectral gap and an inverse for the selfadjoint realization. Pointwise positivity without a uniform lower bound does not imply a uniform spectral gap on a non-compact manifold. An estimate only outside a subset is instead the starting point for an index class with support near that subset.

2.8 Gradings and Callias type operators

A grading on E is a parallel selfadjoint involution γ. Write $E = E^+ \oplus E^-$ for its eigenbundles. We require Clifford multiplication to be odd, $\gamma c(\xi) = -c(\xi)\gamma$; then D is odd and has the matrix form

$$D = \begin{pmatrix} 0 & D^- \\ D^+ & 0 \end{pmatrix}, \qquad (D^+)^* = D^- \quad \text{formally.}$$

This is the structure used for an even index class. Additional Clifford symmetries encode other degrees in the later index theory.

For the following construction assume E is graded and D is odd. We use the graded convention for Callias operators, compatible with the source indicated below.

Definition 2.22. *A* potential *for D is a smooth odd selfadjoint endomorphism Ψ such that*

$$A_\Psi := D\Psi + \Psi D$$

has order zero. The associated Callias type operator *is $B = D + \Psi$. When using it for Fredholm theory one imposes a further uniform positivity condition outside a compact subset.*

The order-zero condition is equivalent to

$$c(\xi)\Psi + \Psi c(\xi) = 0 \qquad \text{for every real } \xi. \tag{2.7}$$

Indeed, applying the Leibniz rule in a local frame gives

$$D\Psi + \Psi D = \sum_j c(e^j)(\nabla^{\mathtt{End}E}_{e_j}\Psi) + \sum_j \big(c(e^j)\Psi + \Psi c(e^j)\big)\nabla^E_{e_j}.$$

Hence, for a potential,

$$A_\Psi = \sum_j c(e^j)(\nabla^{\mathtt{End}E}_{e_j}\Psi), \qquad B^2 = D^2 + A_\Psi + \Psi^2.$$

The endomorphism A_Ψ is selfadjoint. The notation denotes an anticommutator of odd operators, rather than an ordinary commutator. The potential may be unbounded on M; all identities here are on compactly supported smooth sections.

Proposition 2.23. *Suppose that on $M \setminus K$, for a closed subset K, a potential satisfies*

$$\Psi(x)^2 \geq \big(\|A_\Psi(x)\| + a^2\big)\mathtt{id}_{E_x}, \qquad a > 0. \tag{2.8}$$

Then

$$\|Bs\|_{L^2}^2 \geq a^2\|s\|_{L^2}^2, \qquad s \in C_c^\infty(M \setminus K, E).$$

Proof. Formal selfadjointness gives

$$\|Bs\|^2 = \|Ds\|^2 + \int_M \langle (A_\Psi + \Psi^2)s, s\rangle \, d\mathrm{vol}_g.$$

The first term is non-negative and $A_\Psi(x) \geq -\|A_\Psi(x)\|\mathtt{id}$ fibrewise, so (2.8) proves the estimate. □

This mechanism does not require a positive curvature term: the potential itself can force positivity at infinity. With K compact, completeness and the analytic results of the following lectures turn this estimate into Fredholmness. With non-compact K the appropriate invariant is a coarse index with support.

2.8.1 A worked Callias example on the line

On $E = \mathbb{R} \times \mathbb{C}^2$ take the trivial connection and matrices

$$\gamma = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad C = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

They satisfy $C^* = -C$, $C^2 = -\mathtt{id}$, $T^* = T$, $T^2 = \mathtt{id}$, $CT + TC = 0$, and both C and T are odd. Set

$$D = C\frac{d}{dx}, \qquad \Psi = xT.$$

Then $A_\Psi = CT = \mathrm{diag}(-1, 1)$ and $\Psi^2 = x^2\mathtt{id}$. Thus (2.8) holds outside $[-R, R]$ for $R > 1$, with $a^2 = R^2 - 1$. Explicitly,

$$B = \begin{pmatrix} 0 & -\partial_x + x \\ \partial_x + x & 0 \end{pmatrix}, \qquad B^2 = \begin{pmatrix} -\partial_x^2 + x^2 - 1 & 0 \\ 0 & -\partial_x^2 + x^2 + 1 \end{pmatrix}.$$

The chiral equations are elementary:

$$B^+u = 0 \iff u(x) = ae^{-x^2/2}, \qquad B^-v = 0 \iff v(x) = be^{x^2/2}.$$

Only the first solution is square-integrable. For the standard closed L^2 realizations, the harmonic oscillator has compact resolvent, and consequently B^+ is Fredholm with index 1. This last analytic assertion is a preview; the calculation of the potential, the square, and the solutions requires no spectral theorem. The example exhibits a nontrivial index even though the potential forces positivity outside a compact interval.

2.9 Transition to the analytic theory

For a Dirac bundle define

$$L^2(M,E) = \text{completion of } C_c^\infty(M,E) \text{ in } \|s\|^2 = \int_M |s|^2 \, d\mathrm{vol}_g.$$

We initially view D as an operator with dense domain $C_c^\infty(M,E)$. Proposition 2.15 shows symmetry on this domain. Symmetry does not imply selfadjointness: the adjoint may have a larger domain. The next lecture studies the closure, the adjoint, and the role of completeness in proving essential selfadjointness. The same domain questions must be addressed for $D+\Psi$ before interpreting the estimates spectrally.

2.10 Exercises and suggested lecture structure

Problem 2.24 (Symbols and zero-order perturbations)**.** *On the trivial line bundle over $\mathbb{R}$, let $P = i\partial_x + \phi(x)$ for a smooth real function ϕ. Compute $[P,f]$ and P^2. Check that P is of Dirac type and formally selfadjoint, and explain why a nonzero scalar ϕ does not satisfy the anticommutator condition for a Callias potential.*

Problem 2.25 (The Clifford action on forms)**.** *Verify the exterior multiplication and contraction identities used above. Deduce $c(\xi)^2 = -|\xi|^2 \mathtt{id}$ and show that $c(\xi)$ interchanges even and odd forms. On $\mathbb{R}$ identify the matrices of $d+d^*$ on $u(x)+v(x)dx$ with those in the worked Callias example.*

Problem 2.26 (Changing the Clifford connection)**.** *If $\widetilde{\nabla}^E = \nabla^E + A$ is another Clifford connection, show that each $A(X)$ is skew-adjoint and commutes with $c(\xi)$. Prove that the two geometric Dirac operators differ by the selfadjoint endomorphism $\sum_j c(e^j)A(e_j)$.*

Problem 2.27 (Curvature and twisting)**.** *Derive the twisted Weitzenbock formula using $R^{E\otimes W} = R^E \otimes \mathtt{id} + \mathtt{id} \otimes R^W$. If the spin twist curvature term has norm at most C and $\mathrm{scal}_g \geq \kappa > 4C$, find a uniform lower bound for $\|D_W s\|^2/\|s\|^2$ for compactly supported nonzero s.*

Problem 2.28 (A bounded Callias potential)**.** *Replace xT in the line example by $h(x)T$. Compute A_Ψ and B^2. Take $h(x) = \tanh x$, verify uniform positivity outside a sufficiently large interval, and solve the two chiral kernel equations. Hint: the solutions are proportional to $\exp(\mp \int_0^x h(t)\,dt)$.*

Source material. For the geometric setup and its later role in supported index theory, see Bunke–Engel, *The coarse index class with support*, arXiv:1706.06959. For the graded potential convention, compare Bunke–Ludewig, *Coronas and Callias type operators in coarse geometry*, arXiv:2411.01646, Definitions 4.19–4.21. The local symbol calculations, proofs, examples, and exercises above provide the introductory material needed before those index-theoretic constructions.