

NOTES FROM ARSTAFANT

ABSTRACT. Notes from ArStAFANT workshop 2023, held at EPFL between June 5 and June 9, 2023. These notes are incomplete and have not been proofread. Any errors should be assumed to be due to the note-taker.

CONTENTS

1. Oleksiy Klurman, <i>On Elliott's conjecture and applications</i>	2
2. Nina Zubrilina, <i>Root Number Correlation Bias of Fourier Coefficients of Modular Form</i>	5
3. Yu-Chen Sun, <i>On divisor bounded multiplicative functions in short intervals</i>	8
3.1. Introduction	8
3.2. Main results	9
3.3. Sketch of proof for the simplest version	9
3.4. Sketch of proof of Matomaki–Radziwiłł type bound	10
4. Agniva Dasgupta, <i>Second moment of twisted cusp forms along a coset</i>	10
5. Nihar Gargava, <i>Random arithmetic lattices as sphere packings</i>	12
6. James Branch, <i>Explicit images of Shimura's map</i>	14
7. Satyabrata Sahoo, <i>On the solutions of $x^p + y^p = 2^r z^p$, $x^p + y^p = z^2$ over totally real fields</i>	17
8. Radu Toma, <i>On the size of newforms of large level on $\mathrm{PGL}(n)$</i>	17
9. Felicien Comtat, <i>Sup norms of Siegel modular forms and Bocherer conjecture</i>	18
9.1. Motivation	18
9.2. Framework	18
9.3. Fourier coefficients of Siegel modular forms	19
9.4. Bocherer conjecture and idea of proof	20
10. Wing Hong (Joseph) Leung, <i>Short second moment bounds and subconvexity for $\mathrm{GL}(3)$ L-functions</i>	20
11. Soumendra Ganguly	21
12. Phillip Harris, <i>Maximal flat periods for $\mathrm{SL}(n)$</i>	21
13. Yiannis Petridis, <i>Counting and equidistribution</i>	23
14. Michael Yiasemidis, <i>Lattice points in elliptic annuli</i>	25
15. Pranendu Darbar, <i>Asymmetric distributions of the extreme values for cubic L-functions on the 1-line</i>	27
16. Jackie Voros, <i>On the average least negative Hecke eigenvalue</i>	27
16.1. Motivation	27
17. Min Lee, <i>An extension of converse theorems</i>	30
17.1. Modular forms	30
17.2. Petersson trace formula	32
18. Junxian Li, <i>Variations of the Titchmarsh divisor problem</i>	33

19.	Mayank Pandey, <i>L^1-norms of exponential sums with multiplicative coefficients</i>	35
20.	Andrei Seymour–Howell, <i>Rigorous computation of Maass cusp forms of squarefree level</i>	39
21.	Beat Zurbuchen, <i>Sato–Tate laws for Fourier duals of exponential sums</i>	40
21.1.	Sato–Tate	40
22.	Qihang Sun, <i>Sums of Kloosterman sums and their applications</i>	42
23.	Nicolas Potthast, <i>On the asymptotics of elementary abelian extensions over global function fields</i>	45
24.	Farrell Brumley, <i>Recent extensions of Duke’s equidistribution problems</i>	45
25.	Simran Khunger, <i>Extending support for the centered moments of the low lying zeros of cuspidal newforms</i>	50
26.	Gilles Felber, <i>L^2-restriction norm of Siegel modular forms</i>	51
27.	Huimin Zhang, <i>Analytic twists of $\mathrm{GL}_2 \times \mathrm{GL}_2$ automorphic forms and applications</i>	51
	References	51

1. OLEKSIY KLURMAN, *On Elliott’s conjecture and applications*

Let’s talk about some recent progress on some topic of multiplicative number theory, starting with two applications of this recent progress.

One application is the following:

- (1) Take any binary sequence $\{a_n\}_{n \in \mathbb{N}} = \{\pm 1\}^{\mathbb{N}}$. Suppose you measure the discrepancy of this sequence along arithmetic progressions. Then Erdos conjectured that

$$\sup_{n,d} \left| \sum_{k=1}^n a_{kd} \right| = +\infty.$$

This was eventually proved by Tao (2015), following a partially successful polymath project.

- (2) The second application concerns (completely) multiplicative functions f , thus $f(mn) = f(m)f(n)$, that are 1-bounded in the sense that $|f| \leq 1$. Suppose for instance that $f : \mathbb{N} \rightarrow \{\pm 1\}$. Then the problem is to determine which multiplicative functions avoid some sign patterns.
 - (a) Clearly there are no multiplicative functions that take the value -1 from some point on. In other words, there are none that have no $+1$ at all from some point on.
 - (b) But what about functions that never take two values $+1$ in a row from some point on? Consider

$$x^2 - pqy^2 = 1, \quad f(p) = f(q) = -1.$$

Then

$$f(pqy^2) = f(q)f(p)f(y)^2 = 1,$$

$$f(pqy^2 + 1) = f(x^2) = 1.$$

- (c) Lehmer conjectured, and Schur proved using a combinatorially argument, that there are exactly two functions, completely multiplicative,

that avoid three consecutive values $+1$, or equivalently, three quadratic residues in a row.

$$\chi_3^\pm(p) = \begin{cases} 1 & \text{if } p \equiv 1(3), \\ -1 & \text{if } p \equiv -1(3), \\ \pm 1 & \text{if } p = 3. \end{cases}$$

(In the 80's, Hildebrand asked the same question about more general sign patterns, and these questions have been understood.)

(d) Four in a row is a much more difficult question.

Conjecture 1 (Hudson, 1974; Mills, 1965). *There are thirteen functions, in addition to the two mentioned above, that avoid $++++$. They are $\chi_p^\pm$ with modulus $p \in \{5, 7, 11, 13, 53\}$. There are two functions that look like characters mod 4:*

$$\begin{aligned} g_1(p) = g_2(p) &= (-1)^{\frac{p-1}{2}}, \quad p \neq 2. \\ g_1(2) &= -g_2(2) = -1. \\ g_3(p) &= 1, p \neq 2. \\ g_3(2) &= -1. \end{aligned}$$

Theorem 2 (KMV, 2023). *Conjecture 1 is true.*

Let's begin with some thoughts. Let's try to understand

$$\frac{1}{16} \sum_{n \leq N} (1 + f(n))(1 + f(n+1))(1 + f(n+2))(1 + f(n+3)).$$

If we open everything up, then we get a main term plus a k -point correlation sum for each $k \in \{1, 2, 3, 4\}$.

$$\frac{N}{16} + \sum f(n)f(n+1)f(n+2)f(n+3) + \sum (3 \text{ pt}) + \sum (2 \text{ pt}) + \sum_{n \leq N} f(n) + \dots$$

We aim to bound everything but the first term.

Question 3. When is $\sum_{n \leq N} f(n) \gg N$? We recall here that $f : \mathbb{N} \rightarrow \{\pm 1\}$ is completely multiplicative, and takes values in $[-1, 1]$.

Wirsing says that no matter which multiplicative function one takes with the above conditions, one has some cancellation, i.e.,

$$\sum_{n \leq N} f(n) = o(N),$$

unless

$$\sum \frac{1 - f(p)}{p} < \infty,$$

or equivalently,

$$\sum_{f(p)=-1} \frac{1}{p} < \infty.$$

In other words, f is close to 1 in a certain sense on the primes. In that case, the mean value is

$$\mathcal{M}_N \sim N \prod \left(1 - \frac{1}{p}\right) \left(1 + \frac{f(p)}{p} + \dots\right).$$

How about the two and three point correlations? Chowla said that the correlations of Liouville or Mobius always vanish, except when they obviously don't. Here we recall that $\lambda(n) = (-1)^{\Omega(n)}$.

Conjecture 4 (Chowla, 1964). *For all distinct $h_1, \dots, h_k \in \mathbb{N}$, we have*

$$\sum_{n \leq N} \lambda(n + h_1) \lambda(n + h_2) \cdots \lambda(n + h_k) = o(N).$$

Conjecture 5 (Elliott, 1994). *If $f : \mathbb{N} \rightarrow \mathcal{U}$ is a 1-bounded completely multiplicative complex-valued function, then its correlations vanish*

$$\sum_{n \leq N} f(n + h_1) \cdots f(n + h_k) = o(N)$$

unless $f(n)$ is “close” to $n^{it}\chi(n)$ in the sense that for some t and χ , we have

$$\mathbb{D}(f, n^{it}\chi, \infty) < \infty,$$

with notation as below.

Definition 6. We define the squared distance between two multiplicative functions at scale x to be

$$\mathbb{D}^2(f, n^{it}\chi(n), x) = \sum_{p \leq x} \frac{1 - \Re(f(p)\overline{\chi(p)}p^{-it})}{p}.$$

Remark 7. Consider functions f such that

$$\mathbb{D}(f, n^{it}\chi, \infty) = \infty$$

for all t and χ . Ergodic theorists call such functions *aperiodic*.

Remark 8. The largest that the distance $\mathbb{D}^2$ can be is

$$2 \sum_{p \leq x} \frac{1}{p} \sim 2 \log \log x.$$

Matomaki, Radziwill and Tao pointed out that this conjecture is wrong. The class of functions for which it is wrong is now called the MRT class. The idea is to construct a function f that looks like different $n^{it}\chi$'s at different scales, but looks like none of them in particular at all scales. In more detail, we are trying to understand

$$\sum f(n) \overline{f(n+1)}. \tag{1}$$

At scale $p \leq S_m$, we take $f(p) \sim p^{it_m}$. Then (1) is large. At the next scale S_{m+1} , which we take to be much, much larger than S_m , we take a different character. Then $\mathbb{D}(f, n^{it_m}, S_m)$ is small, but for all t and χ , we have $\mathbb{D}(f, n^{it}\chi, \infty) = \infty$. It's key here that this scale be much much larger than S_m .

Theorem 9 (Tao, 2015; Terivainen–Tao, 2018). *(Tao proved something about 2-point correlations, which was generalized with Terivainen to odd-point correlations.) Assume one of the following conditions:*

(a) $k = 2$, f is strongly aperiodic:

$$\inf_{\substack{|t| \leq Ax \\ q \leq (\log x)^A}} \mathbb{D}(f, n^{it}\chi, x) \xrightarrow{x \rightarrow \infty} \infty.$$

Then there exists $\mathcal{X} \subseteq \mathbb{N}$ with density one, i.e., with

$$\lim_{X \rightarrow \infty} \frac{\sum_{\substack{n \in \mathcal{X} \\ n \leq X}} 1}{\sum_{n \leq X} \frac{1}{n}} = 1,$$

such that for all h_1, h_2 distinct, we have

$$\lim_{\substack{x \rightarrow \infty \\ x \in \mathcal{X}}} \frac{1}{x} \sum_{n \leq x} f(n + h_1) \overline{f(n + h_2)} = 0.$$

- (b) k : odd. Suppose for simplicity that $k = 3$ and $f : \mathbb{N} \rightarrow [-1, 1]$ and that f is super strongly aperiodic: for each χ ,

$$\mathbb{D}^2(f, \chi, x) \gg \log \log x.$$

(We can ignore t 's because f is real.) Then the same conclusion as in the previous case is true.

Theorem 10 (KMT, 2023). *Same conditions as above:*

- (a) For $k = 2$, and f aperiodic (rather than strongly aperiodic), the same conclusion holds as in Theorem 9, except that we can only take $\mathcal{X}$ to have upper logarithmic density: same condition as above, but using $\limsup$ rather than $\lim$. (This is necessary; one can otherwise construct counterexamples.)
- (b) For k odd, with f aperiodic, and with upper logarithmic-logarithmic density.

2. NINA ZUBRILINA, Root Number Correlation Bias of Fourier Coefficients of Modular Form

Statistical properties of Fourier coefficients of automorphic forms. Distribution of $a_\pi(p)$. Let's backtrack a bit and give some informal overarching context for the more refined phenomena that we'll discuss later.

We start with the conjecture of Sato and Tate, which describes what we'll call the *horizontal* distribution. This refers to the behavior of Fourier coefficients for a *fixed* arithmetic L -function. To its coefficients $a_L(p)$, we associate a measure μ , given by the Haar measure on a certain group G_L .

Example 11. Given a non-CM elliptic curve $E/\mathbb{Q}$, we have

$$a_p = \sqrt{p}\lambda(p) = 2\sqrt{p}\cos\theta_E(p).$$

Then as $p \rightarrow \infty$, the $\theta_E(p)$ follow the famous Sato–Tate distribution:

$$\mu(\theta) = \frac{2}{\pi} \sin^2 \theta d\theta.$$

An orthogonal picture to this is what we'll refer to as the *vertical* distribution. The way we'll like to think of it is that we have some packet $\mathcal{F}_N$ of automorphic forms π , and we'd like to prescribe that the conductors of the elements of these packets have roughly the same size:

$$\frac{N(\pi)}{N} = 1 + o(1).$$

One example of this would be to take a Hecke basis of modular forms of some fixed weight and level N :

$$\mathcal{F}_N := \Gamma_0(N, k).$$

Then the Satake parameters

$$\{\theta_\pi(p)\} \approx_N \mu_p := p\text{-adic Plancherel}.$$

We won't write it down, but remark that it approaches the Sato–Tate distribution as $p \rightarrow \infty$. This is a theorem of Serre.

How about the expectation? Let's write it as the correlation:

$$\text{Cor}(\mathcal{F}_N, p) := \mathbb{E}_{\pi \in \mathcal{F}_N} a_\pi(p) = \frac{1}{|\mathcal{F}_N|} \sum a_\pi(p).$$

In this vertical averaging, this expectation, in many many cases, will be zero. One can heuristically derive this expectation from the fact that these distributions are symmetric. See the paper of Sarnak–Shin–Templier.

A similar question can be asked back on the horizontal size. If we consider a packet of primes

$$P_X = \{p : p = (1 + o(1))X\},$$

then we can again ask about this correlation function

$$\text{Cor}(P_X, \pi) \sqrt{X} = \text{almost periodic function of } \log X.$$

This will oscillate about the mean. It's a generalization of the Chebyshev bias. A reference is a paper of Rubinstein–Sarnak.

What we'll discuss today is a hybrid phenomenon, where we consider a double correlation

$$\text{Cor}(P_X, \mathcal{F}_N) = \mathbb{E}_{\substack{p \in P_X \\ \pi \in \mathcal{F}_N}} a_\pi(p).$$

We'll take N and X to depend upon each other in some way.

The problem that we'll talk about comes from the world of computer science, historically speaking. It comes from a machine learning paper of He, Lee, Oliver and Pozdnyakov. They fed a bunch of invariants coming from number theory into a machine learning algorithm and tried to guess others. For instance, they considered a family

$$\mathcal{E} = \mathcal{E}_{[N_1, N_2]}^k = \{E/\mathbb{Q} : \text{rank}(E) = r, N(E) \in [N_1, N_2]\}.$$

They fed in Frobenius traces for small primes and tried to guess the rank. They observed the following curious feature:

$$\frac{1}{|\mathcal{E}|} \sum_{E \in \mathcal{E}} a_E(p) = \text{Cor}(\mathcal{E}, p) = \text{"interesting"}.$$

This is best illustrated by a picture.

Picture shows interval between 7000 and 10000. x -axis is the parameter p . Blue line: elliptic curves of rank zero. Red line: elliptic curves of rank one. There's this weird oscillation behavior.

Bottom picture: blue is rank zero, green is rank two. Here it looks sort of like the same picture, just shifted a bit.

This might have gone completely unnoticed, but Sutherland stumbled upon it and performed a bunch of computations. He mentioned a couple things about this phenomenon. You can think about it not as a phenomenon about the rank, but rather about the root number (noting the role played by parity). The second thing

he noticed is that perhaps the correct intervals to take are “geometric intervals”: for $w \in \{\pm 1\}$,

$$\mathcal{E}_M^w = \left\{ E/\mathbb{Q} : w(E) = w, N(E) \in \begin{cases} [M, 2M] \\ [M, (1 + o(1))M]. \end{cases} \right\}$$

That is to say, one can consider either a dyadic interval or a short interval. What Sutherland noticed is that when you switch to these geometric intervals, there’s a funny phenomena that shows up, that we’ll call scale invariance.

The picture shows two intervals at unrelated scales (dyadic case) for which the plots have similar shapes, suggesting that perhaps

$$\text{Cor}(\mathcal{E}_M^w, p) \approx w \mathcal{M}\left(\frac{p}{M}\right).$$

We observe next that the ordering is very important. If we plot instead by Faltings height, naive height, j -invariant, then we just get complete gibberish. It matters that we order by conductor.

The last observation is that, rather than just elliptic curves, we can consider broader families

$$\mathcal{F}_M^w = \{\text{weight two newforms for } \Gamma_0(N) : N \in I_M, \text{degree } d\}.$$

The picture becomes very pretty if we impose no condition on degrees. The theorem we’ll present today is that one can actually write down this function and prove something about it. We give an explicit function. The short intervals are nicer to work with.

Some more pictures. Nice.

Let’s write one line about the proof. It’s basically that

$$\sum_p a_f(p) w(f) = \pm \text{trace}(T(p) \circ W_N).$$

Apply the Selberg trace formula. Unravel it. Get that the above is basically the sum of a bunch of class numbers, depending upon p and N . We apply a theorem of Skoruppa–Zagier. Get that the trace is a sum

$$\sum_{|k| \leq 2\sqrt{P/N}} H_1(k^2 N^2 - 4PN) - (P + 1).$$

Let’s add a few words about how this ties up with the Katz–Sarnak philosophy. One way to think about that philosophy is that if we take a family $\mathcal{F}$ with symmetry type $O(\infty)$, then the 1-level density conjecture says that if we take the parameter X for the prime to be the conductor to some power, say N^a , and we compute this correlation, then as $N \rightarrow \infty$, that correlation should behave as follows:

$$\text{Cor}(\mathcal{F}_N^w, P_X) \xrightarrow{N \rightarrow \infty} \begin{cases} 0 & \text{if } a < 1, \\ -\frac{w}{2} & a > 1. \end{cases}$$

Since we took p to be of size N , the murmurations occur at the phase transition between the gap in asymptotics between the two cases of the 1-level density conjecture.

Maybe two closing comments. We can compute this address using Selberg with harmonic weights, but this removes everything. Another thing is that for elliptic curves or other Galois orbits, this remains mysterious. Lastly, maybe want to advertise Sutherland’s website and IAS talk, which has a lot of interesting data.

The Selberg trace formula has the disadvantage that we can't go very high with N in terms of p . The speaker is working through what happens when one removes the weights. The function might be qualitatively different when you take N to be $p^{1/3}$ or something.

One can do higher weights similarly using the same techniques. Qualitatively it's similar, but looks different.

3. YU-CHEN SUN, *On divisor bounded multiplicative functions in short intervals*

3.1. Introduction. Huxley, 1972: prime number theorem in short intervals: for any $H \geq X^{7/12+\varepsilon}$, we have

$$\sum_{X < n \leq X+H} \Lambda(n) = (1 + o(1))H.$$

Similarly, we have

$$\sum_{X < n \leq X+H} \mu(n) = o(H)$$

and

$$\sum_{X < n \leq X+H} \lambda(n) = o(H).$$

Matomaki and Teravainen recently showed that in the Mobius and Liouville cases, the results hold for all $H \geq X^{0.55+\varepsilon}$, but they cannot prove this for the von Mangoldt function (their arguments are based upon multiplicative properties). Under RH, the above results hold for all $H \geq X^{1/2+\varepsilon}$.

In 2016, Matomaki and Radziwill proved the following breakthrough result:

Theorem 12 (Matomaki–Radziwill). *Let $f : \mathbb{N} \rightarrow [-1, 1]$ be a multiplicative function. For almost all $x \in [X, 2X]$, we have*

$$\frac{1}{h} \sum_{x < n \leq x+h} f(n) - \frac{1}{X} \sum_{X < n \leq 2X} f(n) = o(1)$$

provided that $h \rightarrow \infty$ as $X \rightarrow \infty$.

- For any $\varepsilon > 0$, the interval $[X, X + C(\varepsilon)\sqrt{X}]$ contains X^ε numbers.
- For every integer $h \geq 1$, there exists $c(h) > 0$ such that

$$\frac{1}{X} \left| \sum_{n \leq X} \lambda(n) \lambda(n+h) \right| \leq 1 - c(h).$$

How about the k -fold divisor function in short intervals? Let

$$d_k(n) = \sum_{n=m_1 \cdots m_k} 1.$$

The asymptotic formula for d_k in short intervals is that

$$\sum_{x < n \leq x+H} d_k(n) = (1 + o(1)) P_{k-1}(\log x) H,$$

where $P_k(x)$ is an explicit polynomial of degree k . Here

- $k = 2$ and $x^{131/416+\varepsilon} \leq H \leq x$ (Huxley).
- $k = 3$ and $x^{43/96+\varepsilon} \leq H \leq x$ (Holesnik).
- $4 \leq k \leq 5$ and $x^{\frac{3k-4}{4k}+\varepsilon} \leq H \leq x$ (Huxley).

- $k \geq 6$ and $x^{\frac{11}{20}+\varepsilon} \leq H \leq x$ (Matomaki–Teravainen).

It is natural to ask, can we give an almost all result for divisor bounded functions (or d_k bounded multiplicative functions). Yes, but the general theorem is a bit complicated; we will try to say it at the end of the talk. Let's start with the simple case of just the divisor function d_k .

Theorem 13 (Mangerel, 2023). *If $h_0 \rightarrow \infty$ as $X \rightarrow \infty$ and $h \geq h_0(\log X)^{(k-1)^2}$, then*

$$\frac{1}{h} \sum_{x < n \leq x+h} d_k(n) - \frac{1}{x} \sum_{x < n \leq 2x} d_k(n) = o(\log^{k-1} x)$$

for all but $o(X)$ values of $x \in [X, 2X]$.

3.2. Main results.

Theorem 14 (S. 2023). *Let $X > 0$ be sufficiently large. If $h \geq (\log X)^{(1+\varepsilon)k \log k - k + 1}$, then*

$$\frac{1}{h} \sum_{x < n \leq x+h} d_k(n) - \frac{1}{x} \sum_{x < n \leq 2x} d_k(n) = o(\log^{k-1} x)$$

for all but at most $o(X)$ integers $x \in [X, 2X]$.

For convenience, we will focus on the simplest case $k = 2$, with the length of the interval being

$$(\log X)^{(1+\varepsilon)2 \log 2 - 1}.$$

Here $2 \log 2 - 1 \approx 0.38692$, which should be compared with Mangerel's value 1.

3.3. Sketch of proof for the simplest version. We consider a relatively long average Dirichlet divisor problem. Let

$$\Delta(x) = \sum_{n \leq x} d(n) - x(\log x - 2\gamma + 1),$$

where

$$\gamma = 0.57721 \dots$$

is the Euler–Mascheroni constant. The Dirichlet hyperbola method shows that

$$\Delta(x) = O(X^{1/2}).$$

This implies that for any $h \geq x^{1/2}$, we have

$$\sum_{x < n \leq x+h} d(n) = h \log x + O(h).$$

Proposition 15 (Relatively long average). *Let $X > 0$ be a large number. One has, for all $h \in [X^{1/2}, X]$ and (...)*

For short h , we do instead the following. We first restrict to some subset where we have nice prime factorizations. Let $\varepsilon_0 > 0$ be small. We define

$$P_1 := \exp((\log \log X)^{1/2}), \quad Q_1 := (\log X)^{\varepsilon_0},$$

$$P_2 := \exp((\log X \log X)^2), \quad Q_2 = \exp((\log X \log X)^{100}).$$

Let S_A denote the set of all $n \in (X, 2X]$ having at least one prime factor in each interval $[P_1, Q_1]$ and $[P_2, Q_2]$.

Lemma 16. Let S_A be as above. Then for any $h > 0$, we have

$$\sum_{\substack{x < n \leq x+h \\ n \notin S_A}} d(n) = o(h \log X)$$

for all but $o(X)$ values of $x \in [X, 2X]$.

Next, for relatively short averages, we have to restrict to subsets. Let

$$S_{B,\varepsilon} = \{n \in \mathbb{N} : \Omega(n) \leq 2(1 + \varepsilon) \log \log X\}.$$

Here $\Omega(n)$ denotes the number of prime factors of n .

Lemma 17. Let $X > 0$ be a large real number and $\varepsilon > 0$ be a small real number. Then for any $h > 0$, we have

$$\sum_{\substack{n \notin S_{B,\varepsilon} \\ x < n \leq x+h}} d(n) = o(h \log X)$$

for all but $o(X)$ values of $x \in [X, 2X]$.

[These restriction lemmas follow from multiplicative properties, using Brun–Titchmarsh.]

From the above two lemmas, we can restrict to the set $S_A \cap S_{B,\varepsilon}$.

Proposition 18 (Matomaki–Radziwiłł type bound). *Let $X > 0$ be a large number. Let S_A and S_B be as before, and $S = S_A \cap S_B$. Let*

$$H_\varepsilon = (\log X)^{(1+\varepsilon)2 \log 2 - 1} \leq h \leq h_1 = X^{1/2}$$

for any $\varepsilon > 0$. Then

$$\frac{1}{X} \int_X^{2X} \left| \frac{1}{h} \sum_{\substack{n \in S \\ x < n \leq x+h}} d_k(n) - \frac{1}{h_1} \sum_{\substack{n \in S \\ x < n \leq h_1}} ??? \right| \ll ???.$$

3.4. Sketch of proof of Matomaki–Radziwiłł type bound. We use a mean value theorem for Dirichlet polynomials. Improvement comes from reducing the number of terms, decreasing the diagonal.

Next, we have a Parseval-type bound.

Next, we use Ramaré’s identity to carry out a Matomaki–Radziwiłł type decomposition.

4. AGNIVA DASGUPTA, *Second moment of twisted cusp forms along a coset*

We’ll talk in particular about moments of L -functions. We’ll continue in particular with the theme of short intervals that we’ve already seen in this conference, but in a slightly different context.

L -functions are hard to study individually, but often we study them as a collection, via families of L -functions. Often that is done statistically, via moments of families of L -functions. Thus, let $\mathcal{F}$ be a family of L -functions. We’re looking at expressions like

$$\frac{1}{|\mathcal{F}|} \sum_{f \in \mathcal{F}} |L(\tfrac{1}{2}, f)|^k.$$

More generally, one can look at expressions like

$$\frac{1}{T} \int_0^T |L(\tfrac{1}{2} + it)|^k dt.$$

The goal is to get asymptotic expansions or upper bounds for these.

The L -functions that we'll focus on today will mostly be on $\mathrm{GL}(1)$ (the zeta function or Dirichlet L -functions) or $\mathrm{GL}(2)$ (L -functions associated to cusp forms, possibly twisted by $\mathrm{GL}(1)$). We'll focus on

- the 4th moment for $\mathrm{GL}(1)$, and
- the 2nd moment for $\mathrm{GL}(2)$.

We consider

$$\frac{1}{T} \int_0^T |\zeta(\tfrac{1}{2} + it)|^4 dt \tag{2}$$

or, for the q -aspect,

$$\frac{1}{\phi^*(q)} \sum_{\chi(q)} |L(\tfrac{1}{2}, \chi)|^4. \tag{3}$$

The overarching conjecture is the generalized Lindelöf hypothesis (GLH). What that says is that if you have an L -function, and let's say we fix s at $1/2 + it$, then

$$|L(f, s)| \ll |q(f, s)|^\varepsilon$$

for any fixed $\varepsilon > 0$. If you're unfamiliar with the analytic conductor, then just think of it as something that depends upon all the quantities that we use when working with L -functions, such as the arithmetic conductor q and s . This implies that the moment (2) is $O(T^\varepsilon)$ and that (3) is $O(q^\varepsilon)$. Such moment bounds are called *Lindelöf on average*.

For the zeta function, the first breakthrough (i.e., result with power-savings error term) was due to Heath-Brown (1979), who showed that

$$\frac{1}{T} \int_0^T |\zeta(\tfrac{1}{2} + it)|^4 dt = P_4(\log T) + O(T^{-1/8+\varepsilon}).$$

The GL_2 variant of this was done by Anton Good (1982), who showed that

$$\frac{1}{T} \int_0^T |L(\tfrac{1}{2} + it, f)|^2 dt = P_1(\log T) + O(T^{-2/3+\varepsilon}).$$

It also follows immediately from his work that

$$\int_T^{T+T^{2/3}} |L(\tfrac{1}{2} + it, f)|^2 dt \ll T^{2/3+\varepsilon}.$$

For the q -aspect, Young (2011) showed that

$$\frac{1}{\varphi^*(q)} \sum_{\chi(q)}^* |L(\tfrac{1}{2}, \chi)|^4 = P_4(\log Q) + O(q^{-5/512+\varepsilon}).$$

This was the first power savings estimate for the fourth moment. The GL_2 analogue of this was done by Blomer–Fouvry–Kowalski–Michel–Milicevic–Sawin (2017), who showed that for a fixed GL_2 form f ,

$$\frac{1}{\varphi(q)} \sum_{\chi(q)} |L(\tfrac{1}{2}, f \otimes \chi)|^2 = P_1(\log q) + O(q^{-1/144+\varepsilon}).$$

The speaker has been working on moments over short intervals. We want to look at things like

$$\int_T^{T+H} |L(\tfrac{1}{2} + it)|^k dt$$

where H is significantly smaller than T . Here GLH gives $O(HT^\varepsilon)$. One of the first significant breakthroughs here was due to Iwaniec (1978), who showed that

$$\int_T^{T+T^{2/3}} |\zeta(\tfrac{1}{2} + it)|^4 dt \ll_\varepsilon T^{2/3+\varepsilon}.$$

How about short intervals in the q -aspect? One approach is the following. Say you have a family of automorphic forms $\mathcal{F}$, say a big family. What you want to do is to choose a subfamily where the members are kind of close to one another. That is to say, for π_1 and π_2 in the subfamily, we want the conductor of their convolution

$$C(\pi_1 \otimes \bar{\pi}_2)$$

to be small. For instance, we could consider $L(f, 1/2 + it)$ for $t \in [T, T + H]$, and look at the conductors of the Rankin–Selberg convolutions of those. For the q -aspect, let $\mathcal{F}$ be the family of all characters modulo q , and consider, for given $\alpha \in \mathcal{F}$,

$$\mathcal{F}_0 = \{\alpha \cdot \chi, \chi \pmod{d}, d \mid q, d \ll q\}.$$

For these, we have $C(\pi_1 \otimes \bar{\pi}_2) \asymp d$. Petrow–Young (2019) showed that if $q = p^3$, and if $\alpha \pmod{q}$ is primitive, and we look at a short interval fourth moment like

$$\sum_{\chi(p^2)} |L(\tfrac{1}{2}, \alpha \cdot \chi)|^4 \ll_\varepsilon p^{2+\varepsilon}.$$

Theorem 19 (D. 2023). *Let f be a level 1 cusp form. Let $q = p^3$, where p is an odd prime. Let $\alpha \pmod{q}$ be a given primitive character. Look at*

$$\sum_{\chi(q^{2/3})} |L(f \otimes \alpha \cdot \chi, \tfrac{1}{2})|^2 \ll_\varepsilon q^{2/3+\varepsilon}.$$

Corollary 20 (D. 2019). *Let $q = p^3$, $\alpha(q)^*$, and let f be a level 1 cusp form. Then*

$$\#\left\{\chi : L(f \otimes \alpha \cdot \chi, \tfrac{1}{2}) \neq 0, \chi(q^{2/3})\right\} \gg_\varepsilon q^{2/3-\varepsilon}.$$

Munshi, Singh (2019): $L(\tfrac{1}{2}, f \otimes \alpha \chi) \ll q^{1/3+\varepsilon}$ (Weyl-type subconvexity). Their proof used Munshi’s variant of the δ -method, which doesn’t require averaging over families. Also, their work is a bit more general: they work with $q = p^{3r}$.

5. NIHAR GARGAVA, *Random arithmetic lattices as sphere packings*

Given a lattice Λ , let $r > 0$ and consider the open balls $B_r(v)$, where v ranges over Λ , that are disjoint. Then (Λ, r) is called a lattice sphere packing.

Note that $2r$ can be at most

$$m(\Lambda) = \min_{v \in \Lambda - \{0\}} \|v\|,$$

otherwise some balls will begin to intersect. The goal is to maximize packing density, so take $r = \frac{1}{2}m(\Lambda)$. In that case, the packing density will be equal to

$$\frac{\mu(B_{m(\Lambda)/2}(0))}{\mu(\mathbb{R}^d/\Lambda)}$$

and is independent of scaling. So we somehow find

$$\sup_{\substack{\Lambda \subseteq \mathbb{R}^d \\ \mu(\mathbb{R}^d/\Lambda)=1}} \mu(B_{m(\Lambda)/2}(0)) = \sup_{g \in \mathrm{SL}_d(\mathbb{R})} \frac{1}{2^d} \mu(B_{m(g\mathbb{Z}^d)}(0)),$$

which we use to define our dimensional constant c_d , by requiring that the above be

$$\frac{1}{2^d} c_d.$$

Putting it together, and in more technical words, we have

$$c_d = \sup \{ \mu(B_r(0)) : r > 0, \exists g \in \mathrm{SL}_d(\mathbb{R}) \text{ and } B_r(0) \cap g\mathbb{Z}^d = \{0\} \}.$$

Clearly $c_d \in [0, 2^d]$, so the supremum exists. What is known about c_d ? The exact value is known only for

$$d \in \{1, 2, 3, 4, 5, 6, 7, 8, 24\}.$$

For other d , we want to understand the asymptotic behavior. In this talk, we will only focus on lower bounds. Note that this is only the *lattice packing problem*, not the sphere packing problem, so it was known before the work of the speaker's advisor.

The exact value of the constant c_d is known only for $d \in \{1, 2, 3, 4, 5, 6, 7, 8, 24\}$. For other d , we want to understand the asymptotic behavior. In this talk, we will only focus on lower bounds. (The speaker presented a table of the known lower bounds.)

— $c_d \geq 1$ — Minkowski–Hlawka (1896–1940) — $d \geq 2$ — —

The speaker's result works with probabilistic arguments that lie on $(D \otimes_{\mathbb{Q}} \mathbb{R})^2$, where D is a division algebra over $\mathbb{Q}$. The speaker's result is stronger than Venkatesh's for some small d , but weaker for larger d . Also, it's along a different sparse sequence.

Theorem 21 (G. 2022). *Let D be a finite-dimensional division algebra over $\mathbb{Q}$. Let $\mathcal{O} \subseteq D$ be an order in D , and let $G_0 \subseteq \mathcal{O}$ be a finite group embedded in the multiplicative group of D . Then if $d = 2 \dim_{\mathbb{Q}} D$, we have*

$$c_d \geq \#G_0.$$

To recover Venkatesh's result, take

$$D = \mathbb{Q}(\mu_n), \quad \mathcal{O} = \mathbb{Z}[\mu_n], \quad G_0 = \langle \mu_n \rangle.$$

Hence, this gives

$$c_{2\varphi(n)} \geq n.$$

The cherrypicked sequence of Venkatesh achieves an asymptotic growth of $O(d \log \log d)$. This is achieved by taking K to be the n th cyclotomic field, where

$$n = \prod_{p < N} p.$$

The division algebra construction gives more freedom to cherrypick sequences.

Let's give a taste of how this machinery works. To show that $c_d \geq K$, we must prove the existence of $g \in \mathrm{SL}_d(\mathbb{R})$ such that the origin-centered ball B with $\mu(B) = K$ satisfies

$$g\mathbb{Z}^d \cap B = \{0\}.$$

This is an optimization problem on the space

$$X_d := \{ \Lambda \subseteq \mathbb{R}^d : \mu(\mathbb{R}^d/\Lambda) = 1 \} = \{ g\mathbb{Z}^d : g \in \mathrm{SL}_d(\mathbb{R}) \} \cong \mathrm{SL}_d(\mathbb{R}) / \mathrm{SL}_d(\mathbb{Z}).$$

Probabilistic arguments.

Given $f : \mathbb{R}^d \rightarrow \mathbb{R}$ (measurable, bounded, with compact support), we consider the lattice sum function

$$\Phi_f(\Lambda) = \sum_{v \in \Lambda - \{0\}} f(v).$$

Since we can generate random lattices, we can talk about the expected value of $\Phi_f(\Lambda)$. We do this experimentally.

We are empirically confirming the following theorem of Siegel:

Theorem 22 (Siegel, 1945). *Suppose that $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a compactly-supported bounded measurable function. Then*

$$\int_{X_d} \Phi_f = \int_{\mathrm{SL}_d(\mathbb{R}) / \mathrm{SL}_d(\mathbb{Z})} \left(\sum_{v \in g\mathbb{Z}^d - \{0\}} f(v) \right) dg.$$

But we saw that for any $\Lambda \in X_d$, when f is the indicator function of a ball, we must have

$$\Phi_f(\Lambda) \in \{0, 2, 4, 6, \dots\}.$$

That's because balls are symmetric:

$$v \in \mathrm{supp}(f) \cap (\Lambda - \{0\}) \implies -v \in \mathrm{supp}(f) \cap (\Lambda - \{0\}).$$

If f is the indicator function of a ball of volume $2 - \varepsilon$, then this tells us that for any dimension d ,

$$\int_{X_d} \Phi_f = 2 - \varepsilon.$$

Conclusion: there exists some lattice $\Lambda \in X_d$ such that $\Phi_f(\Lambda) = 0$. Etc.

In the final part of the talk, we'll say a bit about more general spaces of lattices. Can we come up with a general integration formula for some $\mathcal{G}(\mathbb{R})/\Gamma$ for some algebraic $\mathbb{Q}$ -group $\mathcal{G}$ and an arithmetic subgroup Γ . If the group $\mathcal{G}$ is semisimple, we are automatically guaranteed that $\mathcal{G}(\mathbb{R})/\Gamma$ will be a probability space under the natural left-invariant Haar measure. This is a theorem of Borel and Harish-Chandra. In fact, this works for any group with no nontrivial rational characters.

For the higher moment case, what are the orbits of $\mathrm{SL}_d(\mathbb{Z})$ acting on $\mathbb{Z}^{d \times n}$? We work out the case $n < d$.

When n is much less than d , the higher moments start behaving like moments of a Poisson distribution. Rogers used this to prove that $c_d \geq \frac{1}{3}\sqrt{d}$. But he had already obtained in 1947 the stronger lower bound $c_d \geq d \cdot (2e^{-1})$.

6. JAMES BRANCH, *Explicit images of Shimura's map*

In one sentence, Shimura's map is a map from half-integral to integral weight modular forms:

$$S_{k/2} \rightarrow S_{2\lambda}.$$

For us, k and λ will be integers related by the equation

$$k = 2\lambda + 1.$$

We recall that a modular form of integral weight is given by

$$f : \mathbb{H} \rightarrow \mathbb{C}$$

such that

$$f(\gamma\tau) = (c\tau + d)^k f(\tau)$$

for every

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma.$$

We also ask that f vanishes at the cusps of Γ . We call the resulting space $S_k(\Gamma)$.

We recall next that a modular form of half-integral weight is given by something that transforms according to

$$\mathcal{J}(\gamma, \tau) = \underbrace{\varepsilon_d^{-1} \left(\frac{c}{d} \right)}_{\chi^{-1,4}} (c\tau + d)^{1/2},$$

$$\varepsilon_d = \begin{cases} 1 & \text{if } d \equiv 1 \pmod{4}, \\ -1 & \text{if } d \equiv 3 \pmod{4}. \end{cases}$$

We might write it as

$$\sum b(n)q^n \in S_{k/2}(\Gamma, \chi).$$

Theorem 23 (Shimura, 1973). *Let χ be a character modulo $4N$ and let $D > 0$ be squarefree. Let*

$$f = \sum_{n \geq 1} a_f(n)q^n \in S_{k/2}(\Gamma_0(4N), \chi).$$

Then there exists an element

$$g_D(w) = \sum_{n=0}^{\infty} c_D(n)e(nw) \in M_{2\lambda}(\Gamma_0(2N), \chi^2)$$

such that

$$L(g_D, s) = L(\chi_D, s - \lambda + 1) \sum_{n \geq 1} \frac{a_f(n)}{n^s}.$$

In other words, we take

$$B(n) = \sum_{d|n} \chi(d) \chi_{-4}^{\lambda}(d) d^{\lambda-1} b\left(\frac{n^2}{d^2}\right).$$

Question: given some explicit f , can we write the coefficients of g in a nice way?

Theorem 24 (Selberg). *Let $f \in S_{\lambda}(N, \chi)$ be primitive. Consider*

$$f(4\tau)\theta(\tau) \in S_{k/2}(4N, \chi\chi_{-4}).$$

Then

$$\sigma(f(4\tau)\theta(\tau)) = f(\tau)^2 - 2^{\lambda-1}\chi(2)f(2\tau)^2$$

lies in $S_{2\lambda}(2N, \chi^2)$.

Here

$$\theta(\tau) = \sum_{n \in \mathbb{Z}} q^{n^2} \in M_{1/4}(4).$$

Sketch of proof. The coefficient of the product are given by

$$b(n) = \sum_{m \in \mathbb{Z}} a\left(\frac{n-m^2}{4}\right).$$

Then

$$b(n^2) = \sum a_f\left(\frac{n^2-m^2}{4}\right). \quad (4)$$

After $n \mapsto 2m - n$, (4) becomes

$$b(n^2) = \sum_m a_f(n(m-n)).$$

Take f to be a newform. We can then write

$$a_f(n(m-n)) = \sum_{e|(m,n)} \mu(e) e^{\lambda-1} a_f\left(\frac{n}{e}\right) a_f\left(\frac{m-n}{e}\right).$$

We obtain

$$\sum_m \sum_{e|(m,n)}.$$

Then

$$\sum_{d|n} \sum_{e|\frac{n}{d}} \chi(de)(ed)^{\lambda-1} \mu(e) \chi_{0,4}(d) a_{f^2}\left(\frac{n}{ed}\right).$$

Then

$$a_{f^2}(n) = \sum \left(\sum a_f(m) a_f(n-m) \right) q^n.$$

□

Example 25. Let $f = \eta^2 \eta_{11}^2$. Then, since $\chi_{0,44}(2) = 0$, we have

$$\sigma(\eta_4^2 \eta_{44}^2 \theta) = \eta^4 \eta_{11}^4.$$

Next, we introduce h_ψ . Here

$$\eta(\tau) = q^{1/24} \prod_{n \geq 1} (1 - q^n), \quad \eta_{11}(\tau) = \eta(11\tau).$$

Next, let's introduce these h_ψ :

$$h_\psi(\tau) = \sum_{n \in \mathbb{Z}} n^\nu \psi(n) q^{n^2}.$$

Here $\nu = 0$ or $\nu = 1$, with $\psi(-1) = (-1)^\nu$. We have

$$h_\psi \in M_{1/2+\nu}(4r_\psi, \chi_{-1,4}^{-\lambda}).$$

We know that we can do functions of the form $f(4\tau)\theta(\tau)$. What happens if we work with a much more general theta function h_ψ ? That is to say, what happens for

$$f(4r_\psi\tau)h_\psi(\tau)?$$

We have the following:

Theorem 26 (Cipra, 1987). *Let f be as above. Then*

$$\sigma_1(f(4r\tau)h_\psi(\tau)) = g \otimes \psi(\tau) - 2^{\lambda+\nu-1} \chi(2) \psi(2) g \otimes \psi(2\tau)$$

lies in $S_{2(\lambda+\nu)}(2rN_r, \chi^2\psi^2)$, where $N_r := \text{lcm}(N, r)$.

Theorem 27 (Hansen, Naqvi, 2007). *Similar result, but without assuming that ψ is primitive.*

Theorem 28 (Brown, 2013). *Let $f \in M_\lambda(N, \chi)$ be a Hecke eigenform. Then*

(1)

$$\sigma(f(24\tau)\eta_{24}) = (f(\tau)f(6\tau) - f(2\tau)f(3\tau)) \otimes \left(\frac{12}{\cdot}\right)$$

satisfies (...).

What happens when one looks at stuff that is old on the half-integral weight side? For example, look at

$$f(4r_\psi\tau)h_\psi(D\tau),$$

where D is a fundamental discriminant. (...)

Theorem 29 (B. 2022). *Choose any N . Suppose that $f \in S_\lambda^{\text{new}}(N, \chi)$ such that $f\theta \in S_{k/2}(4N, \chi)$. Then*

$$\sigma(f\theta) = U_2 f^2.$$

Here

$$U_2\left(\sum_{n \geq 1} a(n)q^n\right) = \sum_{n \geq 1} a(2n)q^n.$$

Here $U_2 f^2 := (U_2(f))^2$.

7. SATYABRAT SAHOO, *On the solutions of $x^p + y^p = 2^r z^p$, $x^p + y^p = z^2$ over totally real fields*

An elliptic curve E is a nonsingular curve of dimension 1 with a fixed point $O \in E$.

- Any elliptic curve E/K defined over a field is given by a Weierstrass equation with $a_i \in K$.
- Let

$$\Delta_E := -b_2^2 b_8 - 8b_4^3 - 27b_6^2 + 9b_2 b_4 b_6,$$

$$j_E := \frac{c_4^3}{\Delta_E}.$$

8. RADU TOMA, *On the size of newforms of large level on $\text{PGL}(n)$*

Theorem 30 (Toma 2023+). *Under GRH, for a large prime number N , we have*

$$\|\varphi_{\Omega_N}\|_\infty \ll_{\Omega, n, \lambda, \varepsilon} N^{-1/4n+\varepsilon}$$

for all $n \geq 2$.

The actual relevant hypothesis is that

$$\sum_{p \asymp L} |\lambda(p, \varphi)| \gg_\varepsilon L^{3/4-\varepsilon}$$

for L a small power of $N \gg 1$.

Amplifier:

$$A(\varpi) = \sum_{p \asymp L} \lambda(p, \varpi) \overline{\lambda(p, \varphi)}.$$

Atkin–Lehner operator:

$$W_N : z \mapsto \text{diag}(1, \dots, 1, N) z^{-\top}.$$

9. FELICIEN COMTAT, *Sup norms of Siegel modular forms and Bocherer conjecture*

9.1. Motivation. Take G to be a reductive algebraic group over $\mathbb{Q}$, say. Let ϕ be a cusp form for G . In particular, ϕ has rapid decay at cusps. The complement of the cusps is a bounded region, so by continuity, ϕ is bounded, and in particular, L^2 . It's thus an interesting question to relate the L^∞ and L^2 norms.

Problem 1. Bound

$$\frac{\|\phi\|_\infty}{\|\phi\|_2}.$$

Methods:

- (1) Amplified pretrace formula.
- (2) (Steiner) Studying fourth moments via the theta correspondence.
- (3) Bound the Fourier expansion termwise.

The third approach is the subject of today's talk. It motivates the following further question.

Problem 2. How to understand the Fourier coefficients of ϕ ?

9.2. Framework. We denote by

$$\begin{aligned}\mathbb{H}_2 &:= \{z = x + iy \in M_2^{\text{sym}}(\mathbb{C}) : y > 0\}, \\ \text{GSp}_4(R) &:= \{g \in \text{GL}_4(R) : {}^\top g J g = \mu(g) J\}\end{aligned}$$

where

$$J = \begin{pmatrix} 0_2 & 1_2 \\ -1_2 & 0_2 \end{pmatrix}.$$

We also set

$$\text{Sp}_4(R) = \{g \in \text{GSp}_4(R) : \mu(g) = 1\}.$$

Then

$$\text{Sp}_4(\mathbb{R}) \curvearrowright \mathbb{H}_2$$

via the rule

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} Z = (aZ + b)(cZ + d)^{-1}.$$

We say that $F : \mathbb{H}_2 \rightarrow \mathbb{C}$ is a Siegel modular form if it satisfies the transformation property

$$F(\gamma z) = \det(cz + d)^k F(z)$$

for all $z \in \mathbb{H}_2$ and

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Sp}_4(\mathbb{Z}).$$

Fourier expansion:

$$f(z) = \sum_{S \in \mathcal{S}_2} a(S) e^{2\pi i \text{tr}(Sz)},$$

where

$$\mathcal{S}_2 := \left\{ S = \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix} : a, b, c \in \mathbb{Z}, S \geq 0 \right\}.$$

Then

$$\phi_F(z) := (\det y)^{k/2} F(z)$$

is bounded.

9.3. Fourier coefficients of Siegel modular forms.

Conjecture 31 (Reznikov–Saldana). *Take F a Siegel cusp form of weight k . Then*

$$a_F(S) \ll_{\varepsilon, F} (\det S)^{\frac{k}{2} - \frac{3}{4} + \varepsilon}.$$

9.3.1. *Saito–Kurokawa lifts.* F is a Saito–Kurokawa lift if there is a Hecke cusp form of half-integral weight $k - \frac{1}{2}$ in Kohnen’s plus space such that

$$h(z) = \sum_{m>0} c(m) m^{\frac{k}{2} - \frac{3}{4}} e(mz),$$

$$a_F(S) = (\det 2S)^{\frac{k}{2} - \frac{3}{4}} \sum_{d \mid \text{content}(S)} d^{1/2} c\left(\det\left(\frac{2S}{d}\right)\right).$$

If $D < 0$ is a fundamental discriminant, then

$$c(|D|)^2 = L\left(\frac{1}{2}, f \otimes \chi_D\right),$$

where f is the image of h by the Shimura correspondence. This implies that under GRH, we have

$$a_F(S) \ll (\det 2S)^{\frac{k}{2} - \frac{3}{4} + \varepsilon} (\text{content } S)^{1/2}. \quad (5)$$

Using this observation, Blomer proved:

Corollary 32 (Blomer). *If F is a Saito–Kurokawa lift, then*

$$\frac{\|\phi_F\|_{\infty}}{\langle F, F \rangle^{1/2}} \ll k^{3/4 + \varepsilon} \quad \text{under GRH.}$$

Idea of proof. The idea is as follows. Write $z = x + iy$. Then the contribution of $a_F(S)$ is negligible unless Sy is negligible, with eigenvalue

$$\frac{k}{4\pi} + O\left(\sqrt{k} \log k\right).$$

Now count such S and apply (5). □

9.3.2. General case.

Theorem 33 (C., Moyer–Balbesteros?, Saha). *Let k be even. Let F be a Siegel cusp form of weight k that is a Hecke eigenform, but not a Saito–Kurokawa lift. Then under GRH (although maybe GLH suffices), we have*

$$\frac{|a_F(S)|}{\langle F, F \rangle^{1/2}} \ll_{\varepsilon} \frac{(4\pi)^k}{\sqrt{\Gamma(2k-1)}} (\det 2S)^{\frac{k-1}{2} + \varepsilon} \text{content}(S)^{-1/2}.$$

Corollary 34 (C, M-B, S). *Under GRH, we have*

$$\frac{|\phi_F(z)|}{\langle F, F \rangle^{1/2}} \ll (\det Y)^{-1/4} k^{5/4 + \varepsilon}.$$

This $5/4$ improves upon the trivial bound, but not to the same extent as Blomer’s estimate. However, it is stronger higher in the cusp.

9.4. Bocherer conjecture and idea of proof. Let $A \in \mathrm{SL}_2(\mathbb{Z})$. Then F is invariant by the element

$$\begin{pmatrix} TA & 0 \\ 0 & A^{-1} \end{pmatrix} \in \mathrm{Sp}_4(\mathbb{Z}).$$

It follows that

$$a_F({}^tASA) = a_F(S).$$

Thus $a_F(S)$ depends only upon the orbit of S . If d is a fundamental discriminant., then

$$\{S \in \mathcal{S}_2 : \mathrm{disc}(S) = -4 \det S = d\} / \sim \longleftrightarrow \mathrm{Cl}_K, \quad K = \mathbb{Q}(\sqrt{d}).$$

Let Λ be a character of $\mathrm{Cl}_K(M)$. Set

$$R_{L,M}(F, K, \Lambda) = \sum_{S \in \mathrm{Cl}_K(M)} \Lambda^{-1}(S) a_F \left(L \begin{pmatrix} M & 0 \\ 0 & 1 \end{pmatrix} S \begin{pmatrix} M & 0 \\ 0 & 1 \end{pmatrix} \right).$$

Conjecture 35 (Bocherer, D-P-S-S, proved by Furasawa–Morimoto). *We have*

$$\begin{aligned} & \frac{64}{\langle F, F \rangle^{1/2}} |R(F, K, \Lambda)|^2 \\ &= \frac{(4\pi)^{2k-1}}{\Gamma(2k-1)} W(k) |d|^{k-1} \frac{L(\frac{1}{2}, \pi_F \times \mathrm{AI}(\Lambda^{-1}))}{L(1, \pi_F, \mathrm{Ad})} \prod_{p|C(\Lambda)} \mathcal{J}_p(\phi_F^{L,M}, \Lambda_p), \end{aligned}$$

where $\phi_F^{L,M}$ is a certain translate of ϕ_F .

Set

$$J_p(\phi_p^{L,M}, \Lambda_p) = \int_{\mathbb{Q}_p^*} \Lambda^{-1}(t) J_p(\phi^{L,M}, t) dt.$$

Then

$$J_p(\phi, t) = \lim_{k \rightarrow \infty} \int_{M_2^{\mathrm{sym}}(p^{-k}\mathbb{Z})} \Phi_\phi \left(\begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix} t \right) \psi(\mathrm{trace}(Sx))^{-1} dx.$$

10. WING HONG (JOSEPH) LEUNG, *Short second moment bounds and subconvexity for $\mathrm{GL}(3)$ L -functions*

(Joint work with Keshav Aggarwal and Ritabrata Munshi)

Riemann zeta function. Good times.

L -functions. Even better.

Theorem 36 (Aggarwal–L.–Munshi).

$$\int_{t-t^{2/3}}^{t+t^{2/3}} |L(\tfrac{1}{2} + iv, \pi)|^2 dv \ll_{\pi, \varepsilon} t^{3/2-1/4+\varepsilon}.$$

Using a simple argument similar to Good's, we deduce:

Corollary 37.

$$L(\tfrac{1}{2} + it, \pi) \ll_{\pi, \varepsilon} t^{3/4-1/8+\varepsilon}.$$

Remark 38. Pal used the same argument by decomposing the full second moment to get

$$\int_t^{2t} |L(\tfrac{1}{2} + iv, \pi)|^2 dv \ll_{\pi, \varepsilon} t^{3/2-3/1676+\varepsilon}. \quad (6)$$

Remark 39. The speaker is currently working on analogous subconvexity bound for L -functions $\mathrm{GL}(3)$ twisted by characters χ of factorizable modulus $P_1 P_2$.

Remark 40. Ikuyo Kaneko and Maksym Radziwiłł are showing (maybe not yet on the arxiv) something like (6), without decomposing into shorter moments.

Proof idea? Short second moment yields shifted sum problem, with some average over shifts. Ingredients:

- δ -method (“standard recipe”).
- Duality trick.
- “ ∞ -Cauchy Schwarz”, some sort of van der Corput trick.

We want to bound

$$\int_{\mathbb{R}} U\left(\frac{v}{M}\right) |L(\tfrac{1}{2} + it + iv, \pi)|^2 dv.$$

The final optimal choice for subconvexity is $M = t^{2/3}$.

Approximate functional equation yields

$$\sum_{N \ll t^{3/2+\varepsilon}} \frac{1}{\sqrt{N}} \sum_{n \sim N} \lambda(n) n^{-it-iv}.$$

For sketch, focus on $N \sim t^{3/2}$. Squaring and integrating, we arrive at

$$S = \frac{M}{t^{3/2}} \sum_{h \ll t^{3/2+\varepsilon}/M} \sum_{n \sim t^{3/2}} \lambda(n) \overline{\lambda(n+h)} e\left(\frac{th}{2\pi n}\right).$$

Compare with

$$\sum_{h \ll t^{1/2}} \sum_{n \asymp t^{3/2}} \lambda(n) \lambda(n+h).$$

We didn’t know how to do that, but now Ikuya and Maksym have announced that they can.

To study S , we apply the delta method to detect the condition $m = n + h$. For $Q > 1$, get

$$\begin{aligned} & \frac{M}{Q^2 t^{3/2}} \sum_{q \leq Q} \sum_{h \ll t^{3/2+\varepsilon}/M} \sum_{n \asymp t^{3/2}} \lambda(n) \sum_{m \asymp t^{3/2}} \overline{\lambda(m)} \\ & \sum_{a(q)^*} e_q(a(n+h-m)) e\left(\frac{th}{2\pi n}\right) \int U(x) e\left(\frac{(n+h-m)x}{qQ}\right) dx. \end{aligned}$$

For sketch, focus on $q \asymp Q$ and $x \asymp 1$.

Take $Q = \sqrt{M} t^{1/4}$. This corresponds to balancing

$$\frac{th}{2\pi n} \asymp \frac{nx}{qQ}.$$

Poisson on h , Voronoi on m and n .

11. SOUMENDRA GANGULY

12. PHILLIP HARRIS, *Maximal flat periods for $\mathrm{SL}(n)$*

Let $G = \mathrm{SL}(n, \mathbb{R}) = NAK$. We can write

$$\mathfrak{p} = \mathfrak{n} + \mathfrak{a} + \mathfrak{k} = \mathfrak{p} + \mathfrak{k},$$

where $\mathfrak{p}$ is the space of symmetric matrices.

Let $\Gamma < G$ be a discrete subgroup. Set

$$X = \Gamma \backslash G / K.$$

Let $Y \subset X$ be a maximal flat, i.e., a translate gA of the diagonal subgroup. Fix $b \in C_c^\infty(Y)$. Let f_i be an orthonormal basis of Maass forms, with spectral parameter ν_i . The period of interest is then

$$P_i = \int_Y f_i(g) b(g) dg.$$

Pick $\nu \in \mathfrak{a}_{\mathbb{C}}^*$. Sum over all spectral parameters in a ball of radius one around ν :

$$\sum_{\|\Re \nu - \nu_i\| \leq 1} |P_i|^2.$$

Bart Michels (2022): for a closed cone

$$D^{\text{gen}} \subset (\mathfrak{a}^*)^{\text{gen}}$$

(where gen means not singular, i.e., $\langle \nu, \alpha \rangle \neq 0$, and also $\nu \notin \text{span}\{\alpha_1, \dots, \alpha_{n-2}\}$), we have

$$\sum_{\|\Re \nu - \nu_i\| \leq 1} |P_i|^2 \ll_b \beta(\nu) \|\nu\|^{-n+1}.$$

The speaker shows in 2023 a result valid for general spectral parameters:

$$\sum_{\|\Re \nu - \nu_i\| \leq 1} |P_i|^2 \ll_b \beta(\nu) \|\nu\|^{-n+1} \begin{cases} \left(\log \frac{\|\nu\|}{1+|\nu_2+\nu_{n-1}|} \right)^{n-2} & n \neq 4, \\ \left(\log \frac{\|\nu\|}{1+\dots} \right) & \text{if } n = 4. \end{cases}$$

Here we order so that $\nu_1 \leq \dots \leq \nu_n$.

Let

$$k \in C_c^\infty(K \backslash G / K).$$

We can take its support as small as we want.

$$\int_Y \int_Y \sum_i \hat{k}(\nu_i) f_i(x) \overline{f_i(y)} b(x) b(y) dx dy = \int_Y \int_Y k(x^{-1}y) b(x) b(y) dx dy.$$

$$k(z) = \frac{1}{|W|} \int_{\mathfrak{a}^*} \beta(\lambda) \varphi_\lambda(z) \hat{k}(\lambda) d\lambda,$$

$$\varphi_\lambda(g) = \int_{\text{SO}(n)} e^{(i\lambda - \rho)H(kg)} dg,$$

$$H : G \rightarrow \mathfrak{a},$$

$$g \in Ne^{H(g)} K.$$

Theorem 41 (Duistermaat). *There exists $a \in C^\infty(\mathfrak{p})$ so that for all $X \in \mathfrak{p}$, and with $\pi : \mathfrak{p} \rightarrow \mathfrak{a}$ the projection, we have*

$$\varphi_\lambda(e^X) = \int_{\text{SO}(n)} e^{i\lambda\pi(\text{Ad } kX)} a(\text{Ad } kX) dX.$$

So now we can rewrite something as

$$\int_{\mathfrak{a}^*} \beta(\lambda) \hat{k}(\lambda) \int_{\mathfrak{a}} e^{i\lambda\pi(\text{Ad } kX)} \underbrace{a(\text{Ad } kX) b(e^X)}_{C(K,X)} dX d\lambda.$$

13. YIANNIS PETRIDIS, *Counting and equidistribution*

Outline:

- (i) Motivation/example
- (ii) Lattice problems in hyperbolic space. (The above was all beyond 2016.)
- (iii) After 2016
- (iv) Equidistribution

To start, we'll look at the following sum:

$$\sum_{n \leq x-2} a_n r(n) r(n+4),$$

$$a_n = \begin{cases} 1 & \text{if } n \text{ is even,} \\ 1/2 & \text{if } n \text{ is odd.} \end{cases}$$

Here $r(n)$ counts the number of representations of n as a sum of two squares.

The above sum suggests writing

$$n = t^2 + s^2, \quad n + 4 = r^2 + u^2, \quad (r, u, s, t) \in \mathbb{Z}^4.$$

Then

$$1 = \frac{r+s}{2} \frac{r-s}{2} - \frac{t+u}{2} \frac{t-u}{2},$$

$$1 = ad - bc. \tag{7}$$

If n is even, then a, b, c, d will be integers. If n is odd, then r and u (resp. t and s) will have different parities, so by swapping t and s as necessary, we will get an integer solution. That will happen with the swapping half the times.

$$(a, b, c, d) \in \mathbb{Z}^4.$$

$$a^2 + b^2 + c^2 + d^2 = n + 4 \leq x. \tag{8}$$

Write $N(x)$ for the number of quadruples satisfying (7) and (8). Write

$$u(z, w) = \frac{|z - w|^2}{4\Im z \Im w}.$$

Thus, with

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

we have

$$u(\gamma i, i) = \frac{a^2 + b^2 + c^2 + d^2 - 2}{4}.$$

The hyperbolic lattice point problem is to estimate

$$N(x, i, i) = \# \{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) : 4 \cosh u(\gamma i, i) + 2 \leq x \}.$$

More generally,

$$N(x, z, w) = \# \{ \gamma \in \Gamma : 4 \cosh u(\gamma z, w) + 2 \leq x \}.$$

So we have the upper half-plane, and all the points in $i + \mathbb{Z}$. Take a geodesic circle containing i of radius R . Then its area is

$$A(R) = 4\pi \left(\sinh \frac{R}{2} \right)^2 \asymp e^R,$$

while the circumference is

$$C(R) = 2\pi \sinh R \asymp e^R.$$

Heinz Huber (Basel) wrote a set of papers (1969 Math. Annalen) to explain the relation of the problem to automorphic forms. What results is that we have a main term and an error term:

$$N(x, z, w) = M(x, z, w) + E(x, z, w),$$

$$M(x, z, w) = \sum_{1/2 \leq s_j \leq 1} \sqrt{\pi} \frac{\Gamma(s_j - 1/2)}{\Gamma(s_j + 1)} u_j(z) u_j(w) x^{s_j} \sim \frac{\pi}{\text{vol}(\Gamma \backslash \mathbb{H})} x.$$

The s_j correspond to eigenfunctions

$$\Delta u_j + s_j(1 - s_j)u_j = 0, \quad \Delta = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right).$$

Selberg (unpublished notes, 1972) proved that

$$E(x, z, w) = O(X^{2/3}).$$

Patterson (1972) improved the exponent to $3/4$. P. Gunther (1980) improved it further to $2/3$. A. Good (1983) obtained related results to those of Gunther that we'll talk more about later. Conjecturally, one can take $O_\varepsilon(X^{1/2+\varepsilon})$. Phillips–Rudnick established

$$O\left(X^{1/2}(\log \log X)^{1/4-\delta}\right).$$

Let's talk about Good's work. There are three types of subgroups of $\text{SL}_2(\mathbb{R})$:

- elliptic subgroups K ,
- hyperbolic subgroups $\mathcal{H}$, and
- parabolic subgroups P .

Let H, H_1, H_2 be any of the above subgroups. Good was interested in double cosets by these kinds of subgroups:

$$\Gamma \cap H_1 \backslash \Gamma / \Gamma \cap H_2.$$

Chamizo 1996:

$$\frac{1}{X} \int_X^{2X} E(x)^2 dx \ll X^1 \log^2 X.$$

Let's consider some cases.

Elliptic-hyperbolic. $\cosh \rho(z, \ell) = \frac{1}{\cos \theta}$.

With

$$X = \frac{\sinh(t/2)}{\sinh(\mu/2)}.$$

$$\begin{aligned} N(t) &= \# \{ \gamma \in (\Gamma \cap \mathcal{H}) \backslash \Gamma : \rho(\gamma z, \ell) \leq t \} \\ &= \sum_{s_j \leq 1} (\Gamma\text{-factors}) u_j(z) \int_{\ell} u_j X^{s_j} + \underbrace{E(X, \mathcal{H}, z)}_{=O(X^{2/3})}. \end{aligned}$$

The conjecture is that one can take $O(X^{1/2+\varepsilon})$. There's an integral transform here, the one introduced by Huber in his paper in 1988. It's a special case of the Jacobi transform, which covers just about everything in this area of mathematics.

Chatzakos–P 2016:

$$\frac{1}{X} \int_X^{2X} |E(X, \mathcal{H}, z)|^2 dx \ll X^1 \log^2 X.$$

Chatzakos 2019: $O(X^{1/2})$.

Is there any arithmetic interpretation of this counting problem?
Hyperbolic-hyperbolic.

14. MICHAEL YIASSEMIDIS, *Lattice points in elliptic annuli*

Sums of squares.

$$r(n) := \# \{ (a, b) \in \mathbb{Z}^2 : n = a^2 + b^2 \} = 4 \sum_{\substack{d|n \\ d \equiv 1, 3(4)}} (-1)^{\frac{d-1}{2}}.$$

This implies that $p = a^2 + b^2$ if and only if $p \equiv 1(4)$ (Fermat).

For $u, v \geq 0$ integers,

$$r_{u,v}(n) := \# \{ (a, b) \in \mathbb{Z}^2 : ua^2 + vb^2 = n \}.$$

A special case is when $n = p$ is prime, with $u = 1$.

We have

$$\begin{aligned} p = a^2 + 2b^2 &\iff p \equiv 1, 3(8), \\ p = a^2 + 3b^2 &\iff p \equiv 1(3) \quad (\text{or } p = 3), \\ p = a^2 + 14b^2 &\iff \left(\frac{-14}{p} \right) = 1 \text{ and } \exists \text{ solution } x \text{ to } (x^2 + 1)^2 - 8 \equiv 0(p). \end{aligned}$$

For all $v > 0$, there is an irreducible $f_v(x) \in \mathbb{Z}[x]$ such that

$$p = a^2 + vb^2 \iff \left(\frac{-v}{p} \right) = 1 \text{ and } \exists \text{ solution to } f_v(x) \equiv 0(p).$$

A reference is David A. Cox, *Primes of the form $x^2 + ny^2$* , which uses methods such as class field theory and Shimura reciprocity.

How about arithmetic functions?

$$r(n) = \# \{ (a, b) \in \mathbb{Z}^2 : n = a^2 + b^2 \},$$

$$\begin{aligned} \sum_{n=1}^x r(n) &= \# \{ (a, b) \in \mathbb{Z}^2 : a^2 + b^2 \leq x \} \\ &= \pi x + O(\sqrt{x}), \end{aligned}$$

$$\begin{aligned} \sum_{n=1}^x r_{u,v}(n) &= \# \{ (a, b) \in \mathbb{Z}^2 : ua^2 + vb^2 \leq x \} \\ &= \frac{\pi}{\sqrt{uv}} h + E(x, h), \quad E(x, h) = O(\sqrt{x}). \end{aligned}$$

We want to take our error and understand what happens when we average it:

$$\frac{1}{z} \sum_{x=z}^{2z} E(x, h) = \frac{1}{z} \sum_{x=z}^{2z} \left(\sum_{n=x}^{x+h} r_{u,v}(n) - \frac{\pi h}{\sqrt{uv}} \right),$$

and also the variance

$$\frac{1}{z} \sum_{x=z}^{2z} E(x, h)^2 = \frac{1}{z} \sum_{x=z}^{2z} \left(\sum_{n=x}^{x+h} r_{u,v}(n) - \frac{\pi h}{\sqrt{uv}} \right)^2.$$

Four regimes.

- h constant

- $h = h(x) \rightarrow 0$ (Wigman 2005)

$$\sum_{n=x}^{x+h} r_{u,v}(n) = \# \{ (a, b) \in \mathbb{Z}^2 : x \leq ua^2 + vb^2 \leq x+h \}.$$

Analogy between integers and function fields.

Integers	Function Fields
$\mathbb{Z}$	$\mathcal{A} := \mathbb{F}_q[T]$
positives	$A = a_0 + a_1 + \cdots + a_k T^k$
positive	monic
primes	irreducibles
$ n = \mathbb{Z}/n\mathbb{Z} $	$ A = \mathcal{A}/A\mathcal{A} $
$\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$	$\zeta_q(s) = \sum_{A:\text{monic}} A ^{-s}$

We also use the notation, for a polynomial A as above,

$$[A]_k = \begin{pmatrix} a_0 \\ a_1 \\ \dots \\ a_k \end{pmatrix}.$$

We have

$$r_{U,V}(N) = \# \{ (A, B) \in \mathcal{A} : N = UA^2 + VB^2 \},$$

where U and V are fixed monics.

The mean is

$$\frac{1}{q^k} \sum_{\substack{X:\text{monic} \\ \deg X=k}} \sum_{|N-x| < q^h} r_{u,v}(N) = \frac{q^h}{\sqrt{|uv|}}.$$

The variance is

$$\frac{1}{q^k} \sum_{\substack{X:\text{monic} \\ \deg X=k}} \left(\sum_{|N-x| < q^h} r_{u,v}(N) - \frac{q^h}{\sqrt{|uv|}} \right)^2. \quad (9)$$

Solutions to

$$N = UA^2 + VB^2,$$

where

- N is over a “short interval”,
- A, B over intervals of $\deg \leq k$,

are given by

$$\sum_{N,A,B} \mathbf{1}(N - UA^2 - VB^2).$$

The idea is now to take the Fourier expansion of the indicator function. Each of the resulting functions is additive, so we can look separately at the contribution of each parenthetical term. We use

$$\psi : \mathbb{F}_q \rightarrow \mathbb{C}^*,$$

$$\psi(x+y) = \psi(x)\psi(y),$$

$$\mathbf{1}(M) = \sum_{\underline{\alpha} \in \mathbb{F}_q^k} \psi(\underline{\alpha} \cdot [M]_k).$$

This gives the expansion

$$\sum_{\underline{\alpha} \in \mathbb{F}_q^{\kappa-1}} \sum_N$$

]

This gives for the variance (9) the formula

$$\begin{cases} q^k \left(\frac{k}{2} - h \right) \frac{d(uv)}{|uv|^2} \phi(uv) & \text{if } h < \frac{\kappa}{2}, \\ 0 & \text{if } h > \frac{\kappa}{2}. \end{cases}$$

15. PRANENDU DARBAR, *Asymmetric distributions of the extreme values for cubic L-functions on the 1-line*

16. JACKIE VOROS, *On the average least negative Hecke eigenvalue*

16.1. **Motivation.** Recall the Legendre symbol

$$\left(\frac{a}{p} \right) = \begin{cases} 1 & \text{if } a \text{ is a quadratic residue,} \\ -1 & \text{if } a \text{ is not a quadratic residue,} \\ 0 & \text{if } p|a. \end{cases}$$

Totally multiplicative. Quadratic reciprocity:

$$\left(\frac{p}{2} \right) \left(\frac{2}{p} \right) = (-1)^{\frac{p-1}{2} \frac{q-1}{2}}$$

for p and q odd primes.

Question 42. What is

$$n_2(p) := \text{least } n \text{ such that } \left(\frac{n}{p} \right) = -1.$$

Burgess (1957):

$$n_2(p) \ll p^{\frac{1}{4\sqrt{\epsilon}} + \epsilon}.$$

Linnik (1942):

$$\# \{p < x : n_2(p) > x^\epsilon\} \ll_\epsilon 1.$$

Conjecture 43 (Vinogradov). $n_2(p) = O(p^\epsilon)$.

How about the average value, namely

$$\lim_{x \rightarrow \infty} \frac{1}{\pi(x)} \sum_{p < x} n_2(p)?$$

A heuristic: $n_2(p)$ is prime, and

$$\# \left\{ n : \left(\frac{n}{p} \right) = \pm 1, n \in [1, p-1] \right\} = \frac{p-1}{2}.$$

Assume that

$$\text{Prob} \left(\left(\frac{q}{p} \right) = \pm 1 \right) = \frac{1}{2}$$

for p and q prime, independently. Then

$$\text{Prob}(n_2(p) = p_k) = \prod_{j=1}^k \text{Prob} \left(\left(\frac{p_j}{p} \right) = +1 \right) = \frac{1}{2^k}.$$

This is in fact what you get:

Theorem 44 (Erdos 1961).

$$\lim_{x \rightarrow \infty} \frac{1}{\pi(x)} \sum_{p < x} n_2(p) = \sum_{k=1}^{\infty} \frac{p_k}{2^k}.$$

Take

$$f \in S_k^{\text{new}}(\Gamma_0(N)) = S_k^{\text{new}}(N),$$

a normalized eigenform. The set of newforms is finitely-generated. We denote by $H_k^*(N)$ a finite generating set. Each f in this set has the following Fourier expansion:

$$f(z) = \sum_{n \geq 1} \lambda_f(n) n^{\frac{k-1}{2}} e(nz), \quad e(nz) := e^{2\pi i n z}.$$

Here the $\lambda_f(n)$ are Hecke eigenvalues, and they also have a few nice properties much like the Legendre symbol.

- They are (weakly) multiplicative:

$$\lambda_f(nm) = \lambda_f(n)\lambda_f(m)$$

for $(n, m) = 1$.

- We have, e.g.,

$$\lambda_f(p)^2 = 1 + \lambda_f(p^2).$$

- Deligne's bound:

$$|\lambda_f(n)| \leq \tau(n),$$

where $\tau(n)$ denotes the divisor function. Importantly, when n is prime, this says that

$$|\lambda_f(p)| \leq 2.$$

We may thus associate some angle

$$\theta_f(p) \in [0, \pi]$$

such that

$$\lambda_f(p) = 2 \cos \theta_f(p).$$

Question 45. When is the first sign change of $\lambda_f(p)$, namely

$$p_f := \text{least prime such that } \lambda_f(p_f) < 0?$$

Theorem 46 (V). *We have*

$$\lim_{N \rightarrow \infty} \frac{1}{|H_k^*(N)|} \sum_{f \in H_k^*(N)} p_f = \sum_{i=1}^{\infty} \frac{p_i}{2^i},$$

where p_i denotes the i th prime.

Where does this come from? Let's give some evidence or motivation. Let's write

$$n_f := \text{least } n \text{ such that } \lambda_f(n) < 0.$$

(Note that this need not coincide with p_f , because we do not have total multiplicativity.) Then:

- $\lambda_f(n)$ changes sign infinitely often.
- Kohnen, Sengupta, Iwaniec, Kowalski, Lau, Soundararajan, Wu, Matomaki:

$$n_f \ll Q^{3/8}, \quad Q = k^2 N.$$

- Sato–Tate: the numbers $\theta_f(p)$ defined by

$$\lambda_f(p) = 2 \cos \theta_f(p)$$

are equidistributed with respect to the Sato–Tate measure

$$\mu_{ST} = \frac{2}{\pi} \sin^2 \theta d\theta.$$

In particular, half the time the eigenvalue should be positive, the other half, negative.

We will now do a very very quick sketch proof that will maybe shed light on the fixing of the k and the prime condition for n . We will use Sato–Tate, which means that we can say that the size of the set

$$\frac{\#\{f \in H_k^*(N) : \varepsilon_p \lambda_f(p) > 0, p \leq z, p|N\}}{\#|H_k^*(N)|} \rightarrow \mu_{ST}([0, \pi/2])^{\pi_N(z)} = \left(\frac{1}{2}\right)^{\pi_N(z)}, \quad (10)$$

where

$$\pi_N(z) := \#\{\text{primes } p \nmid N\}.$$

Let (ε_p) be a sequence of signs. Then, letting z tend to infinity slowly enough, we deduce that

$$\sum p_f = \sum_{p_i \leq z} p_i \cdot G,$$

where G denotes the numerator on the left hand side of (10).

Lau, Wu 2008:

$$\sum_{f \in H_k^*(N)} \left| \sum_{P \leq p \leq 2P} b_p \frac{\lambda_f(p)}{p} \right|^{2j} \ll k\varphi(N) \left(\frac{384B^2}{P \log P} \right)^j + (kN)^{10/11} \left(\frac{10BQ^{1/10}}{\log P} \right)^{2j}.$$

Here $(b_p)_p$ are real numbers satisfying the bound $|b_p| \leq B$. Basically, we want to count all forms with a sequence of signs that make the summands positive up to some point. We thus take

$$\mathcal{E} := \{f : \varepsilon_p \lambda_f(p) \geq 0, p \in [P, 2P], p \nmid N\}.$$

We can then consider the sum

$$\sum_{f \in \mathcal{E}} \left| \sum_{\substack{P < p \leq 2P \\ p \nmid N}} \frac{\lambda_f(p)^2}{p} \right|^{2j} \leq (\text{apply large sieve}).$$

Then, using that

$$\lambda_f(p)^2 = 1 + \lambda_f(p^2),$$

we arrive at considering

$$\sum_{\substack{P < p \leq 2P \\ p \nmid N}} \frac{1}{p} = \sum_{P \leq p \leq 2P} \frac{1}{p} - \sum_{\substack{P < p \leq 2P \\ p|N}} \frac{1}{p} \geq \frac{\delta}{2 \log P} - \frac{\omega(N)}{p}.$$

(We want to take $P = 2^i z$, where $i = 0, 1, \dots$. But z itself depends upon N , which makes the analysis complicated.)

In the remaining two minutes, we'll explain briefly how we implement this.

Corollary 47. *For N prime, $k \geq 2$ (not fixed), and $z < \beta \leq \log kN$, the set*

$$\{f \in H_k^*(N) : p_f \ll \beta\} \ll kN \exp\left(-C \frac{\beta}{\log \beta}\right),$$

where $C > 0$.

Remark 48. Technically we could take $\omega(N)$ to be P -smooth, but we end up taking P to be very large. We thus don't need N to be prime.

17. MIN LEE, *An extension of converse theorems*

17.1. Modular forms. We start with some notation.

$$\mathrm{GL}_2^+(\mathbb{R}) \curvearrowright \mathbb{H} = \{x + iy : x \in \mathbb{R}, y > 0\}.$$

For $k \in \mathbb{Z}$,

$$(h|_k \gamma) := (\det \gamma)^{k/2} (cz + d)^{-k} h(\gamma z) \quad \text{for } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

ϕ should be a holomorphic function on the upper half-plane, $k \in \mathbb{Z}_{\geq 1}$, $\Gamma_0(n)$, χ .

$$\int_0^1 \phi(x + iy) dx = 0 \quad (\text{cuspidal}),$$

$$\phi|_k \gamma = \chi(d) \cdot \phi.$$

$S_k(N, \chi)$.

$$\phi(z) = \sum_{n=1}^{\infty} a_{\phi}(n) n^{\frac{k-1}{2}} e(nz), \quad e(z) = e^{2\pi i z}.$$

Here we know the Ramanujan conjecture: $a_{\phi}(n) \ll n^{\varepsilon}$.

From such a modular form ϕ we can construct L -functions

$$L(s, \phi) = \sum_{n=1}^{\infty} \frac{a_{\phi}(n)}{n^s} \quad (\Re(s) > 1).$$

$$\begin{aligned} \Lambda(s, \phi) &= \Gamma_{\mathbb{C}} \left(s + \frac{k-1}{2} \right) L(s, \phi) \\ &= i^k N^{\frac{1}{2}-s} \Lambda(1-s, \tilde{\phi}), \end{aligned}$$

where

$$\tilde{\phi} = \phi|_k w_N, \quad w_N = \begin{pmatrix} 0 & -1/\sqrt{N} \\ \sqrt{N} & 0 \end{pmatrix}.$$

This is proved by using that

$$\phi(iy) = (-iy)^k \phi\left(-\frac{1}{Ny}\right)$$

and using the integral representation

$$\Lambda(s, \phi) = 2 \int_0^{\infty} \phi(iy) y^{s+\frac{k-1}{2}} \frac{dy}{y}.$$

Converse theorems address the question: given sequences f_n and $\tilde{f}_n$ indexed by $n \geq 1$ whose associated Dirichlet series satisfy certain conditions, such as

- $f_n, \tilde{f}_n = O(n^{\sigma})$, $\sigma > 0$, and
- analytic continuation and functional equation,

must they come from something in $S_k(N, \chi)$?

Hecke (1936): $N \geq 4$.

$$\{f_n\} \rightsquigarrow f(z) = \sum_{n=1}^{\infty} f_n \cdot n^{\frac{k-1}{2}} e(nz).$$

Then

$$f|_k T = f, \quad T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Also, the condition

$$\tilde{f}(iy) = f|_k w_N(iy)$$

implies that

$$f|_k W_N = f, \quad W_N := w_N T w_N^{-1} = \begin{pmatrix} 1 & 0 \\ N & 1 \end{pmatrix}.$$

These matrices generate $\Gamma_0(N)$ for $N \leq 4$. For $N = 5$, one needs an additional generator. For instance,

$$\Gamma_0(5) = \left\langle T, w_5 T w_5^{-1}, \begin{pmatrix} 2 & -1 \\ 5 & -2 \end{pmatrix} \right\rangle.$$

We can consider the twisted L -functions

$$L(s, f, \psi) := \sum_{n=1}^{\infty} \frac{f_n \psi(n)}{n^s}, \quad \psi : \text{Dirichlet character modulo } q.$$

Weil (1967): if $L(s, f, \psi)$ satisfy analytic continuation and functional equation for all primitive characters ψ modulo q (prime) and all $q \nmid N$, then $f \in S_k(N, \chi)$.

Razar (1977): it is enough to know that $L(s, f, \psi)$ satisfies the above for all characters ψ (not necessarily primitive) with modulus q in $N\mathbb{Z}_{\geq 1}$ satisfying $N \leq q \leq N^2$.

Suppose we know just a single functional equation. Then for $N \geq 5$, the space of solutions, i.e., pairs of sequences $(f_n)_{n \geq 1}$ and $(\tilde{f}_n)_{n \geq 1}$ satisfying the relevant condition, is infinite-dimensional. For an explicit “counterexample” to the naive converse theorem, see the paper of Raphael Steiner (2019).

Conjecture 49 (Farmer, Koutsalios, Lemuell, 2008). *Assume single functional equation and that there is an Euler product, i.e., that*

$$\Lambda(s, f) = \Gamma_{\mathbb{C}} \left(s + \frac{k-1}{2} \right) \prod_{p \nmid N} (1 - f_p p^{-s} + \chi(p) p^{-2s})^{-1}.$$

Then we should have $f \in S_k(N, \chi)$.

Remark 50. By assuming the existence of an Euler product, we can eliminate the known counterexamples.

BBBCMOPSL (2018): consider, instead of an additive twist $\psi(n)$, a twist by the Ramanujan sum

$$c_q(n) = \sum_{a(q)} e \left(n \frac{a}{q} \right).$$

We assume knowledge about such twists for all q with $(q, N) = 1$ and get a converse theorem. We conjecture:

Conjecture 51. *If $L(s, f, c_q)$ have analytic continuation, functional equation, and moderate growth, then for all q with $(q, N) = 1$, we have $f \in S_k(N, \chi)$.*

17.2. Petersson trace formula. Let $H_k(N, \chi) \subset S_k(N, \chi)$ be a basis, and let g be an element of that basis. Write

$$g(z) = \sum_{n=1}^{\infty} \rho_g(n) n^{\frac{k-1}{2}} e(nz).$$

We have $n, m \in \mathbb{Z}_{>0}$,

$$\begin{aligned} & \frac{(4\pi)^{k-1}}{\Gamma(k-1)} \sum_{g \in H_k(N, \chi)} \rho_g(n) \rho_g(m) \\ &= \delta_{n=m} + 2\pi i^{-k} \sum_{q \in N\mathbb{Z}_{\geq 1}} \frac{S_{\chi}(m, n, q)}{q} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{q} \right), \end{aligned}$$

where we recall that

$$S_{\chi}(m, n, q) = \sum_{\substack{u, v(q) \\ uv \equiv 1(q)}} e_q(mu + nv) \chi(u).$$

Rankin–Selberg convolutions:

$$L(s, \phi_1 \times \phi_2) = \zeta^{(N)}(2s) \sum_{m=1}^{\infty} \frac{\lambda_{\phi_1}(m) \overline{\lambda_{\phi_2}(m)}}{m^s}.$$

At $s = 1$, one gets the residue $\langle \phi_1, \phi_2 \rangle$. We obtain

$$\begin{aligned} & \frac{(4\pi)^{k-1}}{\Gamma(k-1)} \sum_{g \in H_k(N, \chi)} \rho_g(n) L(s, f \times \bar{g}) \\ &= f_n n^{-s} \zeta^{(N)}(2s) + 2\pi i^{-k} \zeta^{(N)}(2s) \sum_{q \in N\mathbb{Z}} \frac{1}{q} \sum_{m=1}^{\infty} \frac{f_m S_{\chi}(n, m, q)}{m^s} J_{k-1} \left(\frac{4\pi\sqrt{mn}}{q} \right). \end{aligned}$$

We can write

$$J_{k-1}(4\pi y) = \frac{1}{2\pi i} \int_{(\sigma)} \frac{\Gamma_{\mathbb{C}}(u + \frac{k-1}{2})}{\Gamma_{\mathbb{C}}(-u + \frac{k-1}{2})} y^u \frac{dy}{y}.$$

Venkatesh (2002): for $N = 1$, and assuming finitely many poles of these L -functions, we can recover the converse theorem. Here we assume that

$$\Lambda(s, f, \alpha) = \Gamma_{\mathbb{C}} \left(s + \frac{k-1}{2} \right) L(s, f, \alpha), \quad \alpha = \frac{a}{q}.$$

Theorem 52 (A. Booker, M. Farmer, L 2023). *Let $(f_n)_{n \geq 1}$ and $(\tilde{f}_n)_{n \geq 1}$ such that*

$$L(s, f) := \sum_{n \geq 1} f_n n^{-s}$$

is absolutely convergent for $\Re(s) > 1$. Set

$$\gamma(s) = Q^s \prod_{j=1}^r \Gamma(\lambda_j s + \mu_j),$$

where $\lambda_j > 0$, $\mu_j \in \mathbb{C}$ and $\sum_{j=1}^r \lambda_j = 1$. Let $\alpha = u/q \in \mathbb{Q}$ and $Q > 0$. For $\Re(\mu_j) > -\frac{1}{2}\lambda_j$, assume the functional equation

$$\Lambda(s, f, \alpha) = \gamma(s) \sum_{n=1}^{\infty} \frac{f_n e(n\alpha)}{n^s} = w \cdot \chi(v) q^{1/2-s} \Lambda(1-s, \tilde{f}, -\frac{v}{q}),$$

where $q \in N\mathbb{Z}_{\geq 1}$. Then there exists $k \in \mathbb{Z}_{\geq 1}$ such that

$$f(z) = \sum_{n=1}^{\infty} f_n e(nz) \in M_k(N, \chi).$$

If $k \geq 2$, then it lies in $S_k(N, \chi)$.

18. JUNXIAN LI, *Variations of the Titchmarsh divisor problem*

Titchmarsh divisor problem: find an asymptotic formula for

$$\sum_{p \leq x} \tau(p - a), \quad x \rightarrow \infty. \quad (11)$$

Here p denotes a prime and $a \in \mathbb{Z}$. We know that

$$\sum_{n \leq x} \tau(n) \sim n \log n.$$

We also know that

$$\sum_{p \leq x} 1 \sim \frac{x}{\log x}.$$

So if you believe that the prime factors of a shifted prime behave more-or-less like a shifted integer, then you would think that the main term for the Titchmarsh divisor problem (11) should be roughly x . In reality, there will be arithmetic factors coming up.

$$\sum_{p \leq x} \tau(p - a) = \mathfrak{S}(a)x + E_{\tau}(x).$$

- Titchmarsh (1933): under GRH,
- Linnik (1966): dispersion method, unconditional.
- Halberstam (1967): simplified proof.

$$E_{\tau}(x) \ll \frac{x \log \log x}{\log x},$$

uniform in a .

- Bombieri–Friedlander–Iwaniec (1986):

$$E_{\tau}(x) \ll_{A,a} \frac{x}{(\log x)^A}, \quad A > 0.$$

- Drapeau (2017): under GRH, there exists $\delta > 0$ such that

$$E_{\tau}(x) \ll x^{1-\delta}.$$

Question 53. Do we have

$$E_{\tau}(x) \ll x^{1/2+\varepsilon}?$$

Uniformity in a ? For example,

$$\sum_{p \leq N} \tau(N - p).$$

How many ways can an integer n be written as a sum of a prime and two squares?

$$n = p + x^2 + y^2.$$

This question arises naturally in the study of, e.g.,

$$\sum_{p \leq x} r(n-p).$$

What do we know about such sums? Consider the statement

$$\sum_{p \leq n} r(n-p) \leq \mathfrak{S}(n) \frac{\pi n}{\log n} + E_r(n).$$

- Hooley (1957): established the above under GRH.
- Linnik: unconditional proof.
- Elliott–Halberstam: simplified proof,

$$E_r(n) \ll \frac{n}{(\log n)^{1+\delta}}, \quad \delta \approx 0.04.$$

Theorem 54 (Assing–Blomer–L.). *We have*

$$E_r(n) \ll \frac{n}{(\log n)^A} \quad \text{for all } A > 0,$$

uniformly in $|a| \ll n$. *We have*

$$\sum_{p \leq n} r(n-p) = \mathfrak{S}(n) \text{Li}(n) + O_A \left(\frac{n}{(\log n)^A} \right).$$

Upper bounds for $\sum_{p \leq x} \lambda(p-a)$, where p denotes a prime and $a \in \mathbb{Z}$.

Pitt (2013): there exists $\delta > 0$ such that

$$\sum_{p \leq x} \lambda \left(p + \frac{1}{q} \right) \ll x^{1-\delta}.$$

Assing–Blomer–Li: uniformity in level, shift parameter, weight, spectral parameter.

Recall that $\pi(x) = \#\{p \leq x\}$, $\pi(x, q, a) = \#\{p \equiv a(q)\}$. Then

$$\sum_{q \leq \sqrt{x}/(\log x)^B} \max_{(a,q)} \left| \pi(x, q, a) - \frac{\pi(x)}{\phi(q)} \right| \ll_A \frac{x}{(\log x)^A}$$

for all $A > 0$.

We have

$$\sum_{\frac{\sqrt{x}}{(\log x)^B} < q < \sqrt{x}} \frac{x}{\phi(q) \log(x/q)} \ll \frac{x \log_2 x}{\log x}.$$

Also

$$\sum_{q \sim \sqrt{x}} \left(\pi(x, q, a) - \frac{\pi(x)}{\phi(q)} \right) \ll \frac{x}{(\log x)^A},$$

uniform in $a \ll x^{1+\delta}$; also, you can go up to $x^{1/2+\delta}$, and then all the way up to $x/(\log x)^B$ because of the switching divisor thing?

Leads to

$$\sum_{q_2} \sum_h \sum_{q_1} S(ah(n_1 - n_2) \bar{n}_1, q_2, q_1 n_2).$$

Now apply Kuznetsov and large sieve. This sequence is very sparse when you have large coefficients a . That's why uniformity is a major issue in the original Titchmarsh divisor problem. We overcome this difficulty by factoring out the coefficient

a , in some sense, because of a certain multiplicativity of the Fourier coefficients. But because we have levels and these inverses, it's not at the cusp infinity, but rather at something like the cusp $1/N$.

Study $\sum_p \tau(p^2 + q^2 + 1) - \mathcal{M} + \mathcal{E}$. Improve to arbitrary power of a logarithm savings? How about the analogue for cusp forms, namely $\sum \lambda(p^2 + q^2 + 1)$?

19. MAYANK PANDEY, *L^1 -norms of exponential sums with multiplicative coefficients*

The general problem is to understand the distribution of the magnitude of exponential sums of some arithmetically interesting coefficients $a : \mathbb{N} \rightarrow \mathbb{C}$, namely

$$\left| \sum_{n \leq X} a(n) e(n\alpha) \right|.$$

One approach that gives a lot of information about these is through the moments

$$\int_0^1 \left| \sum_{n \leq X} a(n) e(n\alpha) \right|^s d\alpha.$$

We focus on the L^1 -norm. Here there is comparatively little known. Less structure compare to the even moments.

Why would we want to understand such L^1 -norms?

There's the general problem of understanding the distribution, which one might want for its own sake. It's not inconceivable that it might show up in applications of the circle method. Finally, there's an interesting result of Bourgain:

Theorem 55. *Any finite subset $A \subseteq \mathbb{N}$ has a sumfree subset B with*

$$|B| \geq \frac{1}{3}|A| + \frac{1}{\log|A|} \int_0^1 \left| \sum_{n \in A} e(n\alpha) \right| d\alpha.$$

The best current result (also due to Bourgain) is that the biggest such B has size

$$\geq \frac{1}{3}|A| + \frac{2}{3}.$$

This improves upon a simpler result of Erdos where one just has $|B| \geq (1/3)|A|$ under weaker conditions.

One might hope that sets with small L^1 -norm can be characterized, then manually dealt with, to obtain

$$|B| \geq \frac{1}{3}|A| + \psi(|A|),$$

for some $\psi(X) \rightarrow \infty$. Bloom has announced some results in this direction for general sets.

Let's turn to basic considerations regarding what one can prove. It's not hard to show that

$$\int_0^1 \left| \sum_{n \leq X} e(n\alpha) \right| d\alpha \asymp \log X.$$

This is sharp, due to a (resolved) conjecture of Littlewood:

$$\int_0^1 \left| \sum_{n \leq X} a(n) e(n\alpha) \right| d\alpha \gg \sum_{j \leq k} \frac{|a(n_j)|}{j},$$

where $\{n_1, \dots, n_k\}$ are those $n \leq X$ at which $a(n) \neq 0$. One expects that the L^1 -norm is small only if the coefficients are structured in arithmetic progressions, and the aforementioned work of Bloom makes this precise when the L^1 -norm is very small.

For the case where $a(n)$ is a multiplicative function, here are some examples of results.

Theorem 56 (Balog, Ruzsa). *For fixed $r \geq 2$, we have*

$$\int_0^1 \left| \sum_{n \leq X} 1_{r\text{-free}}(n) e(n\alpha) \right| d\alpha \asymp X^{\frac{1}{r+1}}.$$

Theorem 57 (Vaughan, Goldston). *We have*

$$\sqrt{X} \ll \int_0^1 \left| \sum_{n \leq X} \Lambda(n) e(n\alpha) \right| d\alpha \leq \left(\frac{1}{\sqrt{2}} + o(1) \right) \sqrt{X \log X}.$$

Theorem 58 (Balog–Ruzsa). *For $a(n)$ supported on squarefree n , we have*

$$\int_0^1 \left| \sum_{n \leq X} a(n) e(n\alpha) \right| d\alpha \gg X^{1/8-o(1)} \left(\frac{1}{X} \sum_{n \leq X} |a(n)|^2 \right)^{1/2}.$$

Theorem 59 (P.). *With $\lambda_f(n)$ the normalized Fourier coefficients of an $\mathrm{SL}_2(\mathbb{Z})$ modular form, we have*

$$\int_0^1 \left| \sum_{n \leq X} \lambda_f(n) e(n\alpha) \right| d\alpha = CX^{1/2} + O(X^{1/2-\delta})$$

for some $\delta > 0$ (probably $\delta = 1/100$ suffices).

One of the main results that we'll state today is the following, joint with Maksym Radziwill.

Theorem 60 (P., Radziwill). *Take λ the Liouville function (completely multiplicative, $\lambda(p) = -1$ for all p). We have*

$$\int_0^1 \left| \sum_{n \leq X} \lambda(n) e(n\alpha) \right| d\alpha \gg X^{1/4-o(1)}.$$

- Previous best result due to Balog and Perelli, giving a lower bound of $\exp(c \log X / \log \log X)$.
- Methods apply to the Mobius function, beating the previous Balog–Ruzsa $X^{1/6}$ result lower bound. (Balog–Ruzsa used less information.)

Another one of our results is a structure theorem for the L^1 -norm of an exponential sum with multiplicative coefficients.

Theorem 61 (P. Radziwill). *Suppose that $f : \mathbb{N} \rightarrow \{\pm 1\}$ is completely multiplicative. Suppose that $M \ll X^\eta$ for some absolute $\eta > 0$. Then, if*

$$\int_0^1 \left| \sum_{n \leq X} f(n) e(n\alpha) \right| d\alpha \leq M,$$

then there exist $\chi(q), t \in \mathbb{R}$ with $q(1 + |t|) \ll M^2 \log^{O(1)} \chi$ such that

$$f \approx \{n \mapsto \chi(n) n^{it}\}.$$

While not fully worked out, we expect to make “ $\approx$ ” correspond to pretentiousness. From the lens of $\approx$, this is sharp: for χ of conductor q , and $t \in \mathbb{R}$ with $q(1 + |t|)$ at most a small power of X , we have

$$\int_0^1 \left| \sum_{n \leq X} \chi(n) n^{it} e(n\alpha) \right| d\alpha = (q(1 + |t|))^{1/2} (\dots).$$

The next result follows from similar methods.

Theorem 62. *Let $\lambda_\pi(n_1, \dots, n_{m-1})$ be the Fourier coefficients (or Hecke eigenvalues) of some GL_m cuspidal automorphic representation π . Then we get*

$$\int_0^1 \left| \sum_{n \leq X} \lambda_\pi(1, \dots, 1, n) e(n\alpha) \right| d\alpha \gg X^\delta,$$

where $\delta = \delta(m) \gg 1/m$.

When $m = 2$, this follows from the Jutila/Wilton pointwise bound of $X^{1/2}$, and similarly for $m = 3$ by the bound

$$\sum_{n \leq X} \lambda_\pi(1, n) e(n\alpha) \ll X^{3/4 + o(1)}$$

of Miller, but seems new for general m , where there is no nontrivial L^∞ -bound.

Now we give the sketch of the proof of the result concerning general multiplicative functions. Suppose that f does not pretend to be a character of conductor $\leq Q$ for some $Q \leq X^\delta$. Let

$$S(\alpha) := \sum_{n \leq X} f(n) e(n\alpha).$$

The theorem is then the statement that

$$\|S\|_1 \gg \sqrt{Q}.$$

Define the major arcs

$$\mathfrak{M} = \bigcup_{\substack{q \leq Q \\ (a, q) = 1}} \mathfrak{M}(q, a), \quad \mathfrak{M}(q, a) := (\dots)$$

and the minor arcs $\mathfrak{m}$.

If we have a bound of $X/\sqrt{Q}$ for $S(\alpha)$ on the minor arcs $\mathfrak{m}$, and that

$$\int_{\mathfrak{M}} |S(\alpha)|^2 d\alpha \leq (1 - \sigma)X$$

for some $\sigma \gg 1$. Then we are done, by Cauchy–Schwarz:

$$\int_{\mathfrak{m}} |S(\alpha)|^2 d\alpha \gg X \implies \int_{\mathfrak{m}} |S(\alpha)| d\alpha.$$

Don't have such good minor arc bounds for $S(\alpha)$ from nonpretentiousness, which is insensitive to primes $\sim X$, so we can do no better than saving $1/\log X$. Instead, we decompose as

$$S = S^* + S_{\text{exc}},$$

with

$$\|S_{\text{exc}}\|_2^2 = o_{\delta \rightarrow 0}(X), \quad \sup_{\alpha \in \mathfrak{m}} |S^*(\alpha)| \ll X/\sqrt{Q}.$$

Here S^* is a sum over n with a prime factor in a certain range, and we use Ramare's identity to provide bilinearity to get minor arc bounds. Let

$$\mathcal{E} := \left\{ \alpha : |S_{\text{exc}}(\alpha)| \geq X/\sqrt{Q} \right\}.$$

Something.

Then, because the exceptional set attached to $\alpha \in \mathfrak{m} \cap \mathcal{E}$ has size at least $S^*(\alpha)$, and since we can bound the L^2 -norm of $S(\alpha)$ on $\mathfrak{m} \cap \mathcal{E}$ by that of $S_{\text{exc}}(\alpha)$, then we get the desired lower bound $\|S\|_1 \gg \sqrt{Q}$.

The main part is the L^2 upper bound on the major arcs. By Plancherel and change of variables, we get

$$\int_{\mathfrak{M}} |S(\alpha)|^2 d\alpha \approx \sum_{q \leq Q} \sum_{\chi(q)}^* \int_{|t| \ll Q/q} \left| \sum_{n \sim X} f(n) \chi(n) n^{it} \right|^2 dt.$$

Restricting to n with a prime factor in $[X^{3\delta}, X^{1-3\delta}]$ and applying Turan–Kubilius, there are scales

$$X^{3\delta} \ll N, \quad P \ll X^{1-3\delta}, \quad NP \asymp X$$

such that the above is

$$\ll \log^2 P \sum_{q \leq Q} \sum_{\chi(q)}^* \int_{|t| \ll Q/q} \left| \sum_{n \asymp N} f(n) \chi(n) n^{it} \right|^2 \left| \sum_{p \asymp P} f(p) \chi(p) p^{it} \right|^2 dt.$$

We bound the n -sum in L^∞ via Halasz. We win after using the large sieve on the p -sum: there are more harmonics than the length, so we get a savings of Q^2 from the large sieve.

Let's now give a sketch of the Liouville result. We begin by writing

$$S_X(\alpha) = \sum_{n \leq X} \lambda(n) e(n\alpha).$$

Then S_X varies on scales of size $1/X$, so we have

$$\begin{aligned} \int_0^1 |S_X(\alpha)| d\alpha &\gg \frac{1}{X} \sum_{\substack{p \sim \sqrt{X} \\ (a,p)=1}} \left| S_X\left(\frac{a}{p}\right) \right| \\ &\geq \frac{1}{X} \sum_{p \sim \sqrt{X}} \cdots \end{aligned}$$

For typical $p \sim \sqrt{X}$, we generically expect that (since ζ has nontrivial zeros)

$$p \sum_{n \leq X/p} \lambda(n) \gg p\sqrt{X/p} \asymp X^{3/4},$$

and as X varies, there are Ω results to this effect that have been established.

As for varying X , we write

$$F_X(\alpha) = \sum_{n \leq X} e(n\alpha).$$

We then have that for $y \leq X$,

$$\int_0^1 F_y(\alpha - \beta) S_X(\beta) d\beta = S_y(\beta),$$

and so

$$\|S_y\|_1 \leq \|F_y\|_1 \|S_X\|_1.$$

Since $\|F_y\|_1 \asymp \log y$, we obtain that (...).

We reduce to showing that

$$\frac{1}{X} \cdot X^{1/2} \frac{1}{X} \int_{y \asymp X} \left| p \sum_{n \leq y/p} \lambda(n) + \sum_{n \leq y} \lambda(n) \right| dy \gg X^{1/4-o(1)}.$$

The idea that's used here is to integrate against the kernel $w(y/X)$, where

$$w(u) := \frac{1}{2\pi i} \int_{(2)} \frac{\zeta(s + i\gamma_0)}{s - \frac{1}{2}} u^{-s} e^{s^2} ds,$$

and $\frac{1}{2} + i\gamma_0$ is some zero of ζ . This is based on an idea of Pintz.

With this, we can pick out the oscillation coming from a zero of ζ for the expected Ω result. In particular, on average,

$$p \sum_{n \leq y/p} \lambda(n) + \sum_{n \leq y} \lambda(n) \gg X^{1/2} p^{1/2} \sim X^{3/4}.$$

We thereby obtain the required lower bound

20. ANDREI SEYMOUR-HOWELL, *Rigorous computation of Maass cusp forms of squarefree level*

$S_\lambda(N)$: the space of Maass cusp forms of level N (trivial character) with eigenvalue λ .

Hecke operators T_n , Fourier expansions with coefficients

$$a(n) = \lambda(n), \quad a(-n) = \varepsilon \lambda(n),$$

where $\varepsilon = \pm 1$.

History of computations. In the 1990's, Hejhal developed an algorithm to compute Maass cusp forms, generalized by Stromberg in 2006 to work for general congruence (and non-congruence) subgroups. This argument works very well in practice, but relies on a heuristic argument, and it has not been proven yet to rigorously converge to genuine Maass cusp forms. One gets number that look like Maass forms, but one can't tell if it is an exact Maass form.

Booker-Strombergsson-Venkatesh (2006) developed an algorithm to verify the computations of Maass cusp forms. Originally derived only for level 1, but recent work of Child has generalized this to general level and character.

Booker and Strombergsson (2006) used the Selberg trace formula for computations of Maass cusp forms. However, they focused mainly on proving the non-existence of Maass forms in certain cases.

The main tool we'll use here is the Selberg trace formula. Hecke eigenbasis f_j , with Laplace eigenvalues λ_j and Hecke eigenvalues $a_j(n)$. Let H be a nice (analytic) test function. Then the spectral sum

$$t(n, H) := \sum_j a_j(n) H(\lambda_j)$$

admits a geometric expansion.

Aside to modular forms. Explicit versions of the trace formula have been used to compute basis elements of the spaces of modular forms. Example: PARI command `mfeigenbasis`. Since these spaces are finite-dimensional, one can use Hecke operators and linear algebra to extract Fourier coefficients of basis elements.

In the Maass form case, our spaces become infinite-dimensional. We'll mimic the idea, though, by taking the test function to decay rapidly enough that an essentially bounded number of eigenvalues contribute.

As for details, we define, for a sequence of real numbers $c(m)$ (here $1 \leq m \leq M$) satisfying $c(m) = 0$ if $(m, N) > 1$,

$$Q(c, H) := \sum_{j=1}^{\infty} \left(\sum_{m=1}^M c(m) a_j(m) \right)^2 H(\lambda_j).$$

We can rewrite this using Hecke multiplicativity.

Intuitively, suppose we have some putative numerical approximations $\tilde{\lambda}_j$ and $\tilde{a}_j(m)$ to λ_j and $a_j(m)$. Then we want to choose numbers $c_i(m) = 0$ if $(m, N) > 1$ and

$$\sum_{m=1}^M c_i(m) \tilde{a}_j(m) = \begin{cases} 1 & \text{if } j = i, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\tilde{H}_i(\lambda) = H(\lambda)(\lambda - \tilde{\lambda}_i)^2$. For the verification, we shall prove that there exists a Laplace eigenvalue near $\tilde{\lambda}_i$. For this, we use the definition of Q to compute

$$\varepsilon_i := \sqrt{\frac{Q(c_i, \tilde{H}_i)}{Q(c_i, H)}}.$$

There exists a cuspidal eigenvalue $\lambda \in [\tilde{\lambda}_i - \varepsilon_i, \tilde{\lambda}_i + \varepsilon_i]$.

With the same H as before, we define $\tilde{H}(\lambda) = \lambda H(\lambda)$. Let Q and $\tilde{Q}$ denote the respective matrices of the quadratic forms $Q(c, H)$ and $Q(c, \tilde{H})$. To find the eigenvalues λ_j , we seek solutions to the generalized symmetric eigenvalue problem

$$\tilde{Q}x = \lambda Qx.$$

This gives us numerical approximations to the Laplace eigenvalues. The corresponding eigenvectors we then use as the c_i . This way means we only have to do two matrix diagonalizations per level.

21. BEAT ZURBUCHEN, *Sato–Tate laws for Fourier duals of exponential sums*

21.1. Sato–Tate. Let $E/\mathbb{Q}$ be a non-CM elliptic curve.

Let p be a prime of good reduction for E . Then one can prove using the Lefschetz trace formula that the number of points for the reduction of E modulo p in this finite field is

$$|E_p(\mathbb{F}^n)| = p^k + 1 - p^{k/2} (\alpha_p^n + \alpha_p^{-n}), \quad |\alpha| = 1.$$

$$\Theta_p := \begin{pmatrix} \alpha & 0 \\ 0 & \bar{\alpha} \end{pmatrix} \in \mathrm{SU}(2)^\sharp.$$

Here $\mathrm{SU}(2)^\sharp$ denotes the set of topological conjugacy classes of $\mathrm{SU}(2)$.

Sato–Tate says that for all continuous

$$f : \mathrm{SU}(2)^\sharp \rightarrow \mathbb{C},$$

we have

$$\lim_{N \rightarrow \infty} \mathbb{E}_{\substack{p \leq N \\ \text{good reduction}}} f(\Theta_p) = \int_{\mathrm{SU}(2)} f(g) dg.$$

Tate module: for ℓ a prime, for $p \neq \ell$ of good reduction for E ,

$$T_\ell : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(\mathbb{Q}_\ell).$$

Fix

$$\iota : \overline{\mathbb{Q}_\ell} \xrightarrow{\cong} \mathbb{C}.$$

For each such p ,

$$\Theta_p = (\iota(T_\ell(\mathrm{Frob}_p)))^{ss}.$$

Sato–Tate over $\mathbb{G}_m$? Let p be an odd prime. Let $q = p^m$. Take a prime $\ell \neq p$.

Definition 63. An ℓ -adic local system on $\mathbb{G}_{m, \mathbb{F}_q}$ is a continuous Galois representation

$$\rho : G_{\mathbb{F}_q(t)} \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}_\ell})$$

which ramifies at most at 0 and ∞ .

From now on, ρ is an ℓ -adic local system.

Remark 64. Local systems will give you functions:

$$\begin{aligned} \rho(t) : \mathbb{F}_q^* &\rightarrow \mathbb{C} \\ t &\mapsto \mathrm{trace}(\mathrm{Frob}_t | \rho). \end{aligned}$$

Example 65. (1) Let $\chi : \mathbb{F}_{q^n}^* \rightarrow S^1$, there is a local system h_χ such that

$$h_\chi(t) = \chi(N_{\mathbb{F}_{q^n}/\mathbb{F}_q}(t)).$$

(2) Let $\lambda_2 : \mathbb{F}_p^* \rightarrow S^1$ denote the Legendre symbol, and let $\chi : \mathbb{F}_q^* \rightarrow S^1$ and $\psi : \mathbb{F}_q \rightarrow S^1$ be characters, with ψ nontrivial. Then there is a local system

$$\mathcal{H}(\psi, \chi)$$

such that

$$\mathcal{H}(\psi, \chi) = \sum_{x_1 \cdots x_7 = ty} \psi(\mathrm{trace}_{\mathbb{F}_{q^n}/\mathbb{F}_q}(x_1 \cdots x_7 - y)) \chi(N(x_4 x_5 x_6^{-1} x_7^{-1})) \lambda_2(y).$$

Sato–Tate law. Let ρ be a “normalized” ℓ -adic local system.

Then there exists a compact Lie group $G \subseteq \mathrm{GL}(\rho \otimes_{\mathbb{Q}_\ell} \mathbb{C})$ such that

$$\Theta_t := (\iota(\rho(\mathrm{Frob}_t)))^{ss} \in G^\sharp$$

and for all $f : G^\sharp \rightarrow \mathbb{C}$ continuous, and all $t \in \mathbb{F}_{q^n}^*$, we have

$$\frac{1}{q^n - 1} \sum_{t \in \mathbb{F}_{q^n}^*} f(\Theta_t) = \int_G f(g) dg + O_{f,g}(q^{-n/2}).$$

Example 66. (1) For h_χ , the group G is $\iota(\mathrm{image}(\chi))$.

(2) For $\mathcal{H}(\psi, \chi)$, the monodromy group is G_2 .

Sato–Tate laws for Fourier duals. We can construct a Mellin transform for ρ : for a multiplicative character $\chi : \mathbb{F}_{q^n}^* \rightarrow S^1$,

$$\hat{\rho}(\chi) := \sum_{t \in \mathbb{F}_{q^n}^*} \rho(t) \chi(t).$$

This is not a local system (by any known means).

Sato–Tate (Katz, 2012): for ρ “normalized”, there exists a compact Lie group G , together with a representation $\varphi_\pi : G \rightarrow \mathrm{GL}_n(\mathbb{C})$, such that for all “good” characters χ , there exists a conjugacy class $\Theta_\chi \in G^\sharp$ such that

$$\mathrm{trace}(\Theta_\chi | \pi) = \hat{\rho}(\chi),$$

and for all continuous $f : G^\sharp \rightarrow \mathbb{C}$,

$$\mathbb{E}_{\substack{\chi \in \widehat{\mathbb{F}_{q^n}^*} \\ \chi \text{ is good}}} f(\Theta_\chi) = \int_G f + O(q^{-n/2}).$$

Example 67. There exists a local system ρ_t for each $t \in \mathbb{F}_{p^n}^*$ such that $\hat{\rho}_t(\chi) = \mathcal{H}(\psi, \chi)(t)$.

Theorem 68 (Katz). *For t in a set of full measure, the monodromy group is G_2 .*

The speaker has been able to show the following.

Theorem 69. *The same is true for all t , provided that p is sufficiently large.*

22. QIHANG SUN, *Sums of Kloosterman sums and their applications*

We consider the Kloosterman sums

$$S(m, n, c) := \sum_{d(c)} e_c(ma + nd).$$

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}).$$

Weil bound with no cancellation gives

$$|S(m, n, c)| \leq \sigma_0(c) (m, n, c)^{1/2} c^{1/2}.$$

$$\sum_{c \leq X} \frac{S(m, n, c)}{c} \ll_\varepsilon (m, n)^\varepsilon X^{1/2+\varepsilon}.$$

Kuznetsov: the above is $\ll_{m,n} X^{1/6} \log^{1/3} X$.

Theorem 70 (Sarnak–Tsimmerman, 2009). *The above is*

$$\ll_\varepsilon \left(X^{1/6} + (mn)^{1/6} + (m+n)^{1/8} (mn)^{\vartheta/2} \right) (mnX)^\varepsilon.$$

Conjecture 71 (Linnik–Selberg). *The above is*

$$\ll_\varepsilon (mnX)^\varepsilon.$$

j -invariant:

$$j(z) = q^{-1} + 744 + O(q).$$

$$a_j(n) = \frac{2\pi}{\sqrt{n}} \sum_{c=1}^{\infty} \frac{S(1, n, c)}{c} I_1 \left(\frac{2\pi\sqrt{n}}{c} \right).$$

Generalized Kloosterman sum:

$$\begin{aligned}\nu : I &\rightarrow \mathbb{T}, \\ \nu(-I) &= e^{-\pi i k}.\end{aligned}$$

Then

$$\theta(z) := \sum_{n \in \mathbb{Z}} q^{n^2}, \quad \theta(\gamma z) = \sqrt{cz + d} \underbrace{\left(\frac{c}{d}\right) \varepsilon_d^{-1}}_{\nu_\theta(\gamma)} \theta(z).$$

$$\eta(z) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n),$$

$$z \mapsto z + 1,$$

$$\eta(\gamma z) = \sqrt{cz + d} \nu_\eta(\gamma) \eta(z),$$

$$\eta(z) = \sum_{n=1}^{\infty} a_\eta(n) q^{n-23/24} \rightarrow \tilde{n}.$$

$$S(m, n, c, \iota) := \sum_{d(c)^*} \overline{\nu}(\gamma) e_c(\tilde{m}a + \tilde{n}d).$$

- (1) Weil bound.
- (2) Goldfeld–Sarnak, 1983:

$$\sum_{c \leq X} \frac{S(m, n, c, \iota)}{c} = \sum_{1/2 < s_j < 1} \tau_j(m, n) \frac{X^{2s_j-1}}{2s_j-1} + O_{m,n,I,\nu,k,\varepsilon}(X^{\beta/3+\varepsilon}),$$

where

$$\lambda_j = s_j(1 - s_j).$$

- (3) Uniform bound. Ahlgren–Andersen, 2015:

$$O_{\nu,\varepsilon} \left((X^{1/6} + |\tilde{m}\tilde{n}|^{1/4})(\dots)^\varepsilon \right).$$

For the even multiplier, e.g., ν_j , the exponent gets reduced to

$$\frac{1}{4} - \frac{1}{168}. \tag{12}$$

Andersen and Wu got the better result

$$\frac{1}{4} - \frac{1}{36} \tag{13}$$

by using the subconvexity of L -functions.

For ψ , get

$$\frac{1}{4} - \frac{1}{147}. \tag{14}$$

Next, we turn to the partition functions. Recall that $p(n)$ denotes the number of ways to write n as a sum of positive integers. For example,

$$p(4) = 5$$

because we can write 4 in the following five ways:

$$4, \quad 3+1, \quad 2+2, \quad 2+1+1, \quad 1+1+1+1.$$

(1) We have

$$p(n) \sim \frac{1}{4\sqrt{3n}} e^{\pi\sqrt{\frac{2n}{3}}} + \dots + O(\dots).$$

Rademacher, 1938:

$$p(n) = \frac{2\pi e(-\frac{1}{8})}{(24n-1)^{3/4}} \sum_{c=1}^{\infty} \frac{S(1, 1-n, c, \nu_{\eta})}{c} I_{3/2} \left(\frac{4\pi\sqrt{24n-1}}{24} \right).$$

Estimates:

$$R_1(n, x) = (\dots) \sum_{n>x}$$

satisfies

$$R_1(n, \alpha\sqrt{n}) \ll n^{-1/2-1/168+\varepsilon}.$$

(2) Another case comes from the congruence satisfied by the partition function. The following congruence was conjectured by Ramanujan.

$$p(5n+4) \equiv 0(5).$$

Other congruences were conjectured involving $7n+5$ and $11n+6$.

Dyson, 1994: the rank of the partition

$$\Lambda_1 + \dots + \Lambda_k = n, \quad \Lambda_1 \geq \dots \geq \Lambda_k,$$

namely $\Lambda_1 - k$, is congruent to a modulo 5. Let $N(m, n)$ denote the number of partitions of n of rank m . Then

$$\begin{aligned} R(w, q) &= 1 + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} N(m, n) w^m q^n \\ &= 1 + \sum_{n=1}^{\infty} \frac{q^{n^2}}{\prod_{j=1}^n (1 - wq^j)(1 - w^{-1}q^j)}. \end{aligned}$$

$w = -1$, $R(-1, q) = f(q)$, 3rd mock theta function. Bringmann–Ono, 2006: we can write

$$a_f(n) = \frac{2\pi e(-\frac{1}{8})}{(24n-1)^{1/4}} \sum_{2|c} \frac{S(0, n, c, \psi)}{c} I_{1/2} \left(\frac{4\pi\sqrt{24n-1}}{24} \right),$$

where

$$\frac{\eta(z)^5}{\eta(2z)^2} \in S_{3/2}(2, \bar{\psi}).$$

Ahlgren–Dunn, 2019: with

$$R_2(n, x) = (\dots) \sum_{n>x} \dots,$$

we have

$$R_2(n, \alpha\sqrt{n}) \ll n^{-1/147+\varepsilon}.$$

The speaker's main result (2023) is to generalize the improvements (12) and (14): we get

$$\frac{1}{4} - \frac{1}{147}$$

for the multipliers

$$\left(\frac{D}{\cdot} \right) \nu_{\theta}, \quad \left(\frac{D}{\cdot} \right) \nu_{\eta}.$$

The speaker is trying to improve this further using subconvexity, like in (13).

23. NICOLAS POTTHAST, *On the asymptotics of elementary abelian extensions over global function fields*

Fix a global field F and a finite group G .

Question 72. Are there Galois extensions of F with Galois group G .

Question 73. How many of these extensions are there with discriminant bounded in norm from above by X ?

Question 74. How does the counting function $Z(F, G, X)$, for the number of extensions K/F with Galois group G and discriminant bounded by X , grow as $X \rightarrow \infty$?

Conjecture 75 (Malle 2005). *For a number field F , there exists a constant $c(F, G) > 0$ that describes the asymptotics.*

Wright made a conjecture about the case where the characteristic of F divides $|G|$, but further research showed that the conjecture is wrong.

Lagemann: Distribution of Artin–Schreier extensions.

Theorem 76. *The number of C_p^r -extensions of a global function field F , when counted by discriminant, has asymptotics*

$$Z(F, C_p^r, X) \sim \begin{cases} c(F, C_2^2, e) X^{1/2} \log(X)^3 & \text{for } C_2^2, \\ c(F, C_p^r, e) X^{\frac{1+r(p-1)}{p(p^r-1)}} & \text{otherwise.} \end{cases}$$

In these formulas, we have $X = q^m$, and we consider $m \rightarrow \infty$ with

$$m \equiv e \pmod{L},$$

for a fixed e , with $L = 2$ for C_2^2 and $L = p(p^r - 1)$ otherwise.

24. FARRELL BRUMLEY, *Recent extensions of Duke’s equidistribution problems*

Duke’s theorem is a statement about equidistribution that is solved by appealing to a subconvex bound. What we’d like to do today is to look at some variants of Duke’s theorem which are no longer simply reduced to subconvex bounds on L -functions, but rather require more and more technology in terms of L -functions and also other ingredients such as some classical analytic number theory and some ergodic theory. So we’ll start simple and then ramp up the degree of difficulty, throwing new tools at the problem. Some people already attended a talk of this nature of mine in Oberwolfach a month ago, but the ending of this talk will be different, so there will still be something new for you.

Let’s start with classical Duke’s theorem (or Linnik’s problem). As we have said, there are many variants and we’ll just choose one. The one we’ll choose will have to do with the sphere. We’ll take a squarefree integer $d \in \mathbb{N}$ and we’re going to look at the ways to represent d as a sum of three squares:

$$R_d := \{x = (a, b, c) \in \mathbb{Z}^3 : a^2 + b^2 + c^2 = d\}.$$

We know, by Gauss and Legendre, a criterion for when this set is nonempty:

$$R_d \neq \emptyset \iff d \not\equiv 0, 4, 7(8).$$

We will call d *admissible*, and write

$$d \in \mathbb{D},$$

if it satisfies the above conditions.

Question 77. Does the probability measure μ on the set

$$\frac{1}{\sqrt{d}} R_d$$

equidistribute on the sphere S^2 as $d \rightarrow \infty$?

Remark 78. One can ask the same question in higher dimensions, but dimension three is in some sense the hardest case.

The first progress on this question is due to Linnik and his school, who showed that these probability measures converge weak-star to the uniform measure on the sphere provided that our discriminants satisfy a congruence condition, called a *Linnik condition*. More precisely,

$$\mu_d \rightarrow \mu_{S^2} \quad \text{provided that } -d \in (\mathbb{F}_p^*)^2$$

for some auxiliary choice of prime p . We will denote by $\mathbb{D}(p)$ the integers that satisfy this last condition. You can choose your prime as you wish; it's completely auxiliary and foreign to the problem, but you need to fix it, and once you do, you get equidistribution along that sequence of discriminants. So there is a defect here, and that defect is the choice of auxiliary prime.

So the problem was not solved fully by Linnik. His methods were ergodic-theoretic, and we're not going to say much about them, other than to say that this auxiliary condition is the means by which one can access ergodic theory. In the end, membership in $\mathbb{D}(p)$ is a splitting condition of a certain acting torus, and we want this torus to be non-split, so that it is a non-compact group acting on the solution space, allowing one to work ergodically.

After that, Duke (late 80's, early 90's) showed that one could remove this ergodic condition of Linnik:

$$\mu_d \rightarrow \mu_{S^2}$$

as $d \rightarrow \infty$ in $\mathbb{D}$, effectively. We will talk briefly about the proof of this. As mentioned, this is proved by appealing to subconvexity. What is the bridge to subconvexity? One wants to show that if you test this measure against some continuous function f on S^2 that is orthogonal to constants, then

$$\mu_d(f) \rightarrow 0.$$

We now discuss *orthogonal lattices*. Let

$$Y_0(1) := \mathrm{SL}_2(\mathbb{Z}) \backslash \mathbb{H}^2 = \{ \text{unimodular lattices in } \mathbb{R}^2 \text{ up to rotation} \},$$

write μ_{hyp} for the normalized hyperbolic measure. To each $v \in R_d$, we associate the binary quadratic lattice

$$\Lambda_v := v^\perp \cap \mathbb{Z}^3$$

obtained by taking integral elements of the orthogonal complement of v . This defines a map

$$\begin{aligned} R_d &\rightarrow Y_0(1) \\ v &\mapsto [\Lambda_v]. \end{aligned}$$

Playing the same game, we define

$$\mu_d = \frac{1}{|R_d|} \sum_{v \in R_d} \delta_{[\Lambda_v]}.$$

Linnik tells us that

$$\mu_d \rightarrow \mu_{\text{hyp}}$$

for $d \in \mathbb{D}(p)$, and Duke tells us that this happens for all $d \in \mathbb{D}$. One can ask now for *joint equidistribution*: consider the measure

$$\nu_d = \frac{1}{|R_d|} \sum_{v \in R_d} \delta_{(\frac{v}{\|v\|}, [\Lambda_d])} \quad \text{on } S^2 \times Y_0(1).$$

There was a Bourbaki talk on this last year by Menny Akka, who had wonderful numerics that he showed. The speaker attempts to reproduce one of them on the blackboard. We consider a partition of the fundamental domain into a bulk region L of volume $1/2$, and a complementary part L^c of volume $1/2$, where L^c is a tubular neighborhood of the cusp. Informally, lattices lie in L when they have a short vector. Equidistribution in this context says that the probability measures obtained by partitioning μ_d according to whether a vector v satisfies $[\Lambda_v] \in L$ or not equidistribute on S^2 as $d \rightarrow \infty$.

Theorem 79 (Akka–Einsiedler–Shapira (2019)). *Let $p_1, p_2 > 2$ be distinct. Then*

$$\nu_d \rightarrow \mu_{S^2} \otimes \mu_{\text{hyp}}$$

as $d \rightarrow \infty$ in $\mathbb{D}(p_1) \cap \mathbb{D}(p_2)$.

This may be understood as a classically expressed consequence of a powerful joinings classification result of Einsiedler–Lindenstrauss for higher rank diagonalizable actions. So in contrast to the Linnik problem, where what was underlying the dynamics there was a split rank one torus, here we’re going to have a split rank two torus, coming from the fact that we have fixed two auxiliary primes.

That’s the beginning of my recent work on this topic, with Valentin and others. The first theorem that we would like to present today is with Blomer a few years ago. We use automorphic methods (rather than ergodic theory), but assume GRH.

Theorem 80 (B–Blomer). *Assume GRH. Then*

$$\nu_d \rightarrow \mu_{S^2} \otimes \mu_{\text{hyp}}$$

as $d \rightarrow \infty$ in $\mathbb{D}$.

Let’s give the big picture. The spaces S^2 and $Y_0(1)$ are examples of what we call locally homogeneous quaternionic varieties, which are adelic quotients of the form

$$[G]_K = \underbrace{G(\mathbb{Q}) \backslash G(\mathbb{A})}_{[G]} / K.$$

As an example, S^2 corresponds to the Hamilton quaternions $B_{2,\infty}$, while $Y_0(1)$ corresponds to the matrix algebra M_2 . We may understand $\frac{1}{\sqrt{d}}R_d$ as defining a subset of $[PB_{2,\infty}^\times]_{K_1}$ and $\{[\Lambda_v] : v \in R_d\}$ as defining a subset of $[\text{PGL}_2]_{K_2}$.

- We could also take $G_1 = PB_1^\times$ and $G_2 = PB_2^\times$, where B_1 and B_2 are distinct fixed quaternion algebras.

- Quadratic field extension $E/\mathbb{Q}$ and (varying) embeddings

$$\text{Res}_{E/\mathbb{Q}} \mathbb{G}_m \rightarrow T_E \subset G_j \quad (j = 1, 2).$$

- We consider $\Delta T_E \subset G_1 \times G_2$, noting that $\Delta T_E \subset T_E \times T_E$ is not maximal.

The idea of the proof comes down to a representation-theoretic fact concerning the sequence of inclusions

$$\Delta T_E \rightarrow T_E \times T_E \rightarrow G_1 \times G_2.$$

Each of these inclusions defines a multiplicity one space, and we're going to make use of these intermediate spaces. These problems are difficult because the thing that's not varying (namely, ΔT_E) is kind of small in the thing that's fixed (namely, $G_1 \times G_2$). Making use of this intermediate group allows you to say that

$$\nu_d(f_1 \otimes f_2) = \frac{1}{h_d} \sum_{t \in \text{Cl}_d} f_1(t) f_2(t).$$

You're basically looking at an inner product along the class group. We may apply Parseval to rewrite the above as

$$\frac{1}{h_d} \sum_{\chi \in \text{Cl}_d^\wedge} \hat{f}_1(\chi) \hat{f}_2(\chi^{-1}).$$

These measures are similar to what we had for the Duke problem, but they don't come with a square. So what we get instead is a square-root L -function. What we'll get is an average of square roots of central values:

$$\frac{1}{h_d} \sum_{\chi \in \text{Cl}_d^\wedge} L(\tfrac{1}{2}, f_1 \times \chi)^{1/2} L(\tfrac{1}{2}, f_2 \times \chi)^{1/2}.$$

We're assuming GRH, but if you apply GRH pointwise to bound each one of these central values, then you won't get $o(1)$. What we do instead is to apply absolute values (killing our chances a bit by not considering any possible cancellation between the signs of these two square root L -values) Holowinsky did something similar for the shifted convolution problem in the setting of QUE. Here we're going to take $f_1 \neq f_2$ (because $G_1 \neq G_2$). We will use GRH to approximate

$$\log L(\tfrac{1}{2}, f_i \times \chi)$$

by a short sum of Hecke eigenvalues. We will then use the method of moments as developed by Sound.

That's all we'll say about that. Let's turn now to the *mixing conjecture* of Michel–Venkatesh. We'll try to be brief here, because we do want to talk about new stuff at the end here. Same setup, but with $G_1 = G_2$. The problem is now the existence of, e.g., $\Delta G \subseteq G \times G$.

Let's recall their conjecture in the classical setting. Set $G := \text{PGL}_2$, and let K be maximal. Then $[G]_K = Y_0(1)$. We then have the following diagram.

$$\begin{array}{ccc} Y_0(1) \times Y_0(1) & \longleftarrow & \Delta Y_0(1) \\ \uparrow & & \uparrow \\ \text{CM}_d \times \text{CM}_d & \longleftarrow & \Delta \text{CM}_d. \end{array}$$

We know by Duke that the vertical arrow on the right equidistributes. The way to get an interesting question was introduced by Michel–Venkatesh. Let $[\mathfrak{a}] \in \text{Cl}_d$,

and put

$$\Delta\mathrm{CM}_d([\mathfrak{a}]) = \{(x, [\mathfrak{a}]x) : x \in \mathrm{CM}_d\}. \quad (15)$$

Then for all $\mathfrak{n} \subset \mathcal{O}_E$ such that $[\mathfrak{n}] = [\mathfrak{a}]$, we have the inclusion

$$\Delta\mathrm{CM}_d([\mathfrak{a}]) \subset Y_0(N),$$

where $N := \mathrm{norm} \, \mathfrak{n}$.

Conjecture 81 (Michel–Venkatesh). *Let $d \in \mathbb{N}$ be a fundamental discriminant. For each such d , pick a class $[\mathfrak{a}] \in \mathrm{Cl}_d$ and form the probability measure associate to shifted CM points as in (15). Then we get equidistribution: the class $\Delta\mathrm{CM}_d([\mathfrak{a}])$ equidistributes in $Y_0(1)^2$ if both parameters go to infinite, or more precisely, if*

$$d \rightarrow \infty$$

and

$$q := \min \{ \|\mathfrak{n}\| : [\mathfrak{n}] = [\mathfrak{a}] \} \rightarrow \infty.$$

There are ergodic and automorphic approaches to this question, which each mix ideas from the other subject.

- Khayutin solved the mixing conjecture under a double Linnik condition, involving two split primes p_1, p_2 ; his proof is not altogether ergodic, since it involves analytic techniques and also assumes the absence of a Siegel zero.
- Blomer–B–Khayutin: solve the mixing conjecture under GRH and other hypotheses.

Remark 82. In both of these cases, the ambient group is a product group. In the ergodic approach to these problems, the fact that the ambient group is a product group allows you to access the joinings theorem, as done by Khayutin. But the joinings theorem is a corollary of a more general result of Einsiedler and Lindenstrauss, which is a classification of measures invariant under a higher rank diagonalizable action. It's not as nice as, say, Ratner's theorem, because it requires an additional input, namely positive entropy, which we'll need to verify.

Let's now move to higher rank. Let's move to PGL_3 .

Theorem 83 (ELMV, 2006). *Let*

$$A = \begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ 0 & 0 & 1 \end{pmatrix} < \mathrm{PGL}_3(\mathbb{R}).$$

For a natural number $d \in \mathbb{N}$, let $\mathcal{F}_d$ denote the set of periodic A -orbits on $X_3 := \mathrm{PGL}_3(\mathbb{Z}) \backslash \mathrm{PGL}_3(\mathbb{R})$ of discriminant d . Then $\mathcal{F}_d$ equidistributes on X_3 as $d \rightarrow \infty$.

Proof. The proof really hangs by a thread. It uses, critically, ergodic theory and number theory. It can't get by without one or the other.

- (1) $\mathcal{F}_d$, like in the classical case, is in bijection with the class group Cl_d of a *real* cubic extension $E/\mathbb{Q}$ of discriminant d . This has to do with the fact that A is a split (rank 2) torus, and this is a splitting condition, which is then the doorway to access ergodic methods. But one then needs to establish positive entropy. That's the role of ergodic theory in this paper.

- (2) We use critically that X_3 is non-compact. This allows us to use Eisenstein series and test our measure against them. There, we get an L -function: by a formula of Hecke (proved by direct unfolding), we get that this measure is given by a Dedekind zeta function $\zeta_E(s)$. A subconvex bound now yields entropy positivity. $\square$

We now discuss some work in progress with Matz. We want a similar theorem, but for a different group. Change PGL_3 to GSp_4 . This setting is tantalizing from an arithmetic point of view. However, when you try to move from GL_n to GSp_4 , what goes away is the Hecke formula. The ergodic part thus applies as before, but establishing positive entropy is a bit more mysterious. Suppose we have a maximal anisotropic torus $S \subset \mathrm{GSp}_4$. Then S gives rise to a quartic extension $E/\mathbb{Q}$ equipped with an involution σ , of which F is the subfield E^σ of fixed points. For instance, we could take E to be a quartic CM extension, and F would then be the unique quadratic totally real subfield. We then get

$$S_{E,F} \hookrightarrow \mathrm{GL}_{2,F}^* \hookrightarrow \mathrm{GSp}_4.$$

The first inclusion enjoys a Hecke-like formula, which we can use to establish positive entropy.

25. SIMRAN KHUNGER, *Extending support for the centered moments of the low lying zeros of cuspidal newforms*

Zeros of L -functions near central point and energy levels of heavy nuclei can both be modeled by eigenvalues of random matrices. Famous story of Montgomery and Dyson.

Limiting distribution of random matrix eigenvalues are reasonable to calculate. It's more elusive to establish the distribution of zeros of L -functions.

$H_k^*(N)$: holomorphic cuspidal newforms of weight k and level N .

$H_k^\pm(N)$, $\varepsilon_f = \pm 1$.

$\phi \in \mathcal{S}(\mathbb{R}^n)$, even, $\hat{\phi}$ compactly-supported, $\phi(0) > 0$, $\phi \geq 0$. One-level density

$$D(f, \phi) = \sum_{\gamma_f} \phi \left(\frac{\log R}{2\pi} \gamma_f \right).$$

Here R is the analytic conductor, and the γ_f are the imaginary parts of the zeros of the L -function associated with f .

Definition 84 (Notation for averages).

$$\langle D(f, \phi) \rangle_\pm := \mathbb{E}_{f \in H_{k \pm}(N)} D(f, \phi).$$

We'll be fixing the weight k and looking at the limit as $N \rightarrow \infty$. We'll also look at, e.g.,

$$\lim_{N \rightarrow \infty} \langle (D(f, \phi) - \langle D(f, \phi) \rangle_\pm)^n \rangle_\pm.$$

Random matrix theory analogues. For $U \in \mathrm{U}(M)$ with eigenvalues $e^{i\theta_n}$, and for $\phi \in \mathcal{S}(\mathbb{R})$, we define

$$Z_\phi(U) := \sum_{n=1}^M \sum_{j \in \mathbb{Z}} \phi \left(\frac{M}{2\pi} \cdots \right).$$

We also look at higher powers.

The examples of spectral distributions are

- for orthogonal matrices, $1 + (1/2)\delta_0$,
- for $\mathrm{SO}(\text{even})$, $1 + \sin(2\pi x)/2\pi x$,
- for $\mathrm{SO}(\text{odd})$, $1 - \sin(2\pi x)/2\pi x + \dots$.

26. GILLES FELBER, *L^2 -restriction norm of Siegel modular forms*

Siegel upper half-plane $\mathbb{H}^{(n)} \ni Z = X + iY$, with Z symmetric and $Y > 0$.
Mobius transforms by $\mathrm{Sp}_{2n}(\mathbb{R})$.

Siegel modular forms f of weight k .

$$f(Z) = \sum_{T>0} a_f(T) \det(T)^{\frac{k-3/2}{2}} e(\mathrm{tr}(TZ)).$$

L^2 -restriction norm to the imaginary axis $\mathrm{SL}_2(\mathbb{Z}) \backslash \{Y > 0\}$.

Parseval, spectral expansion over $\mathrm{GL}_2(\mathbb{Z}) \backslash \mathrm{GL}_2(\mathbb{R})$.

27. HUIMIN ZHANG, *Analytic twists of $\mathrm{GL}_2 \times \mathrm{GL}_2$ automorphic forms and applications*

REFERENCES