

✓ Congratulations! You passed!

Grade received 100% Latest Submission Grade 100% To pass 80% or higher

Go to next item

1.

1 / 1 point

$$J(\vec{w}, b) = \frac{1}{m} \sum_{i=1}^m L(f_{\vec{w}, b}(\vec{x}^{(i)}), y^{(i)})$$

The equation shows the cost function $J(\vec{w}, b)$ as the average of loss functions L over m training examples. A blue bracket above $J(\vec{w}, b)$ is labeled with a question mark. A blue bracket below the loss function L is also labeled with a question mark.

In this lecture series, "cost" and "loss" have distinct meanings. Which one applies to a single training example?

☒ Loss

✓ Correct

In these lectures, loss is calculated on a single training example. It is worth noting that this definition is not universal. Other lecture series may have a different definition.

☐ Cost

☐ Both Loss and Cost

☐ Neither Loss nor Cost

2.

1 / 1 point

Simplified loss function

$$L(f_{\vec{w}, b}(\vec{x}^{(i)}), y^{(i)}) = \begin{cases} -\log(f_{\vec{w}, b}(\vec{x}^{(i)})) & \text{if } y^{(i)} = 1 \\ -\log(1 - f_{\vec{w}, b}(\vec{x}^{(i)})) & \text{if } y^{(i)} = 0 \end{cases}$$

$$L(f_{\vec{w}, b}(\vec{x}^{(i)}), y^{(i)}) = -y^{(i)} \log(f_{\vec{w}, b}(\vec{x}^{(i)})) - (1 - y^{(i)}) \log(1 - f_{\vec{w}, b}(\vec{x}^{(i)}))$$

For the simplified loss function, if the label $y^{(i)} = 0$, then what does this expression simplify to?

☐ $\log(1 - f_{\vec{w}, b}(\vec{x}^{(i)})) + \log(1 - f_{\vec{w}, b}(\vec{x}^{(i)}))$

☒ $-\log(1 - f_{\vec{w}, b}(\vec{x}^{(i)}))$

☐ $-\log(1 - f_{\vec{w}, b}(\vec{x}^{(i)})) - \log(1 - f_{\vec{w}, b}(\vec{x}^{(i)}))$

☐ $\log(f_{\vec{w}, b}(\vec{x}^{(i)}))$

✓ Correct

When $y^{(i)} = 0$, the first term reduces to zero.