



**190F**  
Fall 2018

# Foundations of Data Science

## Lecture 14

---

Statistics

# **Announcements**

# **Probability & Simulation**

# Calculation

---

Roll a fair die 4 times.

What is  $P(\text{get at least one } 6)$ ?

---

# Calculation

---

Roll a fair die 20 times. What is  $P(\text{get at least one } 6)$ ?

Three ways to compute it:

- **Calculation:** Use math.
  - **Enumeration:** Count all outcomes.
  - **Estimation:** Randomly sample outcomes. Estimate.
-

# **Statistical Inference & Simulation**

# Terminology

---

- **Statistical Inference:** Making conclusions based on data in random samples
- **Parameter:** A number associated with a population.
- **Statistic:** A number calculated from a sample drawn at random from a population.

A statistic can be used to **estimate** a parameter, or to **test hypotheses** about the process that generated the data.

---

# Simulating a Statistic

---

- Figure out the code to generate *one* value of the statistic
- Create an empty array in which you will collect all the simulated values
- For each repetition of the process:
  - Simulate one value of the statistic
  - Append this value to the collection array
- At the end of all the repetitions, the array will contain all the simulated values

(Demo)

---



# Probability Distribution of a Statistic

---

- Values of a statistic vary because random samples vary
  - “Sampling distribution” or “probability distribution” of the statistic consists of:
    - All possible values of the statistic,
    - and all the corresponding probabilities
  - Can be hard to calculate
    - Either have to do the math,
    - or have to generate all possible samples and calculate the statistic based on each sample
-

# Empirical Distribution of a Statistic

---

- Empirical distribution of the statistic:
  - Based on simulated values of the statistic
  - Consists of all the observed values of the statistic,
  - and the proportion of times each value appeared
- Good approximation to the probability distribution of the statistic *if the number of repetitions in the simulation is large.*

(Demo)

---

# **Jury Selection**

# Swain vs. Alabama, 1965

---

- Talladega County, Alabama
  - Robert Swain, black man convicted of crime
  - Appeal: one factor was all-white jury
  - Only men 21 years or older were allowed to serve
  - 26% of this population were black
  - Swain's jury panel consisted of 100 men
  - 8 people on the panel were black (8%)
-

# Supreme Court Ruling

---

- About disparities between the percentages in the eligible population and the jury panel, the Supreme Court wrote:

*“... the overall percentage disparity has been small and reflects no studied attempt to include or exclude a specified number of [blacks]”*

- The Supreme Court denied Robert Swain’s appeal
-

# Sampling from a Distribution

---

- Sample at random from a categorical distribution:

```
sample_proportions(sample_size, pop_distribution)
```

- Samples at random from the population
- Returns an array containing the distribution of the categories in the sample

(Demo)

---

# **A Genetic Model**

# Steps in Assessing a Model


---

- Come up with a statistic that will help you decide whether the data support the model or an alternative view of the world.
  - Simulate the statistic under the assumptions of the model.
  - Draw a histogram of the simulated values. This is the model's prediction for how the statistic should come out.
  - Compute the observed statistic from the sample in the study.
  - Compare this value with the histogram.
  - If the two are not consistent, that's evidence against the model.
-



# Gregor Mendel, 1822-1884

---



# A Model

---

- Pea plants of a particular kind
  - Each one has either purple flowers or white flowers
  - Mendel's model:
    - Each plant is purple-flowering with chance 75%,
    - regardless of the colors of the other plants
  - Question:
    - Is the model good, or not?
-

# Choosing a Statistic

---

- Start with percent of purple-flowering plants in sample
- If that percent is much larger or much smaller than 75, that is evidence against the model
- ***Distance*** from 75 is the key
- Statistic:  
| sample percent of purple-flowering plants - 75 |
- If the statistic is large, that is evidence against the model

(Demo)

---

# Testing Hypotheses

# Choosing One of Two Viewpoints

---

- Based on data
    - “Chocolate has no effect on cardiac disease.”
    - “Yes, it does.”
    - “This jury panel was selected at random from eligible jurors.”
    - “No, it has too many people with college degrees.”
-

# Estimation

# How many enemy planes?

---



# Assumptions

---

- Planes have serial numbers  $1, 2, 3, \dots, N$ .
- We don't know  $N$ .
- We would like to estimate  $N$  based on the serial numbers of the planes that we see.

## The main assumption

- The serial numbers of the planes that we see are a uniform random sample drawn with replacement from  $1, 2, 3, \dots, N$ .
-



# Discussion question

---

If you saw these serial numbers, what would be your estimate of  $N$ ?

170	271	285	290	48
235	24	90	291	19

**One idea:** 291. Just go with the largest one.

---

# The largest number observed

---

- Is it likely to be close to  $N$ ?
  - How likely?
  - How close?

**Option 1.** We could try to calculate the probabilities and draw a probability histogram.

**Option 2.** We could simulate and draw an empirical histogram.

(Demo)

---

# Verdict on the estimate

---

- The largest serial number observed is likely to be close to  $N$ .
- But it is also likely to underestimate  $N$ .

## **Another idea for an estimate:**

Average of the serial numbers observed  $\sim N/2$

**New estimate:** 2 times the average

(Demo)

---