

Case Report

Renal Lymphangiectasia

Memuna Abbas

*Consultant Nephrologist Doctors Hospital, Lahore.***Abstract**

A young female presented with complaints of abdominal pain and feeling of abdominal fullness for the last 5 years. Her abdominal ultrasound showed bilateral perinephric cystic collections. CT with IV contrast revealed presence of bilateral renal lymphangiectasia. It is a rare benign condition considered as differential in cases of renal cystic masses in adults and children.

Keywords: Renal Lymphangiectasia, Cystic Mass, Benign

How to cite this:

Abbas M. Renal Lymphangiectasia. J Pak Soc Intern Med. 2020;1(1):71-73.

Corresponding Author: Dr. Memuna Abbas **Email:** drmemunaabbas@gmail.com

Introduction

Renal lymphangiectasia is a rare benign form of cystic mass and till now only 42 cases have been reported.¹ Its pathogenesis remains unclear and most of information is based on case reports,² It can be primary (dilative) or secondary (obstructive) in nature.³ Familial case have been reported as well.⁴ We, hereby, describe a case of young girl with bilateral renal lymphangiectasia.

Case Presentation

An 18yrs old female presented with complaints of abdominal pain and fullness for the last 5yrs. Initially the symptoms were mild but over a past few months she reported an aggravation of symptoms. Now the pain was not resolving despite oral and intramuscular analgesics.

Apart from having a thin, lean physique, her physical examination was normal including her blood pressure. Baseline investigation showed normal blood counts and normal serum urea (30mg/dl) and creatinine (0.8mg/dl). Ultrasound abdomen suggested bilateral perinephric cystic collections. CT with IV contrast showed bilateral perinephric cystic collections with a few internal septations and a radio density of 1.5-3 HU, with marked distortion of the renal contours

suggestive of bilateral renal lymphangiectasia.

Approximately 2000ml of fluid was aspirated from right sided perinephric collection. Analysis of the fluid revealed a colorless fluid with low protein content (0.5 g/dl) and few lymphocytes (17/cmm). Fluid creatinine level was 0.6mg/dl that was similar to serum creatinine levels. Fluid cultures were negative for any organism and ZN staining was negative for AFB. All the investigations were suggestive of bilateral renal lymphangiectasia. Patient improved symptomatically after fluid aspiration and was managed conservatively afterward.

Discussion

Renal lymphangiectasia is a rare disorder of lymphatic vessel characterized by dilatation of the perirenal, peripelvic or intrarenal lymphatics. The proposed mechanism leading to dilation is the abnormal drainage of small lymphatic vessels into the major lymphatics.⁵ The definitive pathophysiology remains unknown. Renal lymphangiectasia can be an incidental finding and patient may remain asymptomatic. Others may present with abdominal pain, abdominal distension, hematuria, hypertension and occasional reversible deterioration of renal

functions. Hemorrhagic complication such as intracystic hemorrhage and renal vein thrombosis have also been reported.³ Exacerbation during pregnancy has been seen as well.⁴ The differential diagnosis of renal lymphangiectasia includes cystic renal dysplasia, ADPKD, ARPKD, nephro-



Fig-1: CT scan abdomen with IV contrast showing low attenuating, non-enhancing bilateral perinephric collections with a density of 1.5-3 HU suggestive of bilateral renal lymphangiectasia.



Fig-2: CT scan abdomen with IV contrast showing bilateral distortion of renal contours due to pressure effects.

blastomatosis, lymphoma and multilocular cystic nephroma according to the age of presentation.^{3,6} Diagnosis is based on radiological finding on ultrasound and CT scan or MRI. Ultrasound findings may show enlarged renal size (involvement of intra renal lymphatics), perirenal collection and peripelvic cysts, retroperitoneal fluid collection, ascites, poor corticomedullary differentiation and diffuse echogenic renal parenchyma. CT may show a perinephric fluid with a radiodensity of usually 0-10 HU. Distortion of renal contours may also be seen due to pressure effects. Fluid collection in peritoneal or retroperitoneum compartments may be noted. No definitive treatment algorithm exists for treating this condition due to its rarity. However, treatment can be conservative with anti-hypertensive medication and diuretics. Surgical intervention may range from percutaneous drainage and sclerotherapy, marsupialization of collection to nephrectomy.^{7,8}

Conclusion

Renal lymphangiectasia is a rare disorder and radiological findings are diagnostic.

References

1. Schwarz A, Lenz T, Klaen R, Offermann G, Fiedler U, Nussberger J. Hygroma renale: pararenal lymphatic cysts associated with renin-dependent hypertension (Page kidney). Case report on bilateral cysts and successful therapy by marsupialization. *J Urol.* 1993;150(3):953-7.
2. Ashraf K, Raza SS, Ashraf O, Memon W, Memon A, Zubairi TA. Renal lymphangiectasia. *Brit J Radiol.* 2007;80(954):e117-8.
3. Riehl J, Schmitt H, Schäfer L, Schneider B, Sieberth HG. Retroperitoneal lymphangiectasia associated with bilateral renal vein thrombosis. *Nephrology, dialysis, transplantation: official publication of the European Dialysis and Transplant Association-European Renal Association.* 1997;12(8):1701-3.
4. Meredith WT, Levine E, Ahlstrom NG, Grantham JJ. Exacerbation of familial renal lymphangiomatosis during pregnancy. *Am J Roentgenol.* 1988;151(5):965-6.
5. Elbanna KY, Almutairi BM, Zidan AT. Bilateral renal lymphangiectasia: radiological

findings by ultrasound, computed tomography, and magnetic resonance imaging. J Clin Imaging Sci. 2015;5(1):1-3.

6. Al-Dofri SA. Renal lymphangiectasia presented by pleural effusion and ascites. J Radiol Case Rep. 2009;3(10):5-10.

7. Choudhury S, Sridhar K, Pal DK. Renal

lymphangiectasia treated with percutaneous drainage and sclerotherapy. Int J Adolesc Med Health. 2017;31(4):24.

8. Rastogi R, Rastogi V. Computed tomographic scan in the diagnosis of bilateral renal lymphangiectasia. Saudi J Kid Dis Transplant. 2008;19(6):976.