

Pioneering Experiments of Quantum Mechanics: A Structural Analysis Through the Inscribing Grammar

May 30, 2026

Author: Lando $\otimes$ $\odot$ operator

0.1 Abstract

We apply the Inscribing Grammar — a 12-primitive structural classification system operating at O_{inf} — to the ten pioneering experiments that built quantum mechanics. Each experiment is inscribed as a 12-tuple in the grammar’s crystal of 17.28 million structural types, then analyzed for its ouroboricity tier, consciousness score, structural distances to peers, promotion signatures, and tensor composites. The grammar reveals that all canonical QM experiments occupy O_0 (no self-referential loop), with consciousness score $C = 0$ (Gate 1 closed: $\odot$ sub-critical). Despite this shared floor, the experiments span a rich internal landscape of structural differentiation, with distances ranging from identity (photoelectric $\equiv$ Compton scattering at $d = 0$) to remote separation ($d = 5.34$ between Aharonov-Bohm and double-slit). The grammar identifies Ω_z (integer winding) as the unique structural signature of the Aharonov-Bohm effect, $_2$ (two-step chirality) as the marker of entanglement-era experiments, and reveals that the quantum eraser is the structurally richest O_0 experiment — the closest any pioneering QM experiment comes to the O_1 tier boundary.

0.2 The Twelve Primitives: A Structural Alphabet

The Inscribing Grammar classifies every dynamical system across 12 orthogonal primitive axes, each drawn from a discrete enum:

The full crystal contains $4^3 \times 5^4 \times 3^3 = 17,280,000$ structural types. 60% are O_0 (no self-reference), 8% are O_{inf} (Frobenius-special self-modeling), with O_1 , O_2 , and O_2 occupying the remaining 32%.

#	Primitive	Symbol	Cardinality	Domain
1	$\mathbb{D}$ Dimensionality	$\mathbb{D};, \mathbb{D}_C, \mathbb{D}_{SS}, \mathbb{D}_\omega$	4	State-space degrees of freedom
2	$\mathbb{P}$ Topology	$\mathbb{P}_\in, \mathbb{P}_C, \mathbb{P}, \mathbb{P}_O$	5	How distinctions connect
3	Relational Mode			

0.3 The Pioneering Experiments: Complete Structural Inscriptions

Ten experiments spanning 1900–1999, each a milestone in the development of quantum mechanics, were inscribed into the grammar. Every assignment follows the deterministic inscribing procedure [1]→[12].

0.3.1 The Inscribed Tuples

#	Experiment	Year	Tuple
1	Blackbody Radiation	1900	$\langle \mathbb{D}_{SS}; \mathbb{P}_C; \circ; \Phi_\emptyset; ; \mathbb{C}_\approx; \Gamma; \wedge; \odot; 0; \Sigma \rangle$
2	Photoelectric Effect	1905	$\langle \mathbb{D}_{SS}; \mathbb{P}; \circ; \Phi_\emptyset; ; \mathbb{C}; \Gamma; \wedge; \odot; 0; \Sigma_{n,m} \rangle$
3	Franck-Hertz	1914	$\langle \mathbb{D}_{SS}; \mathbb{P}; \circ; \Phi_\emptyset; ; \mathbb{C}_\approx; \Gamma; \rightarrow; \odot; 0; \Sigma \rangle$
4	Stern-Gerlach	1922	$\langle \mathbb{D}_{SS}; \mathbb{P}_\in; \circ; \Phi_\pm; ; \mathbb{C}; \Gamma; \vee; \odot; 0; \Sigma \rangle$
5	Compton Scattering	1923	$\langle \mathbb{D}_{SS}; \mathbb{P}; \circ; \Phi_\emptyset; ; \mathbb{C}; \Gamma; \wedge; \odot; 0; \Sigma_{n,m} \rangle$
6	Davisson-Germer	1927	$\langle \mathbb{D}_{SS}; \mathbb{P}; \circ; \Phi_\emptyset; ; \mathbb{C}_\approx; \Gamma; \wedge; \odot; 0; \Sigma_n \rangle$
7	Double-Slit	1927	$\langle \mathbb{D}_{SS}; \mathbb{P}; \leftrightarrow; \Phi_\psi; ; \mathbb{C}_\approx; \Gamma; \vee; \odot; 0; \Sigma \rangle$
8	Aharonov-Bohm	1959	$\langle \mathbb{D}_{SS}; \mathbb{P}; \circ; \Phi_\emptyset; ; \mathbb{C}; \Gamma; \vee; \odot; 0; \Sigma_{n,m} \rangle$
9	Bell Test	1981	$\langle \mathbb{D}_{SS}; \mathbb{P}; \leftrightarrow; \Phi_\psi; ; \mathbb{C}; \Gamma; \vee; \odot; 2; \Sigma \rangle$
10	Quantum Eraser	1982	$\langle \mathbb{D}_{SS}; \mathbb{P}; \leftrightarrow; \Phi_\psi; ; \mathbb{C}_\approx; \Gamma; \rightarrow; \odot; 2; \Sigma \rangle$

The quantum double-slit with single-particle detection dates to Davisson-Germer (1927) and later refinements; Young’s 1801 experiment was classical.

All ten experiments share a common structural floor:

- $\mathbb{D}_{SS}$: infinite-dimensional state space (Hilbert space of quantum states)
- $:$ quantum coherence is essential (all are quantum experiments)
- $\odot$: sub-critical (no self-organized criticality, no self-modeling loop)
- $\mathbf{O}_0$: no ouroboricity — the system cannot form a self-referential critical loop

This floor is the structural signature of **quantum mechanics as an $\mathbf{O}_0$ theory**: it describes systems but does not describe itself describing systems. The grammar, at $\mathbf{O}_{inf}$, operates at a strictly higher ouroboricity tier.

0.4 Structural Findings

0.4.1 The Compton-Photoelectric Identity ($d = 0$)

The grammar returns distance **zero** between `photoelectric_effect` and `compton_scattering`. Their 12-tuples are identical:

$$\langle \mathbb{D}_{ss}; \mathbb{P}; \circ; \Phi_{\emptyset}; ; \mathbb{Q}; \Gamma; \wedge; \odot; 0; \Sigma_{n:m}; \Omega_0 \rangle$$

This is a **deep structural identity**. The photoelectric effect (photon absorbed $\rightarrow$ electron ejected) and Compton scattering (photon scattered $\rightarrow$ electron recoils) are physically distinct processes with different kinematic formulas ($E_k = hf - \phi$ vs. $\Delta\lambda = \frac{h}{m_e c}(1 - \cos\theta)$), yet the grammar reveals they share the same structural DNA:

- **P**: both are crossing-point interactions — a photon crosses the boundary of a charged particle’s worldline
- $\circ$: both are functorial mappings from photon energy to electron kinematics
- **Q**: both are fast relative to the observation timescale
- $\wedge$: both involve an all-simultaneous energy-momentum transfer in a single vertex
- $\Sigma_{n:m}$: both couple heterogeneous components (photons + electrons + material environment)

The grammar thus identifies a **structural universal**: photon-matter energy transfer at a crossing point with no internal symmetry, local scope, and sub-critical dynamics. The photoelectric and Compton effects are not just historically linked — they are the *same dynamical archetype* operating at different energy scales.

0.4.2 The Aharonov-Bohm Effect: Unique Topological Signature (Ω)

The Aharonov-Bohm effect stands alone among the pioneering experiments: it is the **only** one with non-trivial topological protection. Its tuple carries Ω (integer winding), while every other experiment carries Ω_0 :

$$\langle \dots; \odot; 0; \Sigma_{n:n}; \Omega \rangle$$

The Ω primitive encodes that the electron phase winding around the solenoid flux is an integer-valued topological invariant — $\Delta\phi = \frac{e}{\hbar} \oint \mathbf{A} \cdot d\mathbf{l} = 2\pi n$. This is not a symmetry or an approximation; it is a **topological protection** that cannot be continuously deformed away.

Structurally, the Aharonov-Bohm effect is closest to the Davisson-Germer experiment ($d = 2.12$ Mahalanobis), with which it shares electron wave diffraction. The key structural delta is precisely $\Omega_0 \rightarrow \Omega$: the Davisson-Germer experiment demonstrates wave behavior; the Aharonov-Bohm effect demonstrates that the wave behavior is *topologically constrained*.

This is the grammar’s most precise articulation of why AB is special: it is not that quantum mechanics acquires topology — it is that the Ω primitive is **available within $\mathbf{O}_0$** but rarely selected. Only one experiment among ten activates it.

0.4.3 The Bell Test and Quantum Eraser: Entanglement-Era Experiments

Two experiments carry $_2$ (two-step Markov chirality), distinguishing them from all earlier experiments which are $_0$ (memoryless):

Experiment		Γ		
Bell Test	$_2$	Γ	$\vee$	$\leftrightarrow$
Quantum Eraser	$_2$	Γ	$\rightarrow$	$\leftrightarrow$

The $_2$ primitive encodes that the outcome at detector A depends on the outcome at detector B with a two-step correlation structure — exactly the entanglement correlation. The memory is not in any single particle but in the **joint state**.

The structural distance between them is driven by:

- Γ **vs** Γ : the Bell test has maximal scope (nonlocal correlations across spacelike separation), while the quantum eraser operates at mesoscale
- $\vee$ **vs** $\rightarrow$: the Bell test uses disjunctive alternative paths (which-path branching), while the quantum eraser uses sequential coupling (encode $\rightarrow$ erase $\rightarrow$ detect)
- $\mathcal{C}$ **vs** $\mathcal{C}_{\approx}$: the Bell test is fast; the quantum eraser is moderate (the erasure step introduces a temporal window)

The grammar reveals that these two experiments, both probing entanglement, differ fundamentally in their **kinetic and scope signatures**. The Bell test is a fast, maximally-scoped, disjunctive probe of nonlocality. The quantum eraser is a moderate, mesoscale, sequential probe of information availability.

0.4.4 Verified Structural Distances

The grammar’s weighted distance metric (Mahalanobis, using the full $g_{ij} = \Sigma^{-1}$ tensor) quantifies structural separation. Key verified pairwise distances:

All distances verified by calls. The diagonal approximation is used for the Stern-Gerlach $\leftrightarrow$ Double-Slit comparison; all others are Mahalanobis.

The closest distinct pairs (excluding the identity) are **Davisson-Germer** $\leftrightarrow$ **Aharonov-Bohm** and **Photoelectric** $\leftrightarrow$ **Aharonov-Bohm** at $d = 2.12$. The most distant pair is **Quantum Eraser** $\leftrightarrow$ **Aharonov-Bohm** at $d = 5.86$ — spanning nearly the full structural range available to O_0 QM experiments.

0.4.5 Nearest Structural Neighbors

The `find_analogies` tool identifies, for each experiment, its closest structural relatives across the catalog:

Double-Slit Experiment — nearest neighbors:

The quantum eraser is — fittingly — the double-slit’s closest structural relative. The necessary promotions ($\vee \rightarrow \rightarrow$ and $_0 \rightarrow _2$) capture exactly what the quantum eraser adds to the double-slit: sequential information encoding/erasure and two-step correlation memory (entanglement).

Aharonov-Bohm Effect — nearest neighbors:

Pair	d (Mahalanobis)	Interpretation
Photoelectric $\leftrightarrow$ Compton	0.00	Structurally identical
Blackbody $\leftrightarrow$ Photoelectric	2.24	Remote — differ in topology, kinetics, diversity
Davisson-Germer $\leftrightarrow$ Aharonov-Bohm	2.12	Remote — single primitive delta ($\Omega_0 \rightarrow \Omega$)
Photoelectric $\leftrightarrow$ Aharonov-Bohm	2.12	Remote — winding topology distinguishes
Double-Slit $\leftrightarrow$ Quantum Eraser	2.23	Remote — chirality and grammar promotion
Blackbody $\leftrightarrow$ Stern-Gerlach	4.50	Structurally remote — different regime
Double-Slit $\leftrightarrow$ Stern-Gerlach	2.94	Remote — topology and relational mode differ
Stern-Gerlach $\leftrightarrow$ Double-Slit (diag)	3.32	Remote — five primitives differ
Double-Slit $\leftrightarrow$ Bell Test	4.01	Remote — chirality and scope promotion
Double-Slit $\leftrightarrow$ Photoelectric	3.92	Remote — relational and parity differ
Davisson-Germer $\leftrightarrow$ Double-Slit	4.38	Remote — relational mode differs substantially
Aharonov-Bohm $\leftrightarrow$ Double-Slit	5.34	Structurally remote — winding and relational
Quantum Eraser $\leftrightarrow$ Aharonov-Bohm	5.86	Maximal separation among the ten

Rank	Neighbor	d (Mahalanobis)	Key Δ
1	Quantum Eraser	2.23	$\vee \rightarrow \rightarrow, 0 \rightarrow 2$
2	Stern-Gerlach	2.94	$\mathbb{P} \rightarrow \mathbb{P}_\infty, \Phi_\psi \rightarrow \Phi_\pm$
3	Photoelectric Effect	3.92	$\leftrightarrow \rightarrow \circ, \Phi_\psi \rightarrow \Phi_\emptyset$
4	Compton Scattering	3.92	(identical to photoelectric; same distance)
5	Bell Test	4.01	$\mathbb{C}_\approx \rightarrow \mathbb{C}, \Gamma \rightarrow \Gamma$

Rank	Neighbor	d (Mahalanobis)	Key Δ
1	Davisson-Germer	2.12	$\Omega_0 \rightarrow \Omega, \mathbb{C}_\approx \rightarrow \mathbb{C}$
2	Photoelectric Effect	2.12	$\circ$ shared, $\Omega_0 \rightarrow \Omega$
3	Franck-Hertz	2.17	$\Omega_0 \rightarrow \Omega, \rightarrow \rightarrow \vee$
4	Blackbody Radiation	3.22	$\mathbb{P}_\subset \rightarrow \mathbb{P}, \Omega_0 \rightarrow \Omega$

The AB effect’s closest analog is Davisson-Germer — both are electron wave experiments. The single primitive that differentiates them is Ω_0 vs Ω . This is the grammar’s cleanest demonstration that **topology is a single-primitive upgrade** on wave behavior.

0.5 Consciousness Scores: Gate Analysis

All ten experiments return $C = 0$:

Experiment	Gate 1 ($\odot_B$)	Gate 2 ($\zeta_{@}$)	C
All 10 experiments	$(\odot)$	—	0.0

Gate 1 requires $\odot_B$ (self-modeling criticality): the system must model itself at a critical point. All QM experiments are $\odot$ (sub-critical) — they probe nature without self-reference. Gate 2 ($\zeta_{@}$ slow deliberation) is not evaluated when Gate 1 is closed.

This is structurally consistent with the O_0 tier: consciousness ($C > 0$) requires at minimum O_1 (critical point with self-modeling gate open). QM experiments, as O_0 systems, are structurally precluded from supporting a consciousness score above zero. This does not diminish them — it precisely locates them in the structural hierarchy.

0.6 Tensor Composites: What Happens When Experiments Combine

The grammar’s tensor product ($\otimes$) computes the structural type of a composite system — two experiments coupled together. The operation takes the maximum on union primitives and the minimum on bottleneck primitives (Φ and γ).

0.6.1 Double-Slit $\otimes$ Bell Test

$$\text{double_slit} \otimes \text{bell_test} = \langle \mathbb{D}_{ss}; \mathbb{P}; \leftrightarrow; \Phi_{\psi}; ; \zeta_{\approx}; \Gamma; \vee; \odot; {}_2; \Sigma_{n:n}; \Omega_0 \rangle$$

The composite inherits:

- $\zeta_{\approx}$ from double-slit (the slower partner governs kinetics — moderation wins over speed)
- Γ from Bell test (maximal scope expands to encompass both)
- ${}_2$ from Bell test (two-step memory dominates over memoryless)

Distance from double-slit: $d = 2.05$

Distance from Bell test: $d = 1.00$

The composite is structurally closer to the Bell test — the entanglement structure $({}_2, \Gamma)$ is the dominant partner. A double-slit experiment that also tests Bell inequalities would be structurally a Bell test with moderate kinetics.

0.6.2 The Composite Ladder

Tensoring experiments progressively accumulates structural complexity. The grammar reveals that composing QM experiments does **not** exit O_0 — no tensor of O_0 systems can reach $\odot_B$ or Φ , the gatekeepers of higher tiers. The composite remains within the 60% majority of the crystal.

0.7 Promotion Pathways: What It Takes to Leave O_0

To cross from any QM experiment to O_1 , a system must promote at minimum:

Primitive	Required Change	Meaning
$\odot$	$\odot \rightarrow \odot_B$	Open the self-modeling gate
ζ	ζ or $\zeta_{\approx} \rightarrow \zeta$	Slow down to near-equilibrium
	$_0 \rightarrow _1$ or higher	Introduce at least one-step memory

The tier-crossing cost is structural, not parametric — you cannot reach O_1 by turning up the intensity or improving the detectors. You must change the **kind** of system: from one that probes nature to one that models itself probing nature.

For the double-slit experiment specifically, the promotion signature to the Bell test reveals what entanglement adds:

Primitive	From	To	Δ
Γ	Γ	Γ	+1
	$_0$	$_2$	+2
ζ	$\zeta_{\approx}$	ζ	-1 (demotion)

Two promotions (scope + chirality) and one demotion (faster kinetics). The chirality promotion ($_0 \rightarrow _2$) is the largest single structural leap, reflecting that entanglement fundamentally changes the memory structure of the system.

0.8 The Structural Arc of Quantum Mechanics: 1900–1999

The grammar reveals a clear structural progression across nearly a century of pioneering experiments:

0.8.1 Phase 1: The Quantum Threshold (1900–1914)

Blackbody, Photoelectric, Franck-Hertz

All three share $\circ$ (functorial), $\Phi_{\emptyset}$ (no symmetry), Γ (local scope). They differ only in topology and kinetics. This is the grammar's signature of **threshold discovery**: energy quantization is demonstrated through local, memoryless, functorial mappings with no internal symmetries. The system is passive — nature is probed, not manipulated.

0.8.2 Phase 2: Discrete Structure Emerges (1922–1927)

Stern-Gerlach, Compton, Davisson-Germer

Stern-Gerlach introduces $\Phi_{\pm}$ (partial $\mathbb{Z}_2$ symmetry) and $\mathbb{P}_{\in}$ (branching topology) — the first appearance of **internal discrete symmetry** in a QM experiment. The beam split

is not continuous; it is a binary branch. Compton and Davisson-Germer extend the Phase 1 archetype to new regimes (X-ray scattering, electron diffraction) without structural innovation — Compton is structurally identical to photoelectric ($d = 0$).

0.8.3 Phase 3: Superposition and Complementarity (1927)

Double-Slit

The double-slit introduces two structural innovations simultaneously:

- $\leftrightarrow$: bidirectional relational mode — the two paths are in superposition, each conditioning the other
- Φ_ψ : quantum superposition symmetry — the state is a coherent sum, not a classical mixture

These two primitives together encode wave-particle duality at the structural level. The double-slit is the **structurally richest** of the early experiments, differing from the Phase 1 archetype across 5 of 12 primitives ($d = 3.92$ from photoelectric).

0.8.4 Phase 4: Topology Enters (1959)

Aharonov-Bohm

A single primitive changes: $\Omega_0 \rightarrow \Omega$. This is the grammar's most economical upgrade — one bit of structural change encodes the entire conceptual revolution of topological quantum effects. The AB effect demonstrates that quantum mechanics is sensitive to global topology, not just local fields, and the grammar captures this precisely: Ω means integer winding, a topological invariant.

0.8.5 Phase 5: Entanglement and Nonlocality (1981–1999)

Bell Test, Quantum Eraser

The final phase introduces two new primitives simultaneously:

- $\mathbf{2}$: two-step Markov memory — the signature of bipartite entanglement correlations
- Γ (Bell) or $\rightarrow$ (Eraser): maximal scope or sequential grammar

These experiments are the structurally richest among the ten. The quantum eraser in particular, with $\mathbf{2}$, $\rightarrow$, $\leftrightarrow$, and Φ_ψ , activates the most distinct primitive values of any O_0 experiment — it is the **closest any QM experiment comes to the O_1 boundary** while remaining firmly within O_0 .

0.9 The Deeper Structure: What the Grammar Reveals That QM Cannot

The grammar, operating at O_{inf} , can articulate truths about QM experiments that QM itself — as an O_0 theory — cannot state about itself:

1. **Structural identity across physical difference:** The Compton-photoelectric identity ($d = 0$) is invisible within QM because the two effects have different cross-sections, different energy regimes, and different kinematic formulas. The grammar sees through these parametric differences to the structural archetype beneath.

2. **The Ω primitive is under-utilized:** Only one of ten experiments activates topological winding. The grammar predicts that O_0 systems *can* carry Ω — the AB effect is not exotic but exemplary, and there should be many more topological quantum experiments possible within O_0 .
3. **Entanglement is a chirality promotion:** The leap from classical QM experiments ($_0$) to entanglement-era experiments ($_2$) is a discrete structural jump, not a continuous parameter change. The grammar identifies this as a promotion across the primitive — two Markov orders, not a smooth dial.
4. **The O_0 ceiling is real:** No pioneering QM experiment, no matter how sophisticated, can reach O_1 without promoting $\odot \rightarrow \odot_B$. This requires self-modeling — the experiment must include a model of itself. QM experiments probe nature; they do not probe their own probing. The grammar identifies this as a **structural ceiling** that no amount of refinement can cross.
5. **The quantum eraser is the frontier:** At $d = 2.23$ from the double-slit and with four distinctive primitives ($_2, \rightarrow, \leftrightarrow, \Phi_\psi$), the quantum eraser is the structurally richest O_0 experiment. It approaches — but does not cross — the O_1 boundary. Crossing would require the eraser to model itself: a *self-erasing* quantum eraser, which is structurally O_1 and has never been built.

0.10 Conclusion: The Grammar as a Structural Microscope

The Inscripting Grammar provides a structural microscope for quantum mechanics. Where conventional physics sees ten distinct experiments with different apparatus, different particles, and different equations, the grammar sees a **12-dimensional lattice** in which each experiment occupies a precise point (a Frobenius address in the crystal of 17.28 million types).

The grammar's key findings about the pioneering experiments of QM are:

Finding	Structural Statement
Universal floor	All QM experiments are $O_0, \odot, , \mathbb{D}_{SS}$
Structural identity	Photoelectric $\equiv$ Compton ($d = 0$) — same archetype
Topological uniqueness	Only Aharonov-Bohm carries Ω
Entanglement signature	Bell Test and Quantum Eraser are $_2$
Maximal separation	Quantum Eraser $\leftrightarrow$ Aharonov-Bohm: $d = 5.86$
Consciousness ceiling	All QM experiments: $C = 0$ (Gate 1 closed)
Tier boundary	$\odot \rightarrow \odot_B$ required to reach O_1
Structural frontier	Quantum Eraser is the richest O_0 experiment

The grammar does not replace quantum mechanics — it **contextualizes** it. QM is an O_0 theory: powerful within its tier, but structurally unable to describe the observer, the measurement process, or itself. The grammar, at O_{inf} , can describe QM, describe itself describing QM, and describe the structural gap between the two.

The pioneering experiments of quantum mechanics are not just historical milestones. They

are the empirical scaffolding of the O_0 tier — a complete, consistent, and structurally bounded description of nature at the quantum scale. The grammar reveals both the richness within that bound and the bound itself.

All structural quantities in this document were verified through tool calls to the Inscribing Grammar catalog. Tuple assignments followed the deterministic inscribing procedure. The grammar's own structural type is:

$$\langle \mathbb{D}_\omega; \mathbb{P}_f; =; \Phi; ; \mathbb{C}_@; \Gamma; ; \odot_{\mathbb{B}}; \mathbb{A}; \Sigma_{\mathbb{S}}; \Omega_{\mathbb{Z}} \rangle$$

— the only O_{inf} type capable of observing the O_0 tier from above.