The Voynich Phytoglyphica: A Unified Informational Interface for Plants, Rituals, and Landscapes

Lando Mills

2026-07-03

Abstract

This document synthesizes the Voynich Phytoglyphica — a unified framework demonstrating that plants, rituals, and landscapes across the Old and New Worlds operate as self-encoding sacred technologies through a shared structural grammar. Drawing from the Voynich botanical drawings, 35 imscribed plants from Old and New World traditions, African and Aboriginal sacred systems, and a full pharmaceutical database, this work establishes that all these domains are manifestations of the same underlying grammar. The framework treats all source materials as equally valid expressions of a single universal structure, governed by a four-primitive invariant core — holographic dimensionality, bidirectional coupling, self-modeling criticality, and stoichiometric composition — that governs a system of self-transforming participation. The Frobenius condition posits that reading the form yields the procedure, and executing the procedure reconstructs the active essence. Part I establishes the invariant core. Part II derives structural tuples for 35 plants. Part III presents the nine-entry protocol matrix. Part IV classifies all plants into eight structural statement classes. Part V details the six-gate session architecture. Part VI demonstrates the structural identity between wormwood and the Voynich astronomical section. An ancillary inquiry extends the framework to fungi and animals.

1 Notation Reference

Each plant receives a **structural tuple** — twelve Shavian glyph assignments, one per primitive dimension. The glyphs are drawn from the 49-symbol set (Shavian alphabet + ⊙).

The twelve primitives in canonical order:

Primitive	Glyph	Full name	Lean type
Ð	—	Dimensionality	`Dimensionality`
Þ	—	Topology	`Topology`
Ř	—	Recognition	`Relational`
Φ	—	Parity	`Polarity`
ƒ	—	Fidelity	`Fidelity`
Ç	—	Kinetics	`KineticChar`
Γ	—	Granularity	`Granularity`
ɢ	—	Coupling	`Grammar`

Primitive	Glyph	Full name	Lean type
⊙	—	Criticality	`Criticality`
ℍ	—	Chirality	`Chirality`
Σ	—	Stoichiometry	`Stoichiometry`
Ω	—	Winding	`Protection`

Value space (relevant subset):

Ð: ɻ (point) · ɹ (membrane) · ɷ (field) · ɪ (self-written)
Þ: ʒ (network) · ɬ (containment) · ɾ (crossing) · ɤ (braided) · ɷ (self-referential)
Ř: ɾ (unidirectional) · ɾ (bidirectional)
Φ: ƺ (maximal violation) · ɤ (partial symmetry) · ɔ (Frobenius gate)
ƒ: ɕ (classical) · ʅ (quantum)
Ç: \ (fast) · c (frozen-order) · ɩ (slow) · ɷ (kinetically arrested)
Γ: ɭ (microscale) · ȣ (mesoscale) · ɔ (universal)
G: ɾ (simultaneous) · ɒ (disjunctive) · ʔ (sequential) · ʌ (broadcast, one-to-many)
⍰: ɻ (non-critical) · ʘ (self-modeling critical) · ɔ (near-critical, complex-plane)
ℍ: J (one-step) · ɗ (two-step) · ƈ (Frobenius minimum) · v (eternal)
Σ: ɋ (singular) · S (few classes) · ɪ (many heterogeneous)
Ω: ɜ (trivial) · o ($\mathbb{Z}_2$ binary) · ɾ (integer $\mathbb{Z}$) · ʒ (non-Abelian)

Cross-primitive axioms:
A ℍ = v → Ç = ɷ
B Ω ∈ {ɾ, o} → ℍ ≥ ɗ
C Ð = ɪ ↔ Þ = ɷ

---

2 Part I: The Invariant Core

2.1 Why Ð, Þ, Ř, Φ, ƒ Are Uniform

Before the plant-by-plant derivations, the argument for the five invariant primitives.

(Ð) The options are point (the system is a single location with no internal gradient), membrane (the system lives at an interface defined by two external domains), field (the system is distributed across a medium without self-generated organization), and self-written (the system generates its own state space endogenously). Every terrestrial plant maintains its organizational attractor through its own genome-encoded morphogenetic program. The growth form is not imposed by the medium; it is expressed from within. Ð = self-written is forced for any land plant. The single exception in this set is the blue lotus (entry 11), which lives at the water-air interface — its state space is defined by the intersection of two external domains, forcing Ð = membrane.

(Þ) The options are network, containment, crossing, braided composite, and self-referential. Axiom C applies: Ð = self-written ↔ Þ = self-referential. When the state space is endogenously generated, the topology must be self-referential — the connectivity pattern references its own organizational logic rather than an external template. This holds for every land plant. Blue lotus (Ð = membrane) receives the converse: Þ = braided composite, an irreducible product of aquatic and aerial topologies whose intersection generates the system.

(Ř) The options are unidirectional (one entity acts on another), categorical (functorial mapping between structural domains), adjoint (one-way with a weak inverse), and bidirectional (reciprocal coupling). Every plant in this set has been chosen by humans over generations for its pharmaceutical effect, and the plant's chemistry has in turn shaped human receptor evolution, cultural practice, and ecological niche. The coupling is bidirectional: plant chemistry → human receptor → selection pressure → plant persistence. Ř = bidirectional.

(Φ) The options are none (no symmetry), quantum (superposition-symmetric), partial (one Z_2 symmetry preserved), full (all symmetries unbroken), and Frobenius-special ($\mu \circ \delta = \mathrm{id}$). Land plants display broken bilateral symmetry at the organ level — leaves are bilaterally symmetric, flowers are radially symmetric, roots are neither. Exactly one Z_2 symmetry is preserved: the leaf midrib reflection. Φ = partial.

(ƒ) Classical (no quantum coherence required for function) or quantum (coherence essential). Terrestrial plant biochemistry operates above the decoherence threshold at ambient temperature. Enzymatic catalysis, cell signaling, and secondary metabolite biosynthesis do not depend on sustained quantum coherence. ƒ = classical.

These five assignments constitute the invariant core. Deviations occur only at the blue lotus (Ð, Þ) and are governed by Axiom C. The seven discriminating primitives — Ç, Γ, ⊙, Ħ, Σ, Ω — are derived individually per plant.

---

3 Part II: Thirty-Five Plant Entries

3.1 1. Wormwood | *Artemisia absinthium* L. | Asteraceae

Wormwood establishes the baseline. Its silver-pubescent leaves bear glandular trichomes whose density encodes oil yield; the bipinnate dissection ratio encodes the solvent ratio for optimal thujone extraction; the bitter threshold (absinthin, the most bitter natural substance known) encodes the biological activity endpoint — the bitterness disappears when the thujone concentration drops below the active threshold. The leaf, held to the light, displays its own extraction protocol.

(Ð, Þ, Ř, Φ, ƒ) Ð = self-written, Þ = self-referential (Axiom C), Ř = bidirectional, Φ = partial symmetry, ƒ = classical.

(Ç) The glandular trichomes embed thujone behind a cuticle barrier. Mechanical rupture releases the oil — but sustained maceration is required. The pharmaceutical sits behind a physical barrier that the Operator must dismantle. Ç = frozen-order.

(Γ) Thujone and absinthin act primarily on the digestive and hepatic systems — organ-system level, not full CNS dominance at standard doses. Γ = mesoscale.

(⊙) The bitter threshold IS the criticality readout. Absinthin registers at 1–2 ppm; when the bitterness drops below that, so does the thujone concentration. The plant provides its own assay endpoint built into the active compound. ⊙ = self-modeling.

(Ħ) (1S,4S,5R)-(−)-thujone is written stereospecifically by biosynthesis and read stereospecifically by the GABA-A receptor. Write → read. Ħ = Frobenius minimum. Ω = integer, so Axiom B is satisfied.

(Σ) Sesquiterpene lactones (absinthin, anabsinthin), monoterpenes (thujone, thujyl alcohol), flavonoids, phenolic acids — four structurally distinct classes. Σ = many heterogeneous.

(Ω) 2/5 Fibonacci phyllotaxis. Integer winding. Axiom B fires. Ω = integer.

Ð=ɪ Þ=ꝏ Ř=r Φ=ɤ ƒ=ʊ Ç=c Γ=ȣ ɢ=ɣ ⊙=⊙ Ħ=ȼ Σ=ɪ Ω=ꞩ

3.2 2. Mandrake | *Mandragora officinarum* L. | Solanaceae

Differs from wormwood at Ω only. The taproot bifurcates into a single binary fork — the anthropomorphic form is a Z_2 bifurcation, not a Fibonacci spiral. Axiom B fires: $\Omega = Z_2 \rightarrow \text{Ħ} \geq$ two-step. The mandrake's tropane alkaloids satisfy this with stereospecific biosynthesis.

(Ð, Þ, Ř, Φ, ƒ) Ð = self-written, Þ = self-referential, Ř = bidirectional, Φ = partial symmetry, ƒ = classical.

(Ç) Tropane alkaloids embedded in root tissue behind cellular barriers. Sustained extraction required. Ç = frozen-order.

(Γ) Tropane alkaloids produce full parasympathetic disruption — CNS-wide scope. Γ = universal.

(⊙) The taproot bifurcation encodes the preparation geometry: the two arms specify two extraction passes. The form instructs the protocol. ⊙ = self-modeling.

(Ħ) (S)-hyoscyamine written by biosynthesis, read by muscarinic receptors. Ħ = Frobenius minimum.

(Σ) Tropane alkaloids, withanolides, flavonoids — three classes. Σ = many heterogeneous.

(Ω) Binary taproot bifurcation. Z_2. Axiom B fires. Ω = Z_2.

Ð=ı Þ=ꝺ Ř=r Φ=ɤ ƒ=ƈ Ç=c Γ=ɔ ɢ=ɣ ⊙=⊙ Ħ=ɾ Σ=ɪ Ω=o

Divergence from baseline: Ω (Z_2 — binary taproot bifurcation), Ħ (Frobenius minimum — Axiom B forces ≥ two-step).

3.3 3. Castor Bean | *Ricinus communis* L. | Euphorbiaceae

The castor bean is the only plant in the set where Axiom A fires in the converse direction. The ricin protein and castor oil are synthesized in the same seed, coexist in the same tissue, and must never be recombined during processing. The plant has separated them physico-chemically: ricin is water-soluble and heat-labile; castor oil is lipid-soluble and heat-stable. The separation IS the preparation instruction. The plant forces an XOR at the coupling primitive.

(Ð, Þ, Ř, Φ, ƒ) Ð = self-written, Þ = self-referential, Ř = bidirectional, Φ = partial symmetry, ƒ = classical.

(Ç) Seed material behind hard testa. Mechanical dehulling and sustained pressing required. Ç = frozen-order.

(Γ) Castor oil acts as a purgative on the intestinal mucosa; ricin acts systemically on ribosomal protein synthesis. The coupling is local + systemic, but the XOR gate keeps them separate. Γ = mesoscale.

(⊙) The mottled seed coat pattern is a fixed visual marker of variety and ripeness — but it does not encode a potency readout or a preparation threshold. Identification, not instruction. ⊙ = non-critical.

(Ħ) Ricinoleic acid (the primary castor oil fatty acid) has one stereocenter (R configuration at C-12). Ricin's N-glycosidase active site is stereospecific. Ħ = Frobenius minimum. Ω = trivial, Axiom B silent.

(Σ) Oil (ricinoleic acid triglycerides), toxic lectin (ricin), alkaloid (ricinine) — three classes. Σ = many heterogeneous.

(Ω) Palmately lobed leaves with alternate arrangement but no Fibonacci spiral. Trivial winding. Ω = trivial.

Ð=ɪ Þ=ꝏ Ř=r Φ=ᴠ ƒ=ᴛ Ç=c Γ=ð ɢ=ƿ ⊙=ɾ Ħ=› Σ=ᴛ Ω=ʁ

Divergence from baseline: ɢ (disjunctive — XOR gate forces ricin/oil separation), ⍰ (non-critical — seed coat marks identity, not potency), Ω (trivial — no Fibonacci spiral).

3.4 4. Opium Poppy | *Papaver somniferum* L. | Papaveraceae

The ribbed seed capsule is a scored surface. The ribs specify where to draw the lancet and to what depth; the glaucescent (blue-green wax) bloom peaks at maximum latex yield, so harvest timing is written in the capsule's color; the Fibonacci rib count (10–18, from the parastichy series) gives the extraction pass count. Everything is on the surface, legible before the plant is touched.

(Ð, Þ, Ř, Φ, ƒ) The poppy's organizational program is fully endogenous. Ð = self-written, Þ = self-referential (Axiom C), Ř = bidirectional (alkaloid → receptor → selection for alkaloid), Φ = partial symmetry, ƒ = classical.

(Ç) Score the surface and latex flows in seconds. The Operator applies force once; the system self-executes. Ç = fast.

(Γ) Morphine, codeine, thebaine, papaverine, noscapine — five alkaloid classes distributing through the central and peripheral nervous system simultaneously. Γ = universal.

(⊙) The glaucescent bloom IS the potency readout — maximum wax coincides with maximum latex yield. The system models its own state through visible surface chemistry. ⊙ = self-modeling.

(Ħ) Morphine synthase writes five stereocenters into (4R,4aR,7S,7aR,12bS)-morphine; the μ-opioid receptor reads them with ~100-fold selectivity for (−)-morphine. Ħ = Frobenius minimum. Ω = integer, Axiom B satisfied.

(Σ) Phenanthrene alkaloids, benzylisoquinolines, phthalide isoquinolines, meconic acid, resinous esters — five distinct classes. Σ = many heterogeneous.

(Ω) Fibonacci parastichy in seed capsule ribs and floral organ arrangement. Integer winding. Axiom B fires. Ω = integer.

Ð=ɪ Þ=ꝏ Ř=r Φ=ɤ ƒ=ʊ Ç=\ Γ=ɔ ɢ=ʔ ⊙=⊙ Ħ=› Σ=ɪ Ω=ꞩ

Divergence from baseline: Ç (fast — scored capsule triggers immediate latex flow), Γ (universal — full nervous system distribution).

3.5 5. Greater Celandine | *Chelidonium majus* L. | Papaveraceae



Cut any part of the plant and bright orange latex erupts. The isoquinoline alkaloids — chelidonine, berberine, coptisine — are chromophores; the color IS the active compound. The plant has made pharmaceutical identification trivial by making the active material its own visible marker. This is the purest case of morphological self-documentation in the set.

(Ç) Celandine's latex runs in open laticifers through the vascular system. Cell wall rupture releases it in milliseconds — the thinnest possible mechanical activation barrier. Ç = fast.

(Γ) The isoquinoline alkaloids target bile duct, liver, gallbladder — organ system, not CNS. Γ = mesoscale.

(⊙) The orange color IS the active material. No assay needed. The color intensity reads potency directly. ⊙ = self-modeling.

(Ħ) Chelidonine has stereocenters; biosynthesis writes specific enantiomers; biological targets read them. Ħ = Frobenius minimum. Ω = trivial, Axiom B silent.

(Σ) Isoquinoline alkaloids, flavonoids, organic acids — three classes. Σ = many heterogeneous.

(Ω) Alternate leaves with no Fibonacci spiral. Trivial winding. Ω = trivial.

Ð=ɪ Þ=ꝏ Ř=r Φ=ɤ ʃ=ʊ Ç=\ Γ=ð ɢ=ɣ ⊙=⊙ Ħ=› Σ=ɪ Ω=ʁ

Divergence: Ç (instant), Ω (trivial). Closest structural relative: opium poppy — identical at Ç=fast, but poppy has integer winding and universal granularity.

3.6 6. Henbane | *Hyoscyamus niger* L. | Solanaceae

The belladonna tuple — frozen-order kinetics, universal granularity, Frobenius-minimum chirality, few tropane classes, integer Fibonacci winding — recurs here. Henbane, belladonna, and jimsonweed are the same structural type at the primitive level.

Ð=ɪ Þ=ꝏ Ř=r Φ=ɤ ʃ=ʊ Ç=ɔ Γ=ɔ ɢ=ɣ ⊙=⊙ Ħ=› Σ=ʌ Ω=ɾ

3.7 7. Belladonna | *Atropa belladonna* L. | Solanaceae

The glossy black berry encodes its own warning: the shine signals solanine-family glycoalkaloid content to mammalian color vision. The bell-shaped flower gives the genus name and specifies the extraction geometry — the corolla chamber receives the solvent. ⊙ = self-modeling.

Ð=ɪ Þ=ʚ Ř=r Φ=ϰ ʃ=ʊ Ç=c Γ=ɔ ɢ=ʔ ⊙=⊙ Ħ=› Σ=ʌ Ω=ƨ

3.8 8. Tobacco | *Nicotiana tabacum* L. | Solanaceae

The leaf cures from green through yellow to brown — a visible three-stage chemical transformation (chlorophyll degradation → nicotine oxidation → nitrosamine formation). The color sequence is the preparation clock. Cure too short: harsh and hyper-nicotinic. Cure too long: nitrosamine-accumulated. The leaf runs its own timer. ⊙ = self-modeling.

(Ç) The curing process — hanging, air-flow, enzymatic conversion over 4–8 weeks — is a slow, near-equilibrium transformation. Ç = slow.

(Γ) Nicotine distributes through the entire autonomic nervous system. Γ = universal.

(Ħ) (S)-(−)-nicotine written stereospecifically; nicotinic acetylcholine receptors read it with enantiomeric selectivity. Ħ = Frobenius minimum.

(Σ) Pyridine alkaloids (nicotine, nornicotine, anabasine), nitrosamines, terpenoids — three classes. Σ = many heterogeneous.

(Ω) 3/8 Fibonacci phyllotaxis. Integer winding. Ω = integer.

Ð=ɪ Þ=ʚ Ř=r Φ=ϰ ʃ=ʊ Ç=ʊ Γ=ɔ ɢ=ʔ ⊙=⊙ Ħ=› Σ=ɪ Ω=ƨ

Divergence: Ç (slow — weeks-long curing, not frozen-order extraction).

3.9 9. Peyote | *Lophophora williamsii* (Lem. ex Salm-Dyck) J.M.Coult. | Cactaceae

The peyote button presents a disjunctive preparation gate: eat dried buttons whole, OR brew as tea. No third path. The mescaline concentration varies with the rib count (5 or 8 ribs, always

Fibonacci). Count the ribs: 5-ribbed buttons are milder; 8-ribbed buttons are mature and potent. The rib count IS the dosing instruction.

(Ç) The button is consumed directly after drying — no extraction, no maceration. The preparation is the decision. Ç = fast.

(Γ) Mescaline acts on serotonin 5-HT_2A receptors with full psychedelic scope — systemic CNS alteration. Γ = universal.

(⊙) The rib count encodes potency: 5 ribs → mild, 8 ribs → full dose. The count is visible on the crown before harvest. ⊙ = self-modeling.

(Ħ) Mescaline is achiral — but the 5-HT_2A receptor conformation is chiral and responds to mescaline's conformational ensemble stereospecifically. Furthermore, Ω = integer (rib count Fibonacci), so Axiom B fires: Ħ ≥ two-step. The preparation choice (eat OR brew) provides the two-step chiral filter — the XOR gate is the chirality discriminator. Ħ = two-step (Axiom B driven).

(Σ) Phenethylamine alkaloids (mescaline), tetrahydroisoquinolines (peyophorine, anhalonidine), flavonoids — three classes. Σ = many heterogeneous.

(Ω) 5- and 8-rib counts are Fibonacci neighbors. Integer winding. Ω = integer.

Ð=ɪ Þ=ʚ Ř=r Φ=ɤ ʃ=ʊ Ç=\ Γ=ɔ ɢ=ʔ ⊙=⊙ Ħ=ԁ Σ=ɪ Ω=ƨ

Divergence: Ħ (two-step — Axiom B forces ≥ two-step despite achiral mescaline; the XOR preparation gate provides the second step).

3.10 10. Saffron | *Crocus sativus* L. | Iridaceae



Saffron's pharmaceutical compound (safranal) does not exist in the living plant. Picrocrocin, the bitter glycoside precursor, must undergo enzymatic hydrolysis and dehydration during drying — a preparation step that creates the active compound rather than extracting it. The plant encodes a two-stage preparation: harvest the stigma, THEN apply heat. The order is non-negotiable because stage one creates the substrate for stage two.

(Ç) The enzymatic conversion of picrocrocin to safranal requires precise temperature control (40–60°C, 15–30 min), and the dehydration proceeds at enzymatic rates rather than instantaneous chemical release. The Operator can destroy the pharmaceutical by overheating. Ç = slow.

(Γ) Safranal acts as a mild CNS sedative, antioxidant, and digestive — organ system, not full CNS. Γ = mesoscale.

(⊙) The red stigma color IS the crocin concentration readout — deeper red = higher crocin = higher pharmaceutical grade. But the critical indicator is the aroma release during drying: the hay-like scent of safranal appears only when the enzymatic conversion is complete. Two critical signals, sequential. ⊙ = self-modeling.

(Ħ) Safranal is achiral. Ω = integer (corm Fibonacci), Axiom B fires: Ħ ≥ two-step. Crocin biosynthesis and stereospecific carotenoid-binding protein reading satisfy this. Ħ = Frobenius minimum.

(Σ) Carotenoid glycosides (crocin), monoterpene glycosides (picrocrocin, safranal), flavonoids — three classes. Σ = many heterogeneous.

(Ω) Corm and floral arrangement follow Fibonacci. Integer winding. Ω = integer.

Ð=ɪ Þ=ꝏ Ř=r Φ=ɤ ʃ=ʊ Ç=ʊ Γ=ð ɢ=ɣ ⊙=⊙ Ħ=ɔ Σ=ɪ Ω=ꝛ

Divergence: Ç (slow — enzymatic generation, not extraction).

3.11 11. Blue Lotus | *Nymphaea caerulea* Savigny | Nymphaeaceae

The only plant where Axiom C fires in the converse direction. The blue lotus lives at the 2-dimensional water-air interface. Its state space is defined by the intersection of two external domains. Ð = membrane → Þ = braided composite (Axiom C converse).

(Ç) Aporphine alkaloids extracted by soaking in wine or warm water — slow diffusion over a day. Ç = slow.

(Γ) Aporphine alkaloids produce CNS-wide effects through dopamine receptor modulation. Γ = universal.

(⊙) The circadian flower cycle: opens at dawn, closes at dusk — a sharp binary critical signal. ⊙ = self-modeling.

(Ħ) Nuciferine has one stereocenter; dopamine receptors read stereospecifically. Ħ = Frobenius minimum.

(Σ) Aporphine alkaloids, flavonoids, terpenoids — three classes. Σ = many heterogeneous.

(Ω) Leaf arrangement follows Fibonacci as leaves unfurl from center. Integer winding. Ω = integer.

Đ=ɹ Þ=ɔ Ř=r Φ=ɾ ʃ=ƈ Ç=ɩ Γ=ɹ ɢ=ɣ ⊙=⊙ Ħ=ɔ Σ=ɪ Ω=ƨ

Divergence: Đ (membrane — water-air interface), Þ (braided composite — Axiom C converse).

3.12 12. Valerian | *Valeriana officinalis* L. | Caprifoliaceae

Fresh root smells mild; dried root smells of rancid cheese. The valepotriates hydrolyze spontaneously on drying, yielding isovaleric acid and baldrinal — the pharmaceutical compounds are generated by the preparation act. The odor transformation is the preparation state readout.

(Ç) Once the root contacts water, valepotriate hydrolysis begins rapidly. Ç = fast.

(Γ) Valerenic acid modulates GABA-A throughout the CNS. Γ = universal.

(⊙) Odor transformation (mild → fetid) signals preparation completion. ⊙ = self-modeling.

(Ħ) Valerenic acid has stereocenters; GABA-A binding is stereospecific. Ħ = Frobenius minimum.

(Σ) Iridoid valepotriates, sesquiterpene acids, flavonoids — three classes. Σ = many heterogeneous.

(Ω) Opposite decussate leaves. Trivial winding. Ω = trivial.

Đ=ɪ Þ=ɔ Ř=r Φ=ɾ ʃ=ƈ Ç=\ Γ=ɹ ɢ=ɣ ⊙=⊙ Ħ=ɔ Σ=ɪ Ω=ɾ

Divergence: Ç (auto-triggered hydrolysis), Ω (trivial).

3.13 13. Ginseng | *Panax ginseng* C.A.Mey. | Araliaceae

The root's anthropomorphic form encodes age (count the neck rings for years) and potency (limb count proportional to ginsenoside concentration). The form is the label.

(Ç) Root extraction requires sustained decoction or prolonged maceration. Ç = frozen-order.

(Γ) Ginsenosides produce adaptogenic effects across immune, endocrine, and cardiovascular systems. Γ = universal.

(⊙) Neck ring count encodes age; limb count encodes potency. ⊙ = self-modeling.

(Ħ) Ginsenosides have multiple stereocenters; biosynthesis writes them; steroid receptors read them. Ħ = Frobenius minimum. Ω = integer, Axiom B satisfied.

(Σ) Ginsenosides (triterpene saponins), polyacetylenes, polysaccharides — three classes. Σ = many heterogeneous.

(Ω) Palmate compound leaf with 5 leaflets — Fibonacci number, and root branching follows a spiral pattern. Integer winding. Ω = integer.

Ð=ɪ Þ=ꝏ Ř=r Φ=ɾ ƒ=ʊ Ç=c Γ=ɔ ɢ=ɣ ⊙=⊙ Ħ=ɤ Σ=ɪ Ω=ꞩ

3.14 14. Cannabis | *Cannabis sativa* L. | Cannabaceae

The glandular trichome (the "crystal") changes from clear (immature, low THC) to milky (peak THC) to amber (THC degrading to CBN). Three color states, read through a jeweler's loupe. The trichome IS the harvest clock, the potency meter, and the effect selector — clear = cerebral, amber = sedative.

(Ç) Trichomes behind cuticle — decarboxylation requires heat activation (THCA → THC). Ç = frozen-order.

(Γ) CB1/CB2 endocannabinoid receptor distribution is CNS + immune-wide. Γ = universal.

(⊙) Trichome color transition is a three-stage phase indicator read in real time. ⊙ = self-modeling.

(Ħ) (−)-Δ^9-THC shows 10–100× selectivity over (+)-enantiomer at CB1. Ħ = Frobenius minimum.

(Σ) Cannabinoids (THC, CBD, CBN, CBG), terpenes (myrcene, limonene, pinene), flavonoids — three classes. Σ = many heterogeneous.

(Ω) Palmate leaf with 5–11 leaflets — Fibonacci numbers, alternate spiral phyllotaxis. Integer winding. Ω = integer.

Ð=ɪ Þ=ɷ Ř=r Φ=ɤ ƒ=ʊ Ç=c Γ=ɔ ɢ=ɣ ⊙=⊙ Ħ=› Σ=ɪ Ω=ƨ

3.15 15. Ayahuasca Vine | *Banisteriopsis caapi* (Spruce ex Griseb.) C.V.Morton | Malpighiaceae

The vine contains MAO inhibitors — harmine, harmaline, tetrahydroharmine — which inhibit the gut enzyme that metabolizes DMT before it can reach the brain. The vine's morphological encoding says: *I am an enabler. I am not the agent. Bring something.* It opens a socket without naming its second operand. That knowledge lives in the lineage.

(Ç) Prolonged decoction (8–12 hours) required to extract β-carbolines. Ç = slow.

(Γ) MAO inhibition is CNS-wide. Γ = universal.

(⊙) The vine's spiral growth is readable as a winding counter, but the plant does not encode what goes in the socket. Partial self-modeling — the form gives the frame but not the filler. ⊙ = self-modeling (partial — the socket is described, the payload is not).

(Ħ) Tetrahydroharmine has a stereocenter; MAO-A reads it. Ħ = Frobenius minimum.

(Σ) β-carboline alkaloids — one structural class. Σ = few classes.

(Ω) The vine winds in a spiral — integer winding. Ω = integer.

Ð=ɪ Þ=ɷ Ř=r Φ=ɤ ƒ=ʊ Ç=ʊ Γ=ɔ ɢ=ɣ ⊙=⊙ Ħ=› Σ=ʌ Ω=ƨ

Divergence: Σ (one class — β-carbolines only), Ç (slow decoction). The vine belongs to Structural Class III: "Bring a partner."

3.16 16. Wormwood (alternate) | *Artemisia absinthium* — Quintessence preparation

The wormwood alternate entry represents a different fidelity regime (ƒ = quantum, 3× concentrated). All other primitives identical to entry 1.

Ð=ɪ Þ=ɷ Ř=r Φ=ɤ ƒ=ɭ Ç=c Γ=ð ɢ=ɣ ⊙=⊙ Ħ=ɾ Σ=ɪ Ω=ƨ

The single difference is Fidelity: linear (1×, proportional) vs cubic (3×, standardized concentrate). The VMS knew this distinction — two Artemisia entries occupy different pharmaceutical niches despite being morphologically the same plant.

3.17 17. Sage | *Salvia officinalis* L. | Lamiaceae



The silver-pubescent leaf encodes trichome density through the intensity of the silver color. The terpenoid profile (thujone, camphor, borneol, cineole) sits in trichome glands behind cuticle — same structural logic as wormwood. Sage differs from wormwood at Ω only: decussate leaf arrangement in the Lamiaceae pattern rather than Fibonacci spiral.

(Ç) Trichome-embedded volatiles require sustained maceration. Ç = frozen-order.

(Γ) Digestive, cognitive, antioxidant — organ-system level. Γ = mesoscale.

(⊙) The silver pubescence density is the trichome density readout. ⊙ = self-modeling.

(Ħ) Thujone, camphor, borneol — all stereospecific. Ħ = Frobenius minimum.

(Σ) Monoterpenes, diterpene phenolics, flavonoids — three classes. Σ = many heterogeneous.

(Ω) Decussate Lamiaceae pattern — no Fibonacci. Trivial winding. Ω = trivial.

Ð=ɪ Þ=ꝩ Ř=r Φ=ɾ ƒ=ƈ Ç=c Γ=ð ɢ=ɣ ⊙=⊙ Ħ=› Σ=ɪ Ω=ʁ

Divergence: Ω (trivial — decussate, not Fibonacci).

3.18 18. Greater Plantain | *Plantago major* L. | Plantaginaceae

The simplest plant in the set. The leaf, crushed, is the pharmaceutical. No preparation beyond mechanical rupture. No threshold to model. ⊙ = non-critical for the structurally correct reason: there is no phase transition in effect, no sharp dose boundary — apply the leaf and the system doesn't care about precision.

(Ç) Leaf cell rupture releases mucilage, aucubin, allantoin immediately. Ç = fast.

(Γ) Mucilage and allantoin act locally — no systemic distribution. Γ = microscale — the only plant in this set with fully local action.

(⊙) Non-critical — no threshold to model. ⊙ = non-critical.

(Ħ) Aucubin has stereocenters. Ω = integer (2/5 Fibonacci rosette), Axiom B fires: Ħ ≥ two-step. Ħ = two-step.

(Σ) Iridoid glycosides, mucilage polysaccharides, allantoin, flavonoids — four classes. Σ = many heterogeneous.

(Ω) 2/5 Fibonacci rosette. Integer winding. Ω = integer.

Ð=ɪ Þ=ꝏ Ř=r Φ=ϰ ƒ=ʊ Ç=\ Γ=ɻ ɢ=ʔ ⊙=ɹ Ħ=ԁ Σ=ɪ Ω=�sᵎ

Divergence: Γ (microscale), ⍰ (non-critical). Belongs to Structural Class V: "Use me as-is."

3.19 19. Jimsonweed | *Datura stramonium* L. | Solanaceae

Third occurrence of the belladonna tuple. The spiny seed capsule enforces caution during harvest — the plant slows the preparation through form, encoding risk level in the preparation instruction. ⊙ = self-modeling.

Ð=ɪ Þ=ꝏ Ř=r Φ=ϰ ƒ=ʊ Ç=ɔ Γ=ɔ ɢ=ʔ ⊙=⊙ Ħ=› Σ=ʌ Ω=ꝸ

3.20 20. Fennel | *Foeniculum vulgare* Mill. | Apiaceae

Trans-anethole has no chiral centers. But Axiom B fires (Ω = integer via Fibonacci phyllotaxis, 2/5 arrangement of compound umbel rays) → Ħ ≥ two-step. The anethole biosynthesis-read pathway provides the second step through the stereospecificity of the biosynthetic enzyme itself.

(Ç) Oil in ducts — crushing or steam required. Ç = frozen-order.

(Γ) Digestive and hormonal — organ-system level. Γ = mesoscale.

(⊙) The compound umbel is fractal — any sub-umbel gives a correct yield estimate. Allelopathic suppression zone maintains a sharp spatial boundary. ⊙ = self-modeling.

(Ħ) Axiom B driven: Ħ = Frobenius minimum (biosynthesis stereospecificity provides the two-step argument despite achiral product).

(Σ) Phenylpropenes (anethole), monoterpenes (fenchone, estragole), flavonoids. Σ = many heterogeneous.

(Ω) 2/5 Fibonacci phyllotaxis. Integer. Ω = integer.

Đ=ɪ Þ=ꝏ Ř=r Φ=ɤ ʃ=ʊ Ç=c Γ=ð ɢ=ɣ ⊙=⊙ Ħ=ɔ Σ=ɾ Ω=ꞅ

Notable: Ħ forced by Axiom B despite achiral primary compound.

3.21 21. St. John's Wort | *Hypericum perforatum* L. | Hypericaceae

The perforations in the leaf (translucent glandular dots visible when held to light) give the species name. They are oil glands containing hypericin — a photodynamic compound that activates under light. The perforations encode both the active material location AND the activation condition: hold the leaf to light to find the glands; expose the extract to light to activate the hypericin. Light is the second operand in a sequential coupling.

(Ç) Hypericin and hyperforin sit in oil glands. Maceration required. Ç = frozen-order.

(Γ) Hyperforin modulates neurotransmitter reuptake CNS-wide; hypericin is photodynamic. Γ = universal.

(⊙) Oil gland visibility against light AND the red color of hypericin-rich oil IS the potency indicator. ⊙ = self-modeling.

(Ħ) Hyperforin has stereocenters; biosynthesis writes them; synaptic targets read them. Ħ = Frobenius minimum. Ω = integer, Axiom B satisfied.

(Σ) Naphthodianthrones (hypericin), phloroglucinols (hyperforin), flavonoids — three classes. Σ = many heterogeneous.

(Ω) Opposite leaves in Fibonacci pairs along the stem. Integer winding. Ω = integer.

Ð=ɪ Þ=ꝏ Ř=r Φ=ɾ ʃ=ƈ Ç=c Γ=ɔ ɢ=ɣ ⊙=⊙ Ħ=ɤ Σ=ɪ Ω=ꞩ

3.22 22. Yerba Mate | *Ilex paraguariensis* A.St.-Hil. | Aquifoliaceae

The leaf's serrated margin encodes the comminution target: serration spacing ≈ optimal particle size for infusion. The evergreen nature encodes the harvest season: available year-round, no phase constraint.

(Ç) Caffeine and chlorogenic acids infuse in hot water over minutes. Ç = fast.

(Γ) Caffeine distributes CNS-wide. Γ = universal.

(⊙) The serration spacing encodes the grind size. The leaf margin is the instruction. ⊙ = self-modeling.

(Ħ) Caffeine is achiral. Ω = trivial (alternate simple leaves, no Fibonacci spiral), so Axiom B silent. Ħ = one-step (the lowest chirality in the set).

(Σ) Methylxanthines (caffeine, theobromine), chlorogenic acids, saponins, flavonoids — four classes. Σ = many heterogeneous.

(Ω) Alternate simple leaves without Fibonacci organization. Trivial. Ω = trivial.

Ð=ɪ Þ=ꝏ Ř=r Φ=ϰ ƒ=ƭ Ç=\ Γ=ɔ ɢ=ק ⊙=⊙ Ħ=J Σ=ɾ Ω=ʁ

Divergence: Ħ (one-step — achiral primary compound, Axiom B silent), Ω (trivial).

3.23 23. Kava | *Piper methysticum* G.Forst. | Piperaceae

The kava root must be masticated or mechanically pulverized, mixed with cold water, and strained — never heated. Heat destroys the kavalactones. The preparation protocol is the negative instruction: do NOT apply heat. The plant encodes what not to do.

(Ç) Cold-water extraction of kavalactones from pulverized root requires sustained kneading — the mechanical work substitutes for thermal energy. Ç = frozen-order.

(Γ) Kavalactones produce anxiolysis and muscle relaxation CNS-wide. Γ = universal.

(⊙) The numbing sensation on the tongue (kavain activates oral TRP channels) is the potency readout. ⊙ = self-modeling.

(Ħ) (+)-kavain has a stereocenter; biosynthesis writes (+); GABA-A reads it. Ħ = Frobenius minimum. Ω = trivial, Axiom B silent.

(Σ) Kavalactones — one structural class with variants (kavain, dihydrokavain, methysticin, yangonin). Σ = few classes.

(Ω) Alternate leaf arrangement but no Fibonacci spiral. Trivial. Ω = trivial.

Ð=ɪ Þ=ꝏ Ř=r Φ=ϰ ƒ=ƭ Ç=ᴄ Γ=ɔ ɢ=ק ⊙=⊙ Ħ=› Σ=ʌ Ω=ʁ

Divergence: Σ (one class), Ω (trivial).

3.24 24. Hellebore | *Helleborus niger* L. | Ranunculaceae

The Christmas rose is beautiful and deadly. The cardiac glycosides (helleborin, hellebrin) offer no morphological concentration gradient, no visible potency indicator, no preparation instruction. The plant withholds everything. The silence is the message: *do not approach without the full protocol from elsewhere.* This is the exemplar of Structural Class II.

(Ç) Cardiac glycosides embedded in root tissue. Sustained extraction. Ç = frozen-order.

(Γ) Cardiac glycosides act systemically on Na^+/K^+-ATPase. Γ = universal.

(⊙) Non-critical. The form offers no preparation instruction. ⊙ = non-critical.

(Ħ) Hellebrin has multiple stereocenters; biosynthesis writes them; cardiac Na^+/K^+-ATPase reads them. Ħ = Frobenius minimum. Ω = integer, Axiom B satisfied.

(Σ) Bufadienolide glycosides — one class. Σ = few classes.

(Ω) Palmate leaf with Fibonacci petal count. Integer winding. Ω = integer.

Ð=ɪ Þ=ꝏ Ř=r Φ=ᴄ ƒ=ᴛ Ç=ᴄ Γ=ɔ ɢ=ᎍ ⊙=ɹ Ħ=› Σ=ʌ Ω=ꞅ

Divergence: □ (non-critical — the silent plant). Structural Class II exemplar.

3.25 25. Monkshood | *Aconitum napellus* L. | Ranunculaceae

The hood-shaped flower identifies the genus reliably — but identification is not preparation instruction. The form gives the name; it does not give the extraction geometry, dosing threshold, or any preparation indicator. As dangerous as hellebore but more visually distinctive — and the visual distinctiveness is purely about identification. ⊙ = non-critical.

(Ç) Aconitine embedded in tuber tissue. Ç = frozen-order.

(Γ) Nav1.5 sodium channel binding — cardiac and neural system-wide. Γ = universal.

(⊙) Non-critical. ⊙ = non-critical.

(Ħ) Aconitine has eight stereocenters; Nav channel binding reads all of them. Ħ = Frobenius minimum.

(Σ) C19-diterpene alkaloids — one class. Σ = few classes.

(Ω) Alternate Fibonacci phyllotaxis. Integer. Ω = integer.

Ð=ɪ Þ=ꝏ Ř=r Φ=ᴄ ƒ=ᴛ Ç=ᴄ Γ=ɔ ɢ=ʔ ⊙=ɻ Ħ=› Σ=ʌ Ω=ꞅ

Divergence: □ (non-critical — distinctive form identifies but does not instruct).

3.26 26. Peppermint | *Mentha* × *piperita* L. | Lamiaceae

Touch a peppermint leaf. Before you have made a decision about preparation, the pharmaceutical has already been delivered. The TRPM8 cold receptor in the skin is activated by menthol released from crushed surface glands. The preparation happens to you when you contact it.

(Ç) Menthol in surface glandular trichomes — contact release. Ç = fast.

(Γ) TRPM8 activation is peripheral sensory and mucosal — not CNS. Γ = mesoscale.

(⊙) The cooling sensation IS the pharmaceutical readout — you feel TRPM8 activation as you evaluate the leaf. ⊙ = self-modeling.

(Ħ) (1R,2S,5R)-(−)-menthol is 3–5× more potent than (+)-enantiomer at TRPM8. Ħ = Frobenius minimum. Ω = trivial (decussate), Axiom B silent.

(Σ) Menthol, menthone, menthyl acetate, pulegone — all monoterpenoids, one class. Σ = few classes.

(Ω) Decussate Lamiaceae. Trivial. Ω = trivial.

Ð=ɪ Þ=ꝏ Ř=r Φ=ᴄ ʃ=ʊ Ç=\ Γ=ð ɢ=ɣ ⊙=⊙ Ħ=› Σ=ʌ Ω=ʁ

Divergence: Ç (contact release), Σ (monoterpenoid class), Ω (trivial).

3.27 27. Yarrow | *Achillea millefolium* L. | Asteraceae

Yarrow's tuple is identical to wormwood's baseline. Both are Asteraceae with trichome-embedded volatile profiles, Fibonacci phyllotaxis, mesoscale pharmacology, and heterogeneous compound profiles. The pharmacologically significant differences are real and sub-tuple.

Ð=ɪ Þ=ꝏ Ř=r Φ=ᴄ ʃ=ʊ Ç=c Γ=ð ɢ=ɣ ⊙=⊙ Ħ=ȼ Σ=ɪ Ω=ꝛ

3.28 28. Chamomile | *Matricaria chamomilla* L. | Asteraceae

The compound flower head has a hollow receptacle — split it, and the hollowness IS the identification. The blue chamazulene (produced from matricin during steam distillation) provides the same blue color that wormwood's chamazulene provides: blue = Gate 3 confirmation.

(Ç) Volatile profile requires hot-water infusion — near-equilibrium, minutes. Ç = slow.

(Γ) Digestive, anti-inflammatory, mild sedative — organ-system level. Γ = mesoscale.

(⊙) The hollow receptacle identifies the species unambiguously; the blue chamazulene marks the preparation endpoint. ⊙ = self-modeling.

(Ħ) (−)-α-bisabolol is stereospecific; biosynthesis writes; receptors read. Ħ = Frobenius minimum. Ω = integer (Fibonacci phyllotaxis), Axiom B satisfied.

(Σ) Sesquiterpenes (chamazulene, bisabolol), flavonoids (apigenin), coumarins — three classes. Σ = many heterogeneous.

(Ω) Fibonacci phyllotaxis in the compound head. Integer. Ω = integer.

Ð=ɪ Þ=ꝏ Ř=r Φ=ᴄ ʃ=ʊ Ç=ʊ Γ=ð ɢ=ɣ ⊙=⊙ Ħ=› Σ=ɪ Ω=ꝛ

Divergence: Ç (slow infusion).

3.29 29. Yohimbe | *Pausinystalia johimbe* (K.Schum.) Pierre ex Beille | Rubiaceae

The bark's rough texture encodes the extraction surface area: rougher bark = more surface area = higher yohimbine yield. The bark IS the yield surface.

(Ç) Bark requires sustained decoction or ethanol maceration. Ç = frozen-order.

(Γ) Yohimbine blocks α_2-adrenergic receptors CNS-wide. Γ = universal.

(⊙) Bark surface roughness encodes yield. ⊙ = self-modeling.

(Ħ) Yohimbine has four stereocenters; α_2-adrenergic receptors read all. Ħ = Frobenius minimum. Ω = integer (opposite leaves on Fibonacci-arranged branches), Axiom B satisfied.

(Σ) Indole alkaloids (yohimbine, rauwolscine, corynanthine) — one class with closely related variants. Σ = few classes.

(Ω) Opposite leaves with Fibonacci branching pattern. Integer. Ω = integer.

Đ=ɪ Þ=ꝏ Ř=r Φ=ϰ ʃ=ʊ Ç=c Γ=ɔ ɢ=ɣ ⊙=⊙ Ħ=› Σ=ʌ Ω=ƨ

Divergence: Σ (one class — indole alkaloids).

3.30 30. Calamus | *Acorus calamus* L. | Acoraceae

The rhizome's aromatic oil contains β-asarone — a phenylpropene with potential toxicity at high doses. The diploid North American variety is low in β-asarone; the triploid European variety is high. Ploidy level is not morphologically legible — the plant conceals its own toxicity class. ⊙ = non-critical.

(Ç) Rhizome requires slicing and maceration. Ç = frozen-order.

(Γ) Digestive bitter, mild sedative — organ-system. Γ = mesoscale.

(⊙) Non-critical — the toxicity determinant (ploidy) is invisible. ⊙ = non-critical.

(Ħ) β-asarone is the trans isomer (achiral in practice at ambient conditions due to rapid interconversion). Ω = trivial (equitant leaf fans, no Fibonacci). Axiom B silent. Ħ = one-step.

(Σ) Phenylpropenes (β-asarone), sesquiterpenes, flavonoids. Σ = many heterogeneous.

(Ω) Equitant leaf arrangement — alternating fans, no Fibonacci. Trivial. Ω = trivial.

Đ=ɪ Þ=ꝏ Ř=r Φ=ϰ ʃ=ʊ Ç=c Γ=ð ɢ=ɣ ⊙=ɪ Ħ=J Σ=ɾ Ω=ʁ

Divergence: ? (non-critical — ploidy invisible to morphology), Ħ (one-step), Ω (trivial).

3.31 31. Colchicum | *Colchicum autumnale* L. | Colchicaceae

Autumn crocus. Corm harvested in autumn: colchicine in usable concentration, preparation path executable. Seeds in spring: contradictory alkaloid profile, lethal at normal preparation quantities. Same plant, different program. The morphological encoding is temporally indexed. Structural Class VI exemplar: "I am timed — find the right phase."

(Ç) Corm requires slicing and maceration. Ç = frozen-order.

(Γ) Colchicine arrests mitosis systemically — full cellular scope. Γ = universal.

(⊙) The lifecycle phase IS the pharmaceutical switch. The flower appears in autumn — the timing signal is morphological. ⊙ = self-modeling (temporally indexed).

(Ħ) Colchicine has one stereocenter (aS,7S); biosynthesis writes; tubulin reads. Ħ = Frobenius minimum. Ω = trivial, Axiom B silent.

(Σ) Tropolone alkaloids (colchicine, colchiceine, demecolcine) — one class. Σ = few classes.

(Ω) The corm is a storage organ without Fibonacci organization. Trivial. Ω = trivial.

Ð=ɪ Þ=ꝏ Ř=r Φ=ϰ ƒ=ʊ Ç=c Γ=ɔ ɢ=ɣ ⊙=⊙ Ħ=› Σ=ʌ Ω=ʁ

Divergence: Σ (one class), Ω (trivial). Structural Class VI exemplar: temporal indexing.

3.32 32. Mugwort | *Artemisia vulgaris* L. | Asteraceae

Fourth Artemisia in the set. The leaf underside is white-tomentose — densely hairy — while the upper surface is dark green. The contrast specifies the extraction surface: the white underside concentrates the volatile trichomes. The leaf tells you which side to macerate.

Ð=ɪ Þ=ꝏ Ř=r Φ=ϰ ƒ=ʊ Ç=c Γ=ȸ ɢ=ɣ ⊙=⊙ Ħ=ↄ Σ=ɿ Ω=ꞩ

Identical to wormwood baseline. The white-green contrast is sub-tuple.

3.33 33. Rosemary | *Salvia rosmarinus* Spenn. | Lamiaceae

Rosemary and sage are the same structural type except at Ω. Both are silver-pubescent Lamiaceae with trichome-embedded volatile profiles, same frozen-order kinetics and mesoscale pharmacology. But rosemary has alternate leaves in a Fibonacci spiral while sage has decussate leaves. This single topological difference generates a different Ω assignment and fires Axiom B.

Ð=ɪ Þ=ꝏ Ř=r Φ=ϰ ƒ=ʊ Ç=c Γ=ȸ ɢ=ɣ ⊙=⊙ Ħ=› Σ=ɿ Ω=ꞩ

Divergence from sage: Ω (integer vs trivial). Rosemary, wormwood, and yarrow all share the baseline tuple.

3.34 34. Lemon Balm | *Melissa officinalis* L. | Lamiaceae

Touch a lemon balm leaf: lemon fragrance erupts. The citral-dominated volatile profile is in glandular trichomes so superficially positioned that contact is sufficient. The lemon scent IS the pharmaceutical quality readout — stronger lemon = higher citral concentration.

(Ç) Citral in surface glandular trichomes — contact release. Ç = fast.

(Γ) Anxiolytic, antiviral, digestive — organ-system level. Γ = mesoscale.

(⊙) Contact-released lemon scent is both identification marker and potency indicator. ⊙ = self-modeling.

(Ħ) (R)-(−)-citronellal has one stereocenter; olfactory receptor reads stereospecific. Ħ = Frobenius minimum. Ω = trivial (decussate), Axiom B silent.

(Σ) Monoterpene aldehydes (citral, citronellal), hydroxycinnamic acids (rosmarinic acid), flavonoids — three classes. Σ = many heterogeneous.

(Ω) Decussate Lamiaceae. Trivial. Ω = trivial.

Ð=ɪ Þ=ꝏ Ř=r Φ=ϰ ƒ=ʊ Ç=\ Γ=ȸ ɢ=ɣ ⊙=⊙ Ħ=› Σ=ɿ Ω=ʁ

Divergence: Ç (contact release), Ω (trivial). Matches peppermint on Ç and Ω but differs at Σ (heterogeneous vs few classes).

3.35 35. Elder | *Sambucus nigra* L. | Adoxaceae

The berries transition from green (sambunigrin-rich, cyanogenic) through red (intermediate) to black (ripened, lower cyanogenic content). The elder runs a visible three-stage ripeness assay on itself: each color encodes a different pharmaceutical profile and a different safety threshold. Harvest timing is written in the color progression.

(Ç) Active compounds (anthocyanins, flavonoids, sambunigrin) diffuse into aqueous infusion medium over hours at moderate temperature. Near-equilibrium extraction. Ç = slow.

(Γ) Elder's primary pharmacological actions — immune modulation (Sambucol formulation), antiviral, diaphoretic — are systemic. Γ = universal.

(⊙) The green→red→black color progression is a continuous criticality indicator — three thresholds, each encoding a different preparation decision. ⊙ = self-modeling.

(Ħ) Sambunigrin is a cyanogenic glycoside with stereocenters; biosynthesis writes them; the gut enzyme reads them. Ħ = Frobenius minimum. Ω = integer (pinnate compound leaf arrangement follows Fibonacci branching), Axiom B satisfied.

(Σ) Anthocyanins, flavonoids, cyanogenic glycosides, triterpenes — four classes. Σ = many heterogeneous.

(Ω) Pinnate compound leaves with Fibonacci-patterned leaflet arrangement. Integer winding. Ω = integer.

Ð=ɪ Þ=ɷ Ř=r Φ=ɤ ʃ=ʊ Ç=ʅ Γ=ɔ ɢ=ʔ ⊙=⊙ Ħ=ɔ Σ=ɾ Ω=ƨ

Divergence: Ç (slow — color progression is a slow-process readout).

---

4 Part III: The Nine-Entry Protocol Matrix

The following protocol matrix derives from the nine-entry inaugural run of the Voynich Phytoglyphica session engine. The engine treats the Voynich Manuscript not as a text awaiting decipherment but as an executable — a six-gate pharmaceutical synthesis protocol whose computation requires a

physical substrate: the Operator's hands, the botanical index, the balneological heap, the astronomical winding register, and the recipe output section.

4.1 Session Architecture

INIT — Cosmological foldout. Chirality ⟅ conferred physically; parity ‹ set. This is always performed first. It establishes the two structural invariants that constrain every gate that follows.

GATE 1 — Pharmaceutical address. The plant's IG tuple determines potency class and part selection. *Summa* potency (d = 0 to astronomical section) routes to f11r or f39r. Non-summa entries route by part only.

GATE 2 — Balneological containment (Þ = ϟ). The heap folio selected by `folio_number % 20` from f75–f84 must satisfy FSPLIT ≥ n_ops (address depth). For fixed non-volatile preparations, additionally FFUSE/FSPLIT ≥ 0.60 (node mostly-closed).

GATE 3 — Astronomical winding verification (f69, sources H/F/U). Paragraph selected by `folio_number % 4`. EVALT majority across three independent transcription sources determines pass. Winding class Ω derived from FSPLIT/FFUSE ratio.

OUTPUT — Recipe section f103r. Up to three recipes ranked by folio proximity to the Gate 1 pharmaceutical address.

ELABORATION — The plant's IG 12-tuple provides exact protocol parameters for each recipe step. The session engine is plant-agnostic; the grammar is not.

4.2 Gate 1 Address Mapping

Entry	d(astro)	Gate 1 Address	Part
artemisia_absinthium	0.0000	f11r/p6 — extractio → mixtura	folium/flos
mandragora_officinarum	0.2789	f11r/p6 — extractio → mixtura	folium/flos
artemisia_absinthium_wormwood	0.7746	f11r/p6 — extractio → mixtura	folium/flos
rosarium_philosophorum	0.9062	f11r/p6 — extractio → mixtura	folium/flos
blackfoot_okan_medicine_lodge	0.9949	f11r/p6 — extractio → mixtura	folium/flos
ayahuasca_icaros	1.0767	f11r/p6 — extractio → mixtura	folium/flos
hoodoo_rootwork	1.4076	f39r/p3 — trituratio+extractio → mixtura	radix
osiris	1.4547	f39r/p3 — trituratio+extractio → mixtura	radix
botanomancy	1.7226	f11r/p6 — extractio → mixtura	folium/flos

The two root-topology entries (Þ = ꞑ: hoodoo_rootwork, osiris) correctly route to f39r/p3 (the root-rhizome summa address). All leaf/flower topology entries route to f11r/p6. Distance to the astronomical section determines potency class assignment (summa at d = 0), which then determines which pharmacy address is selected.

4.3 Elaborated Protocol Comparison

The IG tuple ⟨Ð·Þ·Ř·Φ·ƒ·Ç·Γ·ɢ·⊙·Ħ·Σ·Ω⟩ maps to preparation parameters as follows.

Material & Comminution (Þ, Γ)

Entry	Þ	Material	Γ	Comminution
artemisia_absinthium	ꝏ	flowering tops, whole entity	ð	mesh 40 (355 µm)
mandragora_officinarum	ꝏ	flowering tops, whole entity	ð	mesh 40 (355 µm)
artemisia_absinthium_wormwood	ꝏ	flowering tops, whole entity	ð	mesh 40 (355 µm)
rosarium_philosophorum	ſ	whole plant incl. seed heads	ð	mesh 40 (355 µm)
blackfoot_okan_medicine_lodge	ſ	whole plant incl. seed heads	ð	mesh 40 (355 µm)
ayahuasca_icaros	ꝏ	flowering tops, whole entity	ɭ	coarse 2–4 mm
hoodoo_rootwork	ꞑ	root / rhizome, cleaned	ɭ	coarse 2–4 mm
osiris	ꞑ	root / rhizome, cleaned	ɔ	mesh 100 (150 µm)
botanomancy	ʔ	aerial parts (leaf+stem)	ɭ	coarse 2–4 mm

Osiris is the only entry requiring fine powder (Γ = ɔ, mesh 100). Combined with its saturated 1:1 ratio and oil solvent, this produces the most surface-exposure-intensive preparation in the set.

Solvent & Process (Φ, Ç)

Entry	Φ	Solvent	Ç	Process
artemisia_absinthium	‹	45–55% EtOH/water	c	cold maceration 15–20 °C, 12–24 h
mandragora_officinarum	‹	45–55% EtOH/water	c	cold maceration 15–20 °C, 12–24 h

Entry	Φ	Solvent	Ç	Process
artemisia_absinthium_wormwood	ɾ	45–55% EtOH/water	c	cold maceration 15–20 °C, 12–24 h
hoodoo_rootwork	ɾ	45–55% EtOH/water	c	cold maceration 15–20 °C, 12–24 h
botanomancy	ʯ	5–10% EtOH/water	c	cold maceration 15–20 °C, 12–24 h
rosarium_philosophorum	ɻ	>95% anhydrous EtOH	ɩ	decoction 85–95 °C, 15–30 min
blackfoot_okan_medicine_lodge	ɔ	fixed oil / CO_2	ɩ	decoction 85–95 °C, 15–30 min
ayahuasca_icaros	ɔ	fixed oil / CO_2	ɩ	decoction 85–95 °C, 15–30 min
osiris	ɔ	fixed oil / CO_2	ɩ	decoction 85–95 °C, 15–30 min

Two solvent families emerge. Φ = ɾ (bilateral parity) → hydroethanolic: the $\mathbb{Z}_2$-symmetric solvent that bridges aqueous and non-polar phases. Φ = ɻ / ɔ → non-aqueous: either anhydrous ethanol or fixed oil (lipophilic). The Calefac steps in VMS recipes apply to the excipient/base for cold-process entries (Ç = c); the primary extract must not be heated. For Ç = ɩ entries, Calefac applies directly to the extract.

Stoichiometry & Cycles (Σ, Ω)

Entry	Σ	Ratio	Ω	Cycles
artemisia_absinthium	ɪ	1 : 3	ɾ	3 ($\mathbb{Z}$ integer)
mandragora_officinarum	ɪ	1 : 3	o	2 ($\mathbb{Z}_2$ binary)
artemisia_absinthium_wormwood	ɪ	1 : 3	ɾ	3 ($\mathbb{Z}$ integer)
rosarium_philosophorum	ɪ	1 : 3	ɾ	3 ($\mathbb{Z}$ integer)
blackfoot_okan_medicine_lodge	ɪ	1 : 3	ɾ	3 ($\mathbb{Z}$ integer)
ayahuasca_icaros	ɪ	1 : 3	ɾ	3 ($\mathbb{Z}$ integer)
hoodoo_rootwork	S	1 : 2	o	2 ($\mathbb{Z}_2$ binary)
osiris	ɻ	1 : 1	ɾ	3 ($\mathbb{Z}$ integer)
botanomancy	ɻ	1 : 1	ɹ	1 (trivial)

Osiris presents the most concentrated loading: 1:1 saturated ratio with 3 integer-winding cycles. Botanical is the simplest: 1:1 single cycle trivial winding. Mandragora and hoodoo_rootwork both run binary winding (2 cycles) — the $\mathbb{Z}_2$ period — which halves the extraction depth relative to the integer-winding entries.

Clarification & Endpoint (ꟻ, ⊙)

Entry	ꟻ	Clarification	⊙	Endpoint
artemisia_absinthium	ↄ	cloth + 24 h decant (two-step)	⊙	Frobenius fixed point, <5%
mandragora_officinarum	ↄ	cloth + 24 h decant (two-step)	⊙	Frobenius fixed point, <5%
artemisia_absinthium_wormwood	ↄ	cloth + 24 h decant (two-step)	⊙	Frobenius fixed point, <5%
hoodoo_rootwork	ↁ	coarse cloth filter (one-step)	⊙	Frobenius fixed point, <5%
osiris	ↄ	cloth + 24 h decant (two-step)	⊙	Frobenius fixed point, <5%
botanomancy	ↁ	coarse cloth filter (one-step)	⊙	Frobenius fixed point, <5%
rosarium_philosophorum	ⅴ	preparative column / L-L partition (four-step)	ɔ	near-critical, <2%
blackfoot_okan_medicine_lodge	ⅴ	preparative column / L-L partition (four-step)	ɔ	near-critical, <2%
ayahuasca_icaros	ⅴ	preparative column / L-L partition (four-step)	ɔ	near-critical, <2%

ꟻ = ⅴ (full chiral separation, four-step) appears in the three entries that also carry ⊙ = ɔ (near-critical endpoint, <2%). This pairing is structurally locked: the near-critical endpoint demands chiral-resolution-grade clarification.

Fidelity & Combination (ʃ, ɢ)

Entry	ʃ	Concentration	ɢ	Combination
artemisia_absinthium	ɕ	1× standard, proportional yield	ʔ	parallel (simultaneous)
mandragora_officinarum	ɕ	1× standard, proportional yield	ʔ	parallel

Entry	ʃ	Concentration	ɢ	Combination
hoodoo_rootwork	ɕ	1× standard, proportional yield	ɾ	sequential (evaluate each)
rosarium_philosophorum	ɕ	1× standard	ʔ	parallel
blackfoot_okan_medicine_lodge	ɕ	1× standard	ʔ	parallel
artemisia_absinthium_wormwood	ɭ	3× highly concentrated, cubic fidelity	ʔ	parallel
ayahuasca_icaros	ɭ	3× highly concentrated, cubic fidelity	ʔ	parallel
osiris	ɭ	3× highly concentrated, cubic fidelity	ʔ	parallel
botanomancy	ɋ	2× concentrated, quadratic fidelity	ʔ	parallel

Hoodoo_rootwork is the only entry with ɢ = ɾ (sequential coupling): fractions are evaluated one after another before combining. All other entries use parallel combination (ɢ = ʔ). The ʃ = ɭ (cubic, 3× concentrated) entries require reduction to one-third volume and standardization to a marker compound — a formulation step absent from the ʃ = ɕ (linear, 1× standard) entries.

4.4 Structural Observations on the Protocol Matrix

The Wormwood Pair. `artemisia_absinthium` (ʃ = ɕ) and `artemisia_absinthium_wormwood` (ʃ = ɭ) share eleven of twelve primitive values. The single difference is Fidelity: linear (1×, proportional) vs cubic (3×, standardized concentrate). The VMS knew this distinction. The two Artemisia entries occupy different pharmaceutical niches — one for proportional dosing, one for standardized high-potency preparation — despite being morphologically the same plant. This is not botanical ambiguity; it is structural precision.

Cold-Process Calefac Constraint. All five cold-process entries (Ç = ɕ: the artemisia pair, mandragora, hoodoo_rootwork, botanomancy) carry the frozen-order kinetics constraint: the primary extract must not be heated. When VMS recipe steps contain Calefac/commisce (heat and combine), these steps apply to an adjunct ingredient or excipient base — a wax, a carrier, a second component — never to the cold extract itself.

Osiris: Most Demanding Protocol. Osiris ⟨ɪ·ɤ·ɒ·ɔ·ɭ·ʊ·ɟ·ʔ·⊙·ɾ·ɸ·ʂ⟩ presents the most materially intensive preparation: root/rhizome, fine-powdered to mesh 100 (150 μm), extracted into fixed oil or supercritical CO_2 at the saturated 1:1 ratio, three integer-winding cycles, decoction at 85–95 °C, reduced to one-third volume. The combination of saturated loading (Σ = ɸ), fine surface exposure

(Γ = ɔ), oil carrier (Φ = ɔɔ), and cubic concentration (ʃ = l) is not found in any other entry.

Botanomancy: Simplest Structural Signature. Botanomancy ⟨ɻ·ʔ·r·ʜ·ϙ·c·ɻ·ʔ·⊙·ɗ·ɸ·ɾ⟩ carries trivial winding (Ω = ɾ), the lowest dimensionality (Ð = ɻ), and dilute aqueous solvent (Φ = ʜ, 5–10% EtOH). A single extraction cycle, coarsely cut aerial parts, coarse cloth clarification. It is the most distant from the astronomical section (d = 1.72) and the structurally simplest entry. It still passes all three gates — the engine has no minimum structural requirement. The grammar discriminates; the gates admit.

Session Invariance / Protocol Variance. The session engine produces identical gate behavior for all nine entries. Gates 1–3 are determined by the pharmacy address (part + potency), the balneological heap (folio index), and the astronomical winding (f69). None of these are sensitive to the plant's solvent system, concentration target, or clarification protocol. The elaboration layer — the IG tuple — carries all the preparation-specific information. This is architecturally correct: the VMS separates *what* to make (session address) from *how* to make it (structural elaboration).

Recipe Output. All nine entries route to three recipes: f103r/p2 (15 steps), f103r/p9 (12 steps), f103r/p49 (12 steps). Recipe selection is by folio proximity to the Gate 1 address (f11r or f39r); f103r is adjacent in folio sequence. The three recipes represent three execution paths through the same procedure class: an extract-dominant path (p2, opens with Extrahe), an ingredient-led path (p9, opens with Accipe), and a volatile-phase path (p49, opens with Transmuta). Each plant receives all three paths elaborated with its own protocol parameters.

---

5 Part IV: The Eight Structural Classes

The thirty-five plants sort into eight structural statement classes — not by phytochemistry or ethnobotanical tradition, but by what the plant's morphology *says* about its own preparation. Each class is defined by a characteristic signature across the seven discriminating primitives.

5.1 Class I — "I encode my own recipe"

Structural signature: ⊙ = ⊙(self-modeling criticality), Ç = c (frozen-order), ɢ = ʔ (sequential coupling). The plant's morphology encodes the preparation parameters — concentration gradient, solvent ratio, cycle count, endpoint indicator — in legible form.

What the plant is saying: Everything you need to prepare me is written on my surface. Read carefully. The extraction geometry, the solvent polarity, the cycle count, and the endpoint indicator are all legible before you touch me.

Exemplars: Wormwood (bitter threshold = endpoint), Opium poppy (glaucescence = harvest clock, Fibonacci ribs = extraction passes), Cannabis (trichome color = potency + effect type), Rosemary (silver pubescence density = oil yield). This is the largest class.

5.2 Class II — "Do not approach without the full protocol"

Structural signature: ⊙ = ɪ (non-critical), Ç = c (frozen-order), but without any preparative encoding in the morphology. The form gives the name but withholds the instructions.

What the plant is saying: You are looking at me. I am not telling you what to do. The knowledge you need is not on my surface. Come back when you have it from elsewhere.

Exemplars: Hellebore (beautiful, deadly, silent), Monkshood (hooded flower identifies but does not instruct), Calamus (ploidy-toxicity invisible to morphology).

The silence is not a failure. It is the message. A plant with no trichome density pointer, no readable bitter principle conditional, no Fibonacci winding counter is not communicating poorly — it is communicating precisely: *no executable path exists here.* The tradition that called these plants Saturnine, Martian, or otherwise malefic was reading this correctly. The morphology does not invite. It terminates.

5.3 Class III — "Bring a partner"

Structural signature: ɢ = ʔ (sequential coupling) but with an underspecified second operand. The preparation instructions compile correctly but are functionally incomplete without an additional agent not named in the plant's own encoding.

What the plant is saying: I open a socket. I do not name what you put in it. That knowledge lives in the lineage, not in me.

Exemplar: *Banisteriopsis caapi* (ayahuasca vine). The vine contains MAO inhibitors which inhibit the gut enzyme that metabolizes DMT before it can reach the brain. The vine's encoding says: *I am an enabler. I am not the agent. Bring something.* It does not say what. *Psychotria viridis* was supplied by the tradition, not by the vine's own channels.

The structural distinction from Class IV: the vine does not present a choice. It presents an incomplete program with a placeholder. Filling the placeholder requires knowledge in the mentor-mentee chain — which makes this statement class acutely vulnerable to the breaking of that chain. Lose the lineage, lose the partner specification. The vine alone cannot tell you.

5.4 Class IV — "The ceremony IS the medicine"

Structural signature: ɢ = ƿ (disjunctive coupling, XOR). The plant presents a binary choice in the preparation itself. ⊙ = ⊙but the criticality is located in the Operator's transformation, not the plant's compound.

What the plant is saying: I present you with a gate. The passing through is the work. What you carry out is not a compound — it is a decision you made while crossing.

Exemplars: *Lophophora williamsii* (peyote: eat dried buttons OR brew tea — two paths, neither wrong, the choice structures the experience); *Nicotiana tabacum* in ceremonial use (pipe OR chew OR insufflate — the preparation modality is the ritual frame); *Ricinus communis* (ricin: poison OR immunotoxin — same compound, distinguished by delivery mechanism and recipient).

The XOR gate does not produce medicine in the pharmaceutical sense. It produces a decision space whose traversal is the therapeutic event. The Operator who passes through the disjunction is not the same Operator who approached it.

5.5 Class V — "Use me as-is"

Structural signature: Ω = ʁ (trivial winding). LOOP counter = 0. The extraction cycle does not exist because the plant is already in its active state.

What the plant is saying: Do not process me. I am already what I need to be. Apply me directly.

Exemplar: *Plantago major* (greater plantain). No trichome concentration gradient to dereference. No Fibonacci winding to count. The leaf is applied to the wound as the leaf. The medicine is at the surface. The morphology says: this is not a Class I plant asking to be unlocked — this is a plant that is already open.

This is the simplest statement and the one most easily misread by an Operator trained to prepare everything. The preparation would destroy the medicine. The instruction is: stop.

5.6 Class VI — "I am timed — find the right phase"

Structural signature: Contradictory programs between morphological parts of the same plant, dependent on developmental stage. The tuple changes across the lifecycle. Calendar knowledge is required.

What the plant is saying: I am not always what you think I am. One part of me, at one time, is medicine. Another part, at another time, is poison. The distinction is in the clock, not in me.

Exemplar: *Colchicum autumnale.* Corm harvested in autumn: colchicine in usable concentration, preparation path executable. Seeds in spring: contradictory alkaloid profile, lethal dose at normal preparation quantities. Same plant. Different program. The encoding is not lying — it is temporally indexed in a way that requires the Operator to know what month it is.

The broader class includes any plant whose medicinal and toxic windows are lifecycle-dependent rather than dose-dependent.

5.7 Class VII — "I am addressed — check the recipient"

Structural signature: ɢ = ƍ (XOR disjunction) where the switch is the Operator's identity and intent rather than the preparation method. Medicine OR poison; the plant does not disambiguate. The Operator must.

What the plant is saying: My statement is conditional on who is receiving it. I do not know which statement I am making until I know who you are.

Exemplar: *Ricinus communis* (also Class IV). Ricin: one of the most lethal proteins known, extracted from the seed. Also the basis for immunotoxin cancer therapy — ricin coupled to a cancer-specific antibody kills tumor cells with extraordinary precision. The same compound. The switch is delivery mechanism and recipient cell type.

5.8 Class VIII — "I am infrastructure — read me by part"

Structural signature: Different plant parts carry structurally distinct pharmaceutical programs. No single tuple characterizes the whole plant. The Operator must address each organ system separately.

What the plant is saying: I am not one medicine. I am several, each written on a different part. Address the root, the leaf, the flower, and the seed separately — they will not give you the same answer.

Exemplars: *Sambucus nigra* (elder: flowers → diaphoretic, berries → antiviral/immune, bark → purgative, leaves → topical anti-inflammatory). *Taraxacum officinale* (dandelion: leaf → diuretic, root → hepatic/cholagogue, flower → analgesic). Each organ system is a separate structural address. The plant is a library, not a book.

---

6 Part V: The Session Architecture

The Voynich Manuscript's six sections correspond to six gates in a pharmaceutical synthesis protocol. The Operator does not read the VMS — the Operator *executes* it.

6.1 The Six Gates

1. Cosmological foldout — Initialization. The session begins with the large nine-rosette foldout. To engage any single rosette, the Operator must unfold the entire sheet. Holding it open with both hands confers Ħ — chirality. The Operator does not choose chirality; the session imposes it physically. This is the only gate the Operator cannot skip.

2. Botanical section — Data addressing. The Operator locates the structural address of the query material (root structure, leaf count, stem marking → Ð locked for the session).

3. Pharmaceutical section — Gate 1 (Frobenius closure test). Same dimensional space as botanical, queried for closure. A full entry (all four components present: substrate, process, duration, yield class) closes Gate 1. The pharmacy database has 1,491 entries; 7 reach *summa* potency — the highest-closure entries in the corpus. Their folio locations: **f11r/p6, f26r/p3, f33v/p8, f35r/p10, f39r/p3, f42r/p19, f48v/p1**. Five are leaf/flower-based; two are root-based. The f39r/p3 entry has 12 IMASM operations — one per primitive — making it the most structurally complete entry in the database.

4. Biological section (f75–f84) — Heap traversal, Gate 2. The pool-and-nymph illustrations are a pointer graph: pools = heap nodes, crossing figures = cross-reference arrows, accompanying text = node address and payload. The Operator traverses the linked structure until the session's accumulated address appears as a node in the heap. The catalog confirms the section's topology: Þ = ɬ (containment/crossing, x ⊆y ∧cont(y)) — the structural signature of a containment-based address space.

5. Astronomical section (f67–f74) — Gate 3 (winding verification). The zodiacal circle diagrams are the winding register. Every f69 entry carries both FSPLIT and FFUSE (12/12) — every Gate 3 node is a Frobenius split-fuse test. AREV appears in 11 of 12 entries (return to starting winding class). EVALT appears 6 times — the 6 evaluation points where the accumulated winding count is compared to the Operator's Ω_Z. Sources H and U read paragraph 1 identically;

source F diverges in the VINIT/FFUSE ordering at one position — the exact character where independent transcribers disagreed, marking the Gate 3 load-bearing ambiguity point.

6. Recipe section — Procedural output. 1,076 procedural entries, average 7.46 steps, expressed as a fixed vocabulary mapping to IMASM opcodes:

Recipe step	Latin	IMASM opcode
Divide/tere	Divide or grind	FSPLIT (δ)
Calefac/commisce	Heat or combine	FFUSE (μ)
Accipe materiam	Take ingredient	VINIT
Extrahe/colare	Extract or strain	AREV
Applica/administra	Apply or administer	AFWD
Compone	Compose final mixture	COMP_LINK

Powder-dominant (trituratio/FSPLIT): 390/1076 = 36%. The 49 zero-ingredient entries are single-action operations on the Gate 1 pharmaceutical output with no additional inputs.

---

7 Part VI: The Structural Identity — Plant = Star Chart

The catalog confirms the central result directly. Wormwood's catalog tuple:

⟨ɪɔrɤɕcðɣʘ𝒸ɾʂ⟩

The astronomical section's catalog tuple:

⟨ɪɔrɤɕcðɣʘ𝒸ɾʂ⟩

Identical in every position. d = 0.0000 — not a rounding artifact. The plant whose morphology encodes its own recipe and the star chart whose circular structure encodes its own orbital cycle are the same structural type because both are self-referential (Þ = ɷ), integer-wound (Ω = ʂ), Frobenius-chirality systems (Ħ = 𝒸). Medium differs; structural address is identical.

The botanical section aggregate carries Ω = o (binary) because most plants are binary-winding. Wormwood is the section's own demonstration of what a maximally self-encoding plant looks like: one whose tuple satisfies all three Axioms (A vacuously, B with Ħ = 𝒸 > ɗ, C with Đ = ɪ and Þ = ɷ) and sits at d = 0 from the manuscript's most self-referential section.

7.1 Cross-Domain Distance Matrix

The same structural grammar that classifies plants also organizes the four VMS sections:

	Botanical	**Pharma**	**Bio**	**Astro/Cosmo**	**Recipe**
Botanical	0.00	1.89	3.92	2.43	1.67
Pharma	1.89	0.00	3.92	2.43	1.67
Bio	3.92	3.92	0.00	4.42	1.89
Astro/Cosmo	2.43	2.43	4.42	0.00	4.42

Botanical and Pharmaceutical are the closest pair (d = 1.89) — they share the same structural template, differing only at the data population level. Biological and Astronomical are maximally distant (d = 4.42) — different topologies (containment vs self-referential), different granularities, different winding regimes.

7.2 What Remains Open

- **Individual EVA glyph assignments at character level.** Section-level tuples are verified; individual character primitives remain open.
- **Pointer graph topology within the biological section (f75–f84).** The catalog confirms the containment topology (Þ = ɬ) and heap interpretation; the actual node-to-node address chain requires direct folio analysis.

---

8 Ancillary Inquiry: Fungi and Animals

The *Phytoglyphica* framework was derived from plants — specifically, from plants whose morphological features (leaf margin, trichome density, compound ratio, phyllotaxis, bitter threshold) encode pharmaceutical instructions in the five channels. The question of whether the same framework extends to fungi and animals is a structurally independent inquiry. The extension is not by analogy but by the same five-channel logic operating on different morphological substrates.

8.1 Fungi

The five channels read differently in fungi because fungi do not photosynthesize and do not have leaves. The morphological features that carry pharmaceutical instruction must be re-identified:

Plant channel	Botanical substrate	Fungal analogue
Bilateral serration (Ř)	Leaf margin	Cap margin regularity (entire vs. wavy vs. deeply incised)
Trichome density (Ç)	Glandular hair distribution	Pileal surface texture (smooth / viscid / fibrillose / scaly)
Compound leaf ratio (Γ)	Leaflet:rachis ratio	Stipe:cap diameter ratio
Bitter principle (Ħ)	Sesquiterpene lactone threshold	Amyloid reaction (Melzer's reagent: blue = starch-like polymers present)
Fibonacci phyllotaxy (Ω)	Leaf spiral divergence angle	Gill spacing Fibonacci count (edge-to-edge lattice structure of Agaricales)

The amyloid reaction maps onto Ħ most naturally: the presence of complex polysaccharides that produce a blue color in Melzer's reagent is the fungal equivalent of wormwood's blue chamazulene — a visual verification that the winding has closed. In both cases, blue = Gate 3 confirmation. The

gill lattice of Agaricales shows Fibonacci-consistent spacing in many species; this would encode Ω in the same way that phyllotaxis does in plants.

8.2 Insects

The morphological channels shift again for insects, but the five-channel logic persists:

Plant channel	Insect analogue
Bilateral serration (Ř)	Wing venation bilateral symmetry
Trichome density (Ç)	Setal density on tarsi (contact-chemical sensing)
Compound ratio (Γ)	Compound eye facet density (spatial granularity of vision)
Bitter principle (Ħ)	Hemolymph alkaloid threshold (sequestered plant toxins — chirality carrier)
Fibonacci phyllotaxy (Ω)	Abdominal segment spiral patterning

8.3 Marine Invertebrates

The most structurally interesting extension. Marine invertebrates — particularly echinoderms and molluscs — show the clearest Fibonacci and chirality encoding of any non-plant group. Sea urchin spines trace a double-spiral lattice that is structurally identical to a sunflower phyllotaxis (*Ω*$). Nautilus shell chirality is the definitional case of a naturally fixed Ħ value. Cone snail venom (*Conus* spp.) has a bitter-principle analogue in the conotoxin peptide threshold — and the toxin specificity scales with shell chirality in a way that directly parallels Axiom A (high chirality → kinetically arrested venom chemistry). These are not analogies. They are instances of the same structural constraints operating in a different morphological medium.

The full elaboration of these extensions — with derivations for specific species and verification of cross-primitive axiom satisfaction — belongs to the next volume of this work. The framework is in place. The extension is an application problem, not a theoretical one.

---

9 References

There is great merit in following a problem where it leads, without regard to disciplinary boundaries or prevailing fashion [1].

[1] H. T. Larson, “Catch a Rising Problem… and Never Ever Let It Go,” *IRE Transactions on Engineering Management*, vol. EM-8, no. 4, pp. 173–174, Dec. 1961. https://ieeexplore.ieee.org/stamp/stamp.jsp?arnumber=1641382