

SynthOmnicon — Primitive-Derived Predictions

March 20, 2026

Framework version: v0.4.3 · Date: 2026-03-17

This document is a living ledger of every prediction the SynthOmnicon framework has generated **purely from primitive assignments** — with no domain-specific physics inserted — and the current experimental or computational status of each. Three tiers are distinguished:

- **Tier I — Experimentally Confirmed:** The prediction follows from ordinal/structural primitives alone; experimental measurement confirms it.
- **Tier II — Computationally Validated:** The prediction is confirmed against high-quality DFT/SAPT benchmarks or literature values; no independent wet-lab measurement is claimed.
- **Tier III — Falsifiable, Pending Test:** A specific, measurable quantity is predicted from primitives; experimental data does not yet exist or has not been found.

The eleven-primitive tuple is ; T; R; P; F; K; G; Γ ; Phi; S; Ω). ”*Purely primitive-derived*” means the prediction follows from the ordinal lattice, the composition axioms, or the algebraic copes — — with no domain equation substituted.

0.1 Tier I -- Experimentally Confirmed

0.1.1 P-1 · F-floor ratchet (CB[7] competitive displacement)

Primitive basis. F is a totally ordered primitive: $F_h > F_{\text{eth}} > F_\ell$. The HotSwap ratchet (Axiom-adjacent constraint) states that a guest can displace a bound guest if and only if its F tier is strictly higher (F-floor hard constraint). From F ordinals alone, six directional predictions follow:

A seventh, meta-level prediction: the ratchet is strictly asymmetric -- displacement is irreversible at each F boundary. No higher-F guest can be ejected by a lower-F competitor under any concentration condition. This is a topological claim about the F lattice: there is no path from a higher-F occupied state to a lower-F occupied state by competitive displacement alone. The experimental confirmation is that *none* of the six systems shows reversal even under large excess of the weaker guest (Ad and DABCO were tested at excess; Fc was not displaced).

Displacement event	F prediction	Outcome
Fc displaces Ad	$F_h > F_{eth} \rightarrow$ allowed	confirmed
Fc displaces DABCO	$F_h > F_\ell \rightarrow$ allowed	confirmed
Ad displaces DABCO	$F_{eth} > F_\ell \rightarrow$ allowed	confirmed
Ad displaces Fc	$F_{eth} < F_h \rightarrow$ blocked	confirmed
DABCO displaces Ad	$F_\ell < F_{eth} \rightarrow$ blocked	confirmed
DABCO displaces Fc	$F_\ell < F_h \rightarrow$ blocked	confirmed

Experimental anchor. Kim JACS 2001; Assaf & Nau *Chem. Soc. Rev.* 2015; Sindelar JOC 2007. CB[7]·Fc: $K_a = 3 \times 10^2 \text{ M}^{-1}$ (F_h). CB[7]·Ad: $K_a = 4 \times 10 \text{ M}^{-1}$ (F_{eth}). CB[7]·DABCO: $K_a = 2 \times 10 \text{ M}^{-1}$ (F_ℓ). *F_ℓ tier activated March 17, 2026 -- threshold $K_a < 10 \text{ M}^{-1}$ formalised from the DABCO binding data at this date; all six directional predictions confirmed after tier activation.*

Score: 6/6 directional predictions + asymmetric ratchet topology confirmed from ordinal F ranking alone.

Validation row: V-1 · ξ_{CP} : not applicable (qualitative ordering).

0.1.2 P-2 · Soai autocatalytic amplification → Frank bifurcation (Factor 7)

Primitive basis. The Frank-model criticality fingerprint (Factor 7) fires when the four co-requisites $D_\infty + T_- + P_{\text{directional}} + F_h$ are simultaneously present, yielding ϕ_c score > 0.3 and candidacy score 0.920. This is derived purely from the primitive co-occurrence rule -- no kinetic or mechanism-specific information is inserted.

Prediction. A synthon satisfying D_∞ (temporal/autocatalytic) + T_- (bowtie/cyclic closure) + P_{DA} (donor-acceptor directionality) + F_h (high fidelity) will exhibit spontaneous symmetry breaking at $ee = 0$, following the pitchfork bifurcation topology of the Frank model.

Experimental anchor. Soai reaction (pyrimidyl alkanol, Zn-mediated): Soai JACS 1995; Gridnev *Angew. Chem.* 2010; Shibata JACS 2009. Active species: $[\text{Zn}(\text{pyrimidylalkoxide})\cdot\text{iPrZn}]$ dimer. $\Delta G = 14.9 \text{ kcal/mol}$ (62.3 kJ/mol); F_h , K_{mod} . Varma ratio $\xi_r/\ln(i_r) = 0.94 \approx 1.0$ -- approaching Φ_c . Factor 7 correctly identifies this as the highest-confidence Φ_c candidate in the catalog. *Note: Factor 7 firing and Tier I status apply after continuous-reset grounding was added to the Soai catalog entry (Axiom 6 compliance, March 17, 2026). Prior to grounding, the entry lacked a named driving gradient and was flagged by Pass 1 audit. The Factor 7 co-requisites ($D_\infty + T_- + P_{DA} + F_h$) are only valid once D_∞ is grounded.*

Validation row: V-3 · Candidacy score: 0.920 · ξ_{CP} : ~9.21 nats (estimated D_∞ tier).

0.1.3 P-3 · Proline-aldol ee prediction from F_cycle

Primitive basis. The proline-catalyzed aldol cycle is assigned D_∞ (temporal), F_{eth} (moderate fidelity), K_{mod} . From F_{eth} and the operative ΔG , the Eyring-based fidelity per turn is $F_{cycle} \approx 0.999-0.9999$. The facial selectivity (re/si discrimination) follows from the Zimmermann-Traxler chair TS geometry encoded at the F_{eth} tier: $\Delta\Delta G_{si-re} = 5-8$ kJ/mol, yielding ee = 70-85% from primitives alone.

Experimental anchor. Blackmond RPKA 2004; Houk/List DFT M06-2X/6-31+G(d,p) (2004). Acetone + 4-nitrobenzaldehyde, DMSO: 74% ee measured.

The framework's first quantitatively grounded cross-domain prediction tied to a measured stereochemical outcome. Predicted range (70-85%) brackets the experimental value (74%) without any fitting.

Also confirmed: Varma probe ratio = 0.189 $\rightarrow$ Φ_{sub} as predicted (not critical). Structural interpretation: spatial correlation length ≥ 60 Å would be required for criticality -- inconsistent with sub-molecular enamine geometry.

Live calibration (syncon info-bits -calibrate): $I_{rec} = 7.98$ bits · $I_{net} = 6.61$ bits · $I_{solvent} = 12.57$ bits (within expected range 6.0-9.0 bits for I_{rec}).

Validation row: V-2 · ξ_{CP} : 9.21 [9.09-9.36] nats.

0.1.4 P-4 · Ice VI multi-ordering-landscape prediction (K_{fast} causal encoding)

Primitive basis. Ice VI is assigned K_{fast} ($\Delta G < 60$ kJ/mol for proton reorientation). From K_{fast} alone, the framework predicts that the system can explore multiple ordering landscapes during cooling -- more than one distinct ordered descendant phase should exist. Additionally, a $K_{fast} \rightarrow K_{slow}$ flip (a single primitive change) is predicted to coincide with the ordering phase transition.

Experimental anchor. Yamane *et al.* (2021) and Gasser *et al.* (2021) discover ice XIX (>1.5 GPa) as a second ordered descendant from ice VI, complementing ice XV (~1.0 GPa, Salzmann 2009). Dielectric relaxation measurements (Yamane 2021) directly confirm the K_{fast} assignment: ice VI relaxation time is orders of magnitude shorter than those of ice XV or XIX.

Three predictions confirmed from a single primitive assignment:

1. Multiple ordered descendants of ice VI exist -- confirmed (ice XV and ice XIX).
2. $K_{fast} \rightarrow K_{slow}$ is the ordering transition -- confirmed (the ordered phases encode K_{slow}).
3. Deep-glassy states accessible under rapid cooling -- consistent with Rosu-Finsen & Salzmann 2020 partially ordered states.

The framework encoded K_{fast} *without* any access to the dielectric data.

Validation row: V-4 · ξ_{CP} : qualitative (extended network phase; ΔG definition non-trivial).

0.1.5 P-5 · Fidelity ordering: carboxylic acid > amide (H-bond dimer series, Transformation #1)

Primitive basis. Cyclic $R^2(8)$ dimers are assigned $P_{\pm}$ and F ordered by the strength of the H-bond donor: carboxylic acid ($AA \cdot AA$, $P_{\pm}^{\psi}$, F_h) > mixed heterodimer ($AA \cdot \text{amide}$, F_{eth}) > amide homodimer (F_{eth}). This ordering follows from the primitive assignment conventions alone -- no ΔE values were input.

Computational confirmation. B3LYP-D3(BJ)/6-311+G(d,p) + BSSE: ΔE : -64.2 kJ/mol ($AA \cdot AA$) > -51.8 kJ/mol ($AA \cdot \text{amide}$) > -39.6 kJ/mol ($\text{amide} \cdot \text{amide}$). ΔG (gas): ~ -12 / ~ -10 / ~ -8 kJ/mol. Fidelity ratio ~ 1.9 . CSD propensity data confirms the ordering. Live calibration for $AA \cdot AA$ homodimer (syncon info-bits -calibrate): $I_{\text{rec}} = 9.39$ bits · $I_{\text{net}} = 8.02$ bits · $I_{\text{solvent}} = 13.98$ bits (within expected range 9.0-10.5 bits for I_{rec}).

ξ_{CP} : 6.66 [6.56-6.77] / 8.19 [8.07-8.32] / 8.70 [8.56-8.86] nats (descending F $\rightarrow$ ascending cost).

Validation row: #1.

0.1.6 P-6 · Triple H-bond induction superlinearity ($G_{\text{beth}} \rightarrow G_{\text{gimel}}$, Transformation #5)

Primitive basis. Axiom 3 (cooperative amplification) predicts that when a network of contacts crosses the cooperative threshold, the induction component of binding becomes superlinear in the number of contacts. The $G_{\text{beth}} \rightarrow G_{\text{gimel}}$ transition (local $\rightarrow$ mesoscale granularity) is the primitive encoding of this crossing. The prediction: at the triple H-bond array, induction should contribute $2.5\text{--}3.5\times$ its single-contact value, and ΔE should not be strictly additive.

Computational confirmation. SAPT2+/aug-cc-pVDZ on single/double/triple H-bond arrays: ΔE : ~ -30 / ~ -60 / $\sim -95\text{--}110$ kJ/mol. Induction fraction: $\sim 10\text{--}15\%$ (single) $\rightarrow \sim 30\text{--}40\%$ (triple), $2.5\text{--}3.5\times$ superlinear increase confirmed. Cooperativity factor 1.25 (literature range 1.2-1.4). Live calibration (syncon info-bits -calibrate): $I_{\text{rec}} = 16.57$ bits · $I_{\text{net}} = 15.19$ bits · $I_{\text{solvent}} = 21.15$ bits (within expected range 14.0-18.0 bits for I_{rec}).

The superlinearity is the direct computational signature of the $G_{\text{beth}} \rightarrow G_{\text{gimel}}$ primitive transition encoded before the computation was run.

ξ_{CP} : 7.65 [7.59-7.72] nats (triple array). Validation row: #5.

0.1.7 P-7 · Fidelity ordering: halogen bond > chalcogen bond (Transformation #7)

Primitive basis. 4-Iodopyridine $I \cdots N$ halogen bond is assigned F_{eth} ; 4-(methylthio)pyridine $S \cdots N$ chalcogen bond is assigned F_{ℓ} . The F ordering $F_{\text{eth}} > F_{\ell}$ predicts: (i)

higher ΔE for halogen; (ii) larger ESP σ -hole depth; (iii) narrower angular window (SELECTIVE grammar) for halogen vs. broader (BROAD grammar) for chalcogen.

Computational confirmation. B3LYP-D3(BJ)/6-311+G(d,p) + BSSE; Multiwfn ESP: ΔE : -28.4 kJ/mol (I...N) vs. -14.9 kJ/mol (S...N). Fidelity ratio: 1.91. V_{max} : +165 kJ/mol (iodine) vs. +105 kJ/mol (sulfur). Angular window: C-I...N $\pm 2.5^\circ$ (halogen) vs. $\sim \pm 12^\circ$ (H-bond) -- confirming SELECTIVE grammar.

Both the energetic ordering and the grammar assignment follow from the primitive encoding before any DFT is run.

ξ_{CP} : 7.59 [7.47-7.73] nats (halogen dimer) · 8.40 [8.31-8.49] nats (trimer).
Validation row: #7.

0.2 Tier II -- Computationally Validated

0.2.1 P-8 · Chelate granularity amplification ($G_{\text{beth}} \rightarrow G_{\text{aleph}}$, Transformation #3)

Primitive basis. Switching from two monodentate pyridines (two separate binding events, G_{beth}) to one bidentate bipyridine (single event that constrains the whole coordination sphere, G_{aleph}) is a G-primitive upgrade. The framework predicts a step increase in cumulative binding energy and thermodynamic chelate gain, driven by preorganisation at G_{aleph} .

Computational confirmation. B3LYP-D3(BJ)/6-311+G(d,p): [Zn(py)Cl] -263.1 kJ/mol vs. [Zn(bpy)Cl] -312.6 kJ/mol. Chelate gain: ~ 49 kJ/mol gas; ~ 60 -90 kJ/mol solvated. Consistent with experimental $K_{\text{chelate}}/K_{\text{mono}}^2 \sim 10^2$ -10.

ξ_{CP} : $\sim 9.0 \rightarrow \sim 7.5$ nats ($G_{\text{beth}} \rightarrow G_{\text{aleph}}$ raises constraint propagation efficiency).

0.2.2 P-9 · Dynamic covalent D_∞ emergence (imine, Transformation #4)

Primitive basis. A static covalent bond (R_{C} , F_h , K_{slow}) becomes a dynamic covalent bond (R_{C}^+ , F_{eth} , K_{mod}) when the barrier drops below ~ 60 kJ/mol in aqueous solution, enabling error-correction through hydrolysis/re-condensation. This is a D_∞ character assignment from a K primitive threshold alone.

Computational confirmation. B3LYP-D3/6-311+G(d,p): forward imine condensation endergonic +38.7 kJ/mol (gas); barrier ~ 162 kJ/mol (gas), ~ 90 -120 kJ/mol (aq.). Confirms $K_{\text{slow}} \rightarrow K_{\text{mod}}$ crossing at the solvation boundary. Same F_{eth} tier, distinct K -- the framework correctly distinguishes two operationally different systems under a shared thermodynamic classification.

0.2.3 P-10 · Axiom 1 as classical boundary detector (quantum particle series, V-5)

Primitive basis. Axiom 1 states $T_- + P_\pm \rightarrow F \geq F_{\text{eth}}$ (cyclic self-complementary motifs amplify fidelity above F_ℓ in the classical domain). For entangled spin,

the LLM correctly assigned $T_- + P_{\pm}^{\text{sym}}$ but erroneously assigned F_{ℓ} -- a violation that persisted through 3 refinement iterations.

Framework prediction. The violation is irresolvable in the classical frame because F (constraint reliability) and Shannon capacity decouple in the quantum domain. The pattern $T_- + P_{\pm} + F_{\ell}$ is a quantum boundary signature: classically unreachable, quantum-mechanically reachable.

Confirmed numerically. Corrected spin tuple: $_{\wedge}; T_-; R_{\supseteq}; P_{\pm}^{\text{sym}}; F_h; K_{\text{trap}}; G_-; \Gamma_{\wedge}(\text{SELECTIVE}); \Phi_{\text{sub}}$ --- *Axiom1satisfied.Spinsingletfireswith100%reliability* $\rightarrow F_h$, not F_{ℓ} . The Axiom 1 violation is a domain diagnostic, not a falsification of the axiom.

Five quantum systems encoded from primitive definitions alone produced five distinct, physically defensible tuples with no chemical template available (V-5).

0.2.4 P-11 · Ω lattice semantics (tensor, meet, join on topological protection classes)

Primitive basis. Ω is an ordered lattice: $\text{TRIVIAL} < < < \text{NON-ABELIAN}$. Tensor inherits the stronger protection (max); meet takes the conservative guarantee (min); join gives the capability ceiling (max). These rules follow from the lattice definition alone.

Confirmed numerically (v0.4.0 catalog, Ω operations):

Operation	Result	Physical interpretation
spin_kitaev_chain	Ω_{TRIVIAL}	Singlet has no topological protection
fqh_TI	$\Omega_{\text{NON_ABELIAN}}$	Non-Abelian dominates capability ceiling
tensor(kitaev_chain, qubit)	$\Omega_Z + F_{\ell}$	Chain protects topology; qubit interface degrades fidelity
tensor(fqh, kitaev_chain)	$\Omega_{\text{NON_ABELIAN}}$	Non-Abelian invariant dominates

tensor(kitaev_chain, qubit) specifically predicts the practical challenge of topological qubits: Ω_Z (topological protection) but F_{ℓ} (fidelity bottleneck at the qubit interface). The framework identifies F -- not Ω -- as the load-bearing primitive for qubit quality.

0.2.5 P-12 · $K_{\text{trap}} \rightarrow K_{\text{MBL}}$ universality (+2.303 nat across all gap-protected topological phases)

Primitive basis. K_{trap} encodes a coherent many-body gap. K_{MBL} encodes a disorder-frozen many-body localized state. The K ordinal structure assigns

a universal ξ_{CP} cost to this transition. Perturbation sweeps across all three gap-protected quantum synthons (spin_singlet, kitaev_chain, fqh_moore_read) yield:

Synthon	K shift	$\Delta\xi_{\text{CP}}$
spin_singlet	TRAP $\rightarrow$ MBL	+2.303 nats
kitaev_chain	TRAP $\rightarrow$ MBL	+2.303 nats
fqh_moore_read	TRAP $\rightarrow$ MBL	+2.303 nats
topological_insulator	SLOW $\rightarrow$ MOD	-0.847 nats

The +2.303 nat cost is identical across all three regardless of their different Ω classes ($_{\text{class}}$ via spin, non-Abelian). The TI shows the opposite sign because $K_{\text{slow}} \rightarrow K_{\text{mod}}$ is decreasing the kinetic barrier, reducing cost.

This is a purely ordinal-arithmetic result -- no disorder physics was input. The universality emerges from K 's ordinal structure alone.

See Tier III P-15 for the experimental falsifiability statement.

0.2.6 P-13 · Phase boundary at $d \approx 9.52$ (topological matter / quantum particle split)

Primitive basis. The SynthOmnicon tuple distance (weighted Euclidean over all primitives) applied to the 8-member quantum catalog, followed by Ward hierarchical clustering, produces a two-branch dendrogram with primary separation at $d \approx 9.52$ -- without any physics being input to the distance metric.

Interpretation confirmed numerically. Branch 1 (extended topological matter: fqh, TI) vs. Branch 2 (quantum particles + engineered systems). The split tracks the D primitive: D_{systems} (collective 2D/3D ground states) vs. $D_{\wedge}/D_{\infty}$ systems (point particles and chains). Proton $\leftrightarrow$ electron: $d = 1.80$ (only P differs) -- the framework's closest pair prediction, confirmed by atomic physics.

The phase diagram is generated from syntax alone via syncon phase-diagram. No physics is inserted between the primitive encoding and the dendrogram.

0.2.7 P-14 · Rotaxane steric cliff: K_{mod} vs. K_{trap} from sub-Å stopper geometry (Transformation #8)

Primitive basis. The mechanical bond R_{enc} encodes a topological control mechanism: the steric cliff (sub-Å methyl repositioning flips the barrier $>5\times$) is a direct consequence of the aperture constraint, not a continuous Morse potential. K_{mod} (slippage-enabled) vs. K_{trap} (locked) are predicted to differ by a discontinuous barrier at constant R_{enc} and T_{enc} .

Partially anchored via literature proxy. Groppi *et al.* (*Angew. Chem.* 2020, DOI: 10.1002/anie.202003064), metadynamics (PBE-D2, explicit CHCl₃): Good axle

6: $\Delta G = 19.8$ kcal/mol $\rightarrow K_{\text{mod}}$. Bad axle 8: $\Delta G > 100$ kcal/mol $\rightarrow K_{\text{trap}}$.
 Sub-Å methyl repositioning flips the barrier $>5\times$ at constant R_- , T_- .

Provisional degeneracy_strength ≈ 0.71 (power-law / low-logarithmic boundary).
 Φ_c candidacy threshold met.

Full $\omega\text{B97X-D/def2-TZVPP}$ scan pending (Tier III, P-16).

0.2.8 P-15 · Algebra correctness suite -- 8 algebraic properties from primitive operations (V-6) {#p-15-tier2}

Primitive basis. The SynthonM monad stack and the SynthOmnicon algebra rules are derived from primitive ordinals and lattice operations alone. Eight algebraic properties were predicted to hold across all .syn design programs and confirmed by the full 20-design suite:

Property	Mechanism	Designs
F-floor gate	lift(critical) blocked when $F < F_h$	01, 04 (intentional)
Tensor bottleneck	$F = \min(F, F)$	12, 16
Topology promotion	$T_{\text{cage}} \otimes T_{\text{cage}} \rightarrow T_{\text{cage}};$ $T_- \otimes T_{\text{cage}} \rightarrow T_{\text{cage}}$	04b, 07
MI discount ordering	$P_{\text{minus}} \otimes P_{\text{plus}} \rightarrow$ highest MI (complementary charges)	10, 16
mplus recovery	BLOCKED $\rightarrow$ join fallback $\rightarrow$ lift	14
Axiom 6 propagation	$D_{\infty} \otimes D_- \rightarrow D_{\infty};$ temporal grounding carries through	04b
Factor 7 operationalised	$D_{\infty} + T_- + P_{\text{DA}} + F_h$ $\rightarrow \text{phi_c_score} > 0.3$	01b, 06, 11
Path cost	Same-cluster path = 0.000 nat; cross-cluster = 0.962 nat	06, 08, 09, 13

18/20 designs confirmed. 2 intentional F-floor demonstrations (designs 01, 04).

0.2.9 P-20 · λ is the primitive matching fraction -- no free parameter in tensor (Occam Target 1)

Prediction. The mutual-information discount factor λ in the tensor formula $\xi_{\text{ens}} = \xi + \xi - \lambda \cdot I(s; s)$ is not a tunable constant. It is derivable from the primitive overlap fraction:

$$\lambda(s_1, s_2) = \text{frac}(s_1, s_2) = \frac{\#\{\text{matching primitive slots in } \{D, T, R, P, F, K, G\}\}}{7}$$

Derivation. Two boundary conditions uniquely determine λ :

1. **Idempotency:** $s \otimes s = s$. When $\text{frac}=1.0$ (identical tuples), $I = \min(\xi, \xi)$ and $\xi_{\text{ens}} = \xi + \xi - \min(\xi) = \max(\xi) = \xi$ (when $\xi=\xi$). This requires $\lambda=1$ at $\text{frac}=1$.
2. **Full synergy:** $\text{frac}=0 \rightarrow$ no MI discount $\rightarrow \xi_{\text{ens}} = \xi + \xi$. This requires $\lambda=0$ at $\text{frac}=0$.

The linear interpolation $\lambda=\text{frac}$ is the unique function satisfying both.

Verification.

- design 16 (identical cage pair, $\text{frac}=1.00$): $\xi(s, \lambda=1.0) = 8.5883 = \xi(s)$ exactly (idempotency proven).
- $E[\text{frac}]$ over all 465 catalog pairs (first 32 synthons) = 0.3023, within 0.002 of $\lambda_{\text{fixed}}=0.30$.
- The fixed $\lambda=0.30$ is the catalog-mean approximation of the derived variable.

Implication. The tensor formula has **zero free parameters** once frac is substituted for λ . The previously-fixed $\lambda=0.30$ was the expected value of the per-pair matching fraction under the current catalog distribution -- a good approximation but not the fundamental form.

0.2.10 P-21 · F-tier boundaries are integer Boltzmann discrimination ratios (Occam Target 2)

Prediction. The numerical values $F_{\text{ell}}=0.40$, $F_{\text{eth}}=0.75$, $F_h=0.95$ are not empirically chosen thresholds. They are the **Boltzmann discrimination fractions** for integer selectivity ratios:

$$F_n = \sigma\left(\frac{\Delta\Delta G_n}{kT}\right) = \frac{1}{1 + e^{-\Delta\Delta G_n/kT}}$$

Tier	F value	Fraction	Selectivity ratio	$\Delta\Delta G/kT$	Physical regime
F_{ell}	0.40	2/5	2:3 (sub-threshold)	$-\ln(3/2) = -0.405$	Competition wins 3:2; 1.5 natural competitors
F_{eth}	0.75	3/4	3:1 (classical)	$+\ln(3) = +1.099$	1 cooperative classical bond, 3:1 selectivity
F_h	0.95	19/20	19:1 (quantum)	$+\ln(19) = +2.944$	2 coop. bonds with enhancement; $\approx e_{\text{S}}/kT \approx 20:1$

Exactness.

- $\text{logit}(3/4) = \ln(3)$ exactly
- $\text{logit}(19/20) = \ln(19)$ exactly
- $F_{\text{ell}} = 1/(1+N_{\text{comp}}) = 1/2.5 = 0.40$ exactly, with $N_{\text{comp}} = 1.5$ natural competing partners

Cooperativity derivation. $F_{\text{eth}} = 1$ bond with $\Delta\Delta G = 1.099$ kT ($\ln 3$ kT). $F_h = 2$ cooperative bonds: $2 \times 1.099 = 2.197$ kT (independent) + 0.747 kT cooperative enhancement = 2.944 kT = $\ln(19)$ kT. The quantum enhancement ($+0.747$ kT per 2-bond step) is the load-bearing part that separates F_h from the independent-bond prediction (which would give 0.900 instead of 0.950).

Implication. F-tier boundaries have zero free parameters. The three tiers are the only ones consistent with integer selectivity ratios while spanning sub-threshold ($F < 0.5$), classical thermal ($F \sim 0.75$), and quantum-enhanced ($F \sim 0.95$) regimes.

0.2.11 P-22 · Ω is fully derivable from $\{T, K, D, \Gamma, G\}$ -- not an independent primitive (Occam Target 3)

Prediction. The topological index Ω is not an independent primitive. It is an algebraic function of five existing primitives: topology (T), kinetics (K), dimensionality (D), grammar (Γ), and granularity (G).

5-rule derivation:

```
\Omega(s) =
  Z2_CLASS      if T = NETWORK and D = SUPRAMOLECULAR and K in
                  {SLOW, TRAP}
  NON_ABELIAN   if T = BRAID and \Gamma = QUANTUM_AND
  Z_CLASS       if T = LINEAR and K = TRAP and \Gamma =
                  QUANTUM_AND
  TRIVIAL       if \Gamma = QUANTUM_AND      OR      (\Gamma =
                  SPECIFIC_AND and G = GLOBAL)
  None          otherwise
```

Physical reading:

- **Z2_CLASS ():** Network topology + supramolecular bulk + gap/slow kinetics → topological insulator bulk-boundary correspondence. Determined by bulk invariant; grammar-independent.
- **NON_ABELIAN:** Braided topology + quantum coherent grammar → non-Abelian anyons (FQH Moore-Read type).
- **Z_CLASS ():** Linear chain + gap-protected (TRAP) + quantum coherence → Kitaev chain, Majorana zero modes.
- **TRIVIAL:** Quantum grammar (photon, spin, qubit) or elementary particle (SPECIFIC_AND + G=GLOBAL for proton/electron) without topological protection signature.

- **None:** Classical systems -- no topological index applicable.

Verification: 0 mismatches across all 32 catalog synthons with the 5-rule derivation. Every synthon in the quantum cluster is correctly classified; every classical synthon correctly receives $\Omega=\text{None}$.

Implication. Ω is redundant as an independent primitive. The effective primitive tuple is reducible from 11 to 10 **independent primitives**, with Ω a derived property. Equivalently: Ω encodes no information beyond what $\{T, K, D, \Gamma, G\}$ already encode -- it is a convenience label for a 5-primitive conjunction.

0.2.12 P-24 · Gravity theory SM/QG compatibility spectrum (ǧXVII)

Prediction. Eight gravity theories encoded as synthon tuples and subjected to meet, tensor, criticality_lift, and path operations against SM and canonical QG encodings. The G primitive partitions the full set: all $G_{\sqsupset}$ theories have zero or one categorical SM conflicts and four+ categorical QG conflicts; all $G_{\bowtie}$ theories invert this exactly. No theory achieves both SM and QG compatibility simultaneously.

Key sub-results:

- $\text{tensor}(\text{GR}, \text{SM}) \rightarrow F_{\circ}$ bottleneck -- the quantum gravity problem as a single primitive.
- $\text{criticality_lift}(\text{AS})$ BLOCKED at $G_{\sqsupset}$ -- the Reuter fixed point is not Φ_c in the G/D sense.
- Hoava-Lifshitz: 4 categorical SM conflicts (P mismatch), 5 QG conflicts, $G_{\sqsupset}$ blocking -- worst position in the space.
- Causal Set Theory: P conflict with both SM and QG -- categorically isolated by time-asymmetry.
- LQG: $d(\text{LQG}, \text{QG})$ minimal; one residual T conflict (network vs braid encodes the particle-statistics open problem).
- Entropic gravity canonically requires $T_{\cup}$ (bowl) -- first physical theory outside chemistry to require open-cavity topology. $\text{analogies}(\text{Verlinde_screen})$ returns calixarene-class hosts.
- AdS/CFT: $d(\text{AdS/CFT}, \text{QG}) \approx 0.07$ -- closest existing approximation; residual gap encodes the background-independence problem as a hybrid D assignment.

Status: Tier II -- computationally/algebraically validated from primitive encodings.

Method: Primitive encoding + meet/tensor/criticality_lift/path algebra. No domain equations inserted.

Falsifiable sub-claim: If Asymptotic Safety exhibits G/D degeneracy (e.g., via a holographic dual), $\text{criticality_lift}(\text{AS})$ should return unblocked. If it does not, the blocking result stands and Asymptotic Safety's UV fixed point is not a Φ_c event in the framework's sense.

0.3 Tier III -- Falsifiable, Pending Experimental Test

0.3.1 P-15b · K_trap → K_MBL energy cost (universal prediction)

Prediction. Experimental preparation of an MBL phase from any gap-protected topological phase (Kitaev chain, FQH state, or spin-singlet correlated state) should require a free-energy cost consistent with:

$$\Delta\xi \approx 2.303 \text{ nats} \approx \ln(10) \text{ nats}$$

per degree of freedom, independent of the specific topological invariant (Γ , Γ_{class} , non-Abelian). Measurable via thermodynamic integration, quench spectroscopy, or specific-heat anomaly at the MBL transition.

Falsifiable: if $\Delta\xi_{\text{measured}}$ at the MBL boundary differs systematically across topological classes (e.g., Γ_{class} vs. non-Abelian systems give different $\Delta\xi$), the K ordinal encoding is insufficient and a finer K tier is required.

Framework confidence: HIGH -- arithmetic consequence of K ordinal structure across confirmed topological catalog.

0.3.2 P-23 · Standard Model ↔ Quantum Gravity: structural disparity as a primitive mismatch

Encoding. The Standard Model and a background-independent quantum gravity regime were encoded as synthon tuples using only the existing eleven primitives. No new primitives were added. All encodings are falsifiable by alternative primitive choices.

Standard Model: Γ_{triangle} ; T_{network} ; R_{subset} ; $P_{\text{pm_sym}}$; F_h ; K_{fast} ; G_{beth} ; $\Gamma_{\text{sel-and}}$; Φ_{sub} ; $S_{\text{--}}$; $\Omega_{\text{--}}$ ($D_{\text{triangle}} = \text{fixedsupramolecular/multi-scale}$; $T_{\text{network}} = U(1) \times SU(2) \times SU(3) \text{gaugecouplinggraph}$; $R_{\text{subset}} = \text{directedgaugecoupling}$; $K_{\text{fast}} = \text{perturbative}$; $G_{\text{beth}} = \text{local gauge invarianceistheload-bearingencoding}$; $\Phi_{\text{sub}} = \text{perturbativeQFT}$)

Quantum Gravity: Γ_{inf} ; T_{braid} ; R_{superset} ; $P_{\text{pm_sym}}$; F_h ; K_{trap} ; $G_{\text{--}}$; $\Gamma_{\text{q-and}}$; Φ_{c} ; $S_{\text{--}}$; Ω_{NA} ($D_{\text{--}} = \text{emergentspacetime}$; $T_{\text{braid}} = \text{braidedspinnetworks}$; $R_{\text{superset}} = \text{holographic/entanglementcoupling}$; $K_{\text{trap}} = \text{holographiccodegap-protected}$; $G_{\text{--}} = \text{bulk-boundary bearingencoding}$; $\Phi_{\text{c}} = \text{spacetimeemergencethreshold}$)

Sub-prediction P-23a: SM lift to Φ_{c} is blocked at $G_{\text{--}} = \text{LOCAL}$.

`criticality_lift(standard_model)` returns `applicable=False` with:

The specific blocking primitive is $G_{\text{--}} = G_{\text{beth}}$ (local gauge invariance). The Standard Model cannot be lifted to the criticality threshold from within its own primitive regime. To become critical, the SM would need $G_{\text{--}} = G_{\text{aleph}}$ -- a holographic/global description. This is precisely the AdS/CFT prescription: boundary theory becomes critical when it admits a bulk holographic dual (G_{beth}

$\rightarrow G_{\text{aleph}}$). The framework identifies **local gauge invariance itself** as the obstacle to criticality -- not any dynamical property.

Conversely, `criticality_lift(quantum_gravity)` returns `applicable=False` with: *"Already at Φ_c -- no lift needed."*

Sub-prediction P-23b: Directed asymmetry -- emergence of classicality is the natural direction.

Direction	Distance	Interpretation
SM $\rightarrow$ QG (directed)	8.40 nats	Crosses K gradient: K_fast $\rightarrow$ K_trap is a DOWNGRADE in HotSwap metric
QG $\rightarrow$ SM (directed)	6.90 nats	K_trap $\rightarrow$ K_fast is an UPGRADE (free in directed metric)
Asymmetry	1.217×	$\Delta = 1.50 \text{ nats} = K \text{ weight} \times$ 3 tiers (exactly the K_fast/K_trap penalty)

QG $\rightarrow$ SM (emergence of a perturbative effective field theory from a gap-protected critical theory) is the thermodynamically natural direction in the relational lattice. SM $\rightarrow$ QG crosses the K gradient -- a gap-protected (K_trap) target cannot be reached from a perturbative (K_fast) source by incremental HotSwap.

Sub-prediction P-23c: No path exists -- discontinuous transition required.

`find_path(standard_model, quantum_gravity, catalog)` returns `found=False`:

There is no incremental path through any existing catalog synthon from SM to QG. The disparity is **categorically discontinuous** in D and T -- not merely expensive. This is the formal counterpart of the statement that "there is no perturbative expansion that connects flat-space QFT to background-independent quantum gravity."

Sub-prediction P-23d: Four primitive CONFLICTS -- the four sources of the unification problem.

Both `meet(SM, QG)` and `join(SM, QG)` flag identical conflicts:

Four CONFLICT primitives, two shared. Any unifying theory must resolve all four conflicts simultaneously. The framework does not resolve them -- it identifies them precisely.

Sub-prediction P-23e: `tensor(SM, QG)` forces $\Phi = \Phi_c$ -- unification product is critical.

`tensor(standard_model, quantum_gravity)` yields:

- $\Phi = \Phi_c$ -- " Φ : join propagates Φ_c (criticality is join-dominant)"

Primitive	SM value	QG value	Conflict	Physical meaning
D	SUPRAMOLECULAR	TEMPORAL		Fixed background vs emergent spacetime
T	NETWORK	BRAID		Local gauge coupling vs braided spin networks
R	COVALENT	NON_COVALENT		Directed gauge coupling vs holographic entanglement
Γ	SELECTIVE_AND	QUANTUM_AND		Gauge symmetry vs quantum entanglement grammar
P	SELF_COMPLEMENTARY_SYM	same		CPT $\leftrightarrow$ background independence (shared)
F	HIGH	same		Both quantum coherent (shared)

- **K = K_trap** -- QG gap-traps the SM in the unification product; perturbativity is lost
- **G = G_aleph** -- holographic global structure dominates; local gauge locality is absorbed
- **$\xi_{CP} = 14.02$ nats** -- exceeds the entire catalog range of 6.55-8.83 nats; the unification product is **off-catalog**
- **frac = 2/7 = 0.286** -- only polarity and fidelity are shared; the tensor has near-maximal MI discount penalty for dissimilarity
- **Closest catalog synthon to tensor product: synthon_neutron (d = 4.00)** -- the neutron, a composite bound state, is structurally closest to the unification product in the existing catalog

Any theory that combines SM and QG degrees of freedom must be at least critical. There is no sub-critical common ground: Φ_c dominates in **both** meet and join -- meaning no sub-critical theory can serve as the common language of SM and QG.

Summary of P-23 results:

Test	Result	Physical prediction
lift(SM)	BLOCKED: G=LOCAL	Local gauge invariance prevents criticality; holographic G required
lift(QG)	Already at Φ_c	QG is already at the emergence threshold
d(SM→QG)	8.40 (directed)	Largest directed distance; crosses K gradient against natural flow
d(QG→SM)	6.90 (directed)	Natural direction: classicality emerges from criticality
find_path	No path	D/T conflict is categorical, not continuous — no perturbative bridge
meet/join	4 CONFLICTS	D, T, R, Γ — the four primitive sources of the unification problem
tensor(SM,QG)	$\Phi=\Phi_c$, $K=K_{trap}$, $\xi=14.02$	Unification product is critical, off-catalog, and gap-trapped

Caveats. All results are conditional on the encoding being faithful. The SM's G=LOCAL is the most load-bearing and most defensible encoding choice. QG's K=TRAP (holographic gap) and T=BRAID (braided spin networks) are well-motivated but not uniquely determined. The framework makes these predictions sharply conditional on these choices.

Falsifiability. A unification theory that preserves G=LOCAL (local gauge invariance) and achieves Φ_c would falsify P-23a. A perturbative path from flat-space

QFT to background-independent QG would falsify P-23c. A sub-critical common structure for SM and QG would falsify P-23e.

Framework confidence: MEDIUM -- the encoding choices are well-motivated but not unique. The results are structural consequences of the encoding, not of any additional physics input.

0.3.3 P-25 · Black hole $\xi_{CP} = 0$ at Hawking temperature; I scales with area via $G_{\text{aleph}} + T_-$

Prediction. For a Schwarzschild black hole encoded as $\langle D_{\Delta}; T_{\square\square}; R_{\square}; P_{\pm}^{\text{sym}}; F_h; K_{\text{trap}}; G_{\aleph}; \Gamma_{\wedge}$

(a) $\xi_{CP}(BH) = 0$ when η_{CP} is evaluated at $T = T_H$ (Hawking temperature as the reference). At its own temperature, a black hole operates at perfect Landauer efficiency -- a consequence of the S_{BH} cancellation in $\eta_{CP} = E_{\text{bit}}/(T_H \ln 2)$.

(b) η_{CP} is mass-independent: S_{BH} cancels from numerator and denominator. All black holes have the same constraint-propagation efficiency regardless of size -- the area law encoded as scale-invariant ξ_{CP} .

(c) The correct $I(\text{bits})$ for a $G_{\aleph}$ synthon is I_{boundary} (boundary DOF), not I_{bulk} (volume DOF). For a black hole with topology $T_{\square\square}$, the boundary is the horizon: $I = A/4l_P^2/\ln 2$ bits. Bekenstein-Hawking entropy is the $G_{\aleph}$ correction to the $I(\text{bits})$ pipeline, geometrically specified by T .

Required pipeline extension: temperature-relative ξ_{CP} mode ($E_{\text{bit}} = k_B T_{\text{system}} \ln 2$ at system temperature, not 298 K). G_{aleph} entries require $I \rightarrow I_{\text{boundary}}$ substitution. Both are calibration extensions; the algebraic structure of the framework is unchanged.

Falsification condition: A black hole system whose ξ_{CP} , evaluated at T_H with area-scaling I , is nonzero -- i.e., a physical black hole that violates the holographic bound. This is equivalent to a violation of the Bekenstein-Hawking formula.

Status: Tier III -- derived analytically; experimental confirmation requires a holographic system where T_H is measurable and the entropy-area relation is testable (analogue gravity systems, e.g., sonic black holes with tunable Hawking temperature).

0.3.4 P-26 · Hilbert space factorization failure recovered via P-20 idempotency boundary

Prediction. $\text{tensor}(\text{BH_interior}, \text{BH_exterior})$ at maximal entanglement ($\text{frac} = 1$, all primitives matching between interior and exterior synthons) returns $\xi_{\text{ens}} = \max(\xi_{\text{int}}, \xi_{\text{ext}})$ rather than $\xi_{\text{int}} + \xi_{\text{ext}}$. This is the algebraic signature of Hilbert space factorization failure: combining two maximally entangled $G_{\aleph}$ systems does not add their information.

Derivation. From P-20: $\lambda = \text{frac} = 1$. From the tensor formula: $\xi_{\text{ens}} = \xi_1 + \xi_2 - 1 \cdot \min(\xi_1, \xi_2) = \max(\xi_1, \xi_2)$. No new axioms required -- the idempotency

boundary condition of P-20 already encodes factorization failure as a continuous limit. The interpolation $\xi_{\text{ens}} = (2 - \text{frac}) \cdot \xi$ (for $\xi_1 = \xi_2 = \xi$) smoothly transitions from additive ($\text{frac} = 0$) to non-additive ($\text{frac} = 1$) as entanglement increases.

Residual gap (Type III algebras). The above derivation holds for Type I/II von Neumann algebras (trace-class density operator, countable states). Black hole horizons involve Type III₁ algebras (no trace, no density matrix, only relative entropies well-defined via Tomita-Takesaki modular theory). The framework's I(bits) pipeline must be extended to a relative-entropy form $D_{KL}(\rho \parallel \sigma)$ for Type III systems. This is a calibration extension, not a structural revision: the algebraic result (factorization failure at $\text{frac} = 1$) is correct; only the I computation changes.

Falsification condition: A pair of maximally entangled G_N synthons ($\text{frac} = 1$) for which tensor returns $\xi_{\text{ens}} < \max(\xi_1, \xi_2)$ -- i.e., entanglement somehow reduces the combined information below the maximum of the parts. This would violate the monotonicity of quantum mutual information and is physically excluded.

Status: Tier II/III -- the P-20 derivation is algebraically confirmed (Tier II); the Type III extension and experimental test in an analogue gravity system are Tier III.

0.3.5 P-27 · ER=EPR as R-primitive degeneracy at G_N ; extended Axiom 5

Prediction. If the ER=EPR conjecture (Maldacena-Susskind 2013) is correct, the R primitive is not fully independent at G_N : $R_{\leftrightarrow}$ (mechanical/topological -- wormhole) and $R_{\supseteq}$ (non-covalent/entanglement -- EPR pair) are physically indistinguishable in the G_N regime. The effective R dimension at G_N reduces from 4 to 3 ($R_{\leftrightarrow} \equiv R_{\supseteq}$; $R_{\subseteq}$ and $R_{\dagger}$ remain distinct). The effective independent primitive count at G_N is $10 \rightarrow 8$ (one from G/D degeneracy per Axiom 5; one from R-degeneracy per ER=EPR).

Quantitative form. Under the extended Axiom 5 R-degeneracy: $\text{distance}(\text{ER_bridge}, \text{EPR_pair}) = 0$ at G_N (vs current value of 0.27 with R and D treated as independent). $\text{tensor}(\text{ER}, \text{EPR})$ at full degeneracy: $\text{frac} = 1$, $\xi_{\text{ens}} = \max(\xi_{\text{ER}}, \xi_{\text{EPR}})$ -- they are the same system in dual descriptions, carrying no additional combined information.

Falsification condition (strong form): A physical system that simultaneously exhibits ER bridge geometry (wormhole-like topology with $R_{\leftrightarrow}$ barrier profile -- a steric cliff rather than a smooth Morse curve) and EPR correlations ($R_{\supseteq}$ -- smooth, non-topological entanglement), where the two descriptions give measurably different ξ_{CP} values at G_N . This would disprove ER=EPR directly and confirm that $R_{\leftrightarrow} \neq R_{\supseteq}$ at G_N .

Falsification condition (weak form): Any G_N system where the steric-cliff barrier profile (the operational signature of $R_{\leftrightarrow}$) and the smooth Morse profile (the operational signature of $R_{\supseteq}$) can be independently measured and shown to be distinct. At $G_{\supseteq}$, this is easy (rotaxane vs H-bond). At G_N , the framework predicts no instrument can distinguish them.

The R values that do NOT degenerate at G_N . $R_{\subseteq}$ (covalent -- specific bond

formation) and $R_{\ddagger}$ (catalytic -- transformation of one species by another) remain operationally distinguishable even at G_N , because their distinction does not rely on the presence of a geometric background. Covalent bond formation requires specific orbital overlap regardless of background structure; catalytic recognition requires a distinction between catalyst and substrate that is preserved under holography (the boundary CFT contains both types).

Extended Axiom 5 (proposed formal statement). At G_N and Φ_c : (i) G/D degenerate -- G_N and D_∞ become informationally equivalent (original Axiom 5); (ii) R-degenerate -- $R_{\Leftrightarrow} \equiv R_{\supseteq}$ (ER=EPR; this paper's extension). The framework's distance function must apply a G_{aleph} -conditional metric: $\delta_R(R_{\Leftrightarrow}, R_{\supseteq}) = 0$ when both synthons have $G = G_N$; $\delta_R = 1$ otherwise (standard categorical distance). Implementation: `compute_distance(s1, s2, g_aleph_mode=True)`.

Status: Tier III -- ER=EPR is a conjecture in quantum gravity with strong supporting evidence (Ryu-Takayanagi, firewall paradox resolution) but no proof. The framework's prediction is conditional on ER=EPR; it provides the precise primitive statement of what ER=EPR means in the synthon vocabulary.

0.3.6 P-16 · Rotaxane Φ_c anchor (Transformation #8, full scan)

Prediction. A full constrained-relaxed scan of a DB24C8/dialkylammonium borderline-slippage pseudorotaxane at ω B97X-D/def2-TZVPP + PCM(CHCl) should find: (i) degeneracy_strength ≥ 0.70 (logarithmic class) at the steric-cliff TS (ii) TS spatial correlation length $\xi_r \approx$ axle-crown distance at the steric cliff (3.5-4.5 Å) (iii) The first experimental anchor for Axiom 5 and the Φ_c HotSwap regime

If confirmed, this becomes the canonical Φ_c test case for the criticality-tolerant HotSwap predicted by Axiom 5 (a swap near the steric-cliff locus should be kinetically indistinguishable from a swap at the same F/K primitives away from the cliff).

Falsifiable: degeneracy_strength < 0.50 would disconfirm the Φ_c candidacy and classify the system as non-critical by the framework's own metric.

0.3.7 P-17 · Ice XIX correlation length $<$ ice XV (neutron diffraction)

Prediction. Ice XIX (>1.5 GPa, G_{beth} assigned by the LLM without prompting) has a shorter antiferroelectric ordering correlation length than ice XV (~ 1.0 GPa, G_{gimel}). This follows from the G primitive assignment alone: $G_{\text{beth}} = \text{local}$, $G_{\text{gimel}} = \text{mesoscale}$.

Measurable by: neutron diffraction correlation function analysis on ice XIX and XV under pressure. If correlation lengths are equal or $XV < XIX$, the G assignment convention requires revision.

Framework confidence: MEDIUM -- the G_{beth} vs. G_{gimel} assignment was made by LLM generation from the pressure-dependent O-O distance argument; the prediction is as reliable as that argument.

0.3.8 P-18 · Universality track in tuple space (phase diagram, +2.303 nat displacement)

Prediction. In a metric MDS projection of the quantum synthon catalog, adding a disordered Kitaev chain (K_{MBL} , $T_{\downarrow}$, Ω_Z , all other primitives identical to $kitaev_chain_majorana$) should appear at a **fixed displacement of $d = 2.303$ nats** from $kitaev_chain_majorana$ in the direction shared by all three $K_{\text{trap}} \rightarrow K_{\text{MBL}}$ shifts. This displacement should be parallel for $\text{spin_singlet} \rightarrow \text{disordered_spin}$ and $\text{fqh} \rightarrow \text{disordered_fqh}$.

Measurable by: running syncon phase-diagram with the three MBL synthons added to the catalog and verifying the displacement direction is collinear.

This is the "universality track" in primitive space -- a purely syntactic prediction about catalog geometry.

0.3.9 P-19 · $\chi(T \rightarrow 0) \sim T^{-\gamma}$ divergence from Factor 8 (quantum criticality)

Prediction. Synthons satisfying the Factor 8 fingerprint ($G_{\downarrow} + F_h + K_{\text{trap}} + \neg D_{\infty}$) sit at or near a quantum critical point. Near a QCP, the susceptibility $\chi(T \rightarrow 0) \sim T^{-\gamma}$ with $\gamma > 0$. The framework predicts this divergence for $kitaev_chain$ and fqh_moore_read but not for $\text{topological_insulator_bi2se3}$ (K_{slow} , not K_{trap} ; Factor 8 does not fire).

Measurable by: specific-heat or susceptibility measurements on Kitaev chain candidate materials (e.g., $\alpha\text{-RuCl}_3$, NaIrO_3) vs. BiSe topological insulator. If TI shows no divergence but Kitaev candidates do, Factor 8 is confirmed as a quantum criticality diagnostic. If TI also diverges, Factor 8 under-selects and the $G_{\downarrow} + K_{\text{trap}}$ combination needs refinement.

0.3.10 P-28 · β -sheet scaffolds and active-site mutational tolerance

Primitive basis. The tuple-space distance between active_site ($\langle D_{\wedge}; T_{\boxtimes}; R_{\ddagger}; P_{+-}; F_{\text{H}}; K_{\text{mod}}; G_{\boxminus}; \Gamma_{\wedge} \rangle$) and beta_hairpin ($\langle D_{\wedge}; T_{\boxtimes}; R_{\boxminus}; P_{\pm}^{\text{sym}}; F_h; K_{\text{mod}}; G_{\boxminus}; \Gamma_{\wedge}(\text{SPECIFIC}) \rangle$) is $d = 2.80$ -- the smallest distance between any passive scaffold and the active-site tuple. The α_helix ($T_{\downarrow}$, $\Gamma_{\rightarrow}$) is at $d \geq 4.50$ from active_site . β -sheet topology ($T_{\boxtimes}$) and specificity grammar ($\Gamma_{\wedge}(\text{SPECIFIC})$) are shared between scaffold and catalytic motif; helix topology ($T_{\downarrow}$) and sequential grammar ($\Gamma_{\rightarrow}$) are not.

Prediction. Alanine-scan mutations at active-site residues in β -sheet enzymes (TIM barrel, Rossmann fold, β -propeller) will be tolerated more frequently than equivalent mutations in helix-bundle enzymes (4-helix bundle, up-down bundle, coiled-coil) -- because the scaffold is closer to the active-site tuple and provides structural context that absorbs partial primitive disruptions. The fraction of alanine-scan mutations retaining $> 10\%$ wild-type activity should be higher in β -sheet enzyme classes.

Falsification condition. A systematic alanine-scan dataset (e.g., ProTherm, ProtBank) showing no significant difference in tolerance rates between β -

sheet and helix-bundle enzyme classes, matched for active-site geometry and substrate type.

Status: Tier II -- the distance matrix result is algebraically confirmed (protein_to the experimental correlation requires systematic mutational data analysis.

0.3.11 P-29 · Complex assembly bottleneck at K_{slow} from tensor

Primitive basis. `tensor(active_site, allosteric_domain)` reports K_{slow} as the ensemble bottleneck primitive. The allosteric domain carries K_{mod} ; the protein complex carries K_{slow} ; the tensor result gives $K = \min(K_1, K_2) = K_{\text{slow}}$. The tensor bottleneck rule (validated as P-15, V-6 design suite: design 12, crown[n] F-bottleneck) predicts assembly rate is controlled by the slowest kinetic component.

Prediction. The rate-limiting step in heteromeric protein complex assembly is the association/conformational rearrangement of the allosteric regulatory subunit, not the catalytic subunit. Stopped-flow measurements should show $k_{\text{fold}} > k_{\text{assemble}}$ systematically for assemblies with allosteric regulatory subunits.

Falsification condition. Stopped-flow data showing $k_{\text{assemble}} \approx k_{\text{fold}}$ (assembly not slower than folding) for allosteric enzyme complexes -- implying the tensor bottleneck rule does not transfer from supramolecular chemistry to protein assembly.

Status: Tier II -- tensor bottleneck algebraically confirmed (V-6); protein-domain transfer requires stopped-flow kinetics.

0.3.12 P-30 · Allosteric ON state: multi-timescale R_{ex} spectrum from Φ_c

Primitive basis. The allosteric domain is the only protein structural unit satisfying the G/D degeneracy condition for Φ_c (Varma probe score 0.60). The Primitive Jacobian shows 6 of 8 primitives trigger Axiom 4 violations -- the allosteric domain is axiom-fragile by design. Near-critical systems (gIX) exhibit broad fluctuation spectra because they operate simultaneously at multiple length and time scales; a single-Lorentzian R_{ex} spectrum indicates a single dominant timescale, characteristic of subcritical systems.

Prediction. The conformational exchange spectrum of allosteric domains in the signal-active (ON) state -- measured by CPMG NMR relaxation dispersion -- should show contributions across multiple timescales simultaneously (microsecond to millisecond R_{ex} distribution). The signal-inactive (OFF) state, further from the criticality locus, should show a single dominant Lorentzian or no significant R_{ex} . Test system: CAP protein (cAMP-dependent allosteric transcription factor), whose ON/OFF switch is structurally characterized.

Falsification condition. CPMG data showing a single-Lorentzian R_{ex} spectrum for the ON state of a canonical allosteric protein, indistinguishable from the OFF-state spectrum.

Status: Tier III -- the Φ_c assignment to allosteric domains is primitive-derived; the multi-timescale spectral prediction requires NMR measurement.

0.3.13 P-31 · Directed distance asymmetry: rescue cheaper than aggregation (K-targeting corollary)

Primitive basis. Directed tuple distances (F-floor asymmetry, protein_tests3.py) for all three amyloid systems give rescue cost < aggregation cost:

Disease	Aggregation cost	Rescue cost	Asymmetry
A β	7.20	6.90	0.30
Tau	4.80	3.90	0.90
α -Syn	4.80	3.90	0.90

The asymmetry arises because the F-floor rule blocks upward F transitions ($F_\ell \rightarrow F_h$ in aggregation -- requires discontinuous topology change) while kinetic downgrade ($K_{\text{trap}} \rightarrow K_{\text{fast}}$ in rescue) is not floor-constrained.

Prediction. Kinetic-targeting rescue agents (seeding inhibitors, disaggregases, K -targeting chaperones) outperform thermodynamic denaturants (F -targeting compounds) in fibril dissolution assays. The differential should follow the rank order $A\beta < \text{tau} \approx \alpha\text{-Syn}$, matching the directed-distance asymmetry.

Falsification condition. F -targeting denaturants dissolving fibrils as rapidly as K -targeting disaggregases at matched thermodynamic driving force, or the rank order not following the predicted asymmetry values.

Status: Tier II -- directed-distance asymmetry algebraically confirmed; experimental rank comparison requires head-to-head kinetics.

0.3.14 P-32 · Allosteric-interface chimera: binding retained, Hill coefficient $n_H \rightarrow 1$

Primitive basis. `meet(allosteric_domain, protein_complex)` produces two CONFLICTS: on P (P_{+-} vs. $P_{\pm}^{\text{sym}}$) and on Γ ($\Gamma_{\rightarrow}(\text{SELECTIVE})$ vs. $\Gamma_{\wedge}(\text{SPECIFIC})$). These conflicts encode a structural incompatibility between allosteric signaling grammar (sequential, directional) and interface grammar (simultaneous, symmetric). A chimeric protein at this conflict boundary retains the common primitives ($D_{\wedge\Delta}$, T_{ϵ} , $R_{\supseteq}$, F_{δ} , K_{mod} , G_{J}) -- and therefore binding affinity -- but loses the coherent grammar required for cooperativity.

Prediction. A chimera between an allosteric regulatory domain and a quaternary interface domain retains binding to both cognate partners (K_d unchanged within 10 $\times$) but loses cooperativity: Hill coefficient $n_H \rightarrow 1.0$. ITC should show a normal enthalpic signature but linear binding isotherm; Hill analysis on a cooperative reporter should show n_H collapse.

Falsification condition. A chimera at this conflict boundary retaining $n_H > 1.5$, inconsistent with the meet-conflict prediction of grammar incompatibility.

Status: Tier III -- requires chimera protein engineering and biophysical characteriz

0.3.15 P-33 · Allosteric inhibitors: selectivity advantage from $G_{\downarrow}$ vs. $G_{\uparrow}$

Primitive basis. Orthosteric drugs bind at $R_{\uparrow}$, $G_{\uparrow}$ (local scale -- active-site pocket, shared across enzyme family members). Allosteric drugs bind at $R_{\downarrow}$, $G_{\downarrow}$ (mesoscale -- full allosteric communication path, more divergent across paralogs than the catalytic pocket). By the G primitive's scale-of-control definition: $G_{\downarrow}$ selectivity window = local pocket geometry; $G_{\uparrow}$ selectivity window = domain-level contact network. The domain-level network is more divergent across kinase paralogs than the ATP-binding cleft.

Prediction. For any kinase family with a conserved active site, allosteric inhibitors targeting the regulatory domain ($G_{\downarrow}$) will show higher selectivity (lower off-target activity against family paralogs) than orthosteric inhibitors targeting the ATP-binding site ($G_{\uparrow}$), in a matched selectivity panel (e.g., KINOMEscan, Ambit), even when both compounds have equivalent K_i against the primary target.

Falsification condition. A selectivity panel showing equivalent or superior selectivity for an orthosteric compound vs. its allosteric analog in the same kinase family.

Status: Tier II -- the G primitive assignment is the operative prediction; selectivity panel comparison is available in published kinase inhibitor datasets.

0.3.16 P-34 · $A\beta$ - α -Syn cross-seeding rate $\approx$ homoseeding rate; both faster than tau

Primitive basis. $\text{tuple_distance}(\text{Abeta_fibril}, \text{alphaSyn_fibril}) = 0.00$. Identical primitive encoding across all 8 dimensions: both are $\langle D_{\wedge\Delta}; T_{\epsilon}; R_{\supseteq}; P_{\pm}^{\text{sym}}; F_h; K_{\text{trap}}; G_{\text{N}}; \Gamma_{\wedge}(\text{E}) \rangle$. The same synthon is the same nucleation template -- there is no primitive distinction that would impose a kinetic penalty for heterologous seeding relative to homologous seeding. Tau PHF differs at $d = 2.90$ ($T_{\bowtie}$ vs. T_{ϵ} , $P_{\pm}^{\psi}$ vs. $P_{\pm}^{\text{sym}}$), predicting a cross-seeding barrier.

Prediction. Cross-seeding kinetics ($A\beta$ seeds $\rightarrow$ α -Syn monomer; α -Syn seeds $\rightarrow$ $A\beta$ monomer) should show nucleation lag times and elongation rates approximately equal to homoseeding controls ($A\beta$ seeds $\rightarrow$ $A\beta$ monomer; α -Syn seeds $\rightarrow$ α -Syn monomer). Both should be substantially faster than tau seeding into either $A\beta$ or α -Syn monomer. Measurable by ThT fluorescence kinetics with exogenous seeds at sub-threshold concentration.

Falsification condition. A systematic cross-seeding experiment (matched seed concentration, monomer concentration, buffer) showing cross-seeding lag time $> 3\times$ homoseeding lag time for the $A\beta$ / α -Syn pair.

Status: Tier II -- $d = 0.00$ is algebraically confirmed; the kinetic equality is consistent with published cross-seeding reports (Guo et al. 2013, *Nat. Neurosci.*) but has not been systematically tested with matched homoseeding controls.

0.3.17 P-35 · K -targeting fibril dissolution outperforms F -targeting denaturants

Primitive basis. P-31 establishes the directed-distance asymmetry. Here the corollary is sharpened to a compound-class prediction: F -targeting agents (chaotropes, detergents) must navigate the F -floor to dissolve F_h fibrils, requiring a discontinuous topology change. K -targeting disaggregases (Hsp70+Hsp40+NB) reduce K_{trap} -- a path not floor-constrained. The tuple-space distance from fibril to monomer is 0.90 nat shorter in the K -targeted direction (P-31 table, tau and α -Syn).

Prediction. In a head-to-head dissolution assay at matched thermodynamic driving force, K -targeting disaggregase systems dissolve preformed amyloid fibrils faster than F -targeting chemical denaturants. The rate differential should rank $A\beta < \text{tau} \approx \alpha\text{-Syn}$, following the directed-distance asymmetry: $A\beta$ has the smallest asymmetry (0.30) and thus the smallest predicted advantage.

Falsification condition. GdnHCl or urea at sub-denaturing concentrations dissolving preformed fibrils as rapidly as a chaperone system at matched thermodynamic cost. Or: the dissolution rate differential not following the predicted $A\beta < \text{tau} \approx \alpha\text{-Syn}$ rank.

Status: Tier II -- directed-distance result algebraically confirmed; chaperone vs. denaturant comparison and rank ordering require experimental measurement.

0.3.18 P-36 · Bivalent GNF-2: $T_{\perp} \rightarrow T_{\epsilon}$ closes the gap to the ideal allosteric inhibitor

Primitive basis. GNF-2 encodes $T_{\perp}$ (single branched pocket -- myristoyl binding site only). The ideal allosteric inhibitor encodes T_{ϵ} (distributed contact network). The per-primitive gap analysis (protein_tests4.py) identifies this T mismatch as one of the two primitive gaps between GNF-2 ($d=2.80$ from ideal) and the allosteric ideal. A bitopic inhibitor spanning the myristoyl pocket and a second regulatory site simultaneously shifts $T_{\perp} \rightarrow T_{\epsilon}$, closing half the remaining distance.

Prediction. A bivalent GNF-2 analog engaging the ABL myristoyl pocket and a second allosteric site (e.g., the SH2-kinase linker contact) simultaneously should show: (i) lower ξ_{CP} than GNF-2 monovalent (more Landauer-efficient constraint propagation); (ii) higher kinase selectivity (distributed contact grammar more paralog-discriminating); (iii) slower resistance emergence ($G_{\perp}$ multi-site engagement tolerates single-site mutations). Design direction is analogous to published bitopic type I_j kinase inhibitors.

Falsification condition. A bivalent GNF-2 analog with T_{ϵ} character showing lower selectivity and faster resistance emergence than the GNF-2 monovalent -- contradicting the $T_{\perp} \rightarrow T_{\epsilon}$ primitive upgrade prediction.

Status: Tier III -- requires medicinal chemistry synthesis and SAR characterization.

0.3.19 P-37 · $G_{\perp}$ drugs: binary resistance; $G_{\parallel}$ drugs: incremental resistance

Primitive basis. Imatinib encodes $G_{\perp}$ (local scale): binding grammar operates at the DFG-pocket scale. Any pocket mutation disrupts binding without compensating alternative grammar -- the drug has zero contact redundancy. GNF-2 encodes $G_{\parallel}$ (mesoscale): binding grammar propagates across the allosteric domain. A single myristoyl-pocket mutation changes one node of the contact network but does not destroy full signal propagation. The G primitive's scale-of-control definition directly predicts the resistance trajectory topology.

Prediction. In serial passage resistance evolution experiments: (i) imatinib-resistant clones require complete drug class switching -- pocket mutation abolishes all binding (verified: T315I, E255K → imatinib → ponatinib/asciminib); (ii) GNF-2-resistant clones show incremental resistance -- the first resistance mutation reduces, but does not abolish, sensitivity. The first passage mutation in a GNF-2 serial passage experiment should produce partial resistance (e.g., IC_{50} shift 5-20×), not the binary switch (>1000× shift) characteristic of $G_{\perp}$ drugs.

Falsification condition. GNF-2 showing the same binary resistance profile as imatinib -- a first mutation abolishing all detectable inhibition -- contradicting the $G_{\parallel}$ multi-contact tolerance prediction.

Status: Tier II -- $G_{\perp}$ irreversibility is confirmed by clinical imatinib resistance data; the $G_{\parallel}$ incremental-resistance prediction for GNF-2 is a derived consequence testable by published serial passage protocols.

0.4 Programmable Matter Domain (gXIX) -- P-38 through P-47

Derived from programmable_matter_tests1.py (11 synthon encodings, pairwise distance matrix, meet operations, path algebra, Varma probe, cross-domain analogy) and programmable_matter_tests2.py (Primitive Jacobian, tensor products, DesignPipeline monad). All predictions algebraically confirmed; all await experimental validation.

0.4.1 P-38 -- Programmability pair distance predicts switching energy rank (gXIX)

Primitive basis. The tuple distance between the two states of a programmable material (its "programmability pair distance") measures how many primitives change during a state transition and by how much. This distance should correlate with the switching energy -- the thermodynamic cost of crossing from one state to the other -- because each primitive change corresponds to a real physical difference (F = binding hierarchy, K = kinetic barrier, G = scale of order). Three material classes have cleanly resolved pair distances: $d_{SMP} = 1.70$, $d_{LC} = 3.10$, $d_{colloidal} = 5.10$.

Prediction. The switching energy (measurable as enthalpy: Tg-crossing enthalpy for SMP, N→I transition enthalpy for LC, melting enthalpy of colloidal crystal) follows the same rank: $\Delta H_{SMP} < \Delta H_{LC} < \Delta H_{colloidal \text{ crystal}}$. No domain-specific

thermodynamic equation is inserted between the distance rank and the energy rank prediction -- the ordinal relationship is derived purely from the primitive metric.

Falsification condition. Any experimental rank inversion of switching enthalpies across these three material classes.

Status: Tier II -- algebraically derived from primitive distance matrix; experimental calorimetry values known in literature (DSC for SMP: ~20-40 J/g; LC: ~2-10 J/g; colloidal: ~100+ J/g per particle contact) consistent with prediction but not used as inputs.

0.4.2 P-39 -- Condensate gel rescue requires Γ -targeting or K-targeting; thermal stimulus alone fails (g̃XIX)

Primitive basis. Condensate gel is encoded as $\langle D_{\Delta\wedge}; T_{\text{network}}; R; P_{\text{pm_sym}}; F_h; K_{\text{trap}}; G_N; \Gamma_{\wedge}(\text{BRO})$. The path search from condensate_gel to condensate_liquid is blocked: the F-floor theorem prohibits HotSwap traversal when $F_{\text{src}} = F_h > F_{\text{dst}} = F_{\ell}$ at K_{trap} . No intermediate synthon in the catalog provides a stepping stone. Thermal perturbation acts only on K -- it cannot lower F alone.

Prediction. Gel dissolution requires: (a) a competing binder that lowers F (Γ -targeting agent -- sequesters the gel-forming interaction), or (b) a disaggregase/chaperone that lowers K_{trap} (K-targeting). A stimulus that acts only on temperature (pure thermal) cannot rescue K_{trap} gels at physiological temperatures, because temperature operates on K and cannot by itself change the F -ordering of competing interactions.

Falsification condition. A purely thermal dissolution of a condensate gel (no competing ligand, no chaperone activity) at temperatures compatible with cell viability.

Status: Tier II -- path blocked algebraically; consistent with known biology (Hsp70 disaggregases required for TDP-43/FUS gel rescue; thermal dissolution alone requires denaturing temperatures).

0.4.3 P-40 -- T-conflict programmability pairs predict first-order transitions (g̃XIX)

Primitive basis. The meet operation on a programmability pair identifies which primitives conflict -- primitives that differ between the two states and cannot be reconciled by a shared lower bound. Topology (T) conflicts appear in: LC nematic (T_{linear}) vs isotropic (T_{network}), colloidal crystal ($T_{\text{network_sym}}$) vs fluid (T_{network}). The HotSwap path is blocked for all T -conflicting pairs: "HotSwap requires exact D and T match." Topology is discontinuous -- it cannot be traversed by continuous deformation; any topology change requires a discontinuous jump.

Prediction. All programmability pairs that show T -conflict in their meet operation undergo first-order (discontinuous) transitions: they exhibit latent heat, coexistence regions, and hysteresis. Pairs with no T -conflict (SMP:

$T_{\text{network}} \leftrightarrow T_{\text{network}}$, condensate: $T_{\text{network}} \leftrightarrow T_{\text{network}}$) may be second-order or weakly first-order.

Falsification condition. A programmability pair with T -conflict in its meet that undergoes a continuous (second-order) transition with no latent heat and no coexistence.

Status: Tier II -- algebraically derived; LC $N \rightarrow I$ is known to be weakly first-order (latent heat ~ 2 kJ/mol), colloidal melting is first-order -- both consistent with T -conflict prediction.

0.4.4 P-41 -- Actin-DNA hybrid composites show collective motion only with ATP (ǧXIX)

Primitive basis. Tensor product: $\text{active_gel} \otimes \text{dna_strand_disp.}$ The composite acquires Φ_c from the active gel component (criticality is join-dominant) and $G_{\aleph}$ (global scale from active gel), R (non-covalent recognition from DNA, downgrading the DYNAMIC_CATALYTIC mode). The Φ_c assignment is conditional on the active gel primitive, which requires $D_{\Delta\infty}$ (temporal cycle = ATP hydrolysis). Removing ATP collapses $D_{\Delta\infty} \rightarrow D_{\Delta}$ (no temporal cycle), which changes the composite to a passive DNA network without Φ_c .

Prediction. Actin-DNA composite networks (e.g., kinesin-DNA nanostructures, actin-DNA scaffolds): (1) show collective motion and spatial order absent in either component alone -- emergent from tensor composite; (2) lose collective motion upon ATP depletion faster than DNA scaffold degrades; (3) the threshold ATP concentration for collective motion onset matches Varma criticality score ≈ 0.60 (onset near, but below, full criticality).

Falsification condition. ATP-depleted actin-DNA network shows equivalent spatial order (correlation length, velocity correlations) to ATP-active network.

Status: Tier III -- composite derived from tensor algebra; actin-DNA composites are known but the specific ATP-dependency of collective motion onset has not been tested with this framing.

0.4.5 P-42 -- Maximally versatile programmable matter is generically near-critical (ǧXIX)

Primitive basis. The meet (dynamic floor) of all fluid/dynamic programmable matter states carries Φ_c : $\langle D_{\Delta}; T_{\text{network}}; R; P_{\text{pm_sym}}; F_{\ell}; K_{\text{mod}}; G_{\sqsupset}; \Gamma_{\vee}(\text{BROAD}); \Phi_c \rangle$. This is not a design choice inserted into any individual encoding -- it emerges from the lattice meet of six independently encoded systems. Φ_c is join-dominant in the meet lattice: it propagates whenever at least one component is near-critical.

Prediction. Any programmable matter system engineered to maximize the number of accessible states (highest programmability) will converge to near-critical encoding: $F_{\ell}, K_{\text{mod}}, G_{\text{global}}, \Phi_c$. This is not because designers choose criticality -- it is because the algebra shows Φ_c is the only phase assignment compatible

with the dynamic floor. Consequence: the most versatile programmable matter in biology (cytoplasm, membraneless organelles) operates near criticality -- consistent with measured cortical criticality, now derived from primitive algebra without biological inputs.

Falsification condition. A maximally versatile programmable matter system (demonstrated large state space, global responsiveness) that is provably subcritical in all observables.

Status: Tier II -- lattice algebra result; consistent with growing evidence for criticality in biological PM systems (cortex, cytoplasm); prediction extends to synthetic PM design.

0.4.6 P-43 -- Condensates implement mesoscale allostery with Hill coefficient tied to criticality score (ǧXIX)

Primitive basis. Cross-domain analogy: nearest catalog neighbor to condensate_liquid is allosteric_domain at $d = 2.50$ (closer than any molecular synthon). The shared primitives are: F_ℓ (individually weak contacts), K_{fast} (dynamic exchange), G_J (mesoscale scale of control), Γ_V (promiscuous partner), Φ_c (near-critical). This is the same tuple structure that defines a mesoscale allosteric system: many weak contacts at intermediate scale, dynamic, near-critical, promiscuous.

Prediction. (1) Condensate-mediated signaling shows distance-independent propagation within the droplet -- a perturbation at one face propagates to the opposite face faster than diffusion-limited transport, because $G_\text{J} + \Phi_c$ enables global-from-local propagation. (2) The apparent Hill coefficient of condensate-mediated catalysis equals the Varma criticality score to within 15%: $n_H \approx 0.60/0.50 \approx 1.2$. (3) Condensate dissolution abolishes cooperativity -- n_H drops to 1.0.

Falsification condition. Condensate-mediated signaling shows purely diffusion-limited propagation with no superdiffusive component. Or: n_H in a condensate-mediated reaction is > 2.0 or < 1.0 .

Status: Tier III -- cross-domain analogy; the Hill coefficient prediction is novel and experimentally accessible by comparing condensate vs. dilute-phase catalysis rates.

0.4.7 P-44 -- Colloidal crystals with correct symmetry support topologically protected boundary states (ǧXIX)

Primitive basis. Nearest catalog neighbor to colloidal_crystal is topological_insulator at $d = 2.80$, and synthon_neutronium at $d = 2.80$. The shared structure: F_h (high fidelity collective binding), G_N (global crystalline order), T_{network} (lattice), R (non-covalent). This is the same primitive cluster as a topological insulator. The distance $d = 2.80$ is close enough to predict structural analogy.

Prediction. Colloidal crystals with $d < 3.0$ from topological_insulator in the primitive metric support topologically protected boundary modes -- surface states robust to bulk disorder -- when the interaction design matrix has the

correct band-crossing symmetry (non-trivial Zak phase). The range of colloidal systems predicted to show topological boundary states is defined by $d < 3.0$: any colloidal crystal with F_h , G_N , T_{network} and tuned interaction symmetry qualifies.

Falsification condition. A colloidal crystal with $d < 3.0$ from topological_insulator but no detectable boundary-protected states, even after interaction symmetry tuning.

Status: Tier II -- phenomenon confirmed experimentally for DNA-coated colloidal crystals (Rechtsman 2016 and subsequent); prediction extends to: the primitive distance threshold ($d < 3.0$) as the boundary of the topological analogy.

0.4.8 P-45 -- Topologically closed DNA origami (knots, catenanes) shows polynomial strand-failure sensitivity (ǧXIX)

Primitive basis. DNA origami is nearest to topological_insulator_bi2se3 at $d = 3.70$ -- more distant than colloidal crystal ($d = 2.80$) but still within topological analogy range. DNA origami with $\Omega \neq 0$ (topological closure -- knots, catenanes, Borromean rings) acquires topological protection. In the framework, Ω measures topological robustness: how many strand-failure events are required to destroy structural integrity changes from n (linear: one staple failure is irreversible) to $O(n^2)$ or better (topological: multiple simultaneous failures required).

Prediction. A DNA origami structure with topological closure ($\Omega \neq 0$) shows error-rate scaling as $\sim \exp(-n_{\text{strands}})$ only in open structures. In topologically closed structures, error scaling becomes $\sim n_{\text{strands}}^{-k}$ where k depends on the topological class. Measurable: compare FRET-detected folding yield for open vs. catenated origami as a function of staple strand concentration.

Falsification condition. Topologically closed DNA origami (knots, catenanes) shows identical strand-failure sensitivity to equivalent open origami.

Status: Tier III -- topological DNA origami exists experimentally (Lim et al. 2020) but polynomial vs. exponential error scaling has not been tested as a function of topological class.

0.4.9 P-46 -- Primitive Jacobian identifies dominant design lever: F controls > 40% of d-reduction (ǧXIX)

Primitive basis. The primitive Jacobian $\partial d / \partial \text{primitive}$ measures the sensitivity of programmability pair distance to single-primitive perturbation. Computed for all five PM pairs: F (fidelity) gives the largest $|\Delta d|$ in DNA (-1.10), colloidal (-1.20), SMP (-0.60), LC (-0.60) pairs. K (kinetic character) dominates condensate gel rescue (-1.50). G contributes secondary leverage in DNA (-0.90) and colloidal (-0.80) pairs but is zero for SMP (G_gimel invariant in both states).

Prediction. Engineering the Jacobian-identified primitive alone achieves > 40% of the maximum possible pair distance reduction (full 11-primitive optimization).

Corollary: multi-primitive optimization gives diminishing returns beyond the top two Jacobian-ranked primitives (law of diminishing returns in primitive space). Specific experimental test: vary T_g ($= F$ proxy for SMP) while holding crosslink density (K proxy) and domain size (G proxy) constant -- this alone should halve the switching energy relative to varying crosslink density alone.

Falsification condition. For any PM pair, optimizing the bottom-ranked Jacobian primitive gives equal or larger d-reduction than the top-ranked primitive.

Status: Tier III -- Jacobian computed algebraically; experimental T_g vs crosslink-density vs domain-size optimization comparison is feasible but not yet published with this framing.

0.4.10 P-47 -- Actin-DNA composites cannot achieve a shared locked state: structural integrity requires ATP (gXIX)

Primitive basis. $\text{meet}(\text{active_gel}, \text{dna_origami})$ conflicts on $\{D, R, P, \Gamma\}$. Active gel has $D_{\Delta\infty}$ (temporal cycle = ATP hydrolysis); DNA origami has $D_{\Delta\wedge}$ (molecular + supramolecular, no temporal cycle). These are incompatible at the level of dimensionality: no shared D value exists that satisfies both. Materials whose rigid states conflict on D have no shared locked state in the algebra -- there is no primitive tuple that is both "static DNA scaffold" and "static actin network" simultaneously.

Prediction. Actin-DNA composite materials cannot maintain structural integrity without ATP. The "static DNA scaffold plus actin" design is algebraically incoherent -- actin's structural integrity is encoded in $D_{\Delta\infty}$ (the temporal ATP cycle), so removing ATP collapses the actin component. Specific prediction: actin-DNA hybrids degrade without ATP replenishment faster than equivalent ATP-free DNA origami scaffolds, because actin depolymerisation is part of the actin synthon's D primitive (not an external degradation pathway).

Falsification condition. An actin-DNA composite that maintains structural integrity for > 24 h without ATP, matching performance of equivalent ATP-free DNA origami scaffold under identical conditions.

Status: Tier III -- meet conflict derived algebraically; consistent with known actin biology (G-actin/F-actin equilibrium requires ATP turnover) but specific comparison with DNA origami stability has not been published.

0.5 P-48 -- Condensate Gelation and Amyloid Fibrillization Are the Same Primitive Event; K-Targeting Is the Preferred Therapeutic Strategy for Both

Source. Programmable matter encoding (PROGRAMMABLE_MATTER.md g5); protein domain catalog (PROTEIN_APPLICATIONS.md).

Algebraic basis. $d(\text{condensate_gel}, \text{amyloid}) = 0.00$ -- the programmable matter gel state and the amyloid fibril are identical in all nine primitives:

$$\langle D_{\Delta}; T_{\in}; R; P_{\pm\text{pseudo}}; F_h; K_{\text{trap}}; G_{\text{J}}; \Gamma_{\text{and}}(\text{SELECTIVE}); \Phi_{\text{sub}} \rangle$$

This is not a structural analogy. A distance of zero means the primitive encoding is the same object. The two literatures (condensate biology and amyloid neuroscience) have independently identified the same physical attractor: a trapped, high-fidelity, mesoscale, supramolecular network with self-complementary polarity. The difference is biological context, not physical structure.

Primitive Jacobian result. The Jacobian computed over the full programmable matter catalog assigns the K primitive the highest single-primitive leverage for gel state rescue: targeting K ($K_{\text{trap}} \rightarrow K_{\text{fast}}$) reduces the tuple distance from the locked state to the dynamic floor by more than targeting F alone. This extends directly to amyloid: K -targeting (kinetic remodelling of the trapped state -- disaggregases, chaperones, ATP-dependent unfolding machines) should outperform F -targeting (binding competitors that lower association affinity) as a dissolution strategy.

Prediction 1 -- Mechanistic equivalence. A dissolution agent that successfully converts a condensate gel to the liquid state (reducing $K_{\text{trap}} \rightarrow K_{\text{fast}}$) will show non-trivial cross-reactivity against amyloid fibrils of the same polarity class ($P_{\pm\text{pseudo}}$: $A\beta$, α -Syn, tau), because the primitive target is identical. Agents that act through the F primitive (competitive binding, affinity-based disruption) will not show this cross-reactivity -- they address different positions in the tuple.

Prediction 2 -- Jacobian hierarchy. For any disease system where both condensate gelation and amyloid formation are co-pathological (e.g., TDP-43 in ALS, FUS in FTLD): therapeutic strategies that target K (Hsp70/Hsp104 chaperone axis, disaggregase activity, ATP-dependent remodelling) will outperform strategies that target F (small-molecule binding competitors, antibodies that raise the association barrier) in dissolution efficiency. Ratio of K -strategy to F -strategy dissolution rates should exceed $1.5\times$ from the Jacobian prediction.

Prediction 3 -- Locking asymmetry preservation. The same directed-distance asymmetry that makes programmable matter locking thermodynamically downhill applies to amyloid: the forward rate (monomer $\rightarrow$ fibril, $K_{\text{mod}} \rightarrow K_{\text{trap}}$) will always exceed the reverse rate (fibril $\rightarrow$ monomer, $K_{\text{trap}} \rightarrow K_{\text{mod}}$) under identical conditions, regardless of specific amino acid sequence. The asymmetry is structural, not sequence-specific. Falsified by: any amyloid system where spontaneous dissolution exceeds nucleated aggregation at the same monomer concentration and temperature.

Connection to P-34 and P-35. P-34 established that $A\beta$ and α -Syn share $d = 0.00$ (cross-seeding $\approx$ homoseeding). P-35 established that K -targeting dissolution ranks below F -targeting for the amyloid family. P-48 now links the condensate gelation literature to both: the three predictions form a coherent triangle -- zero-distance identity, K -strategy superiority, locking asymmetry -- all derivable from the same primitive tuple.

Falsification condition. (1) A dissolution agent that targets F (not K)

showing equal or greater cross-reactivity against both condensate gels and amyloid fibrils would falsify Prediction 1. (2) A K-strategy dissolution agent showing no advantage over F-strategy in a head-to-head assay for both target classes would falsify Prediction 2.

Status: Tier II -- zero-distance identity algebraically confirmed; Jacobian hierarchy computed from catalog; consistent with known biology (ATP-dependent disaggregases dissolve both condensates and amyloid; LLPS-to-gel transitions share kinetics with early amyloid nucleation). Experimental head-to-head comparison of K-targeting vs F-targeting agents against matched condensate gel and amyloid fibril targets not yet published.

0.6 P-49 -- Phi_c Categorical Independence: Critical Phase Is a Label, Not a Derived Ordinal

Origin. Decomposition algebra exploration, 2026-03-19. Principal result: `kernel(allosteric_domain, phi_c_probe)` strips F , K , G to their floors, but the $\text{Phi}=\text{CRITICAL}$ field survives because it is a *categorical* field in the synthon dataclass -- not computed from ordinals. Constructing `phi_c_skeleton` ($F=\text{LOW}$, $K=\text{FAST}$, $G=\text{LOCAL}$, $\text{Phi}=\text{CRITICAL}$) is a valid synthon. Catalog proof: `asymptotic_safet` carries $G=\text{LOCAL} + \text{Phi}=\text{CRITICAL}$ -- the UV fixed point of asymptotic safety is *locally* critical without global organisation.

Prediction. Criticality-organising effects should be observable in systems with individually weak interactions ($F=\text{LOW}$, each contact $\sim 1-3k_B T$) if the network topology ($T=\text{NETWORK}$) is correct. Predicted systems: LLPS condensates (weak IDR contacts but near-critical droplet), neuronal avalanches (low synaptic F , global Φ_c), and low-affinity multivalent receptors at crowded membranes. The Phi_c field encodes the *class* of dynamical regime -- not the magnitude of any ordinal.

Falsification condition. A system with $F=\text{LOW}$, $K=\text{FAST}$, $G=\text{LOCAL}$ that cannot in principle support a phase transition would falsify categorical independence. (Note: the prediction is about the encoding grammar -- it does not assert that every $F=\text{LOW}$ system *is* critical, only that criticality is not *forbidden* by low ordinals.)

Status: Tier II -- algebraically exact; catalog proof exists (`asymptotic_safety_reut` consistent with LLPS phenomenology. Systematic ordinal-vs-criticality screen not yet performed.

0.7 P-50 -- Amyloid Formation Algebraically Requires an External High-Fidelity Nucleation Seed

Origin. `cofactor(amyloid_fibril, condensate_liquid) → F-CONFLICT: tensor-min( $F=\text{MEDIUM}$ ,  $B$ ) ≤ MEDIUM < HIGH. A liquid condensate ( $F=\text{MEDIUM}$ ) cannot template amyloid ( $F=\text{HIGH}$ ) by primitive tensor alone. This is not a kinetic`

barrier -- it is a structural impossibility in the primitive algebra: no value of B satisfies $\min(\text{MEDIUM}, B) = \text{HIGH}$.

Prediction. Condensate-to-amyloid conversion must always be gated by an external high-fidelity nucleation event: a seed fibril, a metal ion interface (which elevates local F), a lipid membrane surface, or any other $F=\text{HIGH}$ input. Unseeded liquid condensates of the same protein should show an indefinite lag phase that collapses upon addition of pre-formed fibril seeds. The lag phase duration should be inversely proportional to seed concentration with a saturation plateau at the F -floor threshold.

Falsification condition. A purified liquid condensate that converts to amyloid *de novo* with no detectable nucleation lag, in a system verified to contain no fibril seeds or $F=\text{HIGH}$ surfaces, would falsify this prediction.

Connection to P-34, P-35, P-48. P-34 established $d(\text{A}\beta, \alpha\text{-Syn}) = 0.00$; P-48 established condensate gel and amyloid are the same primitive event. P-50 now algebraically derives the nucleation requirement, completing the triangle: identity (P-34) + K-strategy superiority (P-48) + F-floor nucleation barrier (P-50).

Status: Tier II -- algebraically exact (F-CONFLICT derived); consistent with seeded aggregation literature (Jarrett & Lansbury 1993, Bhak et al. 2009). Matched-control seeded vs. unseeded condensate kinetics under controlled seed-free conditions not yet published as a direct test.

0.8 P-51 -- The Quantization Residual: cofactor(QG, GR) Is the Quantization Operator as a Synthon

Origin. `cofactor(quantum_gravity, general_relativity)` computes residual B such that $\text{GR} \otimes B \approx \text{QG}$. The per-primitive analysis from `decompose_explorations2.py` gives:

Prediction. The D -CONFLICT -- GR's $D=\text{SUPRAMOLECULAR}$ (spatial substrate) vs. QG's $D=\text{TEMPORAL}$ -- is the algebraic form of the *background-dependence problem*: any quantisation scheme that preserves GR's spatial D -component will produce a D -CONFLICT at the quantum gravity level. A background-free theory of quantum gravity must either (a) abandon GR's spatial D -component, or (b) introduce a D -merging primitive not yet in the catalog. Corollary: the quantisation residual $B = \{K_{\text{trap}}, G_{\text{N}}, T_{\text{braid}}, \Phi_c, \Omega_{\text{NA}}\}$ is the minimal primitive tuple that, when tensored with GR, produces the observed structure of QG -- it encodes the complete content of "quantisation" as a single synthon.

Falsification condition. A formulation of quantum gravity that (a) retains GR's spatial D -component and (b) does not produce a D -conflict with the temporal structure of quantum fluctuations would require a revision of the D -primitive taxonomy or the tensor semantics.

Status: Tier III -- algebraically derived from catalog encodings; interpretation consistent with known obstacles to canonical quantisation; not yet experimentally

Primitive	Role	GR value	QG value	Interpretation
K	Contributor	SLOW	TRAP	Path-integral measure freezes high-action histories
G	Contributor	LOCAL	GLOBAL	Quantisation promotes local GR to global entanglement
T	Contributor	NETWORK	BRAID	Smooth spacetime $\rightarrow$ spin-foam / anyonic exchange topology
Phi	Contributor	SUBCRITICAL	CRITICAL	Quantum phase transitions have no classical analogue
Omega	Contributor	None	NON_ABELIAN	Non-Abelian gauge structure / anyonic statistics
D	CONFLICT	SUPRAMOLECULAR	TEMPORAL	GR organises matter in space; QG must organise spacetime — orthogonal D-components

testable in isolation. Cross-reference: P-23 (SM/QG lift blocked), P-24 (G partitions gravity theories), P-27 (ER=EPR extended Axiom 5).

0.9 P-52 -- GNF-2 Drug Combination Strategy: A Supramolecular Network Scaffold Closes the Allosteric Gap

Origin. Drug panel comparison via cofactor(allosteric_domain, drug) for GNF-2, imatinib, and venetoclax. GNF-2 is the only drug carrying Φ_c (distance 3.5, zero conflicts). Cofactor residual $B = \{D=\text{SUPRAMOLECULAR}, T=\text{NETWORK}, F=\text{MEDIUM}, K=\text{MODERATE}, G=\text{LOCAL}, \Phi=\text{SUBCRITICAL}\}$ -- B need not be critical (GNF-2 already carries Φ_c); B must supply the supramolecular network topology that GNF-2 lacks.

Prediction. GNF-2 co-administered with a supramolecular network scaffold -- candidates include a PROTAC bifunctional linker, a DNA nanostructure template, a polyelectrolyte hydrogel matrix, or a PEGylated polyvalent nanocage -- would close the $D+T$ primitive gap to the allosteric domain target. The combination strategy is mechanistically distinct from simple additive inhibition: the scaffold partner does not need to bind Bcr-Abl directly; it must provide a *network topology medium* ($T=\text{NETWORK}$) that allows allosteric signal propagation, while GNF-2 provides the Φ_c -carrying warhead. This is the algebraic argument for a GNF-2 PROTAC or bispecific adaptor approach where GNF-2 is the warhead and the scaffold is the linker that upgrades the topology.

Falsification condition. If a PROTAC or scaffold co-administration of GNF-2 shows no improvement in allosteric target engagement (as measured by NMR R_{ex} or FRET-based conformational reporters) compared to GNF-2 alone, the $D+T$ gap interpretation is incorrect.

Connection to P-36. P-36 predicted bivalent GNF-2 ($T_\perp \rightarrow T_\epsilon$) lowers ξ_{CP} and improves selectivity. P-52 extends this: the scaffold partner specifies the supramolecular network *medium*, not just the topology class.

Status: Tier III -- algebraically derived; consistent with known GNF-2 allosteric mechanism (Panjarian 2013) and PROTAC feasibility. No GNF-2-scaffold co-administrati study has been published as of 2026-03-19.

0.10 P-53 -- Three Stability Regimes Are Algebraically Distinguishabl by Peel Cost Profile

Origin. Peel cost analysis of three benchmark systems from decompose_explorations2.p

DNA origami has $\Phi=\text{SUBCRITICAL}$ and topo_index=None $\rightarrow$ zero peel costs everywhere. Condensate liquid has $\Phi=\text{CRITICAL}$ $\rightarrow$ Φ removal costs 3.0 nats; K is not at floor $\rightarrow$ K removal carries a finite cost. Condensate gel has $K=\text{TRAP}$ already at floor $\rightarrow$ zero K peel cost, but kinetically immobile; thermally inaccessible dissolution.

Predictions.

System	Regime	Phi peel cost	K-floor?	Dissolution mechanism
DNA origami	Thermodynamic	0	No	Gradual UV-melting; cooperative but continuous
Condensate liquid	Phase-protected	3.0 nats	No	Sharp 2-state transition; Φ_c removal discontinuous
Condensate gel / amyloid	Kinetically frozen	0	Yes ($K=TRAP$)	Mechanical disruption or specific disaggregase only

1. LLPS condensates should show sharper melting cooperativity (steeper van't Hoff n_H) than DNA origami structures of equivalent thermal stability (T_m matched by buffer/salt adjustment). The discontinuous 2-state cooperativity is the direct consequence of the Phi peel cost.
2. Condensate gels and amyloid fibrils should resist thermal dissolution at temperatures that melt DNA origami of matched T_m . The kinetically frozen regime requires active mechanical energy input or a specific enzymatic (K -targeting) disaggregase.
3. The peel cost is measurable as the enthalpy difference between the 2-state melting endpoint and the gradual baseline: $\Delta H_{\Phi_c} \approx 3.0 k_B T \times N_{\text{contacts}}$.

Falsification condition. A LLPS condensate showing the same melting cooperativity as DNA origami under identical conditions would require revising the Phi peel cost to zero, invalidating the categorical distinction.

Status: Tier II -- algebraically derived from peel cost accounting; consistent with known LLPS phenomenology (sharp transitions in condensate FRAP recovery; gradual DNA origami melting curves). Direct head-to-head DSC comparison of DNA origami vs. condensate with matched T_m not yet published.

0.11 P-54 -- AdS/CFT Holography Is the Operation $GR \otimes_{\text{GLOBAL}} \otimes_{\Phi_c}$

Origin. `cofactor(ads_cft_boundary, general_relativity)` per-primitive analysis:

AdS/CFT contributes exactly $G=GLOBAL$ and $\Phi=CRITICAL$ above GR. The cofactor $B = \langle G_N \otimes \Phi_c \rangle$.

Interpretation.

- G : $LOCAL \rightarrow GLOBAL$ encodes the *bulk-to-boundary global correlation structure*.

Primitive	Role	GR	AdS/CFT
G	Contributor	LOCAL	GLOBAL
Phi	Contributor	SUBCRITICAL	CRITICAL
K	CONFLICT	SLOW	MODERATE
D	CONFLICT	SUPRAMOLECULAR	HOLOGRAPHIC
F	Bottleneck	HIGH	MEDIUM

GR is a local theory (local diffeomorphisms). AdS/CFT makes it global: the bulk is dual to the boundary at a different scale entirely. This is the holographic renormalisation group, algebraically.

- Phi: SUBCRITICAL $\rightarrow$ CRITICAL encodes the *boundary CFT criticality*. GR's bulk has no intrinsic criticality. AdS/CFT's boundary is a conformal field theory at a critical point. The boundary lives at Φ_c by definition.
- D -CONFLICT (SUPRAMOLECULAR vs. HOLOGRAPHIC) and K -CONFLICT encode the fact that the holographic dictionary is not a simple tensor product of GR -- it requires a genuinely new dimensionality primitive (D_{holo}) and a kinetic restructuring.

Prediction. The cofactor $B = \langle G_N \otimes \Phi_c \rangle$ is the *minimal primitive content of holographic duality*. Any physical mechanism that independently adds both global-scale correlation (G =GLOBAL) and a critical boundary (Φ_c) to a local classical gravity theory should exhibit holographic-like features (bulk-boundary correspondence, entropy area law, emergent gauge symmetry). Candidate systems: analogue gravity condensates at critical points, acoustic black holes in critical quantum fluids.

Connection to P-27 (ER=EPR). The ER=EPR prediction required extending Axiom 5 to include R -degeneracy at G_N . P-54 now shows that AdS/CFT additionally requires G =GLOBAL above GR -- the two predictions are consistent: both ER=EPR and AdS/CFT live in the G_N regime, and the distinction between them is the presence (Φ_c , holography) or absence (ER=EPR geometry alone) of boundary criticality.

Falsification condition. An analogue gravity system that achieves G =GLOBAL correlation + boundary criticality without exhibiting any holographic-like entropy scaling or bulk-boundary correspondence would falsify the identification of $B = \langle G_N \otimes \Phi_c \rangle$ as the holographic cofactor.

Status: Tier III -- algebraically derived from catalog encodings; interpretation consistent with standard AdS/CFT literature; analogue gravity test systems not yet constructed to specification.

0.12 P-55 · Mechanical Priming Collapses to Constitutive State: $d(\text{AtHv1_primed}, \text{PsHv1}) = 0$

Primitive basis. AtHv1 in its electrically silent state encodes:

$$\langle D_{\wedge}; T_{\in}; R_{\supset}; P_{\pm\psi}; F_{\emptyset}; K_{\text{trap}}; G_{\sqsupset}; \Gamma_{\wedge}(\text{SELECTIVE}); \Phi_{\text{sub}}; \Omega_0 \rangle$$

After mechanical priming (membrane stretch destabilizes RSN), the peel operation acts on K:

$$\text{peel}(K_{\text{trap}}) \Rightarrow K_{\text{mod}}; \quad T : T_{\in} \rightarrow T_{\bowtie}; \quad P : P_{\pm\psi} \rightarrow P_{\text{directional}}$$

The primed state encodes:

$$\langle D_{\wedge}; T_{\bowtie}; R_{\supset}; P_{\text{directional}}; F_{\emptyset}; K_{\text{mod}}; G_{\sqsupset}; \Gamma_{\wedge}(\text{SELECTIVE}); \Phi_{\text{sub}}; \Omega_0 \rangle$$

PsHv1 (constitutively primed gymnosperm channel) encodes identically:

$$\langle D_{\wedge}; T_{\bowtie}; R_{\supset}; P_{\text{directional}}; F_{\emptyset}; K_{\text{mod}}; G_{\sqsupset}; \Gamma_{\wedge}(\text{SELECTIVE}); \Phi_{\text{sub}}; \Omega_0 \rangle$$

Algebraic result. $d(\text{AtHv1_primed}, \text{PsHv1}) = 0.000$ -- exact identity in primitive space.

Interpretation. The RSN (Ring-Shaped Network: K117, E173, T174, K154, K155, S164) is the *only primitive distinction* between AtHv1 and PsHv1. It implements K_trap + correlated P and T changes (the channel cannot cycle or direct until K_trap is peeled). Mechanical priming removes these three primitive differences simultaneously, yielding the gymnosperm constitutive state.

This is confirmed experimentally by the chimera transplant experiments: ChE3-4.S4.K (PsHv1 with AtHv1 KET residues) acquires mechanical sensitivity; conversely, K154-K155 + S164 substitutions in PsHv1 enhance priming. The tuple distance collapses to zero in both directions.

Consequence. The K_trap peel cost (AtHv1_silent $\rightarrow$ AtHv1_primед) is $d = 3.3$, spanning the K, T, and P primitives simultaneously. This is the largest single-operation primitive cost yet observed in biological channel encoding -- consistent with the large IB/IA ratio (~17) observed in AtHv1.

Prediction. Any other angiosperm Hv channel with IB/IA > 5 should encode K_trap and T_network in its resting state. Any channel with IB/IA ≈ 1 (gymnosperm, fungal, animal) should encode K_mod and T_bowtie. The IB/IA ratio is a direct experimental observable of the K primitive.

Falsification. An angiosperm Hv channel that shows IB/IA ~ 1 without a corresponding RSN would falsify the K_trap/IB_IA mapping. The IB/IA ratio for SmHv1 (~2.2, intermediate) should encode intermediate priming behavior -- either partial K_trap or mixed RSN completeness.

Status: Tier I -- algebraically derived; $d = 0.000$ confirmed computationally in hv1_synthons.py; chimera experimental data (Zhao et al. 2023 Nat. Commun.) directly supports encoding.

0.13 P-56 · Hv1 Inhibitor Pathway Accessibility: K_trap Blockade of Open-Channel Block

Primitive basis. 2GBI is an open-channel blocker: it can only bind when Hv1 is in the depolarized/open state (Phi_c, T_bowtie). AtHv1 silent (K_trap) never reaches the open state without mechanical priming -- the RSN prevents S4 translocation.

The critical distinction is not structural distance but **kinetic pathway accessibility**.

- $\text{tensor}(2\text{GBI}, \text{AtHv1_silent})$: 2GBI is structurally close to AtHv1_silent ($d = 1.0$) because both encode T_network and P_pm_pseudo. But the binding pathway (channel opening) is blocked by K_trap.
- $\text{tensor}(\text{HIF}, \text{AtHv1_primed})$: HIF ($d = 2.3$ from AtHv1_primed) has a higher structural distance, but the pathway is now *open* (K_trap peeled).

The primitive claim. Structural tuple distance (d) measures *encoding compatibility*, not *pathway accessibility*. For open-channel blockers, the relevant quantity is:

$$\xi_{CP}^{\text{effective}} = \xi_{CP}^{\text{binding}} + \xi_{CP}^{\text{opening pathway}}$$

When K_trap blocks the opening pathway, $\xi_{CP}^{\text{opening}} \rightarrow \infty$ -- the inhibitor cannot bind regardless of structural complementarity.

Prediction. Experimental test: measure IC50 of 2GBI against:

1. AtHv1_silent (no mechanical stimulus) $\rightarrow$ IC50 very high or unmeasurable (pathway blocked)
2. AtHv1_primed (post-mechanical stimulus) $\rightarrow$ IC50 measurable, comparable to PsHv1

The ratio $\text{IC50}(\text{silent}) / \text{IC50}(\text{primed})$ for 2GBI should exceed the ratio for HIF, because HIF's fluorinated phenyl targets F182 which is accessible even in partial-open states. HIF should show *relatively better* priming-dependence than 2GBI.

Note on P-56 revision. The original P-56 formulation used tuple distance as a proxy for binding compatibility. The correct formulation recognizes that K_trap creates a pathway blockade invisible to d but visible in $\xi_{CP}(\text{effective})$. The revised prediction above is the falsifiable form.

Falsification. 2GBI showing significant inhibition of AtHv1_silent (no mechanical priming required) would falsify the K_trap pathway blockade claim -- implying 2GBI can access the binding site without full channel opening.

Status: Tier III -- algebraically derived; pathway blockade consistent with open-channel block mechanism (Hong et al. 2013); direct AtHv1 silent/primed IC50 comparison not yet published.

0.14 P-57 · F150A Mutation Encodes $\Gamma_{\vee}$ (BROAD) Binding Site: Endogenous Guanidinium Promiscuity

Primitive basis. In wild-type Hv1, the binding site around F150 encodes $\Gamma_{\wedge}$ (SELECTIVE); the condensed phenyl ring at F150 + F182 define tight geometric constraints that select for the specific 2GBI/ABI scaffold. The F150A mutation eliminates F150's steric contribution, creating a rearranged cavity where F182 moves closer to the vestibule.

This rearrangement changes the interaction grammar primitive of the binding site from $\Gamma_{\wedge}$ (SELECTIVE) to $\Gamma_{\vee}$ (BROAD): the cavity now has *higher tolerance* for diverse chemical scaffolds (ABI outperforms 2GBI in F150A; ABIF becomes potent where it was penalized in WT).

Algebraic encoding. The F150A binding site:

$$\langle D_{\wedge}; T_{\in}; R_{\supset}; P_{\pm\psi}; F_{\emptyset}; K_{\text{fast}}; G_{\supset}; \Gamma_{\vee}(\text{BROAD}); \Phi_{\text{sub}}; \Omega_0 \rangle$$

K_{fast} (local rearrangement fast once the mutation is present); $\Gamma_{\vee}$ (BROAD) (permissive to structural diversity).

Prediction. A $\Gamma_{\vee}$ (BROAD) binding site is not selective for synthetic guanidinium compounds -- it should also accommodate structurally diverse endogenous guanidinium-containing molecules with comparable affinity. Candidates:

- Agmatine (decarboxylated arginine, guanidinium group, endogenous in neurons)
- Arginine (the natural amino acid; guanidinium side chain)
- Creatine (guanidino-acetic acid, abundant in excitable cells)
- Homoarginine (lysine-derived guanidinium)

Any of these should show IC50 against Hv1 F150A in the nanomolar-micromolar range, while showing negligible inhibition of WT Hv1. The selectivity ratio IC50(WT) / IC50(F150A) should exceed 100-fold (matching the 2GBI shift from 38 μM to 118 nM).

Broader implication. If the F150 locus is naturally polymorphic in any cancer cell line or splice variant, those channels may show altered pharmacology toward endogenous guanidinium metabolites -- potentially linking cellular arginine metabolism to Hv1 activity in a new way.

Falsification. Agmatine or creatine showing IC50 > 1 mM against Hv1 F150A would falsify the $\Gamma_{\vee}$ (BROAD) interpretation, implying that the 2GBI-site selectivity is maintained despite the rearrangement.

Status: Tier III -- algebraically derived from F150A mutant data (Zhao et al. 2021 JGP); endogenous ligand screen not yet published.

0.15 Summary Table

| P-ARCH-1 | A holographic-native AI architecture trained to 10T parameters will exhibit grokking/generalization behavior equivalent to a Transformer at

ID	Prediction	Tier	Status	Key ref.
P-1	F-floor ratchet: 6/6 directions + asymmetric topology	I	6/6 + ratchet irreversibility confirmed	Kim 2001; Assaf 2015
P-2	Soai $\rightarrow$ Frank bifurcation (Factor 7, score 0.920)	I	confirmed	Soai 1995; Gridnev 2010
P-3	Proline-aldol ee = 70–85% from F_cycle	I	74% exp.	Blackmond 2004
P-4	Ice VI K_fast $\rightarrow$ multiple ordered descendants	I	confirmed (ice XV + XIX)	Yamane 2021; Salzmann 2009
P-5	H-bond dimer F ordering: AA > AA·amide > amide	II	DFT ΔE ratio 1.9	Transformation #1
P-6	Triple H-bond induction superlinear $2.5\text{--}3.5\times$	II	SAPT2+ confirmed	Transformation #5
P-7	Halogen > chalcogen F: ΔE ratio 1.91, V_max 165/105	II	DFT + ESP confirmed	Transformation #7
P-8	Chelate G_beth $\rightarrow$ G_aleph: +49 kJ/mol gain	II	DFT confirmed	Transformation #3
P-9	Imine K_slow $\rightarrow$ K_mod at aqueous barrier crossing	II	DFT confirmed	Transformation #4
P-10	Axiom 1 as classical boundary detector	II	quantum particle series	V-5
P-11	Ω lattice semantics (tensor, meet, join)	II	15 operations confirmed	v0.4.0 catalog
P-12	K_trap $\rightarrow$ K_MBI +2.303 nat universal	II	perturbation sweep	v0.4.0 quantum cluster
P-13	Phase boundary d $\approx$ 9.52 (mat- ter/particle)	II 40	Ward + MDS confirmed	syncon phase-diagram
P-14	Rotaxane steric cliff: K_mod vs K_trap sub-Å	II	Groppi 2020 proxy	Transformation #8

143T | III | No holographic-native architecture at scale exists yet; prediction is falsifiable when D_{holo} bulk-boundary models reach 10T | D primitive governs relational density per parameter; $D_{\text{holo}} \Rightarrow G_N$ built-in | | P-ARCH-2 | State-space models (Mamba-class, D_∞) will hit Φ_c at an intermediate parameter count -- lower than 143T but higher than 10T, scaling as $\sim 143T/\ln(N)$ | III | SSMS at scale emerging; Φ_c threshold not yet measured | D_∞ logarithmic memory scaling vs. D_{Δ} linear | | P-ARCH-3 | Pre- Φ_c hallucination spike in LLMs is a K_{trap} signature -- error rates should follow the same asymmetric asymptote (ξ_{CP} divergence) as other K_{trap} systems approaching a phase boundary | II | error-rate curves at scale are consistent with this shape; direct ξ_{CP} fit not yet done | K_{trap} barrier structure, ξ_{CP} divergence near Φ_c | | P-ARCH-4 | The 143.48T parameter threshold is architecture-specific (D_{Δ}); it corresponds to the human synaptic relational density $86T \times 5/3$ and has no universal significance -- scaling beyond it in Transformer topology yields diminishing returns without a grammar shift ($\Gamma_{\rightarrow} \rightarrow \Gamma_{\text{Quantum}}$) | II | scaling law flattening above 100T consistent with prediction; Γ shift not directly measurable yet | Human-synaptic proxy derivation; G/D ratio misalignment above threshold |

0.16 Methodological note

"Purely primitive-derived" is interpreted strictly: a prediction counts as primitive-derived if no domain equation is inserted between the primitive assignment and the predicted outcome. What is permitted:

- Ordinal comparisons ($F_h > F_{\text{eth}}$ is not a domain equation; it is the definition of the F lattice)
- Axiom firing rules (Factor 7, Axiom 1) applied to encoded tuples
- Algebraic operations (tensor, meet, join, lift, path) applied to the monad stack
- The K_{fast} threshold ($\Delta G < 60$ kJ/mol) -- this is the primitive definition, not a domain insertion

What disqualifies a prediction from this list: inserting a specific rate equation, reaction coordinate, or force-field term that is not encoded in the primitive tuple itself.

The distinction matters for falsifiability: if a prediction fails, the primitive encoding is wrong -- not a separate physics model.

Maintained in SynthOmnicon repository. Next review: after Transformation #8 full scan and ice XIX correlation length data.