

The *E. coli* molecular phenotype under different growth conditions

Supplementary material

Mehmet U. Caglar, John R. Houser, Craig S. Barnhart,
Daniel R. Boutz, Sean M. Carroll, Aurko Dasgupta, Walter F. Lenoir,
Bartram L. Smith, Viswanadham Sridhara, Dariya K. Sydykova,
Drew Vander Wood, Christopher J. Marx,
Edward M. Marcotte*, Jeffrey E. Barrick*, Claus O. Wilke*

September 29, 2016

Contents

List of Figures

S1	Summary GO annotations	4
S2	Significantly altered KEGG pathways for mRNA samples in exponential phase tested for glycerol against glucose	5
S3	Significantly altered KEGG pathways for mRNA samples in exponential phase tested for against glucose	6
S4	Significantly altered KEGG pathways for mRNA samples in exponential phase tested for lactate against glucose	7
S5	Significantly altered KEGG pathways for protein samples in exponential phase tested for gluconate against glucose	8
S6	Significantly altered KEGG pathways for protein samples in exponential phase tested for lactate against glucose	9
S7	Significantly altered KEGG pathways for protein samples in stationary phase tested for glycerol against glucose	10
S8	Significantly altered KEGG pathways for protein samples in stationary phase tested for gluconate against glucose	11
S9	Significantly altered KEGG pathways for protein samples in stationary phase tested for lactate against glucose	12
S10	Significantly altered KEGG pathways for mRNA samples in exponential phase tested for low Mg^{+2} levels against base Mg^{+2}	13
S11	Significantly altered KEGG pathways for mRNA samples in exponential phase tested for high Mg^{+2} against base Mg^{+2}	14
S12	Significantly altered KEGG pathways for protein samples in exponential phase tested for high Mg^{+2} against base Mg^{+2}	15
S13	Significantly altered KEGG pathways for mRNA samples in stationary phase tested for high Mg^{+2} against base Mg^{+2}	16
S14	Significantly altered KEGG pathways for mRNA samples in exponential phase tested for high Na^{+1} against base Na^{+1}	17
S15	Significantly altered KEGG pathways for protein samples in exponential phase tested for high Na^{+1} against base Na^{+1}	18
S16	Significantly altered KEGG pathways for mRNA samples in stationary phase tested for high Na^{+1} against base Na^{+1}	19
S17	Significantly altered KEGG pathways for protein samples in stationary phase tested for high Na^{+1} against base Na^{+1}	20
S18	Significantly altered GO annotations associated with molecular functions for mRNA samples in exponential phase tested for glycerol against glucose	21
S19	Significantly altered GO annotations associated with molecular functions for mRNA samples in exponential phase tested for lactate against glucose	22
S20	Significantly altered GO annotations associated with molecular functions for mRNA samples in exponential phase tested for low Mg^{+2} against base Mg^{+2}	23
S21	Significantly altered GO annotations associated with molecular functions for protein samples in exponential phase tested for high Na^{+1} against base Na^{+1}	24
S22	Significantly altered GO annotations associated with molecular functions for protein samples in stationary phase tested for high Na^{+1} against base Na^{+1}	25
S23	Flux Exponential	26

List of Tables

Supplementary Figures

A	mRNA	Protein	
	1. structural constituent of ribosome 2. structural molecule activity		lowMg
			highMg
		1. structural constituent of ribosome 2. structural molecule activity	highNa
	1. structural constituent of ribosome 2. structural molecule activity		glycerol
			gluconate
B			lowMg
			highMg
		1. structural constituent of ribosome 2. structural molecule activity	highNa
			glycerol
			gluconate
			lactate

Figure S1: **Significantly differentially expressed Molecular Functions generated by GO annotations.** For each condition, we show the top-5 differentially expressed MF as determined by either mRNA or protein abundances. (A) exponential phase. (B) stationary phase.




Figure S2: **Significantly differentially expressed KEGG pathways and associated genes with glycerol as carbon source in exponential phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. In figure we show up to 10 most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S3: **Significantly differentially expressed KEGG pathways and associated genes with gluconate as carbon source in exponential phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. In figure we show up to 10 most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S4: **Significantly differentially expressed KEGG pathways and associated genes with lactate as carbon source in exponential phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. In figure we show up to 10 most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S5: **Significantly differentially expressed KEGG pathways and associated genes with gluconate as carbon source in exponential phase, as determined by protein abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. In figure we show up to 10 most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S6: **Significantly differentially expressed KEGG pathways and associated genes with lactate as carbon source in exponential phase, as determined by protein abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. In figure we show up to 10 most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S7: **Significantly differentially expressed KEGG pathways and associated genes with glycerol as carbon source in stationary phase, as determined by protein abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. In figure we show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S8: **Significantly differentially expressed KEGG pathways and associated genes with gluconate as carbon source in stationary phase, as determined by protein abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S9: **Significantly differentially expressed KEGG pathways and associated genes with lactate as carbon source in stationary phase, as determined by protein abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S10: **Significantly differentially expressed KEGG pathways and associated genes with low Mg^{+2} levels in exponential phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S11: **Significantly differentially expressed KEGG pathways and associated genes with high Mg^{+2} levels in exponential phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S12: **Significantly differentially expressed KEGG pathways and associated genes with high Mg^{+2} levels in exponential phase, as determined by protein abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S13: **Significantly differentially expressed KEGG pathways and associated genes with high Mg^{+2} levels in stationary phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S14: **Significantly differentially expressed KEGG pathways and associated genes with high Na^{+1} levels in exponential phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S15: Significantly differentially expressed KEGG pathways and associated genes with high Na^{+1} levels in exponential phase, as determined by protein abundances. The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S16: **Significantly differentially expressed KEGG pathways and associated genes with high Na^{+1} levels in stationary phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S18: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with glycerol as carbon source in exponential phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S19: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with lactate as carbon source in exponential phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S20: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with low Mg^{+2} levels in exponential phase, as determined by mRNA abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S21: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with high Na^{+1} levels in exponential phase, as determined by protein abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S22: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with high Na^{+1} levels in stationary phase, as determined by protein abundances.** The top differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S23: Flux changes with respect to salt stresses in exponential phase. 13 different flux were measured with respect to four different Na and five different Mg²⁺ concentrations. (A) Concentrations with respect to changing Na⁺ concentrations. (B) Concentrations with respect to changing Mg²⁺ concentrations. There is no significant trend of increase or decrease in exponential phase flux measurements with respect to either Mg²⁺ or Na⁺ concentrations after *fdr* correction for multiple testing. Correction. The only significant trend was for labelled CO₂ EC branch at exponential phase with respect to Mg²⁺ concentrations before multiple testing ($P=.02$), but even for this branch, after multiple testing correction corresponding *p* value increases to .4.