

The *E. coli* molecular phenotype under different growth conditions

Supplementary materials

Mehmet U. Caglar*, John R. Houser, Craig S. Barnhart,
Daniel R. Boutz, Sean M. Carroll, Aurko Dasgupta, Walter F. Lenoir,
Bartram L. Smith, Viswanadham Sridhara, Dariya K. Sydykova,
Drew Vander Wood, Christopher J. Marx,
Edward M. Marcotte*, Jeffrey E. Barrick*, Claus O. Wilke*

February 4, 2017

List of Figures

S1	Significantly differentially expressed molecular functions	4
S2	Significantly differentially expressed KEGG pathways for mRNA samples in exponential phase tested for glycerol against glucose	5
S3	Significantly differentially expressed KEGG pathways for mRNA samples in exponential phase tested for gluconate against glucose	6
S4	Significantly differentially expressed KEGG pathways for mRNA samples in exponential phase tested for lactate against glucose	7
S5	Significantly differentially expressed KEGG pathway for protein samples in exponential phase tested for gluconate against glucose	8
S6	Significantly differentially expressed KEGG pathways for protein samples in exponential phase tested for lactate against glucose	9
S7	Significantly differentially expressed KEGG pathway for protein samples in stationary phase tested for glycerol against glucose	10
S8	Significantly differentially expressed KEGG pathway for protein samples in stationary phase tested for gluconate against glucose	11
S9	Significantly differentially expressed KEGG pathways for protein samples in stationary phase tested for lactate against glucose	12
S10	Significantly differentially expressed KEGG pathways for mRNA samples in exponential phase tested for low Mg^{2+} levels against base Mg^{2+}	13
S11	Significantly differentially expressed KEGG pathways for mRNA samples in exponential phase tested for high Mg^{2+} against base Mg^{2+}	14
S12	Significantly differentially expressed KEGG pathways for protein samples in exponential phase tested for high Mg^{2+} against base Mg^{2+}	15
S13	Significantly differentially expressed KEGG pathway for mRNA samples in stationary phase tested for high Mg^{2+} against base Mg^{2+}	16
S14	Significantly differentially expressed KEGG pathway for mRNA samples in exponential phase tested for high Na^+ against base Na^+	17
S15	Significantly differentially expressed KEGG pathways for protein samples in exponential phase tested for high Na^+ against base Na^+	18
S16	Significantly differentially expressed KEGG pathways for mRNA samples in stationary phase tested for high Na^+ against base Na^+	19
S17	Significantly differentially expressed KEGG pathways for protein samples in stationary phase tested for high Na^+ against base Na^+	20
S18	Significantly differentially expressed GO annotations associated with molecular functions for mRNA samples in exponential phase tested for glycerol against glucose	21
S19	Significantly differentially expressed GO annotations associated with molecular functions for mRNA samples in exponential phase tested for gluconate against glucose	21
S20	Significantly differentially expressed GO annotations associated with molecular functions for mRNA samples in exponential phase tested for lactate against glucose	22
S21	Significantly differentially expressed GO annotations associated with molecular functions for protein samples in exponential phase tested for glycerol against glucose	22
S22	Significantly differentially expressed GO annotations associated with molecular functions for protein samples in exponential phase tested for lactate against glucose	23
S23	Significantly differentially expressed GO annotations associated with molecular functions for protein samples in stationary phase tested for glycerol against glucose	23
S24	Significantly differentially expressed GO annotations associated with molecular functions for protein samples in stationary phase tested for gluconate against glucose	24
S25	Significantly differentially expressed GO annotations associated with molecular functions for protein samples in stationary phase tested for lactate against glucose	24
S26	Significantly differentially expressed GO annotations associated with molecular functions for mRNA samples in exponential phase tested for low Mg^{2+} levels against base Mg^{2+} levels	25
S27	Significantly differentially expressed GO annotations associated with molecular functions for mRNA samples in exponential phase tested for high Mg^{2+} levels against base Mg^{2+} levels	25

S28	Significantly differentially expressed GO annotations associated with molecular functions for protein samples in exponential phase tested for low Mg^{2+} levels against base Mg^{2+} levels	26
S29	Significantly differentially expressed GO annotations associated with molecular functions for protein samples in exponential phase tested for high Mg^{2+} levels against base Mg^{2+} levels	26
S30	Significantly differentially expressed GO annotations associated with molecular functions for mRNA samples in stationary phase tested for low Mg^{2+} levels against base Mg^{2+} levels	27
S31	Significantly differentially expressed GO annotations associated with molecular functions for protein samples in exponential phase tested for high Na^+ levels against base Na^+ levels	27
S32	Significantly differentially expressed GO annotations associated with molecular functions for mRNA samples in stationary phase tested for high Na^+ levels against base Na^+ levels	28
S33	Significantly differentially expressed GO annotations associated with molecular functions for protein samples in stationary phase tested for high Na^+ levels against base Na^+ levels	28
S34	Fraction of differentially expressed genes that are found in analyses with or without controlling for doubling time	29
S35	Flux ratios versus ion concentrations	30

Supplementary Figures

A	mRNA		Protein		lowMg	
	1.structural constituent of ribosome ▼▼▼ 2.rRNA binding ▼▼▼ 3.structural molecule activity ▼▼▼ 4.RNA binding ▼▼▼ 5.RNA-dependent ATPase activity ▼▼▼		1.L-lactate dehydrogenase activity ▼▼▼ 2.lactate dehydrogenase activity ▼▼▼			
	1.energy transducer activity ▲ 2.alkanesulfonate transporter activity ▲ 3.oxidoreductase activity ▲▼		1.L-lactate dehydrogenase activity ▼▼▼ 2.lactate dehydrogenase activity ▼▼▼		highMg	
			1.binding ▼ 2.protein binding ▼ 3.structural constituent of ribosome ▼▼▼ 4.small molecule binding ▼ 5.nucleotide binding ▼		highNa	
	1.structural constituent of ribosome ▼▼▼ 2.structural molecule activity ▼▼▼ 3.rRNA binding ▼▼▼ 4.RNA binding ▼▼▼ 5.tRNA binding ▼▼▼		1.ATP binding ▼▼ 2.adenyl ribonucleotide binding ▼▼ 3.adenyl nucleotide binding ▼▼ 4.molecular transducer activity ▼▼▼ 5.oxidoreductase activity, acting on the CH-OH group of donors, quinon		glycerol	
	1.gluconate transmembrane transporter activity ▲ 2.aldonate transmembrane transporter activity ▲				gluconate	
	1.structural constituent of ribosome ▲▲ 2.structural molecule activity ▲▲ 3.rRNA binding ▲▲ 4.RNA binding ▲▲ 5.lactate dehydrogenase activity ▲▲		1.catalytic activity ▲▲ 2.lactate dehydrogenase activity ▲▲ 3.hydrolase activity ▲▲ 4.L-lactate dehydrogenase activity ▲▲		lactate	
	B	mRNA		Protein		lowMg
		1.ATPase activity ▼▼▼ 2.hydrolase activity, acting on acid anhydrides, catalyzing transmembra 3.oligopeptide-transporting ATPase activity ▼▼▼ 4.peptide-transporting ATPase activity ▼▼▼ 5.ATPase activity, coupled ▼▼▼				
						highMg
1.carbohydrate transmembrane transporter activity ▼▼ 2.carbohydrate transporter activity ▼▼ 3.polyol transmembrane transporter activity ▼▼ 4.alcohol transmembrane transporter activity ▼▼ 5.organic hydroxy compound transmembrane transporter activity ▼▼		1.protein binding ▼ 2.binding ▼ 3.structural constituent of ribosome ▼▼▼ 4.ion binding ▼ 5.structural molecule activity ▼▼▼		highNa		
		1.oxidoreductase activity ▼▼ 2.oxidoreductase activity, acting on the aldehyde or oxo group of donors 3.coenzyme binding ▼ 4.oxidoreductase activity, acting on the CH-OH group of donors, quinon 5.cofactor binding ▼		glycerol		
		1.siderophore transmembrane transporter activity ▼▼▼ 2.2,3-dihydroxybenzoate-serine ligase activity ▼▼▼ 3.siderophore transporter activity ▼▼▼ 4.iron chelate transmembrane transporter activity ▼▼▼		gluconate		
		1.lactate dehydrogenase activity ▲▲ 2.oxidoreductase activity, acting on CH-OH group of donors ▲▲ 3.binding ▲▲ 4.L-lactate dehydrogenase activity ▲▲ 5.cofactor binding ▲▲		lactate		

Figure S1: **Significantly differentially expressed molecular functions, as determined by GO annotations.** For each condition, we show the top-5 differentially expressed molecular functions according to either mRNA or protein abundances. Empty boxes indicate that no differentially expressed pathways were found. The arrows next to pathway names indicate the proportion of up- and down-regulated genes among the significantly differentially expressed genes in this pathway. One up arrow indicates that 60% or more of the genes are up-regulated, two arrows correspond to 80% or more genes, and three arrows correspond to 95% or more genes being up-regulated. Similarly, down arrows indicate the proportion of down-regulated genes. (A) Exponential phase. (B) Stationary phase.




Figure S2: **Significantly differentially expressed KEGG pathways and associated genes with glycerol as carbon source, as determined by mRNA abundances in exponential phase.** The top 2 differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S3: **Significantly differentially expressed KEGG pathway and associated genes with gluconate as carbon source, as determined by mRNA abundances in exponential phase.** The top differentially expressed KEGG pathway is shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S4: **Significantly differentially expressed KEGG pathways and associated genes with lactate as carbon source, as determined by mRNA abundances in exponential phase.** The top 3 differentially expressed KEGG pathways are shown along the *y* axis, and the relative fold change of the corresponding genes is shown along the *x* axis. We show up to 10 of the most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S5: **Significantly differentially expressed KEGG pathway and associated genes with gluconate as carbon source, as determined by protein abundances in exponential phase.** The top differentially expressed KEGG pathway is shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S6: **Significantly differentially expressed KEGG pathways and associated genes with lactate as carbon source, as determined by protein abundances in exponential phase.** The top 3 differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway we show up to 15 of the most significantly changing genes.




Figure S7: **Significantly differentially expressed KEGG pathway and associated genes with glycerol as carbon source, as determined by protein abundances in stationary phase.** The top differentially expressed KEGG pathway is shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S8: **Significantly differentially expressed KEGG pathway and associated genes with gluconate as carbon source, as determined by protein abundances in stationary phase.** The top differentially expressed KEGG pathway is shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S9: **Significantly differentially expressed KEGG pathways and associated genes with lactate as carbon source, as determined by protein abundances in stationary phase.** The top 3 differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S10: **Significantly differentially expressed KEGG pathways and associated genes with low Mg^{2+} levels, as determined by mRNA abundances in exponential phase.** The top 2 differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S11: **Significantly differentially expressed KEGG pathways and associated genes with high Mg^{2+} levels, as determined by mRNA abundances in exponential phase.** The top 3 differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S12: **Significantly differentially expressed KEGG pathways and associated genes with high Mg^{2+} levels, as determined by protein abundances in exponential phase.** The top 3 differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S13: **Significantly differentially expressed KEGG pathway and associated genes with high Mg^{2+} levels, as determined by mRNA abundances in stationary phase.** The top differentially expressed KEGG pathway is shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S14: **Significantly differentially expressed KEGG pathway and associated genes with high Na^+ levels, as determined by mRNA abundances in exponential phase.** The top differentially expressed KEGG pathway is shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.




Figure S15: **Significantly differentially expressed KEGG pathways and associated genes with high Na^+ levels, as determined by protein abundances in exponential phase.** The top 10 differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.





Figure S16: **Significantly differentially expressed KEGG pathways and associated genes with high Na^+ levels, as determined by mRNA abundances in stationary phase.** The top 4 differentially expressed KEGG pathways are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed pathways and for each pathway, we show up to 15 of the most significantly changing genes.






Figure S20: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with lactate as carbon source, as determined by mRNA abundances in exponential phase.** The top 5 differentially expressed molecular functions are shown along the *y* axis, and the relative fold change of the corresponding genes is shown along the *x* axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S21: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with glycerol as carbon source, as determined by protein abundances in exponential phase.** The top 10 differentially expressed molecular functions are shown along the *y* axis, and the relative fold change of the corresponding genes is shown along the *x* axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S24: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with gluconate as carbon source, as determined by protein abundances in stationary phase.** The top 4 differentially expressed molecular functions are shown along the *y* axis, and the relative fold change of the corresponding genes is shown along the *x* axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S25: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with lactate as carbon source, as determined by protein abundances in stationary phase.** The top 5 differentially expressed molecular functions are shown along the *y* axis, and the relative fold change of the corresponding genes is shown along the *x* axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S26: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with low Mg^{2+} levels, as determined by mRNA abundances in exponential phase.** The top 10 differentially expressed molecular functions are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S27: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with high Mg^{2+} levels, as determined by mRNA abundances in exponential phase.** The top 3 differentially expressed molecular functions are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S28: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with low Mg^{2+} levels, as determined by protein abundances in exponential phase.** The top 2 differentially expressed molecular functions are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S29: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with high Mg^{2+} levels, as determined by protein abundances in exponential phase.** The top 2 differentially expressed molecular functions are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S30: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with low Mg^{2+} levels, as determined by mRNA abundances in stationary phase.** The top 10 differentially expressed molecular functions are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S31: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with high Na^{+} levels, as determined by protein abundances in exponential phase.** The top differentially expressed molecular functions are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S32: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with high Na^+ levels, as determined by mRNA abundances in stationary phase.** The top 5 differentially expressed molecular functions are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S33: **Significantly differentially expressed GO annotations related with molecular functions and associated genes with high Na^+ levels, as determined by protein abundances in stationary phase.** The top 10 differentially expressed molecular functions are shown along the y axis, and the relative fold change of the corresponding genes is shown along the x axis. We show up to 10 of the most significantly changed molecular functions and for each molecular function, we show up to 15 of the most significantly changing genes.




Figure S34: **Fraction of differentially expressed genes that are found in analyses with or without controlling for doubling time.** Shown are the fractions of genes identified as differentially expressed only when controlling for doubling time (red), only when not controlling for doubling time (blue), or in both cases (green). Combined with the absolute numbers of differentially expressed genes in the various conditions (Figure 5), we can see that the main differences in analyses with or without doubling time arise for protein abundances analyzed with respect to different carbon sources.




Figure S35: **Flux ratios versus ion concentrations.** 13 different flux ratios were measured with respect to four different Na⁺ and five different Mg²⁺ concentrations. (A) Concentrations with respect to changing Na⁺ concentrations. (B) Concentrations with respect to changing Mg²⁺ concentrations. There was no significant trend of increase or decrease in flux ratios with respect to either Na⁺ or Mg²⁺ concentrations (Supplementary Table 13).

List of Supplementary Tables

Supplementary Table S1: Meta information for each sample. Includes information about sample numbers, experiment, the growth time at which the sample was collected, harvest date, number of RNA samples (technical replicates), number of protein samples (technical replicates), batch number, Mg^{2+} and Na^+ concentrations, growth phase, doubling time (mean, $\pm 95\%$ confidence interval, r^2 from the linear fit to OD600 values).

File name: tableS1_meta_data.csv

Supplementary Table S2: Normalized mRNA counts. Includes data for 4197 distinct proteins each for 152 samples.

File name: tableS2_mRNA_normalized_raw_data.csv

Supplementary Table S3: Normalized protein counts. Includes data for 4197 distinct proteins each for 105 samples.

File name: tableS3_protein_normalized_raw_data.csv

Supplementary Table S4: Mean flux ratios for 13 branches each, measured for varying Mg^{2+} and Na^+ concentrations in exponential and stationary phase.

File name: tableS4_fluxData.csv

Supplementary Table S5: Doubling time measurements in exponential phase. Includes the mean, $\pm 95\%$ confidence interval, and r^2 from the linear fit to OD600 values.

File name: tableS5_doubling_times.csv

Supplementary Table S6: z -scores obtained from tests for significant clustering of mRNA counts.

File name: tableS6_clustering_mrna_cophenetic.csv

Supplementary Table S7: z -scores obtained from tests for significant clustering of protein counts.

File name: tableS7_clustering_protein_cophenetic.csv

Supplementary Table S8: Combined results from tests for differential expression for all genes and all distinct tests considered.

The table contains the following information:

- Gene id (ECB number for mRNA and YP number for proteins), and corresponding gene name
- Results from DeSeq2 calculation, including base mean value, log2FoldChange, ifcSE, stat, pvalue, padj.
- Direction of change relative to base level ("1" for increase, "-1" for decrease)
- Data type (mRNA or protein)
- Growth phase
- What is tested; base value and contrast.
- Individual output file name

- Carbon Source, Mg^{2+} and Na^+ levels and growth phase of test data
- Control parameters of the test (batch only or batch plus doubling time)

File name: tableS8_combinedOutputDF_DeSeq.csv

Supplementary Table S9: Filtered version of Supplementary Table S8, retaining only genes with $P < 0.05$ and $\log_2\text{FoldChange} > 2$.

File name: tableS9_combinedDifferentiallyExpressedGenes_DeSeq.csv

Supplementary Table S10: Complete results from DAVID enrichment analysis for KEGG pathways and molecular functions.

File name: tableS10_combinedResultList_DAVID.csv

Supplementary Table S11: List of the additional proteins identified as differentially expressed when controlling for doubling time, tested for different carbon sources. Results are provided for both exponential and stationary phases.

File name: tableS11_changed_protein_carbonSource_ExpSta.csv

Supplementary Table S12: Enriched KEGG pathways and molecular functions based on the genes listed in Supplementary Table S11.

File name: tableS12_changed_DAVID_P05.csv

Supplementary Table S13: Results from linear regressions of flux ratios against ion concentrations (Mg^{2+} and Na^+).

File name: tableS13_flux_vs_conc_Pvalues.csv

Supplementary Table S14: Results from linear regressions of flux ratios against doubling times (Mg^{2+} and Na^+).

File name: tableS14_flux_vs_doublingTime_Pvalues_tog.csv