

# Studies of ERM Models with Correlated Disorder

by Tom Folgmann

Bachelor Thesis Presentation, 2024

# Upfront: Goals.

# Upfront: Goals.

## Foundations

# Upfront: Goals.

## Foundations

- What is ERM?

# Upfront: Goals.

## Foundations

- What is ERM?
  - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?

# Upfront: Goals.

## Foundations

- What is ERM?
  - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
  - Where lies our main theoretical interest?

# Upfront: Goals.

## Foundations

- What is ERM?
  - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
  - Where lies our main theoretical interest?
  - What does randomness mean for the model?

# Upfront: Goals.

## Foundations

- What is ERM?
  - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
  - Where lies our main theoretical interest?
  - What does randomness mean for the model?
- Understanding the action.



# Upfront: Goals.

## Foundations

- What is ERM?
  - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
  - Where lies our main theoretical interest?
  - What does randomness mean for the model?
- Understanding the action.
  - How does it arise?

# Upfront: Goals.

## Foundations

- What is ERM?
  - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
  - Where lies our main theoretical interest?
  - What does randomness mean for the model?
- Understanding the action.
  - How does it arise?
  - How can a Gaussian approach help us out?

# Upfront: Goals.

## Foundations

- What does *generating* with respect to Boltzmann densities mean?

# Upfront: Goals.

## Foundations

- What does *generating* with respect to Boltzmann densities mean?
  - Is there a visual approach to calculations?

# Upfront: Goals.

## Steps forward

# Upfront: Goals.

## Steps forward

- Now, where comes the correlative nature into play?

# Upfront: Goals.

## Steps forward

- Now, where comes the correlative nature into play?
- What did we implement in our research?

# Upfront: Goals.

## Steps forward

- Now, where comes the correlative nature into play?
- What did we implement in our research? (Has this been done before?)



# Upfront: Goals.

## Steps forward

- Now, where comes the correlative nature into play?
- What did we implement in our research? (Has this been done before?)
- What did we **conclude**?

What is ERM?

What is ERM?

Imagine a system of  $N \in \mathbb{N}$  particles.

What is ERM?


Imagine a system of  $N \in \mathbb{N}$  particles.

→ How would you describe their relations?

What is ERM?

Imagine a system of  $N \in \mathbb{N}$  particles.


→ How would you describe their relations?



What is ERM?

Imagine a system of  $N \in \mathbb{N}$  particles.

→ How would you describe their relations?



A system with  $N \in \mathbb{N}$  (related) particles can be described by a mathematical *Graph*.

What is ERM?

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix  $L$* .

What is ERM?

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix*  $L$ .

$$L(G) := D(G) - W(G).$$



Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix*  $L$ .

$$L(G) := D(G) - W(G).$$

- $D(G)$  gives the *degree* of each node: *Number of connected edges*.
-

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix*  $L$ .

$$L(G) := D(G) - W(G).$$

- $D(G)$  gives the *degree* of each node: *Number of connected edges*.
- $W(G)$  encodes the *strength* (and direction) of the connections.

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix*  $L$ .

$$L(G) := D(G) - W(G).$$

- $D(G)$  gives the *degree* of each node: *Number of connected edges*.
- $W(G)$  encodes the *strength* (and direction) of the connections.

$$D(G) := \text{diag}(d), \quad d_i := \#\{e \in E : v_i \in e\},$$

$$W(G) := (w_{ij})_{(i,j) \in [N]^2}, \quad w : [N]^2 \rightarrow \mathbb{R}.$$

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix*  $L$ .

$$L(G) := D(G) - W(G).$$

- $D(G)$  gives the *degree* of each node: *Number of connected edges*.
- $W(G)$  encodes the *strength* (and direction) of the connections.

$$D(G) := \text{diag}(d), \quad d_i := \#\{e \in E : v_i \in e\},$$

$$W(G) := (w_{ij})_{(i,j) \in [N]^2}, \quad w : [N]^2 \rightarrow \mathbb{R}.$$

.. special case is the *Adjacency Matrix*  $A$ , where  $w_{i,j} \in \{0, -1, 1\}$ .

# Definition of the ERM Laplacian Matrix

## Definition of the ERM Laplacian Matrix

In the ERM model the Laplacian matrix is defined as:

$$\tilde{U}(f, r) := \begin{pmatrix} \Sigma(f, 1) & -f_{12} & \dots & -f_{1N} \\ -f_{21} & \Sigma(f, 2) & \dots & -f_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -f_{N1} & -f_{N2} & \dots & \Sigma(f, N) \end{pmatrix} = D(G) - W(G).$$

## Definition of the ERM Laplacian Matrix

In the ERM model the Laplacian matrix is defined as:

$$\tilde{U}(f, r) := \begin{pmatrix} \Sigma(f, 1) & -f_{12} & \dots & -f_{1N} \\ -f_{21} & \Sigma(f, 2) & \dots & -f_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -f_{N1} & -f_{N2} & \dots & \Sigma(f, N) \end{pmatrix} = D(G) - W(G).$$

- *Interaction strength* given by  $f_{ij} \stackrel{\text{m}}{=} f(r_i - r_j)$

## Definition of the ERM Laplacian Matrix

In the ERM model the Laplacian matrix is defined as:

$$\tilde{U}(f, r) := \begin{pmatrix} \Sigma(f, 1) & -f_{12} & \dots & -f_{1N} \\ -f_{21} & \Sigma(f, 2) & \dots & -f_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -f_{N1} & -f_{N2} & \dots & \Sigma(f, N) \end{pmatrix} = D(G) - W(G).$$

- *Interaction strength* given by  $f_{ij} \stackrel{\text{m}}{=} f(r_i - r_j)$
- *Self-interaction* given by  $\Sigma(f, i) \stackrel{\text{m}}{=} \sum_{j \in [N] \setminus \{i\}} f_{ij}$




## How to measure Eigenvalues?

Let  $\Lambda : [n] \rightarrow \sigma_P(\tilde{U}(f, r))$  map bijectively into the *point spectrum* of the ERM Laplacian.


# How to measure Eigenvalues?

Let  $\Lambda : [n] \rightarrow \sigma_P(\tilde{U}(f, r))$  map bijectively into the *point spectrum* of the ERM Laplacian.




# How to measure Eigenvalues?

Let  $\Lambda : [n] \rightarrow \sigma_P(\tilde{U}(f, r))$  map bijectively into the *point spectrum* of the ERM Laplacian.




# How to measure Eigenvalues?

Let  $\Lambda : [n] \rightarrow \sigma_P(\tilde{U}(f, r))$  map bijectively into the *point spectrum* of the ERM Laplacian.




# How to measure Eigenvalues?

Let  $\Lambda : [p] \rightarrow \sigma_P(\tilde{U}(f, r))$  map bijectively into the *point spectrum* of the ERM Laplacian.



## How to measure Eigenvalues?

Let  $\Lambda : [p] \rightarrow \sigma_P(\tilde{U}(f, r))$  map bijectively into the *point spectrum* of the ERM Laplacian.



... results in an (unnormalized) density function


$$E \mapsto \sum_{i \in [n]} \delta_{\Lambda_i}(E) \quad \in \{0, \dots, n\}$$

# The Resolvent Eigenvalue Approximation

.. by an example point  $\Lambda_i$  at  $i \in [n]$ .

# The Resolvent Eigenvalue Approximation


.. by an example point  $\Lambda_i$  at  $i \in [n]$ .





# The Resolvent Eigenvalue Approximation

.. by an example point  $\Lambda_i$  at  $i \in [n]$ .



↪ Usecase is the resolvent with a singularity at  $\Lambda_i$ .

What is ERM?

.. the connection can be found in *Newtonian Physics*:

What is ERM?

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$  map of a particle's position

What is ERM?

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$  map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

What is ERM?

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$  map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

$$\left(\frac{d}{dt}\right)^2 x_i(t) = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot x_j(t), \quad i, j \in [N].$$

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$  map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

$$\left(\frac{d}{dt}\right)^2 x_i(t) = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot x_j(t), \quad i, j \in [N].$$

.. looking at the behaviour with regard to the initial conditions:

$$\left(\frac{d}{dt}\right)^2 \langle x_i(t), x_i(0) \rangle = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot \langle x_j(t), x_i(0) \rangle.$$

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$  map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

$$\left(\frac{d}{dt}\right)^2 x_i(t) = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot x_j(t), \quad i, j \in [N].$$


.. looking at the behaviour with regard to the initial conditions:

$$\left(\frac{d}{dt}\right)^2 \langle x_i(t), x_i(0) \rangle = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot \langle x_j(t), x_i(0) \rangle.$$

This is just  $f_{i,j}$ !  
.. for  $i \neq j$ .


What is ERM?

In a visual approach  $\langle x_i(t), x_i(0) \rangle$  represents:






In a visual approach  $\langle x_i(t), x_i(0) \rangle$  represents:



Implementation of two initial configurations:


In a visual approach  $\langle x_i(t), x_i(0) \rangle$  represents:



Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle = \delta_{ij}$


In a visual approach  $\langle x_i(t), x_i(0) \rangle$  represents:



Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle = \delta_{ij}$
- $\left(\frac{d}{dt}\right) \langle x_i(t), x_j(0) \rangle = 0$

In a visual approach  $\langle x_i(t), x_i(0) \rangle$  represents:




Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle = \delta_{ij}$
- $\left(\frac{d}{dt}\right) \langle x_i(t), x_j(0) \rangle = 0$

.. leads to the *Resolvent representation* (Green's

function): of the ERM  
Laplacian:<sup>5</sup>

In a visual approach  $\langle x_i(t), x_i(0) \rangle$  represents:



Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle = \delta_{ij}$
- $\left(\frac{d}{dt}\right) \langle x_i(t), x_j(0) \rangle = 0$

.. leads to the *Resolvent representation* (Green's

function): of the ERM  
Laplacian:<sup>6</sup>

$$(\mathcal{L}F_{j,i})(s) = \pm \frac{1}{\tilde{U}(f, x^*(t))_{i,j} - \delta_{ij} \cdot \lambda_i^2}.$$

---

<sup>6</sup>With  $x^*(t) = (i \mapsto x_{i(t)})$  and  $F_{j,i}(t) := \langle x_j(t), x_i(0) \rangle$ .

# What does this mean for Eigenvalue measurement?

---

<sup>7</sup>A direct connection can be obtained, see Thesis p. 17.

# What does this mean for Eigenvalue measurement?

Using a physical argument, complex analysis can be utilized for measurement.<sup>8</sup> This leads to the *resolvent*:

# What does this mean for Eigenvalue measurement?

Using a physical argument, complex analysis can be utilized for measurement.<sup>9</sup> This leads to the *resolvent*:



## What does this mean for Eigenvalue measurement?

Using a physical argument, complex analysis can be utilized for measurement.<sup>10</sup> This leads to the *resolvent*:

$$G_{N(\mathbf{p}, z)} = \sum_{(i,j)} \int \pm \frac{1}{\tilde{U}(f, r)_{i,j} - \delta_{ij} \cdot z} \cdot e^{i \cdot \mathbf{p} \cdot (x_i - x_j)} dx.$$

---

<sup>10</sup>A direct connection can be obtained, see Thesis p. 17.

## What does this mean for Eigenvalue measurement?

Using a physical argument, complex analysis can be utilized for measurement.<sup>11</sup> This leads to the *resolvent*:

$$G_{N(\mathbf{p}, z)} = \sum_{(i,j)} \int \pm \frac{1}{\tilde{U}(f, r)_{i,j} - \delta_{ij} \cdot z} \cdot e^{i \cdot \mathbf{p} \cdot (x_i - x_j)} dx.$$

Q: What are we integration over?

---

<sup>11</sup>A direct connection can be obtained, see Thesis p. 17.

What is ERM?

# Where does *Randomness* come into play?

# Where does *Randomness* come into play?

.. by a *slight* modification of functions!

## Where does *Randomness* come into play?

.. by a *slight* modification of functions!

$\Omega \ni \omega \mapsto R(\omega)$ , equivalent to  $x$  from before.

## Where does *Randomness* come into play?

.. by a *slight* modification of functions!

$\Omega \ni \omega \mapsto R(\omega)$ , equivalent to  $x$  from before.

Ev. step	Meaning
$R$	Random variable, abstract
$R(\omega)$	Vector of time dep. pos.
$R(\omega)_i$	$i$ -th particle position, time dep. path
$R(\omega)_i(t)$	Position of $i$ -th particle at time $t$ (fixed for us.)

# The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

# The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} = \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \left( e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, r) - z) \cdot \varphi, \varphi \rangle} \cdot \lambda \right) (d\varphi).$$



# The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} = \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \left( \underbrace{e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, r) - z) \cdot \varphi, \varphi \rangle}}_{\text{Boltzmann density}} \cdot \lambda \right) (d\varphi).$$

# The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} = \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \left( \overbrace{e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, r) - z) \cdot \varphi, \varphi \rangle}}^{\text{Gaussian measure}} \cdot \lambda \right)_{\text{Boltzmann density}} (d\varphi).$$

# The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} = \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \underbrace{\left( \overbrace{e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, r) - z) \cdot \varphi, \varphi \rangle} \cdot \lambda}^{\text{Gaussian measure}} \right)}_{\text{Boltzmann density}} (d\varphi).$$

Value in  $\mathbb{R}$  ..  
(simplification!)

# The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} = \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \underbrace{\left( \overbrace{e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, r) - z) \cdot \varphi, \varphi \rangle}}^{\text{Gaussian measure}} \cdot \lambda \right)}_{\text{Boltzmann density}} (d\varphi).$$

Value in  $\mathbb{R} \dots$   
(simplification!)

This is already a good starting point to understand our *Correlated Disorder* modification!

What is ERM?

.. missing key elements:

---

<sup>12</sup>Expansion to a functional can be argued, see thesis p. 19.

What is ERM?

.. missing key elements:

- The *action* (functional)  $S_{z,R_\omega}$  at a *test point*  $z \in \mathbb{C}$  and a particle position vector  $R_\omega$ .

.. missing key elements:

- The *action* (functional)  $S_{z,R_\omega}$  at a *test point*  $z \in \mathbb{C}$  and a particle position vector  $R_\omega$ .

↪ see the Boltzmann density exponent for implicit def.:<sup>14</sup>

.. missing key elements:

- The *action* (functional)  $S_{z,R_\omega}$  at a *test point*  $z \in \mathbb{C}$  and a particle position vector  $R_\omega$ .

↪ see the Boltzmann density exponent for implicit def.:<sup>15</sup>

$$-\frac{\beta}{2} \cdot S_{z,R_\omega}(\varphi) := -\frac{\beta}{2} \cdot \left\langle \left( \tilde{U}(f, r) - z \right) \cdot \varphi, \varphi \right\rangle$$

---

<sup>15</sup>Expansion to a functional can be argued, see thesis p. 19.



.. missing key elements:

- The *action* (functional)  $S_{z,R_\omega}$  at a *test point*  $z \in \mathbb{C}$  and a particle position vector  $R_\omega$ .

↪ see the Boltzmann density exponent for implicit def.:<sup>16</sup>

$$-\frac{\beta}{2} \cdot S_{z,R_\omega}(\varphi) := -\frac{\beta}{2} \cdot \langle (\tilde{U}(f, r) - z) \cdot \varphi, \varphi \rangle$$

- The *moment generating function*  $Z_{z,R_\omega}[J]$ . It requires the *force field*  $J$ .

---

<sup>16</sup>Expansion to a functional can be argued, see thesis p. 19.

**Definition 2.22.** External Field Shift.

For  $R : \Omega \rightarrow V_{d,N}$  and  $\Phi \in \mathbb{F}_{d,N}$  we define

$$J \mapsto -\frac{1}{2} \cdot S_{z,R_\omega}^{(0)}(\Phi) + \int_{\mathbb{R}^d} J(x) \cdot \Phi(-x) + J(-x) \cdot \Phi(x) \, dx$$

the *field shifted action*  $S_{z,R_\omega}^{(0)}$  by an external field  $J \in \mathcal{S}(\mathbb{R}^d)$ .

**Definition 2.22.** External Field Shift.

For  $R : \Omega \rightarrow V_{d,N}$  and  $\Phi \in \mathbb{F}_{d,N}$  we define

$$J \mapsto -\frac{1}{2} \cdot S_{z,R_\omega}^{(0)}(\Phi) + \int_{\mathbb{R}^d} J(x) \cdot \Phi(-x) + J(-x) \cdot \Phi(x) dx$$

the *field shifted action*  $S_{z,R_\omega}^{(0)}$  by an external field  $J \in \mathcal{S}(\mathbb{R}^d)$ .

$$\hookrightarrow \frac{\delta}{\delta J(x)} S_{z,R_\omega}^{(0)}[\Phi] = \dot{i} \cdot \Phi(-x).$$

# The Generative Operator

Using these tools, an Operator generating  $Z_{z,R_\omega}$  can be deduced:

# The Generative Operator

Using these tools, an Operator generating  $Z_{z,R_\omega}$  can be deduced:

$$Z_{z,R_\omega}[J] = \int_{\mathbb{F}_{d,N}} e\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J] d\Phi$$

# The Generative Operator

Using these tools, an Operator generating  $Z_{z,R_\omega}$  can be deduced:

$$\begin{aligned} Z_{z,R_\omega}[J] &= \int_{\mathbb{F}_{d,N}} e\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J] d\Phi \\ &= \underbrace{\left[ \mathbf{Ex}_{\mathcal{L}_f} \left[ \int_{\mathbb{F}_{d,N}} e\left(S_{z,R_\omega}^{(0)}\Phi\right)[\cdot] d\Phi \right] \right]}_{\text{Generative Part}} [J]. \end{aligned}$$

# The Generative Operator

Using these tools, an Operator generating  $Z_{z,R_\omega}$  can be deduced:

$$\begin{aligned}
 Z_{z,R_\omega}[J] &= \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J]} d\Phi \\
 &= \underbrace{\left[ \mathbf{Ex}_{\mathcal{L}_f} \left[ \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi\right)[\cdot]} d\Phi \right] \right]}_{\text{Generative Part}} [J].
 \end{aligned}$$

Only free theory!

# The Generative Operator

Using these tools, an Operator generating  $Z_{z,R_\omega}$  can be deduced:

$$\begin{aligned}
 Z_{z,R_\omega}[J] &= \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J]} d\Phi \\
 &= \underbrace{\left[ \mathbf{Ex}_{\mathcal{L}_f} \left[ \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi\right)[\cdot]} d\Phi \right] \right]}_{\text{Generative Part}} [J].
 \end{aligned}$$

Only free theory!

This needs explanation.



What is ERM?

.. “Ex” of course is an abbreviation.

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include?

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left( \frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) (\lambda \otimes \widehat{\delta \rho_{R_\omega}})(d\mathbf{p})$$

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left( \frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left( \lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p})$$

$\uparrow$   
 $\varphi \rightarrow \Phi$

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left( \frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left( \lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p})$$

$\rho_R \rightarrow \lambda + \delta \rho_R$   
 $\varphi \rightarrow \Phi$

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left( \frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left( \lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p})$$

for  $\Phi(\mathbf{p}_1) \xrightarrow{\quad} \frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)}$        $\frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \xrightarrow{\quad} \widehat{\delta \rho_{R_\omega}}$

$\rho_R \rightarrow \lambda + \delta \varphi_R$   
 $\varphi \rightarrow \Phi$

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left( \frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left( \lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p})$$

for  $\Phi(\mathbf{p}_1)$  ————— ↑
for  $\Phi(-\mathbf{p}_1 - \mathbf{p}_2)$  ————— ↑

$\rho_R \rightarrow \lambda + \delta \rho_R$   
 $\varphi \rightarrow \Phi$



What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\begin{array}{c}
 \int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left( \frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left( \lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p}) \\
 \begin{array}{ccc}
 \text{for } \Phi(\mathbf{p}_1) \xrightarrow{\quad} \uparrow & & \uparrow \xrightarrow{\quad} \text{for } \Phi(-\mathbf{p}_1 - \mathbf{p}_2) \\
 & & \uparrow \xrightarrow{\quad} \varphi \rightarrow \Phi
 \end{array}
 \end{array}$$

$\rho_R \rightarrow \lambda + \delta \rho_R$   
 $\varphi \rightarrow \Phi$

$$\Phi \mapsto \left( \int_{(\mathbb{R}^d)^2} \Phi(\mathbf{p}_1) \cdot \Phi(-\mathbf{p}_1 - \mathbf{p}_2) \varphi_j \cdot \mu_z(\mathbf{p}_1, \mathbf{p}_2) d\mathbf{p} \right) \cdot e^{(S_{z, R_\omega}^{(0)} \Phi)[J]}.$$

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left( \frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left( \lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p})$$

$\text{for } \Phi(\mathbf{p}_1) \xrightarrow{\quad} \uparrow$ 
 $\text{for } \Phi(-\mathbf{p}_1 - \mathbf{p}_2) \xrightarrow{\quad} \uparrow$

$\rho_R \rightarrow \lambda + \delta \rho_R$   
 $\varphi \rightarrow \Phi$

$$\Phi \mapsto \left( \int_{(\mathbb{R}^d)^2} \Phi(\mathbf{p}_1) \cdot \Phi(-\mathbf{p}_1 - \mathbf{p}_2) \varphi_j \cdot \mu_z(\mathbf{p}_1, \mathbf{p}_2) d\mathbf{p} \right) \cdot e^{(S_{z, R_\omega}^{(0)} \Phi)[J]}.$$

→ Looking at different Taylor expansion terms of  $\exp\left(\int_{(\mathbb{R}^d)^2} \dots d\mathbf{p}\right)$  yields different powers of integral operators.

# Feynman Diagrammatics - Edges

.. conveniently using symmetry in Fourier space:

# Feynman Diagrammatics - Edges

.. conveniently using symmetry in Fourier space:

$$\begin{aligned} \longrightarrow &:= \frac{G_0(\mathbf{p}, z)}{\rho_*} \\ \overset{\gamma}{\sim} &:= \frac{\mathbb{E}((\mathcal{F} \delta \rho_R)(\mathbf{q}) \cdot (\mathcal{F} \delta \rho_R)(-\mathbf{q}))}{\rho_*} \end{aligned}$$

# Feynman Diagrammatics - Edges

.. conveniently using symmetry in Fourier space:


$$\begin{aligned} \longrightarrow &:= \frac{G_0(\mathbf{p}, z)}{\rho_*} \\ \gamma \\ \text{~~~~~} &:= \frac{\mathbb{E}((\mathcal{F} \delta \rho_R)(\mathbf{q}) \cdot (\mathcal{F} \delta \rho_R)(-\mathbf{q}))}{\rho_*} \end{aligned}$$

.. possible connections of these edges are given by *vertices*:

What is ERM?


# Feynman Diagrammatics - Vertices

# Feynman Diagrammatics - Vertices



$$:= \mu_z(\mathbf{p}, \mathbf{q})$$


$\Rightarrow$  Three-point Vertex




$$:= -V(\mathbf{p}, -\mathbf{q}_1)$$

$\Rightarrow$  Four-point Vertex

# Feynman Diagrammatics - Vertices


$$:= \mu_z(\mathbf{p}, \mathbf{q}) \quad \Rightarrow \quad \text{Three-point Vertex}$$


$$:= -V(\mathbf{p}, -\mathbf{q}_1) \quad \Rightarrow \quad \text{Four-point Vertex}$$

.. which completes the set of Feynman rules.




# How can we use diagrammatics?

.. displaying summands in operator expansion.

# How can we use diagrammatics?

.. displaying summands in operator expansion.


Observe **one** loop diagrams:



## How can we use diagrammatics?

.. displaying summands in operator expansion.

Observe **one** loop diagrams:



.. represented diagrams are *irreducible*:  $Z_{z,R_w}[J] \propto \exp\left(\sum_{C \in \mathcal{C}} C\right)$ .

# Integral representations<sup>17</sup>

---

<sup>17</sup>Attention! The terms have been simplified. For more details see Thesis sec. 2.4.2.

# Integral representations<sup>18</sup>

$$\begin{aligned}
 \text{Diagram 1: } \begin{array}{c} \text{A horizontal line with an arrow pointing right, labeled } p \text{ at the left end and } p \text{ at the right end. Above the line is a wavy semi-circular arc. Below the line is a horizontal arrow pointing right, labeled } q. \end{array} &= \frac{G_0(p, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(q - p, z) \cdot \mu_z(p, -q)^2 dq, \\
 \text{Diagram 2: } \begin{array}{c} \text{A horizontal line with an arrow pointing right, labeled } p \text{ at the left end and } p \text{ at the right end. Above the line is a wavy semi-circular arc. Below the line is a horizontal arrow pointing right, labeled } q. \end{array} &= -\frac{2 \cdot G_0(p, z)}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(p - q, z) \cdot \mu_{z(p, -q)} dq, \\
 \text{Diagram 3: } \begin{array}{c} \text{A horizontal line with an arrow pointing right, labeled } q. Above the line is a wavy semi-circular arc. \end{array} &= \frac{1}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(p - q, z) dq.
 \end{aligned}$$

---

<sup>18</sup>Attention! The terms have been simplified. For more details see Thesis sec. 2.4.2.

What is ERM?

# A small note..

What is ERM?

## A small note..

The actual calculation is very tedious.

What is ERM?

## A small note..

The actual calculation is very tedious. Used aspects:



## A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of  $\text{Ex}_{\mathcal{L}_f}$  in *second order*.

## A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of  $\text{Ex}_{\mathcal{L}_f}$  in *second order*.
  - Reduction of  $\mathcal{L}$  to *three point vertex*.

## A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of  $\text{Ex}_{\mathcal{L}_f}$  in *second order*.
  - Reduction of  $\mathcal{L}$  to *three point vertex*.
- Treatment of six point correlator.

## A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of  $\text{Ex}_{\mathcal{L}_f}$  in *second order*.
  - Reduction of  $\mathcal{L}$  to *three point vertex*.
- Treatment of six point correlator.
  - Useful tool: *Isserlis/Wick's Theorem*.

## A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of  $\text{Ex}_{\mathcal{L}_f}$  in *second order*.
  - Reduction of  $\mathcal{L}$  to *three point vertex*.
- Treatment of six point correlator.
  - Useful tool: *Isserlis/Wick's Theorem*.
- Consideration of restrictions towards moment space.

## A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of  $\text{Ex}_{\mathcal{L}_f}$  in *second order*.
  - Reduction of  $\mathcal{L}$  to *three point vertex*.
- Treatment of six point correlator.
  - Useful tool: *Isserlis/Wick's Theorem*.
- Consideration of restrictions towards moment space.
  - Processing linear system of equations for remaining integration parameters.

## A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of  $\text{Ex}_{\mathcal{L}_f}$  in *second order*.
  - Reduction of  $\mathcal{L}$  to *three point vertex*.
- Treatment of six point correlator.
  - Useful tool: *Isserlis/Wick's Theorem*.
- Consideration of restrictions towards moment space.
  - Processing linear system of equations for remaining integration parameters.
- Utilization of vertex' and propagator symmetries.

# What is Correlated Disorder?



# Second Moment of density fluctuations

## Second Moment of density fluctuations

.. previously we used an *a priori* probability density  $R \rightarrow \frac{1}{|V_{d,N}|}$

## Second Moment of density fluctuations

.. previously we used an *a priori* probability density  $R \rightarrow \frac{1}{|V_{d,N}|}$

$\hookrightarrow$  This did not account for the *structure* of our system.

## Second Moment of density fluctuations

.. previously we used an *a priori* probability density  $R \rightarrow \frac{1}{|V_{d,N}|}$

$\hookrightarrow$  This did not account for the *structure* of our system.

Main question to solve:

## Second Moment of density fluctuations

.. previously we used an *a priori* probability density  $R \rightarrow \frac{1}{|V_{d,N}|}$

$\hookrightarrow$  This did not account for the *structure* of our system.

Main question to solve:

**How can we include *structure* in our probability density?**

# The (radial) Particle Distribution Density


# The (radial) Particle Distribution Density

To calculate possibility of finding particles near a given reference  $r_0$  we used the *radial distribution function*

# The (radial) Particle Distribution Density

To calculate possibility of finding particles near a given reference  $r_0$  we used the *radial distribution function*

$$g_\varepsilon(r_0, x_*) = \int_{B_{x_*, \varepsilon}(r_0)} \rho_N^{(2)}(r_0, r) dr,$$






# The (radial) Particle Distribution Density

To calculate possibility of finding particles near a given reference  $r_0$  we used the *radial distribution function*

$$g_\varepsilon(r_0, x_*) = \int_{B_{x_*, \varepsilon}(r_0)} \rho_N^{(2)}(r_0, r) dr,$$

while  $\rho_N^{(2)}$  reflects integration of  $\exp(-\beta \cdot H(r, \cdot))$  for remaining particles.



What is Correlated Disorder?

# How can we use this now?

## How can we use this now?

There is a particular connection between  $g$  and the *static structure factor*  $S_*$ !

## How can we use this now?

There is a particular connection between  $g$  and the *static structure factor*  $S_*$ !

.. namely given by

$$S_*(\mathbf{q}) = 1 + \int_{\mathbb{R}^d} (g_0(\mathbf{r}) - 1) \cdot e^{i \cdot \mathbf{q} \cdot \mathbf{r}} d\mathbf{r}.$$

## How can we use this now?

There is a particular connection between  $g$  and the *static structure factor*  $S_*$ !


.. namely given by

$$S_*(\mathbf{q}) = 1 + \int_{\mathbb{R}^d} (g_0(\mathbf{r}) - 1) \cdot e^{i\mathbf{q}\cdot\mathbf{r}} d\mathbf{r}.$$


↑  
normalized to  $r_0 = 0$

# What does $g_0$ look like?

# What does $g_0$ look like?




# Resulting in the Static Structure Factor





# Resulting in the Static Structure Factor



Where did we implement  
this, what did it change?

Where did we implement this, what did it change?

# Analytical Aspects

Where did we implement this, what did it change?

## Analytical Aspects

We have established a new expectancy for  $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$ :

Where did we implement this, what did it change?

## Analytical Aspects

We have established a new expectancy for  $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$ :

$$\langle \delta\rho_R(\mathbf{q}), \delta\rho_R(-\mathbf{q}) \rangle = \frac{1}{\rho_*} \cdot S_*(\mathbf{q}).$$

Where did we implement this, what did it change?

## Analytical Aspects

We have established a new expectancy for  $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$ :

$$\langle \delta\rho_R(\mathbf{q}), \delta\rho_R(-\mathbf{q}) \rangle = \frac{1}{\rho_*} \cdot S_*(\mathbf{q}).$$

This results in a slight change in Feynman edges:

Where did we implement this, what did it change?

## Analytical Aspects

We have established a new expectancy for  $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$ :

$$\langle \delta\rho_R(\mathbf{q}), \delta\rho_R(-\mathbf{q}) \rangle = \frac{1}{\rho_*} \cdot S_*(\mathbf{q}).$$

This results in a slight change in Feynman edges:

$$\text{~~~~~}\overset{\gamma}{\text{~~~~~}} := \frac{S_*(\mathbf{q})}{\rho_*}$$

Where did we implement this, what did it change?

# Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:



Where did we implement this, what did it change?

## Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:

$$\begin{array}{c} \text{Diagram: A horizontal line with an arrow pointing right, labeled } \mathbf{p} \text{ at the left end and } \mathbf{p} \text{ at the right end. A wavy line (loop) connects the line at a point labeled } \mathbf{q} \text{ below it, forming a semi-circular arc above the line.} \\ \hline \end{array} = \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 \cdot S_*(\mathbf{q}) d\mathbf{q},$$

Where did we implement this, what did it change?

## Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:


$$\begin{array}{c} \text{Diagram: A horizontal line with an arrow pointing right, labeled } p \text{ at the left end and } p \text{ at the right end. A wavy line (representing a disorder) connects the line to itself, forming a loop. The momentum of the wavy line is labeled } q \text{ below it.} \\ \hline \end{array} = \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 \cdot S_*(\mathbf{q}) d\mathbf{q},$$

This also affects the Self-Energy:

Where did we implement this, what did it change?

## Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:


$$= \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 \cdot S_*(\mathbf{q}) d\mathbf{q},$$

This also affects the Self-Energy:

$$\Sigma_{S_*}^{(1)}(\mathbf{p}, z) = \frac{1}{\rho_*} \cdot \int_{\mathbb{R}^d} S_*(\mathbf{q}) \cdot G_0(\mathbf{p} - \mathbf{q}, z) \cdot S_*(\mathbf{q}) d\mathbf{q}.$$

Can we in any way  
compare our results?

Can we in any way compare our results?

# The Approach of Martin-Mayor

Can we in any way compare our results?

## The Approach of Martin-Mayor

Here, a *superposition approximation* was used:

$$\frac{1}{|V_{d,N}|} \cdot \exp(-\beta \cdot U(r)) \approx \frac{1}{|V_{d,N}|} \cdot \exp\left(-\beta \cdot \sum_{i \in [N-1]} u(r_i - r_{i+1})\right)$$

Can we in any way compare our results?

## The Approach of Martin-Mayor

Here, a *superposition approximation* was used:

$$\frac{1}{|V_{d,N}|} \cdot \exp(-\beta \cdot U(r)) \approx \frac{1}{|V_{d,N}|} \cdot \exp\left(-\beta \cdot \sum_{i \in [N-1]} u(r_i - r_{i+1})\right)$$

This approach only considers *direct neighbors* in a chain. Compare:

Can we in any way compare our results?

## The Approach of Martin-Mayor

Here, a *superposition approximation* was used:

$$\frac{1}{|V_{d,N}|} \cdot \exp(-\beta \cdot U(r)) \approx \frac{1}{|V_{d,N}|} \cdot \exp\left(-\beta \cdot \sum_{i \in [N-1]} u(r_i - r_{i+1})\right)$$

This approach only considers *direct neighbors* in a chain. Compare:

$$\exp(-\beta \cdot U(r)) = \exp\left(-\beta \cdot \sum_{(i,j) \in [N]^2} u(r_i - r_j)\right).$$



Can we in any way compare our results?

# The Approach of Martin-Mayor

A spring implementation of  $r \mapsto \frac{\exp(-\beta \cdot \sum \dots)}{|V_{d,N}|}$  is done:

Can we in any way compare our results?

# The Approach of Martin-Mayor

A spring implementation of  $r \mapsto \frac{\exp(-\beta \cdot \sum \dots)}{|V_{d,N}|}$  is done:

$$\ell(r) \approx \frac{f(r)}{|V_{d,N}|} \cdot \exp \left( -\beta \cdot \sum_{(i,j) \in [N]^2} u(r_i - r_j) \right).$$

Can we in any way compare our results?

## The Approach of Martin-Mayor

A spring implementation of  $r \mapsto \frac{\exp(-\beta \cdot \sum \dots)}{|V_{d,N}|}$  is done:

$$\ell(r) \approx \frac{f(r)}{|V_{d,N}|} \cdot \exp \left( -\beta \cdot \sum_{(i,j) \in [N]^2} u(r_i - r_j) \right).$$

This has an explicit approximation built into the spring function!

Can we in any way compare our results?

# The Approach of Martin-Mayor

As a consequence, the bare propagator changes:

Can we in any way compare our results?

## The Approach of Martin-Mayor

As a consequence, the bare propagator changes:

$$G_0(\boldsymbol{p}, z) = \frac{1}{z - \rho_* \cdot (\ell(\mathbf{0}) - S_*(\boldsymbol{p}))} \neq \underbrace{\frac{1}{z - \rho_* \cdot (f(\mathbf{0}) - S_*(\boldsymbol{p}))}}_{\text{Our Approach}}.$$

Can we in any way compare our results?

## The Approach of Martin-Mayor

As a consequence, the bare propagator changes:

$$G_0(\mathbf{p}, z) = \frac{1}{z - \rho_* \cdot (\ell(\mathbf{0}) - S_*(\mathbf{p}))} \neq \underbrace{\frac{1}{z - \rho_* \cdot (f(\mathbf{0}) - S_*(\mathbf{p}))}}_{\text{Our Approach}}.$$

- We explicitly did not approximate the spring function.

## The Approach of Martin-Mayor

As a consequence, the bare propagator changes:

$$G_0(\mathbf{p}, z) = \frac{1}{z - \rho_* \cdot (\ell(\mathbf{0}) - S_*(\mathbf{p}))} \neq \underbrace{\frac{1}{z - \rho_* \cdot (f(\mathbf{0}) - S_*(\mathbf{p}))}}_{\text{Our Approach}}.$$

- We explicitly did not approximate the spring function.
- We did not change the zeroth order term in the propagator.

What did a numerical  
model show?



What did a numerical model show?

# Analytical Foundation

What did a numerical model show?

# Analytical Foundation

We chose a *step function* for the spring mapping:

$$\mathbb{R} \ni r \mapsto f_a^{(num)}(r) = \begin{cases} 1 & \text{if } r < a, \\ 0 & \text{else.} \end{cases}$$

What did a numerical model show?

# Analytical Foundation

We chose a *step function* for the spring mapping:

$$\mathbb{R} \ni r \mapsto f_a^{(num)}(r) = \begin{cases} 1 & \text{if } r < a, \\ 0 & \text{else.} \end{cases}$$

.. resulting in a pair potential ( $d = 3$ ):

# Analytical Foundation

We chose a *step function* for the spring mapping:

$$\mathbb{R} \ni r \mapsto f_a^{(num)}(r) = \begin{cases} 1 & \text{if } r < a, \\ 0 & \text{else.} \end{cases}$$

.. resulting in a pair potential ( $d = 3$ ):


$$V_{d,N} \ni R \mapsto U_a^{(num)}(R) = \sum_{(i,j) \in [N]^2} \begin{cases} \frac{1}{2} \cdot (\|R_i - R_j\| - a)^2 & \text{if } \|R_i - R_j\| < a, \\ 0 & \text{else.} \end{cases}$$

What did a numerical model show?

# Results using the Hypernetted Chain


What did a numerical model show?

# Results using the Hypernetted Chain



What did a numerical model show?

# Results using the Hypernetted Chain



→ No major differences in the velocity of sound noticeable.

What did a numerical model show?

# Concluding terms..



What did a numerical model show?

## Concluding terms..

Adoption of the radial probability density and static structure factor introduced an integral correction in  $c_{\rho_*}$ :

## Concluding terms..

Adoption of the radial probability density and static structure factor introduced an integral correction in  $c_{\rho_*}$ :

$$G \mapsto \left( G_0(\mathbf{p}, 0)^{-1} - \rho_* \cdot \int_{\mathbb{R}^3} S_*(\mathbf{p} - \mathbf{q}) \cdot G(\mathbf{q}, z) \cdot V(\mathbf{q}, \mathbf{p}) d\mathbf{q} \right)^{-1}.$$

## Concluding terms..

Adoption of the radial probability density and static structure factor introduced an integral correction in  $c_{\rho_*}$ :

$$G \mapsto \left( G_0(\mathbf{p}, 0)^{-1} - \rho_* \cdot \int_{\mathbb{R}^3} S_*(\mathbf{p} - \mathbf{q}) \cdot G(\mathbf{q}, z) \cdot V(\mathbf{q}, \mathbf{p}) d\mathbf{q} \right)^{-1}.$$

There is a possibility of extension onto higher loop orders.

## Concluding terms..

Adoption of the radial probability density and static structure factor introduced an integral correction in  $c_{\rho_*}$ :

$$G \mapsto \left( G_0(\mathbf{p}, 0)^{-1} - \rho_* \cdot \int_{\mathbb{R}^3} S_*(\mathbf{p} - \mathbf{q}) \cdot G(\mathbf{q}, z) \cdot V(\mathbf{q}, \mathbf{p}) d\mathbf{q} \right)^{-1}.$$

There is a possibility of extension onto higher loop orders.

$$\exp(-\beta \cdot U(r)) \approx \exp\left(-\beta \cdot (r - \nu)^\perp \cdot A \cdot (r - \nu)\right), (\rho \text{ mediocre})$$

What did a numerical model show?

**Thank you for your attention!**

This presentation was only possible thanks to **Typst**.

What did a numerical model show?

**Thank you for your attention!**

Sources used in this presentation do not differ from the ones in the thesis.

This presentation was only possible thanks to **Typst**.

What did a numerical model show?

# Thank you for your attention!

Sources used in this presentation do not differ from the ones in the thesis. The Presentation, Thesis and Code are available on GitHub.

This presentation was only possible thanks to **Typst**.

# Thank you for your attention!

Sources used in this presentation do not differ from the ones in the thesis. The Presentation, Thesis and Code are available on GitHub.

`unb3rechenbar/BA24-CorDis.git`

This presentation was only possible thanks to **Typst**.