

Studies of ERM Models with Correlated Disorder

by Tom Folgmann

Bachelor Thesis Presentation, 2024

What is ERM?

What is ERM?

Imagine a system of $N \in \mathbb{N}$ particles.

What is ERM?


Imagine a system of $N \in \mathbb{N}$ particles.

→ How would you describe their relations?

What is ERM?

Imagine a system of $N \in \mathbb{N}$ particles.


→ How would you describe their relations?



What is ERM?

Imagine a system of $N \in \mathbb{N}$ particles.

→ How would you describe their relations?



A system with $N \in \mathbb{N}$ (related) particles can be described by a mathematical *Graph*.

What is ERM?

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix L* .

What is ERM?

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix* L .

$$L(G) := D(G) - W(G).$$

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix* L .

$$L(G) := D(G) - W(G).$$

- $D(G)$ gives the *degree* of each node: *Number of connected edges*.
- $W(G)$ encodes the *strength* (and direction) of the connections.

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix* L .

$$L(G) := D(G) - W(G).$$

- $D(G)$ gives the *degree* of each node: *Number of connected edges*.
- $W(G)$ encodes the *strength* (and direction) of the connections.

$$D(G) := \text{diag}(d), \quad d_i := \#\{e \in E : v_i \in e\},$$

$$W(G) := (w_{ij})_{(i,j) \in [N]^2}, \quad w : [N]^2 \rightarrow \mathbb{R}.$$

Fundamentally, the properties of such graphs can be rendered in a *Laplacian Matrix* L .

$$L(G) := D(G) - W(G).$$

- $D(G)$ gives the *degree* of each node: *Number of connected edges*.
- $W(G)$ encodes the *strength* (and direction) of the connections.

$$D(G) := \text{diag}(d), \quad d_i := \#\{e \in E : v_i \in e\},$$

$$W(G) := (w_{ij})_{(i,j) \in [N]^2}, \quad w : [N]^2 \rightarrow \mathbb{R}.$$

.. special case is the *Adjacency Matrix* A , where $w_{i,j} \in \{0, -1, 1\}$.

Definition of the ERM Laplacian Matrix

In the ERM model the Laplacian matrix is defined as

$$\tilde{U}(f, r) := \begin{pmatrix} \Sigma(f, 1) & -f_{12} & \dots & -f_{1N} \\ -f_{21} & \Sigma(f, 2) & \dots & -f_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -f_{N1} & -f_{N2} & \dots & \Sigma(f, N) \end{pmatrix},$$

Definition of the ERM Laplacian Matrix

In the ERM model the Laplacian matrix is defined as

$$\tilde{U}(f, r) := \begin{pmatrix} \Sigma(f, 1) & -f_{12} & \dots & -f_{1N} \\ -f_{21} & \Sigma(f, 2) & \dots & -f_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -f_{N1} & -f_{N2} & \dots & \Sigma(f, N) \end{pmatrix},$$

- *Interaction strength* given by $f_{ij} \stackrel{\text{m}}{=} f(r_i - r_j)$

Definition of the ERM Laplacian Matrix

In the ERM model the Laplacian matrix is defined as

$$\tilde{U}(f, r) := \begin{pmatrix} \Sigma(f, 1) & -f_{12} & \dots & -f_{1N} \\ -f_{21} & \Sigma(f, 2) & \dots & -f_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -f_{N1} & -f_{N2} & \dots & \Sigma(f, N) \end{pmatrix},$$


- *Interaction strength* given by $f_{ij} \stackrel{\text{m}}{=} f(r_i - r_j)$
- *Self-interaction* given by $\Sigma(f, i) \stackrel{\text{m}}{=} \sum_{j \in [N] \setminus \{i\}} f_{ij}$

How to measure Eigenvalues?

Let $\Lambda : [p] \rightarrow \sigma_P(\tilde{U}(f, r))$ map bijectively into the *point spectrum* of the ERM Laplacian.


How to measure Eigenvalues?

Let $\Lambda : [p] \rightarrow \sigma_P(\tilde{U}(f, r))$ map bijectively into the *point spectrum* of the ERM Laplacian.




How to measure Eigenvalues?

Let $\Lambda : [p] \rightarrow \sigma_P(\tilde{U}(f, r))$ map bijectively into the *point spectrum* of the ERM Laplacian.




How to measure Eigenvalues?

Let $\Lambda : [p] \rightarrow \sigma_P(\tilde{U}(f, r))$ map bijectively into the *point spectrum* of the ERM Laplacian.



How to measure Eigenvalues?

Let $\Lambda : [p] \rightarrow \sigma_P(\tilde{U}(f, r))$ map bijectively into the *point spectrum* of the ERM Laplacian.



... results in an (unnormalized) density function


$$E \mapsto \sum_{i \in [p]} \delta_{\Lambda_i}(E) \quad \in \{0, p\}$$

The Resolvent Eigenvalue Approximation

.. by an example point Λ_i at $i \in [p]$.


The Resolvent Eigenvalue Approximation

.. by an example point Λ_i at $i \in [p]$.



The Resolvent Eigenvalue Approximation

.. by an example point Λ_i at $i \in [p]$.



\hookrightarrow Usecase is the resolvent with a singularity at Λ_i .

What is ERM?

.. the connection can be found in *Newtonian Physics*:

What is ERM?

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$ map of a particle's position

What is ERM?

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$ map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

What is ERM?

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$ map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

$$\left(\frac{d}{dt}\right)^2 x_i(t) = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot x_j(t), \quad i, j \in [N].$$

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$ map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:


$$\left(\frac{d}{dt}\right)^2 x_i(t) = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot x_j(t), \quad i, j \in [N].$$

.. looking at the behaviour with regard to the initial conditions:

$$\left(\frac{d}{dt}\right)^2 \langle x_i(t), x_i(0) \rangle = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot \langle x_j(t), x_i(0) \rangle.$$


What is ERM?

In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:




What is ERM?

In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:



Implementation of two initial configurations:


In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:



Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle = \delta_{ij}$


In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:



Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle = \delta_{ij}$
- $\left(\frac{d}{dt}\right) \langle x_i(t), x_j(0) \rangle = 0$

In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:




Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle = \delta_{ij}$
- $\left(\frac{d}{dt}\right) \langle x_i(t), x_j(0) \rangle = 0$

.. leads to the *Resolvent representation* (Green's

function): of the ERM
Laplacian:⁵

In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:



Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle = \delta_{ij}$
- $\left(\frac{d}{dt}\right) \langle x_i(t), x_j(0) \rangle = 0$

.. leads to the *Resolvent representation* (Green's

function): of the ERM
Laplacian:⁶

$$(\mathcal{L}F_{j,i})(s) = \pm \frac{1}{\tilde{U}(f, x^*(t))_{i,j} - \delta_{ij} \cdot \lambda_i^2}.$$

⁶With $x^*(t) = (i \mapsto x_{i(t)})$ and $F_{j,i}(t) := \langle x_j(t), x_i(0) \rangle$.

What is ERM?

Where does *Randomness* come into play?

Where does *Randomness* come into play?

.. by a *slight* modification of functions!

Where does *Randomness* come into play?

.. by a *slight* modification of functions!

$\Omega \ni \omega \mapsto R(\omega)$, equivalent to x from before.

Where does *Randomness* come into play?

.. by a *slight* modification of functions!

$\Omega \ni \omega \mapsto R(\omega)$, equivalent to x from before.

Ev. step	Meaning
R	Random variable, abstract
$R(\omega)$	Vector of time dep. pos.
$R(\omega)_i$	i -th particle position, time dep. path
$R(\omega)_i(t)$	Position of i -th particle at time t

The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} = \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \overbrace{\left(\underbrace{e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, r) - z) \cdot \varphi, \varphi \rangle}}_{\text{Boltzmann density}} \cdot \lambda \right)}^{\text{Gaussian measure}} (d\varphi).$$

The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} = \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \overbrace{\left(\underbrace{e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, r) - z) \cdot \varphi, \varphi \rangle}}_{\text{Boltzmann density}} \cdot \lambda \right)}^{\text{Gaussian measure}} (d\varphi).$$

This is already a good starting point to understand our *Correlated Disorder* modification!

What is ERM?

.. missing key elements:

⁷Expansion to a functional can be argued, see thesis p. 19.

What is ERM?

.. missing key elements:

- The *action* (functional) S_{z,R_ω} at a *test point* $z \in \mathbb{C}$ and a particle position vector R_ω .

.. missing key elements:

- The *action* (functional) S_{z,R_ω} at a *test point* $z \in \mathbb{C}$ and a particle position vector R_ω .

↪ see the Boltzmann density exponent for implicit def.:⁹

.. missing key elements:

- The *action* (functional) S_{z,R_ω} at a *test point* $z \in \mathbb{C}$ and a particle position vector R_ω .

↪ see the Boltzmann density exponent for implicit def.:¹⁰

$$-\frac{\beta}{2} \cdot S_{z,R_\omega}(\varphi) := -\frac{\beta}{2} \cdot \left\langle \left(\tilde{U}(f, r) - z \right) \cdot \varphi, \varphi \right\rangle$$

¹⁰Expansion to a functional can be argued, see thesis p. 19.

.. missing key elements:

- The *action* (functional) S_{z,R_ω} at a *test point* $z \in \mathbb{C}$ and a particle position vector R_ω .

↪ see the Boltzmann density exponent for implicit def.:¹¹

$$-\frac{\beta}{2} \cdot S_{z,R_\omega}(\varphi) := -\frac{\beta}{2} \cdot \langle (\tilde{U}(f, r) - z) \cdot \varphi, \varphi \rangle$$

- The *moment generating function* $Z_{z,R_\omega}[J]$. It requires the *force field* J .

¹¹Expansion to a functional can be argued, see thesis p. 19.

Definition 2.22. External Field Shift.

For $R : \Omega \rightarrow V_{d,N}$ and $\Phi \in \mathbb{F}_{d,N}$ we define

$$J \mapsto -\frac{1}{2} \cdot S_{z,R_\omega}^{(0)}(\Phi) + \int_{\mathbb{R}^d} J(x) \cdot \Phi(-x) + J(-x) \cdot \Phi(x) \lambda(dx)$$

the *field shifted action* $S_{z,R_\omega}^{(0)}$ by an external field $J \in \mathcal{S}(\mathbb{R}^d)$.

Definition 2.22. External Field Shift.

For $R : \Omega \rightarrow V_{d,N}$ and $\Phi \in \mathbb{F}_{d,N}$ we define

$$J \mapsto -\frac{1}{2} \cdot S_{z,R_\omega}^{(0)}(\Phi) + \int_{\mathbb{R}^d} J(x) \cdot \Phi(-x) + J(-x) \cdot \Phi(x) \lambda(dx)$$

the *field shifted action* $S_{z,R_\omega}^{(0)}$ by an external field $J \in \mathcal{S}(\mathbb{R}^d)$.

$$\hookrightarrow \frac{\delta}{\delta J(x)} S_{z,R_\omega}^{(0)}[\Phi] = i \cdot \Phi(-x).$$

The Generative Operator

Using these tools, an Operator generating Z_{z,R_ω} can be deduced:

The Generative Operator

Using these tools, an Operator generating Z_{z,R_ω} can be deduced:

$$Z_{z,R_\omega}[J] = \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J]} d\Phi = \underbrace{\left[\mathbb{E}_{\mathcal{L}_f} \left[\int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi\right)[\cdot]} d\Phi \right] \right]}_{\text{Generative Part}} [J].$$

→ Looking at different Taylor expansion terms yields different integrals.

Feynman Diagrammatics - Edges

.. conveniently using symmetry in Fourier space:

Feynman Diagrammatics - Edges

.. conveniently using symmetry in Fourier space:

$$\begin{aligned} \longrightarrow &:= \frac{G_0(\mathbf{p}, z)}{\rho_*} \\ \overset{\gamma}{\sim} &:= \frac{\mathbb{E}((\mathcal{F} \delta \rho_R)(\mathbf{q}) \cdot (\mathcal{F} \delta \rho_R)(-\mathbf{q}))}{\rho_*} \end{aligned}$$

Feynman Diagrammatics - Edges

.. conveniently using symmetry in Fourier space:


$$\begin{aligned} \longrightarrow &:= \frac{G_0(\mathbf{p}, z)}{\rho_*} \\ \gamma \\ \text{~~~~~} &:= \frac{\mathbb{E}((\mathcal{F} \delta \rho_R)(\mathbf{q}) \cdot (\mathcal{F} \delta \rho_R)(-\mathbf{q}))}{\rho_*} \end{aligned}$$

.. possible connections of these edges are given by *vertices*:


What is ERM?

Feynman Diagrammatics - Vertices

Feynman Diagrammatics - Vertices



Feynman Diagrammatics - Vertices



.. which completes the set of Feynman rules.


How can we use diagrammatics?

.. displaying summands in operator expansion.

How can we use diagrammatics?

.. displaying summands in operator expansion.


Observe **one** loop diagrams:



How can we use diagrammatics?

.. displaying summands in operator expansion.

Observe **one** loop diagrams:



.. represented diagrams are *irreducible*: $Z_{z,R_\omega}[J] = \exp\left(\sum_{C \in \mathcal{C}} C\right)$.

Integral representations¹²

¹²Attention! The terms have been simplified. For more details see Thesis sec. 2.4.2.

Integral representations¹³

$$\begin{aligned}
 \text{Diagram 1} &= \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 d\mathbf{q}, \\
 \text{Diagram 2} &= -\frac{2 \cdot G_0(\mathbf{p}, z)}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{p} - \mathbf{q}, z) \cdot \mu_{z(\mathbf{p}, -\mathbf{q})} d\mathbf{q}, \\
 \text{Diagram 3} &= \frac{1}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{p} - \mathbf{q}, z) d\mathbf{q}.
 \end{aligned}$$

¹³Attention! The terms have been simplified. For more details see Thesis sec. 2.4.2.

What is Correlated Disorder?

Second Moment of density fluctuations

Second Moment of density fluctuations

.. previously we used an *a priori* probability density $R \rightarrow \frac{1}{|V_{d,N}|}$

Second Moment of density fluctuations

.. previously we used an *a priori* probability density $R \rightarrow \frac{1}{|V_{d,N}|}$

\hookrightarrow This did not account for the *structure* of our system.

Second Moment of density fluctuations

.. previously we used an *a priori* probability density $R \rightarrow \frac{1}{|V_{d,N}|}$

\hookrightarrow This did not account for the *structure* of our system.

Main question to solve:

Second Moment of density fluctuations

.. previously we used an *a priori* probability density $R \rightarrow \frac{1}{|V_{d,N}|}$

\hookrightarrow This did not account for the *structure* of our system.

Main question to solve:

How can we include *structure* in our probability density?


The (radial) Particle Distribution Density

The (radial) Particle Distribution Density

To calculate possibility of finding particles near a given reference r_0 we used the *radial distribution function*

$$g_{r_0}(r) = \int_{\mathbb{R}^d} \rho_N^{(2)}(r_0 + r, r) dr,$$

while $\rho_N^{(2)}$ reflects integration of $\exp(-\beta \cdot H(r, \cdot))$ for remaining particles.



What is Correlated Disorder?

How can we use this now?

How can we use this now?

There is a particular connection between g and the *static structure factor* S_* !

How can we use this now?

There is a particular connection between g and the *static structure factor* S_* !


.. namely given by

$$S_*(\mathbf{q}) = 1 + \int_{\mathbb{R}^d} (g_0(\mathbf{r}) - 1) \cdot e^{i\mathbf{q}\cdot\mathbf{r}} d\mathbf{r}.$$

What does g_0 look like?¹⁴

¹⁴Looking at a soft sphere model, see later.


What does g_0 look like?¹⁵



¹⁵Looking at a soft sphere model, see later.

Resulting in the Static Structure Factor

Resulting in the Static Structure Factor



Where did we implement
this, what did it change?

Where did we implement this, what did it change?

Analytical Aspects

Where did we implement this, what did it change?

Analytical Aspects

We have established a new expectancy for $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$:

Where did we implement this, what did it change?

Analytical Aspects

We have established a new expectancy for $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$:

$$\langle \delta\rho_R(\mathbf{q}), \delta\rho_R(-\mathbf{q}) \rangle = \frac{1}{\rho_*} \cdot S_*(\mathbf{q}).$$

Where did we implement this, what did it change?

Analytical Aspects

We have established a new expectancy for $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$:

$$\langle \delta\rho_R(\mathbf{q}), \delta\rho_R(-\mathbf{q}) \rangle = \frac{1}{\rho_*} \cdot S_*(\mathbf{q}).$$

This results in a slight change in Feynman edges:

Where did we implement this, what did it change?

Analytical Aspects

We have established a new expectancy for $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$:

$$\langle \delta\rho_R(\mathbf{q}), \delta\rho_R(-\mathbf{q}) \rangle = \frac{1}{\rho_*} \cdot S_*(\mathbf{q}).$$

This results in a slight change in Feynman edges:

$$\text{~~~~~}\overset{\gamma}{\text{~~~~~}} := \frac{S_*(\mathbf{q})}{\rho_*}$$

Where did we implement this, what did it change?

Analytical Aspects

Where did we implement this, what did it change?


Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:

Where did we implement this, what did it change?

Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:

A Feynman diagram representing a self-energy loop. It consists of a horizontal line with an arrow pointing to the right, labeled with momentum p at both ends. A wavy line (representing a boson) forms a loop above the horizontal line. The horizontal line is labeled with momentum q in the middle. The wavy line is labeled with momentum p at both ends.
$$= \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 \cdot S_*(\mathbf{q}) d\mathbf{q},$$

Where did we implement this, what did it change?

Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:


$$\begin{array}{c} \text{Diagram: A horizontal line with an arrow pointing right, labeled } \mathbf{p} \text{ at the left end and } \mathbf{p} \text{ at the right end. A wavy line (representing a disorder) connects the line to itself, forming a loop. Below the loop, the momentum } \mathbf{q} \text{ is indicated.} \\ \hline \end{array} = \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 \cdot S_*(\mathbf{q}) d\mathbf{q},$$

This also affects the Self-Energy:

Where did we implement this, what did it change?

Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:


$$\text{Diagram} = \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 \cdot S_*(\mathbf{q}) d\mathbf{q},$$

This also affects the Self-Energy:

$$\Sigma_{S_*}^{(1)}(\mathbf{p}, z) = \frac{1}{\rho_*} \cdot \int_{\mathbb{R}^d} S_*(\mathbf{q}) \cdot G_0(\mathbf{p} - \mathbf{q}, z) \cdot S_*(\mathbf{q}) d\mathbf{q}.$$

Can we in any way compare our results?

Where did we implement this, what did it change?

The Approach of Martin-Mayor

Where did we implement this, what did it change?

The Approach of Martin-Mayor

Here, a *superposition approximation* was used:

$$\frac{1}{|V_{d,N}|} \cdot \exp(-\beta \cdot U(r)) \approx \frac{1}{|V_{d,N}|} \cdot \exp\left(-\beta \cdot \sum_{i \in [N-1]} u(r_i - r_{i+1})\right)$$

Where did we implement this, what did it change?

The Approach of Martin-Mayor

Here, a *superposition approximation* was used:

$$\frac{1}{|V_{d,N}|} \cdot \exp(-\beta \cdot U(r)) \approx \frac{1}{|V_{d,N}|} \cdot \exp\left(-\beta \cdot \sum_{i \in [N-1]} u(r_i - r_{i+1})\right)$$

This approach only considers *direct neighbors* in a chain. Compare:

Where did we implement this, what did it change?

The Approach of Martin-Mayor

Here, a *superposition approximation* was used:

$$\frac{1}{|V_{d,N}|} \cdot \exp(-\beta \cdot U(r)) \approx \frac{1}{|V_{d,N}|} \cdot \exp\left(-\beta \cdot \sum_{i \in [N-1]} u(r_i - r_{i+1})\right)$$

This approach only considers *direct neighbors* in a chain. Compare:

$$\exp(-\beta \cdot U(r)) = \exp\left(-\beta \cdot \sum_{(i,j) \in [N]^2} u(r_i - r_j)\right).$$

Where did we implement this, what did it change?

The Approach of Martin-Mayor

Where did we implement this, what did it change?

The Approach of Martin-Mayor

An implementation of $r \mapsto \frac{\exp(-\beta \cdot \sum \dots)}{|V_{d,N}|}$ is done:

$$\ell(r) \approx \frac{f(r)}{|V_{d,N}|} \cdot \exp \left(-\beta \cdot \sum_{(i,j) \in [N]^2} u(r_i - r_j) \right).$$

Where did we implement this, what did it change?

The Approach of Martin-Mayor

An implementation of $r \mapsto \frac{\exp(-\beta \cdot \sum \dots)}{|V_{d,N}|}$ is done:

$$\ell(r) \approx \frac{f(r)}{|V_{d,N}|} \cdot \exp \left(-\beta \cdot \sum_{(i,j) \in [N]^2} u(r_i - r_j) \right).$$

This has an explicit approximation built into the spring function!

Where did we implement this, what did it change?

The Approach of Martin-Mayor

Where did we implement this, what did it change?

The Approach of Martin-Mayor

In a consequence, the bare propagator changes:

Where did we implement this, what did it change?

The Approach of Martin-Mayor

In a consequence, the bare propagator changes:

$$G_0(\mathbf{p}, z) = \frac{1}{z - \rho_* \cdot (\ell(\mathbf{0}) - S_*(\mathbf{p}))} \neq \underbrace{\frac{1}{z - \rho_* \cdot (f(\mathbf{0}) - S_*(\mathbf{p}))}}_{\text{Our Approach}}.$$

Where did we implement this, what did it change?

The Approach of Martin-Mayor

In a consequence, the bare propagator changes:

$$G_0(\mathbf{p}, z) = \frac{1}{z - \rho_* \cdot (\ell(\mathbf{0}) - S_*(\mathbf{p}))} \neq \underbrace{\frac{1}{z - \rho_* \cdot (f(\mathbf{0}) - S_*(\mathbf{p}))}}_{\text{Our Approach}}.$$

- We explicitly did not approximate the spring function.

Where did we implement this, what did it change?

The Approach of Martin-Mayor

In a consequence, the bare propagator changes:

$$G_0(\mathbf{p}, z) = \frac{1}{z - \rho_* \cdot (\ell(\mathbf{0}) - S_*(\mathbf{p}))} \neq \underbrace{\frac{1}{z - \rho_* \cdot (f(\mathbf{0}) - S_*(\mathbf{p}))}}_{\text{Our Approach}}.$$

- We explicitly did not approximate the spring function.
- We did not change the zeroth order term in the propagator.

What did a numerical
model show?

What did a numerical model show?

Analytical Foundation

What did a numerical model show?

Analytical Foundation

We chose a *step function* for the spring mapping:

$$\mathbb{R} \ni r \mapsto f_a^{(num)}(r) = \begin{cases} 1 & \text{if } r < a, \\ 0 & \text{else.} \end{cases}$$

What did a numerical model show?

Analytical Foundation

We chose a *step function* for the spring mapping:

$$\mathbb{R} \ni r \mapsto f_a^{(num)}(r) = \begin{cases} 1 & \text{if } r < a, \\ 0 & \text{else.} \end{cases}$$

.. resulting in a pair potential

Analytical Foundation

We chose a *step function* for the spring mapping:

$$\mathbb{R} \ni r \mapsto f_a^{(num)}(r) = \begin{cases} 1 & \text{if } r < a, \\ 0 & \text{else.} \end{cases}$$

.. resulting in a pair potential


$$V_{d,N} \ni R \mapsto U_a^{(num)}(R) = \sum_{(i,j) \in [N]^2} \begin{cases} \frac{1}{2} \cdot (\|R_i - R_j\| - a)^2 & \text{if } \|R_i - R_j\| < a, \\ 0 & \text{else.} \end{cases}$$

What did a numerical model show?

Results using the Hypernetted Chain


What did a numerical model show?

Results using the Hypernetted Chain



What did a numerical model show?

Results using the Hypernetted Chain



→ Sadly no major differences in the velocity of sound noticeable.