

Studies of ERM Models with Correlated Disorder

by Tom Folgmann

Bachelor Thesis Presentation, 2024

Upfront: Goals.

Upfront: Goals.

Foundations

Upfront: Goals.

Foundations

- What is ERM?

Upfront: Goals.

Foundations

- What is ERM?
 - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?

Upfront: Goals.

Foundations

- What is ERM?
 - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
 - Where lies our main theoretical interest?

Upfront: Goals.

Foundations

- What is ERM?
 - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
 - Where lies our main theoretical interest?
 - What does randomness mean for the model?

Upfront: Goals.

Foundations

- What is ERM?
 - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
 - Where lies our main theoretical interest?
 - What does randomness mean for the model?
- Understanding the action.

Upfront: Goals.

Foundations

- What is ERM?
 - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
 - Where lies our main theoretical interest?
 - What does randomness mean for the model?
- Understanding the action.
 - How does it arise?

Upfront: Goals.

Foundations

- What is ERM?
 - How can a simple model be build on the idea of (repulsively) interacting (finite) particles?
 - Where lies our main theoretical interest?
 - What does randomness mean for the model?
- Understanding the action.
 - How does it arise?
 - How can a Gaussian approach help us out?

Upfront: Goals.

Foundations

- What does *generating* with respect to Boltzmann densities mean?

Upfront: Goals.

Foundations

- What does *generating* with respect to Boltzmann densities mean?
 - Is there a visual approach to calculations?

Upfront: Goals.

Steps forward

Upfront: Goals.

Steps forward

- Now, where comes the correlative nature into play?

Upfront: Goals.

Steps forward

- Now, where comes the correlative nature into play?
- What did we implement in our research?

Upfront: Goals.

Steps forward

- Now, where comes the correlative nature into play?
- What did we implement in our research? (Has this been done before?)

Upfront: Goals.

Steps forward

- Now, where comes the correlative nature into play?
- What did we implement in our research? (Has this been done before?)
- What did we **conclude**?

What is ERM?

What is ERM?

Imagine a system of $N \in \mathbb{N}$ particles.

What is ERM?


Imagine a system of $N \in \mathbb{N}$ particles.

→ How would you describe their relations?

What is ERM?

Imagine a system of $N \in \mathbb{N}$ particles.


→ How would you describe their relations?



What is ERM?

Imagine a system of $N \in \mathbb{N}$ particles.

→ How would you describe their relations?



A system with $N \in \mathbb{N}$ (related) particles can be described by a mathematical *Graph*.

Definition of the ERM Laplacian Matrix

Definition of the ERM Laplacian Matrix

In the ERM model the Laplacian matrix is defined as:

$$\tilde{U}(f, r) := \begin{pmatrix} \Sigma(f, 1) & -f_{12} & \dots & -f_{1N} \\ -f_{21} & \Sigma(f, 2) & \dots & -f_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -f_{N1} & -f_{N2} & \dots & \Sigma(f, N) \end{pmatrix} = D(G) - W(G).$$

Definition of the ERM Laplacian Matrix

In the ERM model the Laplacian matrix is defined as:

$$\tilde{U}(f, r) := \begin{pmatrix} \Sigma(f, 1) & -f_{12} & \dots & -f_{1N} \\ -f_{21} & \Sigma(f, 2) & \dots & -f_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -f_{N1} & -f_{N2} & \dots & \Sigma(f, N) \end{pmatrix} = D(G) - W(G).$$

- *Interaction strength* given by $f_{ij} \stackrel{\text{m}}{=} f(r_i - r_j)$

Definition of the ERM Laplacian Matrix

In the ERM model the Laplacian matrix is defined as:

$$\tilde{U}(f, r) := \begin{pmatrix} \Sigma(f, 1) & -f_{12} & \dots & -f_{1N} \\ -f_{21} & \Sigma(f, 2) & \dots & -f_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ -f_{N1} & -f_{N2} & \dots & \Sigma(f, N) \end{pmatrix} = D(G) - W(G).$$


- *Interaction strength* given by $f_{ij} \stackrel{\text{m}}{=} f(r_i - r_j)$
- *Self-interaction* given by $\Sigma(f, i) \stackrel{\text{m}}{=} \sum_{j \in [N] \setminus \{i\}} f_{ij}$

How to measure Eigenvalues?

Let $\Lambda : [k] \rightarrow \sigma_P(\tilde{U}(f, r))$ map bijectively into the *point spectrum* of the ERM Laplacian.


How to measure Eigenvalues?

Let $\Lambda : [k] \rightarrow \sigma_P(\tilde{U}(f, r))$ map bijectively into the *point spectrum* of the ERM Laplacian.




How to measure Eigenvalues?

Let $\Lambda : [k] \rightarrow \sigma_P(\tilde{U}(f, r))$ map bijectively into the *point spectrum* of the ERM Laplacian.



How to measure Eigenvalues?

Let $\Lambda : [k] \rightarrow \sigma_P(\tilde{U}(f, r))$ map bijectively into the *point spectrum* of the ERM Laplacian.



... results in an (unnormalized) density function


$$E \mapsto \sum_{i \in [k]} \delta_{\Lambda_i}(E) \quad \in \{0, \dots, k\}$$

The Resolvent Eigenvalue Approximation

.. by an example point Λ_i at $i \in [k]$.


The Resolvent Eigenvalue Approximation

.. by an example point Λ_i at $i \in [k]$.



The Resolvent Eigenvalue Approximation

.. by an example point Λ_i at $i \in [k]$.



\hookrightarrow Usecase is the resolvent with a singularity at Λ_i .

What is ERM?

.. the connection can be found in *Newtonian Physics*:

What is ERM?

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$ map of a particle's position

What is ERM?

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$ map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

What is ERM?

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$ map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

$$\left(\frac{d}{dt}\right)^2 x_i(t) = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot x_j(t), \quad i, j \in [N].$$

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$ map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

$$\left(\frac{d}{dt}\right)^2 x_i(t) = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot x_j(t), \quad i, j \in [N].$$

.. looking at the behaviour with regard to the initial conditions:

$$\left(\frac{d}{dt}\right)^2 \langle x_i(t), x_i(0) \rangle = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot \langle x_j(t), x_i(0) \rangle.$$

.. the connection can be found in *Newtonian Physics*:

$[N] \ni i \rightarrow (x_i : \mathbb{R}_{>0} \rightarrow \mathbb{R}^d)$ map of a particle's position

.. using our Laplacian ERM Matrix and Newtonian dynamics:

$$\left(\frac{d}{dt}\right)^2 x_i(t) = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot x_j(t), \quad i, j \in [N].$$


.. looking at the behaviour with regard to the initial conditions:

$$\left(\frac{d}{dt}\right)^2 \langle x_i(t), x_i(0) \rangle = -\tilde{U}(f, i \mapsto x_i(t))_{i,j} \cdot \langle x_j(t), x_i(0) \rangle.$$


This is just $f_{i,j}$!
.. for $i \neq j$.

What is ERM?

In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:




In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:



Implementation of two initial configurations:


In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:



Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle|_{t=0} = \delta_{ij}$


In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:



Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle|_{t=0} = \delta_{ij}$
- $\left(\frac{d}{dt}\right) \langle x_i(t), x_j(0) \rangle|_{t=0} = 0$

In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:




Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle|_{t=0} = \delta_{ij}$
- $\left(\frac{d}{dt}\right) \langle x_i(t), x_j(0) \rangle|_{t=0} = 0$

.. leads to the *Resolvent representation* (Green's

function): of the ERM Laplacian:⁵

In a visual approach $\langle x_i(t), x_i(0) \rangle$ represents:



Implementation of two initial configurations:

- $\langle x_i(t), x_j(0) \rangle|_{t=0} = \delta_{ij}$
- $\left(\frac{d}{dt}\right) \langle x_i(t), x_j(0) \rangle|_{t=0} = 0$

.. leads to the *Resolvent representation* (Green's

function): of the ERM
Laplacian:⁶

$$(\mathcal{L}F_{j,i})(s) = \pm \frac{1}{\tilde{U}(f, x^*(0))_{i,j} - \delta_{ij} \cdot \lambda_i^2}.$$

⁶With $x^*(t) = (i \mapsto x_{i(t)})$ and $F_{j,i}(t) := \langle x_j(t), x_i(0) \rangle$.

What does this mean for Eigenvalue measurement?

⁷A direct connection can be obtained, see Thesis p. 17.

What does this mean for Eigenvalue measurement?

Using a physical argument, complex analysis can be utilized for measurement.⁸ This leads to the *resolvent*:

What does this mean for Eigenvalue measurement?

Using a physical argument, complex analysis can be utilized for measurement.⁹ This leads to the *resolvent*:

$$G_N(\mathbf{p}, z) = \sum_{(i,j)} \int \pm \frac{1}{\tilde{U}(f, r)_{i,j} - \delta_{ij} \cdot z} \cdot e^{i \cdot \mathbf{p} \cdot (r_i - r_j)} dr.$$

⁹A direct connection can be obtained, see Thesis p. 17.

What does this mean for Eigenvalue measurement?

Using a physical argument, complex analysis can be utilized for measurement.¹⁰ This leads to the *resolvent*:

$$G_N(\mathbf{p}, z) = \sum_{(i,j)} \int \pm \frac{1}{\tilde{U}(f, r)_{i,j} - \delta_{ij} \cdot z} \cdot e^{i \cdot \mathbf{p} \cdot (r_i - r_j)} dr.$$

Q: What are we integrating over?

¹⁰A direct connection can be obtained, see Thesis p. 17.

What is ERM?

Where does *Randomness* come into play?

Where does *Randomness* come into play?

.. by a *slight* modification of functions!

Where does *Randomness* come into play?

.. by a *slight* modification of functions!

$\Omega \ni \omega \mapsto R(\omega)$, equivalent to x from before.

Where does *Randomness* come into play?

.. by a *slight* modification of functions!

$\Omega \ni \omega \mapsto R(\omega)$, equivalent to x from before.

Ev. step	Meaning
R	Random variable, abstract
$R(\omega)$	Vector of time dep. pos.
$R(\omega)_i$	i -th particle position, time dep. path
$R(\omega)_i(t)$	Position of i -th particle at time t (fixed for us.)

The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} \propto \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \left(e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, R_\omega) - z) \cdot \varphi, \varphi \rangle} \cdot \lambda \right) (d\varphi).$$

The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} \propto \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \left(\underbrace{e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, R_\omega) - z) \cdot \varphi, \varphi \rangle}}_{\text{Boltzmann density}} \cdot \lambda \right) (d\varphi).$$

The measurement of Eigenvalues


Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} \propto \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \left(\overbrace{e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, R_\omega) - z) \cdot \varphi, \varphi \rangle}}^{\text{Gaussian measure}} \cdot \lambda \right)_{\text{Boltzmann density}} (d\varphi).$$

The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:


$$\left(\tilde{U}(f, r) - z\right)_{ij}^{-1} \propto \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \underbrace{\left(\overbrace{e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, R_\omega) - z) \cdot \varphi, \varphi \rangle}}^{\text{Gaussian measure}} \cdot \lambda \right)}_{\text{Boltzmann density}} (d\varphi).$$

Value in \mathbb{R} ..
(simplification!) 

The measurement of Eigenvalues

Eigenvalue measurement can be done via the resolvent. Gaussian representation gives us:

$$\left(\tilde{U}(f, r) - z \right)_{ij}^{-1} \propto \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \underbrace{\left(e^{-\frac{\beta}{2} \cdot \langle (\tilde{U}(f, R_\omega) - z) \cdot \varphi, \varphi \rangle} \cdot \lambda \right)}_{\text{Boltzmann density}} \underbrace{(d\varphi)}_{\text{Gaussian measure}}.$$

Value in $\mathbb{R} \dots$ (simplification!) 

This is already a good starting point to understand our *Correlated Disorder* modification!

What is ERM?

.. missing key elements:

¹¹Expansion to a functional can be argued, see thesis p. 19.

What is ERM?

.. missing key elements:

- The *action* (functional) S_{z,R_ω} at a *test point* $z \in \mathbb{C}$ and a particle position vector R_ω .

.. missing key elements:

- The *action* (functional) S_{z,R_ω} at a *test point* $z \in \mathbb{C}$ and a particle position vector R_ω .

↪ see the Boltzmann density exponent for implicit def.:¹³

.. missing key elements:

- The *action* (functional) S_{z,R_ω} at a *test point* $z \in \mathbb{C}$ and a particle position vector R_ω .

↪ see the Boltzmann density exponent for implicit def.:¹⁴

$$S_{z,R_\omega}(\varphi) := -\frac{\beta}{2} \cdot \left\langle \left(\tilde{U}(f, r) - z \right) \cdot \varphi, \varphi \right\rangle$$

¹⁴Expansion to a functional can be argued, see thesis p. 19.

.. missing key elements:

- The *action* (functional) S_{z,R_ω} at a *test point* $z \in \mathbb{C}$ and a particle position vector R_ω .

↪ see the Boltzmann density exponent for implicit def.:¹⁵

$$S_{z,R_\omega}(\varphi) := -\frac{\beta}{2} \cdot \left\langle \left(\tilde{U}(f, r) - z \right) \cdot \varphi, \varphi \right\rangle$$

- The *moment generating function* $Z_{z,R_\omega}[J]$. It requires the *force field* J .

¹⁵Expansion to a functional can be argued, see thesis p. 19.

What is ERM?

Action Functional and Densitys

Action Functional and Densitys

A clever modification leads to Dirac integration:

Action Functional and Densitys

A clever modification leads to Dirac integration:

$$S_{z, R_\omega}(\varphi) \propto \sum_{i,j} \iint \Phi_\varphi(x) \cdot \Phi_\varphi(y) \cdot \mathcal{U}(x, y) \left(\delta_{R_\omega(i)} \otimes \delta_{R_\omega(j)} \right) (d(x, y))$$

Action Functional and Densities

A clever modification leads to Dirac integration:

$$S_{z, R_\omega}(\varphi) \propto \sum_{i,j} \iint \Phi_\varphi(x) \cdot \Phi_\varphi(y) \cdot \mathcal{U}(x, y) \left(\delta_{R_\omega(i)} \otimes \delta_{R_\omega(j)} \right) (d(x, y))$$

$\uparrow \qquad \qquad \qquad \rho_{R_\omega} = \sum_{i \in [N]} \delta_{R_\omega(i)} \qquad \qquad \qquad \uparrow$

Action Functional and Densities

A clever modification leads to Dirac integration:

$$S_{z, R_\omega}(\varphi) \propto \sum_{i,j} \iint \Phi_\varphi(x) \cdot \Phi_\varphi(y) \cdot \mathcal{U}(x, y) \left(\delta_{R_\omega(i)} \otimes \delta_{R_\omega(j)} \right) (d(x, y))$$

$\uparrow \quad \quad \quad \rho_{R_\omega} = \sum_{i \in [N]} \delta_{R_\omega(i)} \quad \quad \quad \uparrow$

.. coming from ..

$$\left(\tilde{U}(f, r) - z \right)_{ij}^{-1} \propto \int_{\mathbb{R}^d} \varphi_i \cdot \varphi_j \left(e^{S_{z, R_\omega}(\varphi)} \cdot \lambda \right) (d\varphi).$$

The Generative Operator

Using these tools, an Operator generating Z_{z,R_ω} can be deduced:

The Generative Operator

Using these tools, an Operator generating Z_{z,R_ω} can be deduced:

$$Z_{z,R_\omega}[J] = \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J]} d\Phi$$

The Generative Operator

Using these tools, an Operator generating Z_{z,R_ω} can be deduced:

$$Z_{z,R_\omega}[J] = \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J]} d\Phi$$

Possible due to

$$\rho_R \rightarrow \lambda + \delta\rho_R$$

The Generative Operator

Using these tools, an Operator generating Z_{z,R_ω} can be deduced:

$$\begin{aligned}
 Z_{z,R_\omega}[J] &= \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J]} d\Phi \\
 &\quad \uparrow \\
 \text{Possible due to } \rho_R \rightarrow \lambda + \delta\phi_R &= \underbrace{\left[\mathbb{E}_{\mathbf{x}_{\mathcal{L}_f}} \left[\int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi\right)[\cdot]} d\Phi \right] \right]}_{\text{Generative Part}} [J].
 \end{aligned}$$

The Generative Operator

Using these tools, an Operator generating Z_{z,R_ω} can be deduced:

$$\begin{aligned}
 Z_{z,R_\omega}[J] &= \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J]} d\Phi \\
 &\quad \uparrow \qquad \qquad \qquad \downarrow \text{Only free theory!} \\
 \text{Possible due to } \rho_R \rightarrow \lambda + \delta\phi_R &= \underbrace{\left[\text{Ex}_{\mathcal{L}_f} \left[\int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi\right)[\cdot]} d\Phi \right] \right]}_{\text{Generative Part}} [J].
 \end{aligned}$$

The Generative Operator

Using these tools, an Operator generating Z_{z,R_ω} can be deduced:

$$Z_{z,R_\omega}[J] = \int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi + S_{z,R_\omega}^{(int)}\Phi\right)[J]} d\Phi$$

Possible due to $\rho_R \rightarrow \lambda + \delta\phi_R$

$$= \left[\text{Ex}_{\mathcal{L}_f} \left[\underbrace{\int_{\mathbb{F}_{d,N}} e^{\left(S_{z,R_\omega}^{(0)}\Phi\right)[\cdot]} d\Phi}_{\text{Generative Part}} \right] \right] [J].$$

Only free theory!

This needs explanation.

What is ERM?

.. “Ex” of course is an abbreviation.

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include?

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

What is ERM?


.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left(\frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) (\lambda \otimes \widehat{\delta \rho_{R_\omega}})(d\mathbf{p})$$

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left(\frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left(\lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p})$$

$\rho_R \rightarrow \lambda + \delta \rho_R$


.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left(\frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left(\lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p})$$

$\rho_R \rightarrow \lambda + \delta \rho_R$
 $\varphi \rightarrow \Phi$

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left(\frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left(\lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p})$$

for $\Phi(\mathbf{p}_1) \xrightarrow{\quad} \frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)}$ $\xrightarrow{\quad} \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)}$ $\xrightarrow{\quad} \lambda \otimes \widehat{\delta \rho_{R_\omega}}$

$\rho_R \rightarrow \lambda + \delta \varphi_R$
 $\varphi \rightarrow \Phi$

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left(\frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) \left(\lambda \otimes \widehat{\delta \rho_{R_\omega}} \right) (d\mathbf{p})$$

for $\Phi(\mathbf{p}_1)$ ————— \uparrow for $\Phi(-\mathbf{p}_1 - \mathbf{p}_2)$ ————— \uparrow

$\rho_R \rightarrow \lambda + \delta \rho_R$
 $\varphi \rightarrow \Phi$

What is ERM?

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left(\frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) (\lambda \otimes \widehat{\delta \rho_{R_\omega}})(d\mathbf{p})$$

for $\Phi(\mathbf{p}_1)$ ↑ ↑
 for $\Phi(-\mathbf{p}_1 - \mathbf{p}_2)$ ↑

$\rho_R \rightarrow \lambda + \delta \rho_R$
 $\varphi \rightarrow \Phi$

$$\Phi \mapsto \left(\int_{(\mathbb{R}^d)^2} \Phi(\mathbf{p}_1) \cdot \Phi(-\mathbf{p}_1 - \mathbf{p}_2) \cdot \mu_z(\mathbf{p}_1, \mathbf{p}_2) d\mathbf{p} \right) \cdot e^{(S_{z, R_\omega}^{(0)} \Phi)[J]}.$$

.. “Ex” of course is an abbreviation. Q: What does it include? To keep things simple, just “Integrals”:

$$\int_{(\mathbb{R}^d)^2} \mu_z(\mathbf{p}_1, \mathbf{p}_2) \cdot \left(\frac{\delta}{\delta \hat{J}(-\mathbf{p}_1)} \circ \frac{\delta}{\delta \hat{J}(\mathbf{p}_1 + \mathbf{p}_2)} \right) (\lambda \otimes \widehat{\delta \rho_{R_\omega}})(d\mathbf{p})$$

$\text{for } \Phi(\mathbf{p}_1) \xrightarrow{\quad} \uparrow \quad \quad \quad \uparrow \xrightarrow{\quad} \text{for } \Phi(-\mathbf{p}_1 - \mathbf{p}_2)$

$\rho_R \rightarrow \lambda + \delta \rho_R$
 $\varphi \rightarrow \Phi$

$$\Phi \mapsto \left(\int_{(\mathbb{R}^d)^2} \Phi(\mathbf{p}_1) \cdot \Phi(-\mathbf{p}_1 - \mathbf{p}_2) \cdot \mu_z(\mathbf{p}_1, \mathbf{p}_2) d\mathbf{p} \right) \cdot e^{(S_{z, R_\omega}^{(0)} \Phi)[J]}.$$

→ Looking at different Taylor expansion terms of $\underbrace{\exp\left(\int_{(\mathbb{R}^d)^2} \dots d\mathbf{p}\right)}_{\text{Ex}_{\mathcal{L}_f}}$ yields different powers of integral operators.

What is ERM?

On a broad scale: Dyson Series

On a broad scale: Dyson Series

The generative aspect applies to the resolvent G . This leads to the *Dyson Series* (Fixed Point Equation):

On a broad scale: Dyson Series

The generative aspect applies to the resolvent G . This leads to the *Dyson Series* (Fixed Point Equation):

$$G = \left((G_0)^{-1} - \Sigma(G) \right)^{-1}.$$

On a broad scale: Dyson Series

The generative aspect applies to the resolvent G . This leads to the *Dyson Series* (Fixed Point Equation):

$$G = \left((G_0)^{-1} - \Sigma(G) \right)^{-1}.$$

Hereby $G_0(p, z) = \left(z - \rho_* \cdot \left(\hat{f}(0) - \hat{f}(p) \right) \right)^{-1}$ is the *bare propagator* and $\Sigma(G)$ the *self-energy*.

Feynman Diagrammatics - Edges

.. conveniently using symmetry in Fourier space:

Feynman Diagrammatics - Edges

.. conveniently using symmetry in Fourier space:

$$\begin{aligned} \longrightarrow &:= \frac{G_0(\mathbf{p}, z)}{\rho_*} \\ \text{~~~~~} &:= \frac{\mathbb{E}((\mathcal{F} \delta \rho_R)(\mathbf{q}) \cdot (\mathcal{F} \delta \rho_R)(-\mathbf{q}))}{\rho_*} \end{aligned}$$

Feynman Diagrammatics - Edges

.. conveniently using symmetry in Fourier space:


$$\begin{aligned} \longrightarrow &:= \frac{G_0(\mathbf{p}, z)}{\rho_*} \\ \text{~~~~~} &:= \frac{\mathbb{E}((\mathcal{F} \delta \rho_R)(\mathbf{q}) \cdot (\mathcal{F} \delta \rho_R)(-\mathbf{q}))}{\rho_*} \end{aligned}$$

.. possible connections of these edges are given by *vertices*:


What is ERM?


Feynman Diagrammatics - Vertices

Feynman Diagrammatics - Vertices



Feynman Diagrammatics - Vertices


$$:= \mu_z(\mathbf{p}, \mathbf{q}) \quad \Rightarrow \quad \text{Three-point Vertex}$$


$$:= -V(\mathbf{p}, -\mathbf{q}_1) \quad \Rightarrow \quad \text{Four-point Vertex}$$

.. which completes the set of Feynman rules.


How can we use diagrammatics?

.. displaying summands in operator expansion.

How can we use diagrammatics?

.. displaying summands in operator expansion.


Observe **one** loop diagrams:



How can we use diagrammatics?

.. displaying summands in operator expansion.

Observe **one** loop diagrams:



.. represented diagrams are *irreducible*: $Z_{z,R_\omega}[J] \propto \exp\left(\sum_{C \in \mathcal{C}} C\right)$.

Integral representations¹⁶

¹⁶Attention! The terms have been simplified. For more details see Thesis sec. 2.4.2.

Integral representations¹⁷

$$\begin{aligned}
 \begin{array}{c} \text{Diagram 1: A horizontal line with an arrow pointing right. Above the line, a wavy line forms a semi-circle connecting the start and end points. The start point is labeled } p \text{ and the end point is labeled } p. \text{ Below the line, the label } p - q \text{ is centered.} \end{array} &= \frac{G_0(p, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(q - p, z) \cdot \mu_z(p, -q)^2 dq, \\
 \begin{array}{c} \text{Diagram 2: A horizontal line with an arrow pointing right. Above the line, a wavy line forms a semi-circle connecting the start and end points. The start point is labeled } p - q \text{ and the end point is labeled } p. \end{array} &= -\frac{2 \cdot G_0(p, z)}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(p - q, z) \cdot \mu_z(p, -q) dq, \\
 \begin{array}{c} \text{Diagram 3: A horizontal line with an arrow pointing right. Above the line, a wavy line forms a semi-circle connecting the start and end points. The start point is labeled } p - q \text{ and the end point is labeled } p - q. \end{array} &= \frac{1}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(p - q, z) dq.
 \end{aligned}$$

¹⁷Attention! The terms have been simplified. For more details see Thesis sec. 2.4.2.

What is ERM?

A small note..

What is ERM?

A small note..

The actual calculation is very tedious.

What is ERM?

A small note..

The actual calculation is very tedious. Used aspects:

A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of $\text{Ex}_{\mathcal{L}_f}$ in *second order*.

A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of $\text{Ex}_{\mathcal{L}_f}$ in *second order*.
 - Reduction of \mathcal{L} to *three point vertex*.

A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of $\text{Ex}_{\mathcal{L}_f}$ in *second order*.
 - Reduction of \mathcal{L} to *three point vertex*.
- Treatment of six point correlator.

A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of $\text{Ex}_{\mathcal{L}_f}$ in *second order*.
 - Reduction of \mathcal{L} to *three point vertex*.
- Treatment of six point correlator.
 - Useful tool: *Isserlis/Wick's Theorem*.

A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of $\text{Ex}_{\mathcal{L}_f}$ in *second order*.
 - Reduction of \mathcal{L} to *three point vertex*.
- Treatment of six point correlator.
 - Useful tool: *Isserlis/Wick's Theorem*.
- Consideration of restrictions towards moment space.

A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of $\text{Ex}_{\mathcal{L}_f}$ in *second order*.
 - Reduction of \mathcal{L} to *three point vertex*.
- Treatment of six point correlator.
 - Useful tool: *Isserlis/Wick's Theorem*.
- Consideration of restrictions towards moment space.
 - Processing linear system of equations for remaining integration parameters.

A small note..

The actual calculation is very tedious. Used aspects:

- Integral Operator representation of $\text{Ex}_{\mathcal{L}_f}$ in *second order*.
 - Reduction of \mathcal{L} to *three point vertex*.
- Treatment of six point correlator.
 - Useful tool: *Isserlis/Wick's Theorem*.
- Consideration of restrictions towards moment space.
 - Processing linear system of equations for remaining integration parameters.
- Utilization of vertex' and propagator symmetries.

What is Correlated Disorder?

Second Moment of density fluctuations

Second Moment of density fluctuations

.. previously we used an *a priori* probability density $R \rightarrow \frac{1}{|V_{d,N}|}$

Second Moment of density fluctuations

.. previously we used an *a priori* probability density $R \rightarrow \frac{1}{|V_{d,N}|}$

\hookrightarrow This did not account for the *structure* of our system.

Second Moment of density fluctuations

.. previously we used an *a priori* probability density $R \rightarrow \frac{1}{|V_{d,N}|}$

\hookrightarrow This did not account for the *structure* of our system.

Main question to solve:

Second Moment of density fluctuations

.. previously we used an *a priori* probability density $R \rightarrow \frac{1}{|V_{d,N}|}$

\hookrightarrow This did not account for the *structure* of our system.

Main question to solve:

How can we include *structure* in our probability density?

The (radial) Particle Distribution Density


The (radial) Particle Distribution Density

To calculate possibility of finding particles near a given reference r_0 we used the *radial distribution function*

The (radial) Particle Distribution Density

To calculate possibility of finding particles near a given reference r_0 we used the *radial distribution function*

$$g_\varepsilon(r_0, x_*) = \int_{B_{x_*, \varepsilon}(r_0)} \rho_N^{(2)}(r_0, r) dr,$$




The (radial) Particle Distribution Density

To calculate possibility of finding particles near a given reference r_0 we used the *radial distribution function*

$$g_\varepsilon(r_0, x_*) = \int_{B_{x_*, \varepsilon}(r_0)} \rho_N^{(2)}(r_0, r) dr,$$

while $\rho_N^{(2)}$ reflects integration of $\exp(-\beta \cdot H(r, \cdot))$ for remaining particles.



What is Correlated Disorder?

How can we use this now?

How can we use this now?

There is a particular connection between g and the *static structure factor* S_* !

How can we use this now?

There is a particular connection between g and the *static structure factor* S_* !

.. namely given by

$$S_*(\mathbf{q}) = 1 + \rho_* \cdot \int_{\mathbb{R}^d} (g_0(\mathbf{r}) - 1) \cdot e^{i \cdot \mathbf{q} \cdot \mathbf{r}} d\mathbf{r}.$$

How can we use this now?

There is a particular connection between g and the *static structure factor* S_* !


.. namely given by

$$S_*(\mathbf{q}) = 1 + \rho_* \cdot \int_{\mathbb{R}^d} (g_0(\mathbf{r}) - 1) \cdot e^{i\mathbf{q} \cdot \mathbf{r}} d\mathbf{r}.$$


↑
normalized to $\mathbf{r}_0 = 0$

What does g_0 look like?


What does g_0 look like?



Resulting in the Static Structure Factor



Resulting in the Static Structure Factor



Where did we implement
this, what did it change?

Where did we implement this, what did it change?

Analytical Aspects

Where did we implement this, what did it change?

Analytical Aspects

We have established a new expectancy for $\delta\rho_R(\boldsymbol{q}) \cdot \delta\rho_R(-\boldsymbol{q})$:

Where did we implement this, what did it change?

Analytical Aspects

We have established a new expectancy for $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$:

$$\langle \delta\rho_R(\mathbf{q}), \delta\rho_R(-\mathbf{q}) \rangle = \frac{1}{\rho_*} \cdot S_*(\mathbf{q}).$$

Where did we implement this, what did it change?

Analytical Aspects

We have established a new expectancy for $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$:

$$\langle \delta\rho_R(\mathbf{q}), \delta\rho_R(-\mathbf{q}) \rangle = \frac{1}{\rho_*} \cdot S_*(\mathbf{q}).$$

This results in a slight change in Feynman edges:

Where did we implement this, what did it change?

Analytical Aspects

We have established a new expectancy for $\delta\rho_R(\mathbf{q}) \cdot \delta\rho_R(-\mathbf{q})$:

$$\langle \delta\rho_R(\mathbf{q}), \delta\rho_R(-\mathbf{q}) \rangle = \frac{1}{\rho_*} \cdot S_*(\mathbf{q}).$$

This results in a slight change in Feynman edges:

$$\text{~~~~~} := \frac{S_*(\mathbf{q})}{\rho_*}$$

Where did we implement this, what did it change?

Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:

Where did we implement this, what did it change?

Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:

$$\begin{array}{c} \text{Diagram: A horizontal line with an arrow pointing right, labeled } p \text{ at the left end and } p \text{ at the right end. A wavy line (representing a disorder) connects the line at a point labeled } p-q \text{ below it, forming a loop above the line.} \\ \hline \end{array} = \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 \cdot S_*(\mathbf{q}) d\mathbf{q},$$

Where did we implement this, what did it change?

Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:

$$\begin{array}{c} \text{Diagram: A horizontal line with an arrow pointing right, labeled } p \text{ at the left end and } p \text{ at the right end. A wavy line (representing a disorder) connects the line at a point labeled } p-q \text{ below it, forming a loop above the line.} \\ \hline = \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 \cdot S_*(\mathbf{q}) d\mathbf{q}, \end{array}$$

This also affects the Self-Energy:

Where did we implement this, what did it change?

Analytical Aspects

From this the Integrands of the irreducible diagrams gain a factor:

$$\begin{array}{c} \text{Diagram: A horizontal line with an arrow pointing right, labeled } p \text{ at the left end and } p \text{ at the right end. A wavy line (representing a disorder) connects the line to itself, forming a loop. The momentum of the loop is labeled } p - q. \end{array} = \frac{G_0(\mathbf{p}, z)^2}{\rho_*} \cdot \int_{\mathbb{R}^d} G_0(\mathbf{q} - \mathbf{p}, z) \cdot \mu_z(\mathbf{p}, -\mathbf{q})^2 \cdot S_*(\mathbf{q}) d\mathbf{q},$$

This also affects the Self-Energy:

$$\Sigma_{S_*}^{(1)}(\mathbf{p}, z) = \frac{1}{\rho_*} \cdot \int_{\mathbb{R}^d} S_*(\mathbf{q}) \cdot G_0(\mathbf{p} - \mathbf{q}, z) \cdot V(\mathbf{q}, \mathbf{p})^2 d\mathbf{q}.$$

Can we in any way
compare our results?

Can we in any way compare our results?

The Approach of Martin-Mayor

Can we in any way compare our results?

The Approach of Martin-Mayor

Here, a *superposition approximation* was used:

$$\frac{1}{|V_{d,N}|} \cdot \exp(-\beta \cdot U(r)) \approx \frac{1}{|V_{d,N}|} \cdot \exp\left(-\beta \cdot \sum_{i \in [N-1]} u(r_i - r_{i+1})\right)$$

Can we in any way compare our results?

The Approach of Martin-Mayor

Here, a *superposition approximation* was used:

$$\frac{1}{|V_{d,N}|} \cdot \exp(-\beta \cdot U(r)) \approx \frac{1}{|V_{d,N}|} \cdot \exp\left(-\beta \cdot \sum_{i \in [N-1]} u(r_i - r_{i+1})\right)$$

This approach only considers *direct neighbors* in a chain. Compare:

Can we in any way compare our results?

The Approach of Martin-Mayor

Here, a *superposition approximation* was used:

$$\frac{1}{|V_{d,N}|} \cdot \exp(-\beta \cdot U(r)) \approx \frac{1}{|V_{d,N}|} \cdot \exp\left(-\beta \cdot \sum_{i \in [N-1]} u(r_i - r_{i+1})\right)$$

This approach only considers *direct neighbors* in a chain. Compare:

$$\exp(-\beta \cdot U(r)) = \exp\left(-\beta \cdot \sum_{(i,j) \in [N]^2} u(r_i - r_j)\right).$$

Can we in any way compare our results?

The Approach of Martin-Mayor

A spring implementation of $r \mapsto \frac{\exp(-\beta \cdot \sum \dots)}{|V_{d,N}|}$ is done:

Can we in any way compare our results?

The Approach of Martin-Mayor

A spring implementation of $r \mapsto \frac{\exp(-\beta \cdot \sum \dots)}{|V_{d,N}|}$ is done:

$$\ell(r) \approx \frac{f(r)}{|V_{d,N}|} \cdot \exp \left(-\beta \cdot \sum_{(i,j) \in [N]^2} u(r_i - r_j) \right).$$

Can we in any way compare our results?

The Approach of Martin-Mayor

A spring implementation of $r \mapsto \frac{\exp(-\beta \cdot \sum \dots)}{|V_{d,N}|}$ is done:

$$\ell(r) \approx \frac{f(r)}{|V_{d,N}|} \cdot \exp \left(-\beta \cdot \sum_{(i,j) \in [N]^2} u(r_i - r_j) \right).$$

This has an explicit approximation built into the spring function!

Can we in any way compare our results?

The Approach of Martin-Mayor

As a consequence, the bare propagator changes:

Can we in any way compare our results?

The Approach of Martin-Mayor

As a consequence, the bare propagator changes:

$$G_0(\mathbf{p}, z) = \frac{1}{z - \rho_* \cdot (\hat{\ell}(\mathbf{0}) - \hat{\ell}(\mathbf{p}))} \neq \underbrace{\frac{1}{z - \rho_* \cdot (\hat{f}(\mathbf{0}) - \hat{f}(\mathbf{p}))}}_{\text{Our Approach}}.$$

Can we in any way compare our results?

The Approach of Martin-Mayor

As a consequence, the bare propagator changes:

$$G_0(\mathbf{p}, z) = \frac{1}{z - \rho_* \cdot (\hat{\ell}(\mathbf{0}) - \hat{\ell}(\mathbf{p}))} \neq \underbrace{\frac{1}{z - \rho_* \cdot (\hat{f}(\mathbf{0}) - \hat{f}(\mathbf{p}))}}_{\text{Our Approach}}.$$

- We explicitly did not approximate the spring function.

The Approach of Martin-Mayor

As a consequence, the bare propagator changes:

$$G_0(\mathbf{p}, z) = \frac{1}{z - \rho_* \cdot (\hat{\ell}(\mathbf{0}) - \hat{\ell}(\mathbf{p}))} \neq \underbrace{\frac{1}{z - \rho_* \cdot (\hat{f}(\mathbf{0}) - \hat{f}(\mathbf{p}))}}_{\text{Our Approach}}.$$

- We explicitly did not approximate the spring function.
- We did not change the zeroth order term in the propagator.

What did a numerical
model show?

What did a numerical model show?

Analytical Foundation

What did a numerical model show?

Analytical Foundation

We chose a *step function* for the spring mapping:

$$\mathbb{R} \ni r \mapsto f_a^{(num)}(r) = \begin{cases} 1 & \text{if } r < a, \\ 0 & \text{else.} \end{cases}$$

What did a numerical model show?

Analytical Foundation

We chose a *step function* for the spring mapping:

$$\mathbb{R} \ni r \mapsto f_a^{(num)}(r) = \begin{cases} 1 & \text{if } r < a, \\ 0 & \text{else.} \end{cases}$$

.. resulting in a pair potential ($d = 3$):

Analytical Foundation

We chose a *step function* for the spring mapping:

$$\mathbb{R} \ni r \mapsto f_a^{(num)}(r) = \begin{cases} 1 & \text{if } r < a, \\ 0 & \text{else.} \end{cases}$$

.. resulting in a pair potential ($d = 3$):


$$V_{d,N} \ni R \mapsto U_a^{(num)}(R) = \sum_{(i,j) \in [N]^2} \begin{cases} \frac{1}{2} \cdot (\|R_i - R_j\| - a)^2 & \text{if } \|R_i - R_j\| < a, \\ 0 & \text{else.} \end{cases}$$

What did a numerical model show?

Results using the Hypernetted Chain

What did a numerical model show?

Results using the Hypernetted Chain




→ Physical applicability at $\rho_* > 225$.

What did a numerical model show?

Results using the Hypernetted Chain

What did a numerical model show?

Results using the Hypernetted Chain




→ $S_* \neq 0$ yields slightly higher amplitudes.

What did a numerical model show?

Results using the Hypernetted Chain

What did a numerical model show?

Results using the Hypernetted Chain



→ No *major* differences in the velocity of sound noticeable.

What did a numerical model show?

Concluding terms..

Concluding terms..

Adoption of the radial probability density and static structure factor introduced an integral correction in c_{ρ_*} :

Concluding terms..

Adoption of the radial probability density and static structure factor introduced an integral correction in c_{ρ_*} :

$$G \mapsto \left(G_0(\mathbf{p}, 0)^{-1} - \rho_* \cdot \int_{\mathbb{R}^3} S_*(\mathbf{p} - \mathbf{q}) \cdot G(\mathbf{q}, z) \cdot V(\mathbf{q}, \mathbf{p}) d\mathbf{q} \right)^{-1}.$$

Concluding terms..

Adoption of the radial probability density and static structure factor introduced an integral correction in c_{ρ_*} :

$$G \mapsto \left(G_0(\mathbf{p}, 0)^{-1} - \rho_* \cdot \int_{\mathbb{R}^3} S_*(\mathbf{p} - \mathbf{q}) \cdot G(\mathbf{q}, z) \cdot V(\mathbf{q}, \mathbf{p}) d\mathbf{q} \right)^{-1}.$$

There is a possibility of extension onto higher loop orders.

Concluding terms..

Adoption of the radial probability density and static structure factor introduced an integral correction in c_{ρ_*} :

$$G \mapsto \left(G_0(\mathbf{p}, 0)^{-1} - \rho_* \cdot \int_{\mathbb{R}^3} S_*(\mathbf{p} - \mathbf{q}) \cdot G(\mathbf{q}, z) \cdot V(\mathbf{q}, \mathbf{p}) d\mathbf{q} \right)^{-1}.$$

There is a possibility of extension onto higher loop orders.

$$\exp(-\beta \cdot U(r)) \approx \exp\left(-\beta \cdot (r - \nu)^\perp \cdot A \cdot (r - \nu)\right), (\rho \text{ mediocre})$$

Thank you for your attention!

This presentation was only possible thanks to **Typst**.

Thank you!

Thank you for your attention!

Sources used in this presentation do not differ from the ones in the thesis.

This presentation was only possible thanks to **Typst**.

Thank you!

Thank you for your attention!

Sources used in this presentation do not differ from the ones in the thesis. The Presentation, Thesis and Code are available on GitHub.

This presentation was only possible thanks to **Typst**.

Thank you!

Thank you for your attention!

Sources used in this presentation do not differ from the ones in the thesis. The Presentation, Thesis and Code are available on GitHub.

`unb3rechenbar/BA24-CorDis.git`

This presentation was only possible thanks to **Typst**.