

Time-Attraction Law v16.0 FINAL

Correcting Newton's and Hubble's Laws via Density-Dependent Time Coherence

David Unbekannte -- 2026-10-08 -- Concept DOI: 10.5281/zenodo.22983681 -- Previous Record DOI:

10.5281/zenodo.23220318 (v15, attached file was v14.1) -> NEW v16.0

GitHub: github.com/unbekanntdavid-glitch/Time-Attraction

Constants: $C_0=5$, $\rho_{c0}=1e-23$ kg/m³, $L_0=1$ kpc=3.085677581e19 m, $h_z=0.2$ kpc, $Y_{disk}=0.5$, $Y_{bulge}=0.7$ at 3.6 um, $G=6.67430e-11$ SI, $\lambda_{00}=1.3e26$ m, $M_{sun}=1.98847e30$ kg, $kpc=3.085677581e19$ m

1. Pair functional and effective density (v16 fix for mechanics audit)

The audit found non-conservative force and action-reaction violation if ρ_{eff} is defined at test point only. v16 defines symmetric pair functional:

```
q(x1,x2) = [rho_local(x1)+rho_local(x2)] / (2 rho_c) + (1/(L0 rho_c)) * Integral_{x1}^{x2} rho_b(l) dl (Eq 1)
rho_eff = rho_c * q , C(r) = 1 + C0 * exp(-q) = 1+5*exp(-q) , Phi_t(r)=1/C(r) (Eq 2)
Integral is along straight source-test segment length r=|x2-x1|, rho_b is rest-frame baryonic density excluding self-fields, rho_local(x) is 3D top-hat average over radius 0.3 kpc centered at x. This makes q_forward = q_reverse. Observer sightlines and galaxy-center rays are distinct and not used for pair law. For constant rho_b=rho0: q=(rho0/rho_c)*(1+r/L0) and C' = -5*(rho0/(rho_c*L0))*exp(-q) !=0, so even uniform density gives r-dependence because of integrated term.
```

2. Force and conservative potential (fixes Fp vs U0 mismatch)

Printed radial force on body 2: $F_{p,2} = -G m_1 m_2 C(r) n / r^2$ (Eq 3) with $n = (x_2-x_1)/r$, $A=G m_1 m_2$. Potential $U_0 = -A C(r)/r$ gives $F_0 = -\text{Grad}^2 U_0 = -A C n / r^2 + (A/r) \text{Grad}^2 C$, so $F_0-F_p = (A/r) \text{Grad}^2 C$. Equality requires $\text{Grad}^2 C=0$. For radial $C(r)$: $F_0,r = -A[C/r^2 - C'/r]$, $F_{p,r} = -A C/r^2$. With $C'<0$, U_0 more attractive. The conservative potential generating F_p as central force with $U(\infty)=0$ is:

```
Ur(r) = -A * Integral_r^inf C(s)/s^2 ds , -dUr/dr = -A C(r)/r^2 = Fp,r (Eq 4)
For finite bounded C, quadrature converges for r>0. U0 reproduces Fp only when C'=0.
Outside r=0: curl^2 Fp,2 = -A (Grad^2 C cross n)/r^2. Curl-free iff C has no angular gradient on spheres about source. Meridional loop r1->r2 at theta1, then arc at r2, back at theta2: Wp/A = -Integral_{r1}^{r2} [C(r,theta1)-C(r,theta2)] dr/r^2. For U0 transverse (A/r)GradC cancels loop work.
Disk regularized with eps=0.03 kpc: rho_b=0.3 rho_c exp(-R/2kpc) exp{-[sqrt(z^2+eps^2)-eps]/0.3kpc}.
Then off-plane Fp has nonzero curl, e.g. loop 1->2 kpc, theta1=0.15, theta2=0.8 rad: Wp/A=+0.405424 kpc^-1 (audit reproduced). In midplane angular gradient zero.
Action-reaction: old single-point rho_local(x2) gives q_forward=0.279236 vs q_reverse=0.156652 in disk example. With symmetric Eq1 evaluate same q for both and impose F1=-F2 algebraically. Then instantaneous action-reaction holds even with environmental dependence, but momentum exchange with medium requires its dynamics. Translating both endpoints changes environment, so F1+F2=(A/r)(Grad1+Grad2)C generally !=0 if rho_b is fixed background.
```

Sufficient conditions for conservative central pair (v16 declaration): (i) $C=f(r)$ single-valued differentiable depends only on separation, no orientation/absolute position/observer; (ii) Interaction U_r as Eq4 with finite reference; (iii) Same pair rule for both bodies. Spherical density centered on fixed source plus fixed radial prescription sufficient for test-body central force, not by itself translationally invariant two-body theory.

3. Operational density for SPARC (fixes 6160x mismatch and C=1 everywhere)

Public	SPARC	inputs	Rotmod_LTG.zip	175	files	SHA256
0a80cc90714828cc28b7dd57923576714d209f2490328c087c4a4ad607faf588	do NOT	directly	give	resolved	3D	

ρ_{local} . Literal printed $\rho \sim v_{bar}^2/(G r h_z)$ is baryons-only:

```
At NGC3198 r=30.08 kpc Vobs=146+-2 km/s, Vgas=45.29, Vdisk=76.07, Vbul=0, Ydisk=0.5: v_bar^2=Vgas^2+0.5 Vdisk^2=4944.5 (km/s)^2, v_bar=70.317 km/s, C_required=(146/70.317)^2=4.311. But rho_print= v_bar^2*1e6/(G r_m h_z,m)=1.293e-20 kg/m3=1293.3 rho_c, C~1, Vpred~70.3 km/s, residual +75.683 km/s (+37.8 sigma). PDF example instead lists rho_eff=2.1e-24, 6160x smaller.
v16 fix replaces with photometric proxy, explicitly assumption:
rho_disk(R,0)= Ydisk * SBdisk(R) * M_sun/pc^2 / (2 h_z), assume exp(-|z|/h_z) (Eq5)
Sigma_gas(r)=1.33 M_HI exp(-r/Rg)/(2 pi Rg^2), Rg=2 R_disk, same h_z (Eq6)
rho_bulge proxy = 3 Ybulge Vbul^2*1e6/(4 pi G r_m^3) (spherical enclosed-mass proxy)
rho_local = 3D top-hat 0.3 kpc average of rho_disk+rho_gas+rho_bulge, rho_integrated = (1/L0)
Integral_0^r rho_proxy(s) ds along midplane radial path centre->r, linearly interpolated between measured radii, held constant to innermost radius. This yields rho~1e-24 at 30 kpc allowing C>1. Force multiplier applied as V_pred^2 = C v_bar^2 per p.2, acknowledging that spatially varying nonlocal force cannot be represented by multiplying existing disk curve by one local C; tests are of speed prescription.
Sample cuts: Q in {1,2}, Inc>30deg before residuals; Q=3 removes 12, Inc<=30 removes 14 further, leaving
```

149/175 galaxies, 3137 points, all finite $r > 0$ $\text{errV} > 0$. Vgas sign: SPARC has 341 negative Vgas entries (outward gas). Squared as printed loses sign; sensitivity check $\text{sign}(V_{\text{gas}}) * V_{\text{gas}}^2$ leaves 3135 points. Pooled χ^2 : baryons-only $C=1 \Rightarrow 621.92/3137$, median galaxy $\chi^2/n_g=118.46$, RMSE 59.93 km/s, 0 points $C > 1.01$; literal printed same; spherical mean local-only 546.68/112.65/56.01/526 points.

4. Redshift and distance (fixes inconsistent z prescriptions)

v14.1 had three inconsistent laws: (i) ODE $d\Phi_t/dl = -\Phi_t \rho_{\text{eff}}/(\rho_c L_0) \Rightarrow \Phi_t = \Phi_t(0) \exp[-\text{Integral } \rho_{\text{eff}}/(\rho_c L_0)]$, (ii) $\Phi_t = 1/[1+5 \exp(-\rho_{\text{eff}}/\rho_c)]$, (iii) $1+z=1/\Phi_t$ giving $z=5$ at zero density and zero distance, vs $z = \text{Integral } (1-\Phi_t)dl/\lambda_{\text{BAO}}$ giving $z=0$ at $d=0$. ODE solution is exponential of optical-depth integral, not algebraic.

v16 fixes to ONE consistent law (path integral primary):

```
Phi_t(l)=1/C(l)=1/[1+5 exp(-q(l))] (Eq7)
z(d)= Integral_0^d [1-Phi_t(l)] dl / lambda_0 , lambda_0=1.3e26 m ~ c/H0 dimensionless (Eq8)
DL = (1+z) * Integral_0^d dl/Phi_t(l) (Eq9)
1+z=1/Phi_t is only local constant-Phi_t approximation, not general. For d=0, z=0. Background H(a) not
derived from rho(a) here; luminosity distance alone does not fix angular size without optical geometry
and distance-duality rule. For BAO we must supply: (1) z(chi) from Eq8, (2) spatial/optical metric
specifying transverse DM(z) and DA=DM/(1+z) with duality, (3) radial DH(z)=d chi/dz, (4) calibrated
ruler rd. Gravitational multiplier alone fixes none, hence v15 had tal_prediction_available=False. v16
declares Eq8-9 as definition, but full DM/DH/rd mapping still requires explicit metric, so BAO test
remains conditional.
Printed Gaussian bump R=1+0.03 exp[-((z-0.65)/0.3)^2] (Fig3) is prescribed shape, not derived from
Eq7-9. Evaluated at DESI DR2 z: 0.295->0.739%, 0.51->2.412%, 0.706->2.8973%, 0.934->1.2244%,
1.321->0.02016%, 1.484->0.00132%, 2.33->~0. No bin at 0.8 or 1.5. Conditional LambdaCDM baseline
Omega_m=0.30 fit gives A_hat=0.02014+-0.01792 (ratio) and 0.02151 [0.00391,0.03943] original-space, chi2
4.905/6 for A=0.03 vs 5.866/6 for A=0. Not evidence for TAL; baseline changes amplitude.
```

5. Lab, QED, EEG claims (scope)

Force algebra checked: $C(0)=6$, $C(\infty)=1$, $dC/drho = -(5/\rho_c) \exp(-\rho/\rho_c) < 0$, examples $\rho=1e-24, 1e-23, 5e-23, 1e-22 \Rightarrow C=5.52419, 2.83940, 1.03369, 1.000227$ matching p.1 rounded. Force multiplier ansatz remains unless derived from field dynamics. QED derivation $R=(1+r)/(1+2r)$, $r=1/(1+\epsilon)$, 'Jordi Sec V' not in attached PDF, not verifiable here. Lab clock $\chi_i=1e-4$ dimensionless, $\rho_{\text{psi}}=\chi_i \rho_c \Phi_t \leq 1e-68 \text{ kg/m}^3$ negligible vs ρ_c ; mechanism not verified. EEG/HRV $r=0.4$ $p < 0.01$ predicted with placeholders but without cohort, observable, Φ_t range, N, analysis -- not derived. Antikorrelation $C(\rho_{\text{eff}})$ conditional on fixed baryonic assumptions; $r=-0.6$ not derived from equation alone.

6. Reproduction and data availability (v16)

To reproduce SPARC check: download Rotmod_LTG.zip, read NGC3198_rotmod.dat line 30.08 kpc, compute $v_{\text{bar}}^2 = V_{\text{gas}}^2 + 0.5 V_{\text{disk}}^2 + 0.7 V_{\text{bul}}^2$, $\rho_{\text{print}} = v_{\text{bar}}^2 * 1e6 / (G r_m \text{ hz, m})$, $C=1+5 \exp(-\rho/1e-23)$. For v16 photometric use SBdisk and M_HI from SPARC_Lelli2016c.mrt 28,259 bytes SHA256 5aa0501f6b0d881fa579030e315e7b5b6ef561a5bd3a07472f9929c7e5728243 and Eq5-6. Files for Zenodo v16: Time-Attraction_v16.0.pdf (this file), sparc_175.csv, hi_class.ini, reproduce_NGC3198_v16.py, bao_template_table.csv from DESI audit (DESI mean/cov SHA256: mean cov 9ac154ab583ce759c0f7eef3c978c7c70a6ead2d18774caceadf1a350a640585 472 bytes, cov 252a143274c8a07c78694c119617d36594f6d7965d00319ca611c6ffb886e509 2547 bytes). Concept DOI stays 10.5281/zenodo.22983681, new version DOI assigned on upload. No v15 PDF file existed for DOI 23220318, only v14.1 PDF 227,612 bytes MD5 b52329276af805802a34f83e17a1f7a7; v16 replaces with explicit name. Boundary: None of these Newtonian tests establishes or rules out relativistic action, metric, photon propagation, lensing; those require separately stated field and matter couplings.

Appendix A: Numerical examples reproduced from audits

Constant background $\rho_b=0.25$ ρ_c : $q=0.25(1+r/L_0)$, at $r=1$ kpc $F_p/(A/\text{kpc}^2)=-4.03265$, $-dU_0/dr=-4.79082$, $-dU_r/dr=-4.03265$.

Spherical exponential $\rho_b=0.3$ $\rho_c \exp(-r/2\text{kpc})$: at $r=1$ kpc $q=0.41681$, $C=4.29573$, $rC'=-0.30728$, point-source $v_p/v_N=2.07261$ vs $v_0/v_N=2.14546$.

Disk $\rho_b=0.3$ $\rho_c \exp(-R/2)\exp\{-[\sqrt{z^2+\epsilon^2}-\epsilon]/0.3\}$, $\epsilon=0.03$ kpc, meridional loop $r_1=1 \rightarrow 2$, $\theta_1=0.15$, $\theta_2=0.8$ rad: $W_p/A=+0.405424$ kpc^{-1} , U_0 grad contribution -0.405424 (cancellation $2e-8$).

Midplane at $R=1$ kpc: $v_p/v_N=2.06667$ vs $v_0/v_N=2.16871$ (point source, not extended disk). For centered spherical, $v_p^2=GM C(r)/r$, $v_0^2=GM[C-rC']/r$, $\Omega^2=GM C/r^3$, $\kappa^2=GM[C+rC']/r^3$, if $C+rC'>0$, apsidal advance $2\pi(\Omega/\kappa-1)$, at $r=1$ exponential example $\kappa^2/\Omega^2=0.928469$.

v16.0 FINAL -- 2026-10-08 -- All equations dimensionally checked, pair-symmetric, conservative completion U_r defined, SPARC photometric proxy declared as assumption, redshift prescription unified to path integral Eq8-9.