

A minimal equipment one-pot RT-LAMP assay for detecting reverse transcriptase activity

December 26, 2016

1 Introduction

There are a number of chronic diseases including cancers, neurological diseases, autoimmune diseases, and immune deficiencies with a suspected retroviral etiology [1]. In addition, tests for known retroviruses with a high viral diversity such as HIV are very specific and could miss divergent HIV strains [2][3]. Patients looking for an explanation of their chronic disease with a suspected retroviral etiology face significant challenges in diagnosis due to political impositions. While assays for detecting reverse transcriptase activity are available to researchers, the US government makes it illegal for patients to access these assays as the tests have not been approved for clinical diagnosis. While a reverse transcriptase activity assay by itself cannot diagnose a specific retrovirus, it can provide supporting evidence to look further with more specific investigations.

A patient who would like to investigate the presense of reverse transcriptase activity must either find a researcher who would be willing to run the test, travel to a country whose government allows such testing, perform social engineering to have a sample processed by pretending to be a researcher, or purchase the equipment and supplies and run tests themselves. While all avenues should be explored to maximize the chances of success, this paper explores the last option.

2 Design

design constraints balance minimal design/development time, minimal equipment requirements, adequate performance.

2.1 Sample Collection

minimal sample volume, blood drop from lancet

2.2 Lysis

[4]

[5]

[6]

2.3 Reaction

References

- [1] C. Voisset, R. A. Weiss, and D. J. Griffiths, "Human rna rumor viruses: the search for novel human retroviruses in chronic disease," *Microbiology and Molecular Biology Reviews*, vol. 72, no. 1, pp. 157–196, 2008.
- [2] I. Bártolo and N. Taveira, *HIV-1 Diversity and Its Implications in Diagnosis, Transmission, Disease Progression, and Antiretroviral Therapy*. INTECH Open Access Publisher, 2012.

- [3] L. M. Luft, M. J. Gill, and D. L. Church, “Hiv-1 viral diversity and its implications for viral load testing: review of current platforms,” *International Journal of Infectious Diseases*, vol. 15, no. 10, pp. e661–e670, 2011.
- [4] A. Malmsten, “Reverse transcriptase activity assays for retrovirus quantitation and characterization,” 2005.
- [5] S. Karamohamed, M. Ronaghi, and P. Nyrén, “Bioluminometric method for real-time detection of reverse transcriptase activity,” *oligonucleotides*, vol. 2, p. 5, 1998.
- [6] K. A. Curtis, D. L. Rudolph, and S. M. Owen, “Rapid detection of hiv-1 by reverse-transcription, loop-mediated isothermal amplification (rt-lamp),” *Journal of virological methods*, vol. 151, no. 2, pp. 264–270, 2008.