An I-band Period–Luminosity Distance to the Globular Cluster Kestrel-7 from 42 RR Lyrae Stars

Bianca Azartash
University of Oklahoma Libraries, Norman, OK, USA

Draft of 30 October 2026

DRAFT PREPRINT — MOCK RESEARCH PROJECT. All data are synthetic and Kestrel-7 is a fictional cluster. Cited works marked as fictional in references.bib do not exist.

Abstract

We present V- and I-band time-series photometry of 42 RR Lyrae stars in the recently discovered globular cluster Kestrel-7 (Haddad et al., 2021). Using Lomb–Scargle periodograms and fourth-order Fourier fits, we identify 28 fundamental-mode (RRab) and 14 first-overtone (RRc) pulsators. The mean RRab period, $\langle P_{ab} \rangle = 0.593$ d, places Kestrel-7 in the Oosterhoff gap. A free fit to the I-band period–luminosity relation gives a slope of -1.23 ± 0.09, consistent with recent calibrations. Adopting the parallax-based calibration of Okafor and Lindqvist (2023) and a reddening of $E(B-V) = 0.08$ (Chen and Abara, 2024), we derive a true distance modulus of $\mu_0 = 15.11 \pm 0.01$ (stat) ± 0.07 (sys) mag, corresponding to 10.5 ± 0.3 kpc.

1 Introduction

RR Lyrae stars are low-mass, core-helium-burning pulsators that trace old stellar populations. Although their V-band absolute magnitudes depend only weakly on period, a well-defined period–luminosity (PL) relation appears at redder wavelengths, where the relation is also less sensitive to metallicity and interstellar extinction (Okafor and Lindqvist, 2023). This makes I-band RR Lyrae photometry an efficient distance indicator for globular clusters.

Kestrel-7 was identified as a compact overdensity behind the southern Galactic bulge, with a horizontal-branch distance of 11 ± 2 kpc (Haddad et al., 2021). Variability among horizontal-branch stars in the discovery data motivated the dedicated time-series campaign presented here. Our goals are to (i) build a clean RR Lyrae census, (ii) determine the cluster's Oosterhoff type, and (iii) measure a precise PL distance.

2 Observations and Data Reduction

We obtained 62 epochs in each of V and I over a 118-day baseline. Point-spread-function photometry was performed on all frames and calibrated to a standard system. Each light curve was inspected and epochs flagged by the reduction pipeline (cosmic rays, bad columns; 25 of 5208 measurements) were removed. The resulting photometry is provided in `data/raw/kestrel7_photometry.csv`. Typical uncertainties are 0.01–0.02 mag.




Figure 1: Example phased V- and I-band light curves for one RRab and one RRc star, plotted over two cycles.

3 Analysis

Periods were determined from the V-band data with a Lomb–Scargle periodogram (Lomb, 1976; Scargle, 1982) over 0.2–1.0 d. We inspected folded light curves at the strongest peaks to rule out daily aliases, a common failure mode for nightly sampling (Petrov, 2019). Phased light curves in each band were fit with a fourth-order Fourier series, and mean magnitudes were computed by intensity-averaging the fitted model (Petrov, 2019). Example light curves are shown in Figure 1.

Stars with $P < 0.43$ d and sinusoidal light curves were classified as RRc. Their periods were fundamentalized with $\log P_F = \log P + 0.127$. The resulting period–amplitude (Bailey) diagram (Figure 2) shows two clean sequences with no ambiguous objects.

We fit $\langle I \rangle = a \log P_F + b$ by least squares, and separately derived μ_0 by fixing the slope and zero point to the calibration $M_I = -1.13 \log P_F + 0.47$ (Okafor and Lindqvist, 2023), correcting for $A_I = 0.15$ mag. Statistical uncertainties were estimated with 2000 bootstrap resamples. The analysis code is `src/rrlyrae_pl.py`, and a step-by-step record is in `notebooks/kestrel7_analysis.ipynb`.

4 Results

Our final catalog contains 28 RRab and 14 RRc stars (Table 1). The mean RRab period is $\langle P_{ab} \rangle = 0.593$ d and the mean RRc period is $\langle P_c \rangle = 0.347$ d, with an RRc fraction of 0.33.

The free-slope PL fit (Figure 3) gives $a = -1.23 \pm 0.09$ and $b = 15.70 \pm 0.03$ with an rms scatter of 0.043 mag, consistent with the intrinsic scatter of the calibrating relation. With the slope fixed to the calibration, we find $\mu_0 = 15.11 \pm 0.01$ (stat) ± 0.07 (sys) mag, or $d = 10.5 \pm 0.3$ kpc.

5 Discussion

Our distance agrees with, and is roughly six times more precise than, the horizontal-branch estimate of Haddad et al. (2021). The error budget is dominated by the systematic terms: the calibration zero point (0.05 mag) and the reddening (0.02 mag in $E(B-V)$, or about 0.04 mag in A_I).




Figure 2: Bailey (period–amplitude) diagram. RRab and RRc stars form two distinct sequences.

Table 1: Summary of derived properties of Kestrel-7.

Quantity	Value
RRab / RRc	28 / 14
$\langle P_{ab} \rangle$	0.593 d
$\langle P_c \rangle$	0.347 d
PL slope (free)	-1.23 ± 0.09
PL rms scatter	0.043 mag
A_I	0.15 mag
μ_0	15.11 ± 0.01 (stat) ± 0.07 (sys)
Distance	10.5 ± 0.3 kpc

With $\langle P_{ab} \rangle = 0.593$ d, Kestrel-7 falls in the 0.58–0.62 d range that is sparsely populated among Galactic halo clusters (Moreau et al., 2022). Its RRc fraction (0.33) is also intermediate between typical Oosterhoff I and II values. Oosterhoff-intermediate clusters are often associated with accreted systems, which makes Kestrel-7 a promising target for spectroscopic metallicity and kinematic follow-up.

6 Conclusions

We have identified 42 RR Lyrae stars in Kestrel-7 and used their I-band PL relation to measure a distance of 10.5 ± 0.3 kpc. The cluster is Oosterhoff-intermediate. Planned future work includes spectroscopic metallicities, which will allow us to apply a metallicity-dependent PL calibration.

Acknowledgments

This draft is part of a mock research project created for teaching and documentation exercises. No real telescope time was used.




Figure 3: I-band period–luminosity relation for Kestrel-7. RRc periods are fundamentalized. The line shows the free least-squares fit.

References

L. Chen and D. Abara. A high-resolution reddening map toward the southern galactic bulge from red clump giants. *Journal of Synthetic Astrophysics*, 13:412–430, 2024. FICTIONAL. papers/Chen$_A$bara$_2$024$_b$ulge$_r$eddening$_m$ap.pdf.

R. Haddad, S. Tanaka, and M. Ruiz. Discovery of kestrel-7: a faint, reddened globular cluster behind the galactic bulge. *Proceedings of the Mock Astronomical Society*, 48:233–241, 2021. FICTIONAL. papers/Haddad$_T$anaka$_R$uiz$_2$021$_K$estrel7$_d$iscovery.pdf.

N. R. Lomb. Least-squares frequency analysis of unequally spaced data. *Astrophysics and Space Science*, 39: 447–462, 1976.

J. Moreau, K. Adeyemi, and P. Sørensen. The oosterhoff dichotomy in halo globular clusters revisited. *Journal of Synthetic Astrophysics*, 11:57–80, 2022. FICTIONAL. papers/Moreau$_e$tal$_2$022$_O$osterhoff$_d$ichotomy.pdf.

N. Okafor and E. Lindqvist. An i-band period–luminosity calibration for galactic field rr lyrae stars from trigonometric parallaxes. *Journal of Synthetic Astrophysics*, 12:101–118, 2023. FICTIONAL. papers/Okafor$_L$indqvist$_2$023$_R$RLyrae$_P$L$_c$alibration.pdf.

A. Petrov. Fourier decomposition of rr lyrae light curves: Practical methods and common pitfalls. *Monthly Notes of Synthetic Photometry*, 7:22–39, 2019. FICTIONAL. papers/Petrov$_2$019$_F$ourier$_d$ecomposition$_m$ethods.pdf.

J. D. Scargle. Studies in astronomical time series analysis. ii. statistical aspects of spectral analysis of unevenly spaced data. *The Astrophysical Journal*, 263:835–853, 1982.