




DISTRICT AGRICULTURE CONTINGENCY PLAN DACP - 2019



KOLHAN DIVISION



**BIRSA AGRICULTURAL UNIVERSITY, KANKE, RANCHI, JHARKHAND
AND**

CENTRAL RESEARCH INSTITUTE FOR DRY LAND AGRICULTURE (CRIDA), HYDERABAD



BIRSA AGRICULTURAL UNIVERSITY, RANCHI

P.O.: Kanke, Ranchi-834006, Jharkhand, India



FOREWORD

Dr. Parvinder Kaushal
Vice-Chancellor

Agriculture in Jharkhand state has always been dependent on the mercy of South West monsoon. Production and productivity of crops in Rainfed agriculture of Jharkhand fluctuate year to year as per the time of monsoon arrival, amount of precipitation and distribution. The climate change in recent years has further augmented the problem keeping the farmers in great dilemma with respect to appropriate timing of agricultural operations and selection of right kind of crops and varieties for different land use situations. Even livestock rearing and forestry

activities are being affected due to climatic vagaries in Jharkhand. In this scenario of climatic change and climatic variability, farmers of the region need to have different sets of well defined farming options for different climatic and land situations. There was a need of having District Agriculture Contingency Plans (DACP) for all the districts, based on the probable climatic conditions and existing land situations.

In view of the prevailing agro-climatological constraints and on the initiative of ICAR-CRIDA, contingency plans for all the 24 districts of Jharkhand have been prepared by the Contingency Cell of Birsa Agricultural University under the technical guidance of Dr. D.N. Singh (Chairman, Contingency Cell) and Dr. A. Wadood (Nodal officer, Contingency Cell) involving scientists of all the Zonal Research Stations, Krishi Vigyan Kendras, concerned government officials and farmers. The District Agriculture Plans, so prepared, are fool proof documents published in 5 volumes (one each for 5 administrative divisions of Jharkhand having contingency plans for all the districts of the division concerned).

I hope the DACPs will serve as a ready reckoner and will be of great help to the farmers and other stakeholders in planning and carrying out the agricultural activities in a weather based and climate resilient manner which would certainly give sustenance to Jharkhand agriculture. I congratulate Dr. D. N. Singh, Dr. A. Wadood and entire team for their untiring efforts for bringing out such excellent and useful publication. I have all appreciation for ICAR-CRIDA for this initiative and also for financial support.

(Parvinder Kaushal)

Citations

Patron
Dr. Parvinder Kaushal
Vice-Chancellor, BAU

Co-Patron
Dr. K.V. Rao
Co-ordinator, Contingency Cell,
ICAR - CRIDA, Hyderabad

Chairman, Contingency Cell, BAU
Dr. D.N. Singh
Director Research, BAU

Nodal Officer, Contingency Cell, BAU
Dr. A. Wadood
Chairman, Department of Agrometeorology and Environmental Science

Members of Contingency Cell, BAU
Associate Directors, ZRSs & Heads, KVKS

• *Compilation and Editing* •
Dr. A. Wadood and Sri Binod Kumar
Department of Agrometeorology and Environmental Science
Birsa Agricultural University, Kanke, Ranchi

**DEPARTMENT OF AGROMETEOROLOGY AND ENVIRONMENTAL SCIENCE
BIRSA AGRICULTURAL UNIVERSITY, RANCHI**



PREFACE



Dr. A. Wadood
Chairman

The 28th state of India, Jharkhand came into existence on November 15th, 2000. The state has a total geographical area of 79.71 lakh hectares with a net cultivated area of about 28 lakh hectares (35.12%). The state is maintaining fairly a good forest cover of slightly over 23 lakh hectares. The remaining land falls under barren, cultivable waste, pasture and other categories. Jharkhand state falls in 7th Agroclimatic zone (out of 15 Agroclimatic zones as classified by planning commission), 8th Meteorological Subdivision (out of 36 meteorological subdivisions of India) and in 11th -13th Agro-Ecological zones (out of 20 AgroEcozones of India). For the ease of better agricultural planning based on climate and soil/land features India has been divided into 127 Agroclimatic zones (AZ1 -AZ127). The state of Jharkhand has 3 agroclimatic zones AZ57, AZ58 and AZ59, the corresponding agroclimatic sub zones are Central and North Eastern Plateau sub-zone (sub-zone IV), Western Plateau sub-zone (sub-zone V) and South Eastern Plateau sub-zone (sub-zone VI).

Jharkhand state being different from many other states of the country with respect to topography, physiography and climatic conditions needs special emphasis on water management, particularly

the rain water management in conjunction with underground water management. The state enjoying nature's favour receives fairly a high amount of annual rainfall, on an average 1234.3 mm annually, of which more than 80% is instantly lost through high speed surface and sub-surface run-off causing loss of fertile soil and gully formation. With this much of annual rainfall 79.71 lakh hectares land area of Jharkhand receives about 111 TMC rain water annually. In absence of proper and adequate water management practices crop failure during rainy season in case of prolonged dry spells, virtually no crop during rabi season and drinking/domestic water crisis in the months of March-mid June have become a common feature in Jharkhand. Mainly for the water related limitations, the Agriculture in the state of Jharkhand becomes a gamble, almost solely depending on the mercy of monsoon rainfall. Delay in monsoon onset, frequent dry spells during crop season (early and mid season droughts/dry spells) initiated the need of District Agriculture Contingency Plans (DACP). Hence, with the joint efforts of ICAR and BAU-DACPs were prepared for Jharkhand state in 2010-12. In view of the marked climate change and variations in monsoon rainfall in recent years the existing DACPs required to be updated precisely. Accordingly, Ministry of Agriculture, Govt. of India and ICAR-CRIDA requested to setup a Contingency Cell in BAU to update existing DACPs and to prepare DACPs for newly constituted districts. Birsa Agricultural University constituted a Contingency Cell at BAU with Director Research as Chairman and Dr. A. Wadood as nodal officer along with all Associate Directors of 3 Zonal Research Stations and programme coordinators/Heads of all KVks.

As proposed by ICAR-CRIDA, through DR. K.V. Rao, convenor Agriculture Contingency Cell, CRIDA an MoU was signed between ICAR-CRIDA and BAU on 12.06.2017. ICAR-CRIDA released a fund of Rs 3 lakhs (in two installments) to meet the expenses on DACP updation/modification.

Contingency Cell of BAU organised a series of meetings were conducted at ZRSs (on 20.04.2017 at ZRS Darisai, on 29.04.2017 at ZRS Chianki and on 23.06.2017 at ZRS Dumka) in which most of the stakeholders (ZRS and KVK scientists, officials of state department and farmers) of altogether 9 districts took part in day long discussions. Hard copies of existing DACPs were handed over to programme coordinators/heads of KVks of the rest districts to go through and give appropriate suggestions/advices.

A 3 days meeting was convened in the Department of Agrometeorology and Environmental Science (AMES), BAU on 25-27 September, 2017 having detailed discussions on contingency plans with active participation of Dr. A. Wadood, Dr. D. N. Singh, Dr. Ramesh Kumar, Dr. Pragyan Kumari, Sri Binod Kumar, Sri Sanjiv Kumar, Dr. BK Agarwal, Dr. PB Saha, Dr. MK Barnwal, Dr. Sudhir Kr. Jha, Dr. Shankar Kumar Singh, Dr. Ranjay Kr. Singh, Dr. Pramod Kumar, Dr. Devkant Prasad, Dr. Aarti Beena Ekka, Dr. Amrit Kr. Jha, Sri LK Das, and Dr. Ansar Ahmad. All the DACPs of 24 districts were updated/modified.

Again a 13-days brain storming exercise was done on 18-30 December, 2017 at AMES with a smaller expert-group consisting of Dr. MK Barnwal, Dr. Majid Ansari, Dr. Amrit Kr. Jha, Dr. Pramod Kumar, Dr. Ansar Ahmad, Dr. PK Singh, Dr. SN Karamakar, and Dr. Krishna Prasad along with Dr. A. Wadood and Sri Binod Kumar. All the DACPs were minutely discussed and modified in the light of discussions and obtained inputs. Soft copies of DACPs were sent to Dr. KV Rao and received back with his comments for printing. Contributions of all the scientists, farmers and supports received from ICAR-CRIDA (financial as well as technical) are gratefully acknowledged. Sri Binod Kumar (RA GKMS, Darisai) deserves special appreciation for his consistent efforts in making the DACPs in the present form.

The DACPs have been prepared/updated/modified for different conditions of monsoon and farm situations prevailing in different districts in a little different format as provided by ICAR-CRIDA to suit the conditions of the state. The DACPs have been published in 5 volumes, one for each administrative Divisions comprising the districts under each Division.


(A. Wadood)

DIRECTORATE OF RESEARCH
BIRSA AGRICULTURAL UNIVERSITY, RANCHI



MESSAGE



Dr. D. N. Singh
Director Research

Jharkhand state, carved out from undivided Bihar, came into existence on 15th Nov. 2000 and became 28th state of this country. The state occupies 46% of the geographical area of undivided Bihar. The geographical area of Jharkhand state is 79.7 lakh ha, out of that 38 lakh ha is cultivable but cultivation is done only in 28 lakh ha which is 35.13% of the total geographical area of the state. The state is predominantly a rainfed and dry land system of agriculture is prevailing because only 12% area is under assured irrigation. The land is of different type starting from Tanr to Don III, Don II and Don I. Nearly 60% of the area under cultivation is acidic and organic carbon ranges from 0.3 to 0.5 % coupled with poor water holding capacity of the soils. Because of these two characteristics (rainfed agriculture and acidity of soil) the cropping intensity of state is hardly 125%, the lowest in the country.

As per classification of world bank aided National Agricultural Research project (NARP), the entire country has been divided into 15 agroclimatic zones, out of which Bihar and Jharkhand falls under zone VII (Eastern hill and Plateau region). Again, the zone VII has been sub divided into six subzones of which zone I, II and III are falling in Bihar whereas subzone IV, V and VI is in the Jharkhand. Subzone IV is largest subzone consisting of 15 districts of the state and also known as a central north eastern plateau, whereas subzone V is consisting of seven districts and known as Western Plateau. The subzone VI is smallest zone of the state and consists of only 3 districts, known as South Eastern Plateau sub-zone.

Considering the above characteristics of agro ecological conditions existing in the state of Jharkhand the contingency crop planning is very important in view of increasing the cropping intensity as well as productivity and profitability per unit area per unit time. To meet the above challenges diversification of crops is recommended which is the need of time. In uplands, farmers are cultivating direct seeded Rice which needs to be gradually replaced with drought tolerant crops like finger millet, soybean and black gram etc, these crops could be easily and successfully grown in uplands in place of rice. These crops, being climate resilient ones, are not only helpful in providing better yield under drought like situations but are also useful in providing nutritional security as well. Further, Don III (medium upland) is also required to be diversified where transplanting of rice must be replaced with pigeonpea, maize, sorghum and pearl millet with a caution that sowing of above crop has to be done after following the ridge and furrow and broad bed furrow method. This will also provide food, nutrition and livelihood security. The hybrid rice cultivation needs to be promoted further in Don II (medium land) and Don I (low land) for harvesting better yield of rice under changing climatic condition. Cultivation of vegetables and flowers are required to be promoted under poly house condition which may enable for production of off season flower and vegetable for fetching good prices from the market which would ultimately be helpful in increasing the profitability and income of the farmers.

While congratulating Dr. Wadood and his team for preparing such useful documents I urge the farmers and other stakeholders of respective districts of Jharkhand to make fullest use of the District Agriculture Contingency Plans (DACP) for mitigating the drought and drought like situations prevailing in the state and to help farmers in doubling their income in years to come.

(D. N. Singh)



DISTRICT AGRICULTURE CONTINGENCY PLAN DACP - 2019



KOLHAN DIVISION


**East Singhbhum - Saraikela Kharsawan
- West Singhbhum**



BIRSA AGRICULTURAL UNIVERSITY, KANKE, RANCHI, JHARKHAND
AND
CENTRAL RESEARCH INSTITUTE FOR DRY LAND AGRICULTURE (CRIDA), HYDERABAD

EAST SINGHBHUM DISTRICT

Sl. No.	CONTENTS	Page No.
1.	District Agriculture profile and land use pattern	1
	PART-I	
	CONTINGENCY PLAN FOR KHARIF	
2.	Contingency plan for 2 weeks delay in monsoon arrival (onset in 4th week of June) A1, A4, A5. Upland A2, A6. Midland A3, A7. Lowland Contingency plan for 4 weeks delay in monsoon arrival (onset in 2nd week of July) B1, B4, B5. Upland B2, B6. Midland B3, B7. Lowland Contingency plan for 6 weeks delay in monsoon arrival (onset in 6th week of July) C1, C4. Upland C2, C5. Midland C3, C6. Lowland	2-11 11-20 20-26
	PART-II	
3.	A. Contingency Plan for normal monsoon onset followed by 15-20 days dry spell A1. Upland A2. Midland A3. Lowland	27-28
4.	B. Contingency plan for mid season drought Upland B1. At vegetative phase B2. At Flowering/Fruiting stage Midland B3. At vegetative phase B4. At Flowering/Fruiting stage Lowland B5. At vegetative phase B6. At Flowering/Fruiting stage	28-30
5.	C. Contingency plan for Late season drought/Terminal drought (Early withdrawal of monsoon) At fruiting/pre physiological maturity stage C1.1. Upland C1.2. Midland C1.3. Lowland	31
	PART-III	
6.	A. Unusual rains : Continuous high rainfall in a short span leading to water logging Crop management Pest and disease management	32-34
7.	B. Extreme weather events (Hail storm, Heat wave, Cold wave, Frost CONTINGENCY PLAN FOR RABI	34-35
8.	1. Sowing window information 2. Contingency measures for field crops grown with residual moisture under rainfed condition 2(A) Optimal residual moisture 2A.1 Upland 2A.2 Midland 2A.3 Lowland 2 (B) Less than optimal soil moisture (25 % less than normal-Deficit of 20-40 % rainfall) 2B.1 Upland 2B.2 Midland 2B.3 Lowland	36 36-40
	CONTINGENCY STRATEGIES FOR LIVESTOCK, POULTRY AND FISHERIES	
9.	1. Livestock a) Before the event b) During the event c) After the event 2. Poultry a) Before the event b) During the event c) After the event 3. Fisheries a) Before the event b) During the event c) After the event	41-48





District Agriculture profile

Agro-Climatic Zone	AZ - 59		
Agro Ecological Sub Region (ICAR)	Eastern plateau (chotanagpur) And Eastern Ghats, Hot Subhumid Eco-Region (12.3)		
Agro-Climatic Zone (Planning Commission)	Eastern Plateau And Hills Region (VII)		
Agro Climatic Zone (NARP)	South Eastern Plateau Sub Zone - VI		
Meteorological Subdivision	8 th		
List all the districts falling under the NARP Zone (>50% area falling in the zone)	East Singhbhum, Saraikela-Kharsawan, West Singhbhum		
Geographic coordinates of district headquarters	Latitude		Altitude
	22° 12'35" N - 23° 01' 02" N		86° 03' 59"E - 86°53'28" E 198m
Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/ RRTTS	Zonal Research Station (ZRS), Darisai, Birsa Agricultural University, Ranchi		
Mention the KVK located in the district with address	Krishi Vignan Kendra, Darisai, Vill-Kharia, PO. Barakhurshi Distt. East Singhbhum-832 304		
Name and address of the nearest Agromet Field Unit (AMFU, IMD) for agro-advisories in the Zone	ZRS, Darisai, Eas Singhbhum-832 304		

Land use pattern of the district (area: '000 ha)

Geographical area	Cultivable area	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable wasteland	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
556.697	161.577	124.846	138.649	2.242	40.156	5.416	46.763	86.654	37.048



CONTINGENCY PLAN FOR KHARIF

PART-I

A Monsoon/Weather Situation: 2 Weeks Delay (Onset: 4th Week of June) – Early Season Drought

A1. Major Farming Situation/Land Situation: Upland High rainfall, shallow iron rich, light textured sandy & acidic soil.	
Normal Crop/cropping system	Upland Rice, Pigeonpea, Pigeonpea + Maize, Greengram, Blackgram, Maize+ Creeper vegetables
Suggested Contingency measures	
a) Change in crop/cropping system	
<p><u>Discard Rice crop</u> <u>Sole</u> Pigeonpea, Maize, Soybean, Groundnut , Blackgram, Sorghum, Finger Millet <u>Intercrop</u> Pigeonpea + Groundnut (1:2) , Pigeonpea + Maize (1:1), Pigeonpea + Lady's finger(1:2)/Greengram (1:2), Maize+ Creeper vegetables (Sponge Gourd) (1:2) <u>Horticulture Crop</u> Vegetable : Amaranthus leaf/ Coriander leaf/ Early Cabbage/ Cauliflower/ French bean/ Creepers Variety Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250) Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Vivek hybrid 9 (80) Soybean- JS 9752 (100), Birsa soybean 1 black(120-125), Birsa safed soybean 2 (105-110), RKS 18 Groundnut- Birsa mungfali 3, 4, Girnar 3 Blackgram- Birsa urd 1 (75-80), PU 19 (70-75), WBU 109 (70-75), Sorghum- MP chari, CSV 1616 Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28 Greengram-HUM 16, SML 668 <u>Vegetable crops</u> Lady's finger- Varsa uphar, Hybrid- Sonal, Sarika Coriander- Pant haritima, Rajendra swati Cabbage- Golden acer, Early drumhead Cauliflower-Pusa katki, Pusadipali, Early synthetic, Pusa him jyoti, Pant subhra, Endum early Pusa hybrid 1 Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata Cucurbits- Bitter gourd- Pusa domausami, Bottle gourd-PusaMeghdoot, Coimbtor long green, local, Arka harit Sponge gourd- Long green,Long white Ridge gourd- Swarn manjari, Swarn uphar, Satputia, Red Pumpkin- Arka chandan, Arka suryamukhi</p>	
b) Agronomic measures	
<ul style="list-style-type: none"> • Summer deep ploughing with Mouldboard or disc • Dova construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations. • RD Spacing • Zero tillage practices • Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing • RDF and in case of Intercropping reduce 1/3rd dose for intercrop • Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin@ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables • Bund construction for unbundled upland • Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables • Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables • Inter-cropping to meet the consequences of occasional Drought. • Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed. 	



- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite application for pulses and oilseed @ 3-5 q/ha in furrow at the time of sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha, PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure

c) Remarks on Implementation

- Linkage with RKVY, ATMAS, and NFSM
- Vermicomposting through KVKS ATMAS and NHM
- Goatry and poultry rearing through KVKS, ATMAS and Veterinary Dept of. Govt. and BAU for livelihood support.
- Awareness about balanced use of fertilizers to increase their fertility, productivity and sustainability
- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology.
- Awareness for more and more use of organic manures, bio-pesticides for organic cultivation with IFS (eight components linkages)
- Upland- 15-20 % upland area should be covered with orchard

1. Mango based orchard-

Variety- Amrapali (30 June-5 July), Mallika (15-20 June regular bearer), Sunder langra(15-20 May)
Spacing- 5 m X 5m

i) Recommended package of Practices- Intercrops

- Mango + Papaya (Filler crop for two years) + Blackgram (rainy)/ Chickpea
- Mango + Custard apple (for 10 years and renovate or remove after 10 years) + Blackgram/Chickpea

Variety- Langra (15 June)/Bombay green(15 May)/ Himsagar (20-25 May irregular bearer),

Spacing- 10 m X 10m

ii) Recommended package of practices

- Mango + Guava(Up to 10 years as filler) + Papaya (Less than 3 years) + Blackgram-Chickpea/Lentil
- Mango + Lemon + Papaya + Rabi pulses/vegetables
- Mango + Custard apple + Papaya + Blackgram – Pea/Chickpea/Lentil/ Vegetables

2. Guava base orchard-

Variety- Arka Mridula, Pant Prabhat, Allahabad safeda, L 49

Spacing- 5m X 5m

Recommended package of practices- Intercrops

- Guava + Papaya (For 3 years) + Blackgram-Chickpea
- Guava + Custard apple + Blackgram/Soybean- Pea/Vegetables

3. Ber Based Orchard –

Variety- Banarsi, Karakka, Gola, Apple ber

Spacing- 5m X 5m

Recommended package of practices Intercrops

Ber + Custard apple + Sesame/Blackgram- Toria/Linseed/Safflower

N.B.- Cucurbits, beans or any creeper or climber vegetable should be avoided

- Field crops having height more than one meter should be avoided such as Pigenpea, Maize, Sorghum
- After 3-5 years when shading effects started shade loving crops like ginger, Turmeric, OI or leafy vegetables should be grown
- In citrus leaf minor and aphid susceptible crops should be avoided
- Aphid should be managed of mustard /toria taken in citrus orchard

4. Cashewnut based orchard for Kolhan region

5. Cassava should be grown for the requirement as feed for pig animals

- 6. Moringa should also be grown as fodder or vegetable purpose on upland main field bunds as shelter belt/ wind break. Every year pruning and thinning should be followed for bushy look.

A2. Major Farming Situation/Land Situation: Midland High rainfall, shallow iron rich, light textured sandy & acidic soil

Normal Crop/cropping system	Rice-fallow, Finger millet, Pigeonpea, Sorghum, Maize, Pigeonpea + Sorghum Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures



a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var. –Arize 6444 (Gold), Arize Tej (Gold), PAC 801, PAC 807

Don 3

Ridge and Furrow method or raise bed broad furrow: Replace Rice with cereal/ Pulse/ Vegetable

Cereal – Maize/ Sorghum

Pulse- Pigeonpea + Lady's Finger(1:2) / Groundnut(1:2)/ Soybean(1:2)/ Finger millet (1:1)

Vegetable-Amaranthus leaf/Coriander leaf/ Lady's Finger/ Cowpea/ Dolichos bean/

Variety

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 – Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Groundnut- Birsa mungfali 3, 4, Girnar 3

Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149

Vegetable crops

Coriander- Pant haritima, Rajendra swati

Lady's finger- Pusa A 4, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati, Swarn harit

Dolichos bean-Swarna utkrist, Swarna rituwar

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INP
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P_2O_5 + 1 kg K_2O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P_2O_5 : K_2O (Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P_2O_5 + 40 K_2O /ha ((Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K_2O
- DSR-Use plastic drum seeder rice tools
- Follow RDF and
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /ltr water, Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/ltr. water; Blackgram and Greengram-Leaf minor- Monocrotophos @ 1ml/ltr. water, Mosaic- Methyl Demeton @ 1.5 ml/ltr. water; S vegetables- Nursery management Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/ltr. water.

c) Remarks on Implementation

- A campaign through RKVY, ATMAs, NFSM, KVKS, NHM and other State Govt. line departments are needed to be launched through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.



- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

A3. Major Farming Situation/Land Situation: Lowland High rainfall, shallow iron rich , light textured sandy and acidic soil

Normal Crop/cropping system	Rice -Fallow
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2</u> in <u>Don 1</u> DSR(Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days)	
Transplanting (Hybrid rice varieties) Var-Arize 6444 (Gold), Uday 111, PHB 71, 26P52, 25P25	
b) Agronomic Measures	
<ul style="list-style-type: none"> • Staggered Nursery raising by MAT/ DAPOG method • Follow community based nursery raising • Follow RDF,INPM • Use Post emergence weedicide • Use early to mid early duration of rice variety. • Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² • Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice • Topdressing above mentioned dose 10-15 days after sowing • In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling • Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm • Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering. • DSR-Use plastic drum seeder rice tools • Use of post weedicide • Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/l. water Gundhi bug, leaf folder and BPH -Quinolophos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /lt water 	

4 (c) Remarks on Implementation

Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme

Supply of Plastic drum seeder through line departments

Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.

Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.

Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates

Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation.

A4. Major Farming Situation/Land Situation: Upland High rainfall, shallow depth, light textured sandy & acidic soil.

Normal Crop/cropping system	Upland Rice , Pigeonpea, Pigeonpea + Maize, Greengram, Blackgram, Maize+ creeper vegetables
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Rice crop	
<u>Sole</u>	
Pigeonpea, Maize, Soybean, Groundnut , Blackgram, Sorghum, Finger Millet	
<u>Intercrop</u>	
Pigeonpea + Groundnut (1:2) , Pigeonpea + Maize (1:1), Pigeonpea + Lady's finger(1:2)/Greengram (1:2), Maize+ Creeper vegetables (Sponge Gourd) (1:2)	
<u>Horticulture Crop</u>	
Vegetable : Amaranthus leaf/ Coriander leaf/ Early Cabbage/ Cauliflower/ French bean/ Creepers_ Variety	
Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250)	



Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Vivek hybrid 9 (80)
Soybean- JS 9752 (100), Birsa soybean 1 black(120-125), Birsa safed soybean 2 (105-110), RKS 18
Groundnut- Birsa mungfali 3, 4, Girnar 3
Blackgram- Birsa urd 1 (75-80), PU 19 (70-75), WBU 109 (70-75),
Sorghum- MP chari, CSV 1616
Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28
Greengram-HUM 16, SML 668
Horticulture Crop

Vegetable :Coriander leaf/ Early Cabbage/ Cauliflower/ French bean, Cowpea/ Creepers
Variety

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250)
Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Vivek hybrid 9 (80)
Soybean- JS 9752 (100), Birsa soybean 1 black(120-125), Birsa safed soybean 2 (105-110), RKS 18
Groundnut- Birsa mungfali 3, 4, Girnar 3
Blackgram- Birsa urd 1 (75-80), PU 19 (70-75), WBU 109 (70-75),
Sorghum- MP chari, CSV 1616
Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28
Lady's finger- Varsa uphar, Hybrid- Sonal, Sarika
Greengram-HUM 16
SML 668

Vegetable crops

Coriander- Pant haritima, Rajendra swati
Cabbage- Golden acer, Early drumhead
Cauliflower-Pusa katki, Pusadipali, Early synthetic, Pusa him jyoti, Pant subhra, Endum early Pusa hybrid 1
Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata
Cucurbits-
Bitter gourd- Pusa domausami,
Bottle gourd-PusaMeghdoot, Coimbtor long green, local, Arka harit
Sponge gourd- Long green, Long white
Ridge gourd- Swarn manjari, Swarn uphar, Satputia,
Red Pumpkin- Arka chandan, Arka suryamukhi

b) Agronomic measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbundled upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite application for pulses and oilseed @ 3-5 q/ha in furrow at the time of sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure



c) Remarks on Implementation

- Linkage with RKVY, ATMAs, and NFSM
- Vermicomposting through KVKS ATMAs and NHM
- Goatry and poultry rearing through KVKS, ATMAs and Veterinary Dept of. Govt. and BAU for livelihood support.
- Awareness about balanced use of fertilizers to increase their fertility, productivity and sustainability
- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology.
- Awareness for more and more use of organic manures, biopesticides for organic cultivation with IFS (eight components linkages)
- Upland- 15-20 % upland area should be covered with orchard

1. Mango based orchard-

Variety- Amrapali (30 June-5 July), Mallika (15-20 June regular bearer), Sunder langra(15-20 May)

Spacing- 5 m X 5m

i) Recommended package of Practices- Intercrops

a) Mango + Papaya (Filler crop for two years) + Blackgram (rainy)/ Chickpea

b) Mango + Custard apple (for 10 years and renovate or remove after 10 years) + Blackgram/Chickpea

Variety- Langra (15 June)/Bombay green(15 May)/ Himsagar (20-25 May irregular bearer),

Spacing- 10 m X 10m

ii) Recommended package of practices

a) Mango + Guava(Up to 10 years as filler) + Papaya (Less than 3 years) + Blackgram-Chickpea/Lentil

b) Mango + Lemon + Papaya + Rabi pulses/vegetables

c) Mango + Custard apple + Papaya + Blackgram – Pea/Chickpea/Lentil/ Vegetables

2. Guava base orchard-

Variety- Arka Mridula, Pant Prabhat, Allahabad safeda, L 49

Spacing- 5m X 5m

Recommended package of practices- Intercrops

a).Guava + Papaya (For 3 years) + Blackgram-Chickpea

b).Guava + Custard apple + Blackgram/Soybean- Pea/Vegetables

3. Ber Based Orchard –

Variety- Banarsi, Karakka, Gola, Apple ber

Spacing- 5m X 5m

Recommended package of practices Intercrops

Ber + Custard apple + Sesame/Blackgram- Toria/Linseed/Safflower

N.B.- Cucurbits, beans or any creeper or climber vegetable should be avoided

- Field crops having height more than one meter should be avoided such as Pigenpea, Maize, Sorghum
 - After 3-5 years when shading effects started shade loving crops like ginger, Turmeric, OI or leafy vegetables should be grown
 - In citrus leaf minor and aphid susceptible crops should be avoided
 - Aphid should be managed of mustard /atoria taken in citrus orchard
4. Cashewnut based orchard for Kolhan region
5. Cassava should be grown for the requirement as feed for pig animals
6. Moringa should also be grown as fodder or vegetable purpose on upland main field bunds as shelter belt/ wind break. Every year pruning and thinning should be followed for bushy look.

A5. Major Farming Situation/Land Situation: Upland Less rainfall, shallow depth, light textured sandy & acidic soil

Normal Crop/cropping system	Upland Rice, Pigeonpea, Maize, Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Discard Rice Crop

Sole crop

Pigeonpea, Maize, Soyabean, Groundnut, Blackgram, Sorghum, Finger millet

Intercrop

Pigeonpea + Lady's Finger (1:2), Pigeonpea + Sorghum(1:1), Pigeonpea + Groundnut (1:2), Pigeonpea + Maize (1:1)

Horticulture Crop

Vegetable :Radish/Early Cabbage/ Cauliflower/ French bean / Cowpea

Variety

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250)

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Vivek hybrid 9 (80)



Soybean- JS 9752 (100), Birsa soybean 1 black(120-125), Birsa safed soybean 2 (105-110), RKS 18

Groundnut- Birsa mungfali 3, 4, Girnar 3

Blackgram- Birsa urd 1 (75-80), PU 19 (70-75), WBU 109 (70-75),

Sorghum- MP chari, CSV 1616

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28

Lady's finger- Varsa uphar, Hybrid- Sonal, Sarika

Greengram-HUM 16, SML 668

Vegetable crops

Radish- Pusa chetki (summer), Pusa deshi, Kashi hansi, Jaunpur/ Pusa himani, Japanese white, Pusa roshni

Coriander- Pant haritima, Rajendra swati

Cabbage- Golden acer, Early drumhead

Cauliflower-Pusa katki, Pusadipali, Early synthetic, Pusa him jyoti, Pant subhra, Endum early Pusa hybrid 1

Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata

Cucurbits-

Bitter gourd- Pusa domausami,

Bottle gourd-PusaMeghdoot, Coimbatore long green, local, Arka harit

Sponge gourd- Long green, Long white

Ridge gourd- Swarn manjari, Swarn uphar, Satputia,

Red Pumpkin- Arka chandan, Arka suryamukhi

Cowpea- bushy- CP 4, Arka garima, Pusa komal

b) Agronomic measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathypor or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite application for pulses and oilseed @ 3-5 q/ha in furrow at the time of sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure

c) Remarks on Implementation

- Linkage with RKVY , ATMAS, and NFSM
 - Vermicomposting through KVVs ATMAS and NHM
 - Goatry and poultry rearing through KVVs, ATMAS and Veterinary Dept of. Govt. and BAU for livelihood support.
 - Awareness about balanced use of fertilizers to increase their fertility, productivity and sustainability
 - A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology.
 - Awareness for more and more use of organic manures, biopesticides for organic cultivation with IFS (eight components linkages)
 - Upland- 15-20 % upland area should be covered with orchard
1. Mango based orchard-
Variety- Amrapali (30 June-5 July), Mallika (15-20 June regular bearer), Sunder langra(15-20 May)
Spacing- 5 m X 5 m



- i) Recommended package of Practices- Intercrops
a) Mango + Papaya (Filler crop for two years) + Blackgram (rainy)/ Chickpea
b) Mango + Custard apple (for 10 years and renovate or remove after 10 years) + Blackgram/Chickpea
Variety- Langra (15 June)/Bombay green(15 May)/ Himsagar (20-25 May irregular bearer),
Spacing- 10 m X 10m
- ii) Recommended package of practices
a) Mango + Guava(Upto 10 years as filler) + Papaya (Less than 3 years) + Blackgram-Chickpea/Lentil
b) Mango + Lemon + Papaya + Rabi pulses/vegetables
c) Mango + Custard apple + Papaya + Blackgram – Pea/Chickpea/Lentil/ Vegetables
2. Guava base orchard-
Variety- Arka Mridula, Pant Prabhat, Allahabad safeda, L 49
Spacing- 5m X 5m
Recommended package of practices- Intercrops
a) Guava + Papaya (For 3 years) + Blackgram-Chickpea
b) Guava + Custard apple + Blackgram/Soybean- Pea/Vegetables
3. Ber Based Orchard –
Variety- Banarsi, Karakka, Gola, Apple ber
Spacing- 5m X 5m
Recommended package of practices Intercrops
Ber + Custard apple + Sesame/Blackgram- Toria/Linseed/Safflower
- N.B.- Cucurbits, beans or any creeper or climber vegetable should be avoided
- Field crops having height more than one meter should be avoided such as Pigenpea, Maize, Sorghum
 - After 3-5 years when shading effects started shade loving crops like ginger, Turmeric, Ol or leafy vegetables should be grown
 - In citrus leaf minor and aphid susceptible crops should be avoided
 - Aphid should be managed of mustard /atoria taken in citrus orchard
4. Cashewnut based orchard for Kolhan region
5. Cassava should be grown for the requirement as feed for pig animals
6. Moringa should also be grown as fodder or vegetable purpose on upland main field bunds as shelter belt/wind break. Every year pruning and thinning should be followed for bushy look.

A6. Major Farming Situation/Land Situation: Midland Less rainfall, shallow depth, light textured sandy & acidic soil.

Normal Crop/cropping system	Rice, Pigeonpea, Maize Pigeonpea + Maize , Pigeonpea + Sorghum
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var. – Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 27P31, DRRH 2

Don 3

Ridge and Furrow method or raise bed broad furrow: Replace Rice with cereal/ Pulse/ Vegetable
Cereal – Maize/ Sorghum

Pulse- Pigeonpea + Lady's Finger (1:2) / Groundnut (1:2)/ Soybean (1:2)/ Finger millet(1:1)

Vegetable-Radish/ Lady's Finger/ Cowpea/ Dolichos bean

Variety

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 – Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Groundnut- Birsa mungfali 3, 4, Girnar 3

Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149

Vegetable crops

Lady's finger- Pusa A 4, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati, Swarn harit

Dolichos bean-Swarna utkrist, Swarna rituwar



4 (b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INP
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O
- DSR-Use plastic drum seeder rice tools
- Follow RDF and
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /lit water, Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVks, NHM and other State Govt. line departments are needed to be launched through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

A7. Major Farming Situation/Land Situation: Lowland Less rainfall, shallow depth , light textured sandy and acidic soil

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	

Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of Don 2 in Don 1
 DSR(Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), MTU-1001, MTU 1010, Transplanting (Hybrid rice varieties) Var. – Arize 6444 (Gold), Uday 111, PHB 71, 26P52, 25P25, 27P31, 27P36

b) Agronomic Measures

- Staggered Nursery raising by MAT/ DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use Post emergence weedicide
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm



- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water Gundhi bug, leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /ltr water

c) Remarks on Implementation

- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation.

B. Monsoon/Weather Situation: 4 Weeks Delay (Onset: 2nd Week of July) – Early Season Drought

B1. Major Farming Situation/Land Situation: Upland High rainfall, shallow iron rich, light textured sandy & acidic soil.	
Normal Crop/cropping system	Upland Rice , Pigeonpea, Pigeonpea + Maize, Maize, Greengram and Blackgram
Suggested Contingency measures	
a) Change in crop/cropping system	
<u>Discard Rice Crop</u>	
<u>Sole</u>	
Maize, Sorghum, Blackgram, Groundnut, Soybean, Finger millet	
<u>Intercrop</u>	
Pigeonpea + Groundnut (1:2), Pigeonpea + Maize (1:1), Pigeonpea + Sorghum (1:1)	
<u>Horticulture Crop</u>	
Vegetable- Tomato/ Brinjal/ Chili/ Amaranthus leaf/ Lobia/ French bean/ Creepers	
<u>Fodder Crop</u>	
Sorghum, Maize	
<u>Variety</u>	
Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), VMH 4106 (Sweet corn hybrid), Sorghum- CSV 20-110-20, CSV 1616	
Blackgram- Birsa urd 1 (75-80), PU 35 (70-75), Uttara (75-80 small grain)	
Groundnut- Birsa mungfali 3, 4, Girnar 3	
Soybean- JS 9752 (100), JS 335, Birsa safed soybean 2 (105-110), RAUS 5	
Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67	
<u>Vegetable crops</u>	
Tomato- Swarn samridh, Pusa hybrid 1 Suraksha	
Brinjal- Banaras giant, Swarn pratibha, Swarn mani, hybrid-Swarn shakti , Vijay, Swarna sampada 6	
Chili- Spices- Andhrajyoti, Pusasadabahar, Jwala, California wonder, Bharat	
Cowpea- bushy- Pusa komal, Pusa barsati Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal,	
<u>Cucurbits-</u>	
Bitter gourd- Pusa domausami,	
Bottle gourd-PusaMeghdoot, Coimbtor long green, local, Arka harit	
Sponge gourd- Pusa chikni, Long green,Long white	
Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha,	
Red Pumpkin- CO 1, CO 2, Arka suryamukhi	
<u>Fodder crop</u>	
Sorghum- Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid	
Maize- African tall, JS-1006 and Vijaya composite.	
b) Agronomic Measures	
<ul style="list-style-type: none">• Summer deep ploughing with Mouldboard or disc• Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.	

- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbundled upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha, PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demoton @ 1.5 ml/t. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demoton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water, Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/lit. water or Cartap hydrochloride @ 2 gm/lit. water, hairy caterpillar –Quinolphos 1.5ml /lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuran 3G @ 3 gm/m² before 10 days of transplanting followed by application of Tricoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- Linkage with RKVY , ATMAs and NFSM
- Vermicomposting Awareness through KVks, ATMAs and NHM
- Backyard Goatry and poultry rearing Awareness campaign through KVks, ATMAs and Veterinary Dept of. Govt. and BAU for livelihood support.
- A special programme is needed to be launched in such areas on priority basis to motivate the farmers to adopt improved technology for stress management through ATMAs, KVks, Govt Dept., NGOs
- Campaign for Awareness of crop-weather insurance to meet the losses due to drought/cyclone like weather vagaries.

B2. Major Farming Situation/Land Situation: Midland High rainfall, shallow iron rich, light textured sandy & acidic soil

Normal Crop/cropping system	Rice, Finger millet, Pigeonpea, Sorghum, Maize, Pigeonpea + Sorghum, Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR



Transplanting (Hybrid Rice varieties) Var.- Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807 Don 3

Ridge and Furrow method or raise bed broad furrow: Replace rice with Cereal/Pulses/ vegetable/ Fodder crop
Cereal – Maize/Sorghum

Pulses- Pigeonpea+ Lady's finger (1:2)/Blackgram (1:2)/Soybean (1:2)/ Groundnut(1:2)

Vegetables- Lady's Finger/ Cowpea/ Dolichos bean/

Fodder Crop

Brachiaria grass/ Rice bean (Moth bean)/ Maize / Cowpea

Variety

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 – Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Blackgram- Birsa urd 1 (75-80), PU 31 (70-75), WBU 109 (70-75)

Soybean- Birsa soybean 1 black(120-125), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Groundnut- Birsa mungfali 3, 4, Girnar 3

Vegetable crops

Lady's finger- Pusa A 4, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati, Swarn harit

Dolichos bean-Swarna utkrist, Swarna rituwar

Fodder crop

Maize- African tall, JS-1006 and Vijaya composite.

Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility.
- Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.

- Gap filling and resowing should be done if mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
 - Weed control by applying pre-emergence 5-6 DAS (Pendimethalin) or Post-emergence 18-28 DAS (Bispyribac).
 - Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/lit. water or Cartap hydrochloride @ 2 gm/lit. water, hairy caterpillar –Quinolphos 1.5ml /lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water.
- Rice pest and disease management -Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /lt water. Termite- Methyl parathion dust @ 25 kg/ha

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be awared through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

B3. Major Farming Situation/Land Situation: Lowland High rainfall, shallow iron rich , light textured sandy and acidic soil

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	

Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of Don 2 in Don 1

DSR(Improved Rice variety) var.-- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days)

Transplanting (Hybrid rice varieties)Var.-Arize 6444 (Gold), Uday 111, PHB 71, 26P52, 25P25

b) Agronomic Measures

- Staggered Nursery raising by MAT/ DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use Post emergence weedicide
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /lt water

c) Remarks on Implementation

- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation

B4. Major Farming Situation/Land Situation: Upland High rainfall, shallow depth, light textured sandy and acidic soil.

Normal Crop/cropping system	Upland Rice, Pigeonpea, Maize, Pigeonpea + Maize (Mixed) Sorghum
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Discard Rice crop

Sole crop

Pigeonpea, Blackgram, Greengram, Soybean , Groundnut, Sorghum

Intercrop

Pigeonpea + Blackgram/ Greengram (1:2), Pigeonpea + Groundnut (1:2), Pigeonpea + Maize (1:1)

Pigeonpea + Sorghum (1:1), Maize + Lady's finger (1:2)

Horticulture Crop

Vegetable- Coriander leaf/ Early Cabbage/ Cauliflower/ French bean, Cowpea/ Creepers

Variety

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), ICPH 2671 (200)

Blackgram- Birsa urd 1 (75-80), PU 35 (70-75), Uttara (75-80 small grain)

Greengram- HUM 16, IPM-02-03-60-65, SML 668

Soybean- JS 9752 (100), JS 335, Birsa safed soybean 2 (105-110), RAUS 5

Groundnut- Birsa mungfali 3, 4, Girnar 3

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), LG 32-81 –Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Vegetable crops-

Coriander- Pant haritima, Rajendra swati

Cabbage- Golden acer, Early drumhead

Cauliflower-Pusa katki, Pusadipali, Early synthetic, Pusa him jyoti, Pant subhra, Endum early Pusa hybrid 1

Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata

Cucurbits

Bitter gourd- Pusa domausami,

Bottle gourd-PusaMeghdoot, Coimbtor long green, local, Arka harit

Sponge gourd- Long green,Long white

Ridge gourd- Swarn manjari, Swarn uphar, Satputia,

Red Pumpkin- Arka chandan, Arka suryamukhi

Fodder crop

Sorghum- Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

Maize- African tall, JS-1006 and Vijaya composite.

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables

- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like: Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/lit. water or Cartap hydrochloride @ 2 gm/lit. water, hairy caterpillar -Quinolhos 1.5ml /lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuran 3G @ 3 gm/m² before 10 days of transplanting followed by application of Tricoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- Linkage with RKVY , ATMAs and NFSM
- Vermicomposting Awareness through KVks, ATMAs and NHM
- Backyard Goatry and poultry rearing Awareness campaign through KVks, ATMAs and Veterinary Dept of. Govt. and BAU for livelihood support.
- A special programme is needed to be launched in such areas on priority basis to motivate the farmers to adopt improved technology for stress management through ATMAs, KVks, Govt Dept., NGOs
- Campaign for Awareness of crop-weather insurance to meet the losses due to drought/cyclone like weather vagaries.

B5. Major Farming Situation/Land Situation: Upland Less rainfall, shallow depth, light textured sandy and acidic soil.	
Normal Crop/cropping system	Upland Ric, Pigeonpea, Maize, Pigeonpea + Maize (Mixed)
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Rice crop <u>Solecrop</u> Pigeonpea, Blackgram, Greengram, Soybean, Groundnut, <u>Intercrop</u> Pigeonpea + Groundnut(1:2), Pigeonpea + Maize 1:1), Pigeonpea + Sorghum (1:1) <u>Horticulture Crop</u> Vegetable- Radish/Early Cabbage/ Cauliflower/ French bean / Cowpea (Lobia) <u>Fodder Crop</u> Cowpea/Sorghum/ Maize/ Blackgram <u>Variety</u> Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), ICPH 2671 (200) Blackgram- Birsa urd 1 (75-80), PU 35 (70-75), Uttara (75-80 small grain) Greengram- HUM 16, IPM-02-03-60-65, SML 668 Soybean- JS 9752 (100), JS 335, Birsa safed soybean 2 (105-110), RAUS 5	



Groundnut- Birsa mungfali 3, 4, Girnar 3

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), LG 32-81 –Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Vegetable crops

Radish- Pusa chetki (summer), Pusa deshi, Kashi hansi, Jaunpur/ Pusa himani, Japanese white, Pusa roshni
Coriander- Pant haritima, Rajendra swati

Cabbage- Golden acer, Early drumhead

Cauliflower-Pusa katki, Pusadipali, Early synthetic, Pusa him jyoti, Pant subhra, Endum early Pusa hybrid 1

Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata

Cowpea- bushy- CP 4, Arka garima, Pusa komal

Fodder Crop

Cowpea- UPC-5286, GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Sorghum Sudan hybrid

Maize- African tall, JS-1006 and Vijaya composite

Blackgram- Uttara (75-80)

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin@ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbundled upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/l. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/l. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/l. water, Mosaic- Methyl Demeton @ 1.5 ml/l. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/l. water, Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/l. water or Cartap hydrochloride @ 2 gm/l. water, hairy caterpillar –Quinolphos 1.5ml / lt. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt; vegetables- Nursery management- Application of carbofuran 3G @ 3 gm/m² before 10 days of transplanting followed by application of Tricoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/l. water



c) Remarks on Implementation

- Linkage with RKVY , ATMAs and NFSM
- Vermicomposting Awareness through KVKS, ATMAs and NHM
- Backyard Goatry and poultry rearing Awareness campaign through KVKS, ATMAs and Veterinary Dept of Govt. and BAU for livelihood support.
- A special programme is needed to be launched in such areas on priority basis to motivate the farmers to adopt improved technology for stress management through ATMAs, KVKS, Govt Dept., NGOs
- Campaign for Awareness of crop-weather insurance to meet the losses due to drought/cyclone like weather vagaries.

B6. Major Farming Situation/Land Situation: Midland Less rainfall, shallow depth , light textured sandy and acidic soil

Normal Crop/cropping system	Rice, Pigeonpea, Maize Pigeonpea + Maize, Pigeonpea + Sorghum, Rice-vegetable
-----------------------------	---

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 27P31, DRRH 2

Don 3

Ridge and Furrow method or raise bed broad furrow along the slope :_Replace rice with Cereal/Pulses/ vegetable/ Fodder crop

Cereal – Maize/Sorghum

Pulses- Pigeonpea+ Lady's Finger/Blackgram/Soybean/ Groundnut/Maize

Vegetables- Lady's Finger/ Cowpea/ Dolichos bean/

Fodder Crop -

Rice bean (Moth bean)/ Maize / Cowpea/ Sweet Sorghum

Variety

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 – Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Blackgram- Birsa urd 1 (75-80), PU 31 (70-75), WBU 109 (70-75)

Soybean- Birsa soybean 1 black(120-125), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Groundnut- Birsa mungfali 3, 4, Girnar 3

Maize- Birsa makka (Vikash) 2 (75-80), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)

Vegetable crops

Lady's finger- Pusa A 4, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati, Swarn harit

Dolichos bean-Swarna utkrist, Swarna rituwar

Fodder crop

Maize- African tall, JS-1006 and Vijaya composite.

Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbundled upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables



- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility.
- Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like: Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder-Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water, Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/lit. water or Cartap hydrochloride @ 2 gm/lit. water, hairy caterpillar -Quinolphos 1.5ml /lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuran 3G @ 3 gm/m² before 10 days of transplanting followed by application of Tricoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water.
- Rice pest and disease management -Gundhi bug, leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /lt water. Termite- Methyl parathion dust @ 25 kg/ha

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be awarded through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

B7. Major Farming Situation/Land Situation: Lowland Less rainfall, shallow depth, light textured sandy and acidic soil	
Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2</u> in <u>Don 1</u>	
DSR(Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), MTU-1001, MTU 1010, Transplanting (Hybrid rice varieties) Var.– Arize 6444 (Gold), Uday 111, PHB 71, 26P52, 25P25, 27P31, 27P36	
<u>Vegetable</u>	
Cucurbits-	
Bitter gourd- Arka hait, Pusa domausami,	



Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, Local, Arka harit
 Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green, Long white
 Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,
 Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi

b) Agronomic Measures

- Staggered Nursery raising by MAT/ DAPOG method
- Follow community based nursery raising
- Follow RDF, INPM
- Use Post emergence weedicide
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water, Gundhi bug, leaf folder and BPH -Quinolophos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /lt water

c) Remarks on Implementation

- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation

C. Monsoon/Weather Situation: 6 Weeks Delay (Onset: 6th Week of July) – Early Season Drought

C1. Major Farming Situation/Land Situation: Upland High rainfall, shallow iron rich, light textured sandy and acidic soil.	
Normal Crop/cropping system	Upland Rice, Pigeonpea, Maize, Pigeonpea + Maize
Suggested Contingency measures	
a) Change in crop/cropping system	
<u>Discard Rice Crop</u> <u>Sole crop</u> Raised bed or ridge and furrow method: Niger, Horsegram, Sorghum, Blackgram, Soybean, Finger millet, Pigeonpea, Lady's finger, Maize <u>Intercrop</u> Pigeonpea + Horsegram (1:2), Pigeonpea + Maize (1:1) <u>Horticulture crop</u> Vegetable –Lady's Finger, Ridge Gourd, Bottle Gourd, Cowpea <u>Fodder Crop</u> Deenanath grass/ Guinea grass/ Sadabahar/ Stylo hemata/ Chara badam/ Anjan grass/ Hybrid napier/ Cactus/ Sweet Sorghum Variety Niger- Birsa niger 1, 2 and 3 (95-105) Horsegram- Birsa kulthi1 (90-95) Sorghum- CSV 20-110-20, MP chari, Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75) Soybean- R 518 (110), JS 9752 (100), RKS 18, RAUS 5 Finger millet- A 404, BM 2, BM 3 (BBM 10) Pigeonpea- Birsa Arhar (200-220), Asha (200-220) Lady's finger- Varsa uphar, Hybrid- Sonal, Sarika Maize- Birsa makka (Vikash) 2 (75-80), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85) <u>Fodder crop</u> Sorghum-HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid	

b) Agronomic Measures

- Top dressing of urea and DAP after receipt of the rain for all crops
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10-15 kg/ha
- Replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Soybean, Sesame, Horsegram , Niger, Cowpea, Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour
- Follow mulch after cultural operations to control the weeds in vegetables.
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges
- Foliar application of 2 % DAP or 0.5 to 1 % potassium chloride (KCl) +0.3 % Boric acid or 2% urea at pre-flowering and flowering stage in pulses and vegetables
- 2 % DAP spray for pulses.
- Use antitranspirants : Stomatal closure (Growth hormones like ABA, Ethrel, TIBA, succinic acid, ascorbic acid and Cycocel (CCC), Reflectant (Calcium bicarbonate, Lime water) Thin film (Hexadecanol (Higher alcohols) Cetyl alcohol, Methanol
- Acidic soils should be reclaimed by application of soil ameliorants.
- Follow integrated pest management.
- Weed control by applying pre-emergence 5-6 DAS (Pendimethalin) or Post-emergence 18-28 DAS (Bispyribac)
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lit. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit.

c) Remarks on Implementation

- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology for stress management through ATMAS, KVks, Govt Dept., NGOs and others. Soybean and fodder crops may be promoted.
- Promote Knowingness about climate resilient agriculture at district, block, panchayat and village level through involvement of KVks, ATMAS, DAO, NGO's and other State Agril. Govt line departments.
- Awareness of mechanization and Supply of Mouldboard and disc chisel/harrow through govt. scheme on subsidized way.
- Promote for double their income by curtailing cost of cultivation by introduction of early duration crops variety.
- Campaign for Awareness programme about crop-weather insurance

C2. Major Farming Situation/Land Situation: Midland High rainfall, shallow iron rich , light textured, sandy and acidic soil

Normal Crop/cropping system	Rice, Finger millet, Pigeonpea, Sorghum, Maize, Pigeonpea + Sorghum,Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 110, 111

Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 25P25, 27P36

Don 3

Ridge and Furrow method or raise bed broad furrow: Replace Rice with Pulses and cereals/ Vegetable/ Fodder crop Pulses and cereal-

Pigeonpea/ Soybean/ Maize/ Sorghum

Vegetable- Lady's finger/Tomato,/ Brinjal, creepers/Chili

Fodder Crop-Blackgram/ Greengram/ Cowpea/ Sorghum/ Maize

Variety

Pigeonpea- Birsa Arhar (200-220), ICPH 2671 (200)

Soybean-JS 9752 (100),

Birsa safed soybean 2 (105-110), RAUS 5

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Pusa HM 9(AQH 9), P3544, LG 32-81 –Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Sorghum- CSV 20-110-20, MP chari, CSV 1616



Vegetable crops

Lady's finger- Sonal, Sarika

Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Brinjal- Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali,

Cucurbits-

Bitter gourd- Arka hait, Pusa domausami,

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbur long green, Local, Arka harit

Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green, Long white

Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,

Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi

Chili- Spices- AndhraJyoti, Pusasadabahar, NP 46, Jwala, KA 2, California wonder, Chinese giant, Yellow wonder, Bharat

Fodder crop

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain)

Greengram- HUM 16, IPM-02-03-60-65, SML 668 (summer)

Cowpea- GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1

Maize- African tall, JS-1006 and Vijaya composite

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF, INPM
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P_2O_5 + 1 kg K_2O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @ 300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P_2O_5 : K_2O (Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P_2O_5 + 40 K_2O /ha ((Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K_2O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/l; Gundhi bug, leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /l water; Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/l; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/l., Mosaic- Methyl Demeton @ 1.5 ml/l; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lt.

c) Remarks on Implementation

- Campaign for awareness improved technology through RKVY, ATMAS, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be at different district, block, panchayat and village level
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency crops through Lamps within one months.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

C3. Major Farming Situation/Land Situation: Lowland High rainfall, shallow iron rich , light textured sandy & acidic soil

Normal Crop/cropping system	Rice
Suggested Contingency measures	



a) Change in crop/cropping system

Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of Don 2 in Don 1

DSR(Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), MTU-1001, MTU 1010, Transplanting (Hybrid rice varieties) Var.-PAC 801,25P25, 27P31 Arize 6444 (Gold), PHB 71, 26P52, 27P36 Fodder crop -In case of fallow (Late heavy rainfall)- Allis grass/ Para grass/ Arundino Grass

4 (b) Agronomic Measures

- Staggered Nursery raising by MAT/ DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use Post emergence weedicide
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² at 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water; Gundhi bug,leaf folder and BPH -Quinolophos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /ltr water

c) Remarks on Implementation

- Awareness programme of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments in case of DSR
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency mid early rice varieties through Lamps within one month Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation
- Contingency technology Awareness programme through KVKS, ATMAS, NGOs and DAOs
- Achieve maximum fallow area in case of late drought and suggest to go for cultivation of early duration rabi and fodder crops.

C4. Major Farming Situation/Land Situation: Upland Less rainfall, shallow depth , light textured sandy and acidic soil.

Normal Crop/cropping system	Upland Rice, Pigeonpea + Sorghum, Pigeonpea, Maize
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Discard Rice Crop

Sole crop

Raised bed or ridge and furrow method: Niger, Horsegram, Sorghum, Blackgram, Soybean, Finger millet, Pigeonpea, Lady's finger, Maize

Intercrop

Pigeonpea + Horsegram (1:2), Pigeonpea + Maize (1:1)

Horticulture crop

Vegetable –Lady's Finger, Ridge Gourd, Bottle Gourd, Cowpea

Fodder Crop

Deenanath grass/ Guinea grass/ Sadabahar/ Stylo hemata/ Chara badam/ Anjan grass/ Hybrid napier/ Cactus/ Sweet Sorghum

Variety

Niger- Birsa niger 1, 2 and 3 (95-105)

Horsegram- Birsa kulthi1 (90-95)

Sorghum- CSV 20-110-20, MP chari,

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75)

Soybean- R 518 (110), JS 9752 (100), RKS 18, RAUS 5



Finger millet- A 404, BM 2, BM 3 (BBM 10) Pigeonpea- Birsa Arhar (200-220), Asha (200-220)

Lady's finger- Varsa uphar, Hybrid- Sonal, Sarika

Maize- Birsa makka (Vikash) 2 (75-80), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85)

Fodder crop

Sorghum-HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

b) Agronomic Measures

- Top dressing of urea and DAP after receipt of the rain for all crops
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10-15 kg/ha
- Replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Soybean, Sesame, Horsegram , Niger, Cowpea, Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour
- Follow mulch after cultural operations to control the weeds in vegetables.
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges
- Foliar application of 2 % DAP or 0.5 to 1 % potassium chloride (KCl) +0.3 % Boric acid or 2% urea at pre-flowering and flowering stage in pulses and vegetables
- 2 % DAP spray for pulses.
- Use antitranspirants : Stomatal closure (Growth hormones like ABA, Ethrel, TIBA, succinic acid, ascorbic acid and Cycocel (CCC), Reflectant (Calcium bicarbonate, Lime water) Thin film (Hexadecanol (Higher alcohols) Cetyl alcohol, Methanol
- Acidic soils should be reclaimed by application of soil ameliorants.
- Follow integrated pest management.
- Weed control by applying pre-emergence 5-6 DAS (Pendimethalin) or Post-emergence 18-28 DAS (Bispyribac)
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit.water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology for stress management through ATMAS, KVKS, Govt Dept., NGOs and others. Soybean and fodder crops may be promoted.
- Promote Knowingness about climate resilient agriculture at district, block, panchayat and village level through involvement of KVKS's, ATMAS, DAO, NGO's and other State Agril. Govt line departments.
- Awareness of mechanization and Supply of Mouldboard and disc chisel/harrow through govt. scheme on subsidized way.
- Promote for double their income by curtailing cost of cultivation by introduction of early duration crops variety.
- Campaign for Awareness programme about crop-weather insurance

C5. Major Farming Situation/Land Situation: Midland High rainfall, shallow sandy soil

Normal Crop/cropping system	Rice, Pigeonpea, Maize,Pigeonpea + Maize ,Pigeonpea + Sorghum
-----------------------------	---

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 110, 111, MTU 1001, Transplanting (Hybrid Rice Varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 25P25, 27P36
Don 3

Ridge and Furrow method or raise bed broad furrow: Replace Rice with pulses and cereals/ Vegetable/ Fodder crop Pulses and cereal-Pigeonpea/ Maize/ Sorghum

Vegetable- Lady's finger/Tomato,/ Brinjal, Cucurbits/Chili

Fodder Crop

Blackgram/ Greengram/ Cowpea/ Sorghum/ Maize

Vegetable crop

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika



Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Hybrid- Swarn sampada, Swarn samridh, Pusa hybrid 1 Suraksha Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6

Cucurbits-

Bitter gourd- Arka hait, Pusa domausami,

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, local, Arka harit Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white

Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,

Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi

Chili- Spices- Andhraiyoti, Pusasadabahar, NP 46, Jwala, KA 2, California wonder, Chinese giant, Yellow wonder, Bharat

Fodder crop

Blackgram- WBU 109 (70-75), Uttara (75-80)

Greengram- HUM 16, IPM-02-03-60-65

Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

Maize- African tall, JS-1006 and Vijaya composite.

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P_2O_5 + 1 kg K_2O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P_2O_5 : K_2O (Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P_2O_5 + 40 K_2O /ha ((Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K_2O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/l; Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /lt water; Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/t; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/l., Mosaic- Methyl Demeton @ 1.5 ml/l; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lt.

c) Remarks on Implementation

- Campaign for awareness improved technology through RKVY , ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be at different district, block, panchayat and village level
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency crops through Lamps within one months.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.



C6. Major Farming Situation/Land Situation: Lowland Less rainfall, shallow depth , light textured sandy and acidic soil	
Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2 in Don 1</u>	
DSR(Improved Rice variety) Var.- Shabhagi, IR 64-Drt 1, Abhishek (120 days), MTU 1010, BVD 203, BVS 1 Transplanting (Hybrid rice varieties) Var.-PAC 801,25P25, 27P31 Arize 6444 (Gold), PHB 71, 26P52, 27P36 Fodder crop : In case of fallow (Late heavy rainfall) Para grass	
b) Agronomic Measures	
<ul style="list-style-type: none">• Staggered Nursery raising by MAT/ DAPOG method• Follow community based nursery raising• Follow RDF,INPM• Use Post emergence weedicide• Use early to mid early duration of rice variety.• Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²• Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice• Topdressing 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² at 10-15 days after sowing• In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling• Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm• Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.• DSR-Use plastic drum seeder rice tools• Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lt; Gundhi bug,leaf folder and BPH -Quinolophos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /lt water	
c) Remarks on Implementation	
<ul style="list-style-type: none">• Awareness programme of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme• Supply of Plastic drum seeder through line departments in case of DSR• Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.• Supply of improved and hybrid seeds of contingency mid early rice varieties through Lamps within one month Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates• Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation• Contingency technology Awareness programme through KVKS, ATMAS, NGO's and DAO's• Achieve maximum fallow area in case of late drought and suggest to go for cultivation of early duration rabi and fodder crops.	



PART-II

**Monsoon/Weather Situation: Normal onset followed by 15-20 days dry spell after sowing
(Early Season Drought-Normal onset)**

A1. Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils	
Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
Suggested Contingency measures	
a) Change management Cultivate drought tolerant promising non paddy crops like pigeonpea, blackgram, greengram, rice bean, finger millet, guar, sesame, soyabean, sorghum, pearl millet, sweet potato, castor and vegetables like radish, tomato, brinjal, creeper bean, chili, lady's finger wherever possible in place of upland rice	
<ul style="list-style-type: none"> • Maximum use of organic manures for early seedling vigour along with RDF (N:P₂O₅:K₂O) • Recommend to resow with subsequent rains for better plant stand. • When damage is Less than 30 per cent then go for Gap filling in all upland crops • When damage is More than 50 per cent then go resowing in all upland crops • Removing excess plants where are over crowded,to reduce crop stand to conserve soil moisture • Water spraying during evening and early morning 	
b) Soil nutrient & moisture conservation measures <ul style="list-style-type: none"> • Avoid top dressing of Urea during dry spell and wait till downpour • Go for in-situ moisture conservation • One hand weeding followed by hoeing and simultaneous eartingup after 20 DAS is highly recommended in all upland crops. 	
c) Remarks on Implementation Awareness for Construction of rain water harvesting structures for recycling of water during dry spell like DOVAS through SHG or on subsidized basis through State Govt.schemes.	
A2. Major Farming Situation/Land Situation: MID LAND Sandy loam soils	
Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change management <u>Don 2</u> <ul style="list-style-type: none"> • If possible, go for staggered raising of nursery in rice crop • If possible, raise community nursery of rice at a reliable water source to save time for further delay. • In case, if rice population is less than 40-50 percent, gap filled by retransplanting the rice crop and for more than 50 per cent mortality use fresh seeding for fresh transplanting. • Follow gap filling by removing seedlings from profuse tillers to have a uniform distribution of same aged plants • For termite and disease management in nursery spray Indofil M 45 and Chlorpyriphos @ 0.2 per cent life saving irrigation • DSR on receipt of rain by using Paddy drum seeder or • High yielding varieties- follow transplanting while, Improved varieties – follow DSR • In case of DSR- Use sprouted seeds in plastic drum seeder with increased seed rate by 20-25 per cent for good crop stand • Late transplanted rice during early season drought results in the occurrence of sheath rot and grain discoloration diseases. • Follow pre emergence and post emergence weedicide to disturb/check the crop-weed competition for nutrient • Provide life saving and protective irrigation to over aged seedling in nursery through dovas (harvested rain water). Also, take care of blast disease in nursery and avoid using urea in nursery. • Strengthen the bunds to check the drainage holes and seepage loss in transplanted and direct sown medium land rice regularly <u>Don3</u> <ul style="list-style-type: none"> • Follow raised bed broad furrow or Ridge and furrow method for Maize/ Pigeonpea/ Lady's finger/ Blackgram/ Soybean • Adopt surface mulching with crop residue or tree loppings of <i>Glyricidia</i> wherever possible. If farm waste is not available, use blade to form a thin layer of soil mulch to avoid cracks • Life saving irrigation • In case of transplanting of over aged seedling (35-45 days), increase number of seedling per hill (5-6 seedling/hill) 	

b) Soil nutrient & moisture conservation measures

- Dry seeding of rice with application of pre and post emergence weedicide in over aged seedlings (>25 DOS)
- Split application of Urea fertilizer
- Foliar spray of 2% KNO_3 or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells

c) Remarks on Implementation

Awareness for Construction of rain water harvesting structures for recycling of water during dry spell like DOVAS through SHG or on subsidized basis through State Govt.schemes.

A3. Major Farming Situation/Land Situation: LOW LAND Sandy clay loam soils

Normal Crop/cropping system	Rice
-----------------------------	------

Suggested Contingency measures

a) Change management

- If possible, go for staggered nursery raising in rice crop
- If possible, raise community nursery of rice at a reliable water source to save time for further delay.
- In case, if rice population is less than 40-50 percent, gap filled by retransplanting the rice crop and for more than 50 per cent mortality use fresh seeding for fresh transplanting.
- Follow gap filling by removing seedlings from profuse tillers to have a uniform distribution of same aged plants
- Prefer mid early rice variety instead of late variety
- Use pre and post emergence weedicide
- Over aged seedling should be top cut and treat the seedlings root by Dursban/Chlorpyriphos @ 5 ml per lt water and transplant immediately after treated seedlings with 2 per cent Urea solution
- In case of transplanting over aged seedling (35-45 days), increase number of seedling per hill (5-6 seedling/hill)
- In fallow land go for cultivation of mid early duration rice variety through DSR @ 70-80 Kg/ha

b) Soil nutrient & moisture conservation measures

- Split application of Urea fertilizer
- Foliar spray of 2% KNO_3 or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells

c) Remarks on Implementation

Awareness for Construction of Ponds, check dam through water shed management & MNREGA scheme through SHG or on subsidized basis through State Govt. schemes.

B. Monsoon/Weather Situation: Mid season drought (long dry spell, consecutive 2 weeks rainless (<2.5 mm) period

B1. At vegetative phase

B1.1. Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils

Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
-----------------------------	--

Suggested Contingency measures

a) Change management

- Use organic mulches such as tree leaves, straw and other available crop residue to conserve soil moisture
- Avoid top dressing of fertilizers till sufficient moisture is available in soil
- Use reflectant or antitranspirant like Kaolin @ 3-5 kg/100 lt or
- In pulses, at weekly interval foliar spray of KCl @ 0.5- 1 % + 100 ppm Boric acid followed by foliar spraying of 2 percent urea during evening time
- Spray wax emulser
- Manual weeding followed by hoeing for germinating weeds.
- For termite and leaf folder control spraying or drenching of Chlorpyriphos @ 2ml/lt water and for all pulses and cereals.
- For leaf folder control in Maize (Stem borer) and Pigeonpea apply Carbofuran 3 G @ 12 Kg/acre or Phorate 10 G @ 4 kg/acre or Quinolphos @ 1 ml/lt water in Maize for leaf folder
- Also, spray @ 20/40/60 ppm CaCl_2 in pulses
- Vegetables- Foliar spray of water with 2 per cent KCl + 100 ppm Boron
- Tomato- Foliar spray of CaCl_2 @ 20/40/60 ppm
- Gap filling may be done with pigeonpea to maintain adequate plant stand.
- For termites in pigeonpea, maize and other standing cereal crops which can be controlled by soil drenching with chlorpyriphos 20 EC @ 2 ml/lt water or by adding Chlorpyriphos 1.5% dust @ 8–10 kg/ha or Carbofuran 3G @ 12 kg or Phorate 10 G @ 4 kg. acre before final land preparation and also control Gallmidge



- In greengram and blackgram, cowpea, bean and lady's finger the spread of YMV by insect vector may increase. Hence, to control insect vectors spray Dimethoate @ 1ml/lit. water or Imidacloprid 4 ml/10 lit. water twice at 10 days interval
- In groundnut crop termites and white grub incidence is expected to be more. Methods suggested in rice may be followed to reduce the pest infestation.
- Incidence of leaf miner in groundnut may increase which can be managed by spraying Monocrotophos 36 SL or Triazophos 40 EC @ 1 ml/lit. water twice at fortnight intervals.
- Under dry condition incidence of mites is expected to be more in vegetable crops which can be brought down by spraying of dicofol @ 2 ml/lit water.
- Early and mid season drought favours disease like brown spot of rice, bacterial wilt of brinjal and other vegetables

b) Soil nutrient & moisture conservation measures

Foliar spraying of DAP @ 2 per cent along with Boric acid @ 0.3 per cent. Also, spray Urea @ 1 per cent Provide micro- irrigation with drip for wide spaced crops such as chilies and vegetables and Sprinklers for groundnut, maize and vegetables wherever ground/ surface water is available.
Go for life saving and protective irrigation from constructed dovas.

c) Remarks on Implementation

Promote construction of Rain water harvesting structure watershed programme and MNREGA

B2. At flowering/ fruiting stage

B2.1. Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils

Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
-----------------------------	--

Suggested Contingency measures

a) Change management

- Maize- Harvest it for fodder use
- Pulses- and vegetables- At 2-3 days interval spraying of water followed by 2 per cent KCl + 100 ppm Boron during evening time is recommended.
- In case of groundnut maturing in the month of September which can be harvested after providing light irrigation through dovas to lose the soil.

b) Soil nutrient & moisture conservation measures

Go for life saving and protective irrigation from constructed DOVAS.

c) Remarks on Implementation

Promote for the construction of Rain water harvesting structure watershed programme and MNREGA

B3. At vegetative phase

B3.1. Major Farming Situation/Land Situation: MID LAND Sandy loam soils

Normal Crop/cropping system	Rice
-----------------------------	------

Suggested Contingency measures

a) Crop management

Don 2

Manual weeding followed by hoeing for germinating weeds

Take care of mealy bug and termite attack which are more prevalent in dry weather .

Top dressing should be followed only after receipt of rain .

No urea should be top dressed until receipt of rainfall in rice crop.

For BPH, dusting field bunds and around with Carbaryl (Savin)4% or malathion 5% @ 10 – 12 kg/acre

Don 3

One manual weeding for germinating weeds

Apply 4 Kg N/acre in sorghum and oilseed crops soon after receipt of rains.

In pigeonpea, if the drought affected plants to recuperate with the revival of the rains, spray 2 to 3% urea after the foliage is wetted with the rains.

Foliar application of Sulphur @ 1ppm to mitigate the stress condition in oilseed is necessary after receipt of rainfall

Apply post emergence weedicide for controlling weeds in oilseed (Groundnut) to undisturb the pegging process.

During 40-45 DAS, if there is a severe moisture stress, thinning may be done in kharif sorghum and pearl millet.

b) Soil nutrient & moisture conservation measures

- Foliar spray of KCl or ZNSO₄ @ 2 per cent
- Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells
- Life saving irrigation through dovas, well, ponds, check dams and bora bandh

c) Remarks on Implementation

Promote for the construction of Rain water harvesting structure watershed programme and MNREGA



B4. At flowering/ fruiting stage

B4.1 Major Farming Situation/Land Situation: MID LAND Sandy loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
Don 2 and Don 3	
<ul style="list-style-type: none"> • Life saving irrigation with harvested water • Spray of urea @ 1-2 percent • Drought condition during the month of August-September onwards shall result in severe incidence of foliar blast and brown spot diseases in rice. It is advised to spray Tricyclazole (Tilt) @ 6 g/ 10 lt. water or Casugamycin @ or Kasu B @ 2 ml/lt. water twice at 10 days intervals during drought period. 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Foliar spray of KCl or ZNSO₄ @ 2 per cent • Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells • Life saving irrigation through dovas, well, ponds, check dams and bora bandh 	
c) Remarks on Implementation	
Promote for the construction of Rain water harvesting structure watershed programme and MNREGA	

B5. At vegetative phase

B5.1. Major Farming Situation/Land Situation: LOW LAND Sandy clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<ul style="list-style-type: none"> • Foliar spray of 2 per cent KCL followed by 1-2 per cent Urea. • Weeding should be done • Drought makes the crop vulnerable to sheath rot and sheath blight diseases. Maintenance of field sanitation followed by twice spraying at 10 days interval with validamycin 2-3 ml/lt water or Tricyclazole @ 6g/10 lt. water or carbendazim @ 2 g/lt water are advised. • Life saving irrigation 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Foliar spray of Foliar spray of Urea @ 2 per cent • Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells • Life saving irrigation through dovas, wells, ponds, check dams and bora bandh 	
c) Remarks on Implementation	
Awareness for Construction of Ponds, check dam through water shed management & MNREGA scheme through SHG or on subsidized basis through State Govt.schemes.	

B6. At flowering/ fruiting stage

B6.1. Major Farming Situation/Land Situation: LOW LAND Sandy clayloam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<ul style="list-style-type: none"> • Drought condition during flowering and fruiting and onwards shall result in severe incidence of foliar blast and brown spot diseases in rice. It is advised to spray Tricyclazole (Tilt) @ 6 g/ 10 lt or Casugamycin @ or Kasu B @ 2 ml/lt. water twice at 10 days intervals during drought period. • Life saving irrigation • During drought, attack of gundhi bug shall be more. Apply Quinolphos or Monocrotophos @ 1-2 ml per lt. water. 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Weeding and foliar spray of urea @ 2 per cent • Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells • Life saving irrigation through dovas, well, ponds, check dams and bora bandh 	
c) Remarks on Implementation	
Promote for the construction of Rain water harvesting structure watershed programme and MNREGA	



C. Monsoon/Weather Situation: Terminal drought (Early withdrawal of monsoon)

C1. At fruiting/pre physiological maturity stage

C1.1. Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils

Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
Suggested Contingency measures	
a) Change management	
<ul style="list-style-type: none"> • Life saving irrigation to vegetables through stored moisture from constructed DOVA • If not possible to make survival harvest it for fodder use 	
b) Rabi Crop planning	
<ul style="list-style-type: none"> • Cultivation of Niger, Horsegram, Toria, linseed as relay/paira cropping • In case of availability of irrigation, go for cultivation of early Potato and pea (early Arkel group) • Prepare kachha check dam or Bora Bandh for Water conservation • Mid early variety of radish cultivation is recommended 	
c) Remarks on Implementation	
Promote for the construction of Farm ponds through watershed management programme and MNREGA	

C1.2. Major Farming Situation/Land Situation: MID LAND Sandy loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<u>Don 2</u>	
<ul style="list-style-type: none"> • At milking , soft and dough stage spray KCL @ 2 per cent • In case of gundhi bug attack found more than ETL(>2 gundhibug /m²), spray Chlorpyriphos dust or Monocrotophos @ 1 ml/lit. water • If possible go for life saving irrigation • Late season drought generally results in outbreak of foliar, node, collar or neck blast of rice depending on the stage of crop. 	
<u>Don 3</u>	
Instead of grain purpose crops like sorghum, pearl millet, maize, cowpea, black and greengram that can be harvested for fodder use	
b) Rabi crop planning	
<ul style="list-style-type: none"> • Ensure for all inputs required for rabi season in advance. • In case of failure of kharif crops prefer sowing of pre rabi catch crops like, Toria, Niger, Horsegram, blackgram, sesame linseed in uplands to medium lands 	
c) Remarks on Implementation	
Promote construction of Rain water harvesting structure watershed programme and MNREGA	

C1.3. Major Farming Situation/Land Situation: LOW LAND Sandy loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<ul style="list-style-type: none"> • Life saving irrigation. • The land should be tilled properly in case <i>kharif</i> crop fails sow <i>rabi</i> crops like safflower, pigeonpea in sept-Oct (Short duration) • Spray KCL @ 2 per cent followed by urea @ 2 per cent • Mid early rice crop may be harvested at Physiological maturity • Cultivate vegetables like Tomato, Brinjal, Capsicum, Shimla mirch, Broccoli, Cabbage and Cauliflower, green pea and potato as per suitability near and around tributaries 	
b) Rabi crop planning	
Prefere early sowing of wheat, Mustard, Chickpea,linseed and lentil as sole or intercrop Wheat + Chickpea (4:2) Wheat+ Mustard (4:3)	
c) Remarks on Implementation	
Promote construction of Rain water harvesting structure watershed programme and MNREGA	

PART-III

A. Unusual rains: Continuous high rainfall in a short span leading to water logging

Suggested Contingency measures
a) Crop management
Pigeonpea /Sorghum/Pearl millet
Vegetative stage- Prefer ridge and furrow method of sowing. Ensure for proper drainage through channel. Collect runoff water in Dovas for further use.
Flowering stage- Ensure for proper drainage through channel. Collect runoff water in Dovas for further use.
Crop maturity stage- No such situation at the time of maturity
Post harvest- After Sun drying follow grading and storing
Blackgram and other pulses/Oilseeds
Vegetative stage- Follow Ridge and furrow sowing
Ensure for proper drainage through channel
Collect runoff water in Dovas for further use
Avoid application of fertilizer
Flowering stage- Ensure for proper drainage through channel
Collect runoff water in Dovas for further use
Avoid application of fertilizer
Prophylactic measure for jassid and YMV
Crop maturity stage-
Post harvest-
Rice
Vegetative stage- Safe disposal of excess water from rice field. Bund repairing and strengthen. Application of insecticides in the afternoon hours is preferred seeing the weather condition or after spraying weather should remain rain free for at least 4-5 hrs. Retransplant to maintain plant population in case of mortality more than 50 %
In partially damaged crop, allow to withstand upright. Flood occurs due to heavy storm in mid and lowland which when recedes probability of occurrence of swarming caterpillar on field bunds and around of rice crop is more. So, when it crosses the Economic Threshold Limit (ETL) i.e., one larva / hill then spray the crop with Chlorpyriphos/ Triazophos/ Profenophos @ 2 ml/lt water or dust the crop with Quinalphos @ 1.5% D @ 10kg/ acre. To prevent migration of larvae from one field to other, bunds should be heavily dusted with the dust formulation mentioned above. In partially ponded field, rice caseworm and in general leaf folder attack is expected. If 1-2 cases or folded leaves/hill is seen spray the crop with Monocrotophos / Chlorpyriphos @ 1 ml/lt water or with Cartap Hydrochloride 50 SP / Fipronil 5 SP @ 200 g/acre. Rain storms during kharif may result in severe occurrence of bacterial leaf streak and bacterial blight in rice. It is advised to spray the crop immediately after every rain spell with streptocycline @ 1g/10 lt water or plantomycin @ 1g/lt water or bacterinol @ 2g/lt. water. Control snail occurrence by Acaricide
Flowering stage- Safe disposal of excess water from rice field. Bund repairing and strengthen. Avoid application of fertilizer. Flood occurs due to heavy storm in mid and lowland which when recedes probability of occurrence of swarming caterpillar, BPH and cut worm on field bunds and around of rice crop is more. So, when it crosses the Economic Threshold Limit (ETL) i.e., one larva / hill then spray Chlorpyriphos/ Triazophos/ Profenophos @ 2 ml/lt water or dust the crop with Quinalphos @ 1.5% D @ 10kg/ acre. To prevent migration of larvae from one field to other, bunds should be heavily dusted with the dust formulation mentioned above. In partially ponded field, rice caseworm and in general leaf folder attack is expected. If 1-2 cases or folded leaves/hill is seen spray the crop with Monocrotophos / Chlorpyriphos @ 1 ml/lt water or with Cartap Hydrochloride 50 SP / Fipronil 5 SP @ 200 g/acre. Unusual and heavy rain during kharif may result in severe occurrence of bacterial leaf streak and bacterial blight in rice. It is advised to spray the crop immediately after every rain spell with streptocycline @ 1g/10 lt water or plantomycin @ 1g/lt water or bacterinol @ 2g/lt. water. Control snail occurrence by Acaricide.
Crop maturity stage- Provide drainage for fast removal of water from the field to favour harvesting
Post harvest- Protect the grain from rain and store it after sun drying for 2-3 days
Maize
Vegetative stage- Prefer ridge and furrow method of sowing. Ensure for proper drainage through channel. Earthingup after downpour. At Knee stage apply thimate 10 G @ 4-6 grains in whirl
Flowering stage- Ensure for proper drainage through channel. At flowering and silking stage for ant attack apply dust on silks @ 0.5 g / cob
Crop maturity stage- Provide drainage for fast removal of water from the field to favour harvesting
Post harvest- Protect grains from rain and store it after sun drying for 2-3 days



Horticulture

Vegetative stage- Prefer ridge and furrow method for sowing and proper drainage. Ensure for proper drainage through water ways. Collect runoff water in Dovas for further use. Soil drenching Carbofuran 3G @ 3 g/lt water against insects. In case of web formation with leaves apply (Nuvan)DDVP @ 1 ml/lt water as a fumigant

Flowering stage- Apply hormone to prevent flower drop. Ensure for proper drainage. Take precaution against wilting and fruit rot. In Tomato and Brinjal-drenching Bavisting @ 2 ml/lit. water + Streptocycline @ 1-2 g/lit water. In Cauliflower -In case of Incidence of collar rot -Spraying of Saaf (Metalaxyl + Mancozeb) @ 2 g/lit water solution. Drainage of excess water. In Lady's finger- YVMV- Spray insecticide followed by fungicide. Soil drenching Carbofuran 3G @ 3 g/lt water against insects. In case of web formation with leaves apply (Nuvan) DDVP @ 1 ml/lt water as a fumigant

Crop maturity stage- Take precaution against wilting and fruit rot. For wilting– Soil drenching with Bavistin @ 2 ml/lit + Streptocycline @ 1-2 g/lit water. In YMVM- Insecticide followed by fungicid

Post harvest- Immediate harvest and safe disposal of produce

Vegetables- (Cucurbits, Tomato/ Brinjal/ Cauliflower/ Cabbage/ Lady's finger/Dolichos bean/Amaranthus leaf/ Cariander leaf/Radish)

Vegetative stage- Sowing on ridge and drainage through furrow. Prophylactic measures against pest and diseases.

Damaged twigs and leaves may be removed and follow fungicide spraying and stacking

Flowering stage- Apply hormone to prevent flower drop. Ensure for proper drainage. Take precaution againts wilting and fruit rot. In Tomato and Brinjal-drenching Bavisting @ 2 ml/lit

+ Streptocycline @ 1-2 g/lit water. In Cauliflower -In case of Incidence of collar rot -Spraying of Saaf (Metalaxyl + Mancozeb) @ 2 g/lit water solution. Drainage of excess water. In Lady's finger- YVMV- Spray insecticide followed by fungicide. Provide support through stacking

Crop maturity stage- Take precaution against wilting and fruit rot. In Wilting– Soil drenching with Bavistin @ 2 ml/lit + Streptocycline @ 1-2 g/lit water. In YMVM- Insecticide followed by fungicide

Provide support through stacking.

Post harvest- Immediate harvest and sell produce safely in the market

b) Pest and disease management

Rice

Vegetative stage- Sheath blight- Hexaconazole @ 1ml/lt wate. Blast- Tricyclazole @ 6 g/10 lt water

Flowering stage- Sheath blight- Hexaconazole @ 1ml/lt water. Blast- Tricyclazole @ 6 g/10 lt water. Falsesmut-Nativo @ 4g/10 lt water

Crop maturity stage- False Smut - Control- Nativo @ 4g/10 lt water or Propiconazole + Tricyclazole 52.5 SE @ 1ml/lt water. In case of grain discolourness (Grain blast). Spray Tricyclazole @ 6 ml / 10 liter water

Post harvest- Store grains after proper sun drying to minimize the incidence of stored grain pest

Maize

Vegetative stage- Stem borer Control- Carbofuran 3 G @ 12 Kg/acre or Phorate 10G@ 4 kg/acre

Flowering stage- Sheath blight Control- Hexaconazole1-2 ml/lt water

Vegetables- (Cucurbits, Tomato/ Brinjal/ Cauliflower/ Cabbage/ Lady's finger/Dolichos bean/Amaranthus leaf/ Cariander leaf/Radish)

Vegetative stage- Before sowing apply in soil, Carbofuran 3 G @2-3 g/m². Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt. water streptocycline or 2-3 g/lit. water plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lit water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lt. water against downy mildew diseases of cucurbits.

Flowering stage- Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt. water streptocycline or 2-3 g/lit. water plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lit water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lt. water against downy mildew diseases of cucurbits. YVM Control- Carbofuran 3G @ 3 or Phorate 10 G @ 1 g/m² followed by any fungicide

Crop maturity stage- Stop spraying 1 week before harvesting

Post harvest- Harvest and sell produce in the market



French bean-

Vegetative stage- Rust disease Control- Mancozeb 2g/ lt water. Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt. water streptocycline or 2-3 g/lt. water plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lt water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lt. water against downy mildew diseases of cucurbits.

Flowering stage- Take care of pod borer and aphid attack. Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt. water streptocycline or 2-3 g/lt. water plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lt water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lt. water against downy mildew diseases of cucurbits.

Crop maturity stage- Stop spraying 1 week before harvesting

Post harvest- Harvest and sell produce in the market

B. Extreme Weather Events

Suggested Contingency measures

Hail storm

Seedling / nursery stage- Vegetable nursery should be raised in poly house or make proper arrangement of low height Polly tunnels in open area or cover with plastic sheet or thatching should be done

Vegetative stage- In vegetables-Remove damages parts immediately and apply insecticide followed by fungicide as prophylactic measures. Follow fertilization through foliar as well as broadcasting

Reproductive stage- n vegetables- Remove damaged parts immediately and apply insecticide followed by fungicide as prophylactic measures. Follow fertilization through foliar as well as broadcasting for proper fruiting

At harvest- Safely sell in the market after grading for immediate returns

Heat Wave

Wheat Chickpea/pea

Seedling / nursery stage- For protection from heat and cold wave there is intervention to sow the rabi crops in between 2nd week of October to 2nd week of November to protect theirs vegetative phase from ground/radiation frost results from cold wave/wind chill injury and reproductive phase from terminal heat stress on Mustard, Chickpea, Wheat, Lentil, Linseed and pea crops. Life saving irrigation

Vegetative stage- Timely sown crop never face heat stress while very late sown(January) crop face heat stress hence only one option is to provide life saving irrigation and water spray during evening time frequently at 2-3 days intervals. Take care of termite attack by spraying Chlorpyriphos @ @ 1 ml/lt and drenching @ 3-5 ml/lt water

In Chickpea because of high soil and ambient temperature (> 35 °C) favours the dry root rot disease starts during flowering/reproductive stage (spraying Captan or thiram or carbendazim or ridomil MZ or Saaf @ 1,5-2 g/lt water)

Reproductive stage- To minimize the terminal heat stress during the month of March and April the only and only way is to provide frequent protective irrigation irrespective of theirs stages (Life saving irrigation). Take care of termite attack by spraying Chlorpyriphos @ @ 1 ml/lt. water and drenching @ 3-5 ml/lt water. In Chickpea because of high soil and ambient temperature (> 35 °C) favours the dry root rot disease starts during flowering/ reproductive stage (spraying Captan or thiram or carbendazim or ridomil MZ or Saaf @ 1,5-2 g/lt water)

At harvest- Frequent irrigation should be provided to meet the evaporative losses.

Tomato/Brinjal/ Lady's finger/Cucurbits

Seedling / nursery stage- Due to heat stress wilting and mortality is more hence frequent irrigation and cover the nursery with mulch(Straw/leaves)

Vegetative stage- Due to heat stress wilting and mortality is more hence frequent irrigation and cover the nursery with mulch(Straw/leaves)

Reproductive stage- Drying of flower- Spray PCOA. Follow mulching after irrigation

At harvest- Immediate harvest after irrigation and shift it to safer place

Cold wave

Wheat

Seedling / nursery stage- Cold environment during tillering or branching stage favours more number of tillers in wheat and more branching in mustard, chickpea, lentil and linseed crops which prospects for high yield but it is detrimental for potato, tomato, brinjal, pea, creeper vegetables and fruits. Irrigation. Balanced fertilizer application.



Foliar spray of nutrients

Vegetative stage- Light irrigation. Mulching with crop residue \ weeds. Fertilizer application

Reproductive stage- Irrigation, fertilizer application

At harvest- N/A

Pigeonpea/Mustard/Linseed/Chickpea/pea

Seedling / nursery stage- In Mustard because of cool weather aphid insects attack is more prominent (spraying Rogor (Dimethoate) @ 2 ml or or Monocrotophos 36 EC @ 1 ml /lt water during evening time is advised).

In linseed Alternaria blight (For blight spray Double dose (Iprodione 25 % WP + Carbendazim 25 % WP) @ 2 g per lt. water) and powdery mildew (prophylactic spraying of Sulfex @ 3 g or Kairathen 1 ml per lt water twice at weekly interval during evening time) disease are more common. For powdery mildew in pea (spraying Calixin (Tridemorph 80 % EC @ 5 ml per 10 lt water twice are highly recommended).

In Chickpea-Cold and wet environment (High humidity) during seedling stage cause collar rot, black root rot, wet rot, Pythium root and seed rot in Chickpea, while in potato, pea and tomato favours late blight (spraying of Krilaksil or Ridomil MZ chemical@ 1.5 g per liter water), powdery mildew (spraying newly emerged fungicide Double dose (Iprodione 25 % WP + Carbendazim 25 % WP) 2 g per lt water twice at weekly interval) and bacterial wilt, leaf spot and canker (spraying carbendazim @ 2g/lt water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP) diseases in respective vegetable crops. Anthracnose in cucurbitaceous species. Vegetative stage- Provide light irrigation. Follow mulching with crop residue \ weeds\straw\leaves. In Mustard because of cool weather aphid insects attack is more prominent (spraying Rogor (Dimethoate) @ 2 ml or or Monocrotophos 36 EC @ 1 ml /lt water during evening time is advised)

Reproductive stage- Pigeonpea- During flowering and pod formation stage attack of Pod borer/sucking bug, mites, blister beetle insects as well as sterility disease may occur more (spraying Profenophos 50 EC, methomyl 40 SP or monocrotophos 36 SL kill the larvae but as the webs protect them from contact insecticides hence along with contact insecticides, mixing of fumigant insecticide such as DDVP @ 0.5 ml/lt. water is required to make the larvae come out from the web. For Mites and Aphids, Dimethoate 30 EC @ 2ml/l and acaricides such as Dicofol 18.5 EC @ 2.5 ml/lt. water , for Blister beetle synthetic pyrethroids such as Cypermethrin 10 EC @ 1.0 ml/lt. water or Lamda cyhalothrin 5 EC @ 1.0 ml/lt. water; for sterility mosaic Dicofol 18.5 EC 2.5 ml or Oxydemeton methyl 25 EC or Dimethoate 30 EC 2.0 ml/lt. water on alternate row twice at an interval of 10 days are recommended).

Vegetables

Seedling / nursery stage- Raising seedling in Poly house, re sowing if damage is more. Provide shelter belt (Wind break) at appropriate spacing with Sisham, Ghamhar. Provide irrigation and mulching with straw and leaves

Vegetative stage- Provide light irrigation. Follow mulching with crop residue \ weeds\straw\leaves. Disease and pest control, care for chilling injury or replanting

Reproductive stage- Drying of flower- Spray PCOA. Follow mulching after irrigation

At harvest- Grading and safely dispose produce in the marketing

Frost

Wheat

Seedling / nursery stage- N/A

Vegetative stage- Provide light irrigation. Follow mulching with crop residue \ weeds\straw\leaves

Pigeonpea

Seedling / nursery stage- Exposure of crop to smoke by burning waste material during night time

Vegetative stage- Exposure of crop to smoke by burning waste material during night time, Light sprinkler irrigation

Reproductive stage- Exposure of crop to smoke by burning waste material during night time, Light sprinkler irrigation

At harvest- Exposure of crop to smoke by burning waste material during night time, Light sprinkler irrigation

Tomato & Potato and Horticultural crops (fruit)

Seedling / nursery stage- Create smoke around the field by using waste materials or set afire with used mobile oil in north-west or west-north direction towards incoming cold waves. Use polythene or bamboo hoogli in small horticultural /nursery/cash vegetable crops during morning hour and remove it during daytime. In Perennial or Horticulture crop (Fruit) also frequent irrigation followed by mulching, thatching , creating smoke screen s and lighting of fire should be practiced in availability of irrigation facility

Vegetative stage- Earthing up, Irrigation and create smoke around the field by using waste materials or set a fire with used mobile oil in north-west or west-north direction towards incoming cold waves. Use polythene or bamboo hoogli in small horticultural /nursery/cash vegetable crops during morning hour and remove it during daytime. In Perennial or Horticulture crop (Fruit) also frequent irrigation followed by mulching, thatching , creating smoke screen s and lighting of fire should be practiced

Reproductive stage- Immediate harvesting and disposal

At harvest- Harvest in dry weather

Cyclone- Not applicable

CONTINGENCY PLANS FOR RABI

1. Sowing window information

Land type	Cropping system	Crop name	Optimum sowing window (Please mention along with week)
1.Upland	Maize/ Groundnut-/Finger Millet –Vegetable/ Toria/Mustard / Linseed	Toria, Mustard,Linseed Vegetable- Tomato, Brinjal, Round melon (Tinda),Radish (Under limited Source of water)	Toria- 3rd week of September – 4 th week of September Mustard- 1 st week of October – 4 th week of October Linseed- 1 st week of October- 3 rd week of October Vegetables (Tomato, Brinjal, Radish)- 1 st week of October- 4 th week of November
2.Mid Land	Rice –Wheat Rice –Chickpea Rice –Mustard Rice –Lentil Rice-Linseed Rice-Vegetable	Rainfed (Zero tillage)- Barley, Mustard, Chickpea, Lentil, Linseed, Dolichos bean (September sown) Irrigated- (Zero tillage) Barley/Wheat Potato, Vegetables (Cauliflower, Tomato, Cabbage, Brinjal, Round melon (Tinda), Radish Cucurbits (Pumpkin) Fodder Crop- Oat, Maize, lathyrus	Barley/Wheat – 3 rd week of October – 2 nd week of November Mustard- 1 st week of October – 4 th week of October Chickpea – 2 nd week of October – 2 nd week of November Lentil- 3 rd week of October- 2 nd week of November Linseed- 1 st week of October – 4 th week of October Potato- 4 th week of October -2 nd Week of November Vegetables- 1 st week of October – 4 th week of November Dolichos bean- 1 st week of August – 2 nd week of September Fodder-2 nd week of October – 2 nd week of November
3.Low Land	Rice –Wheat Rice- Linseed (Paira) Rice-Vegetable	Linseed/ lathyrus (Paira cropping), Wheat Vegetables (Onion Fodder Crop- Oat, Maize, lathyrus	Linseed- 4 th week of October – 2 nd week of November Wheat- Timely- 1 st week of November- 3 rd week of December, Late Sown - 1 st week of December- 4 th week of December Lathyrus- 4 th week of October – 2 nd week of November Fodder-2 nd week of November – 2 nd week of December

2. Contingency measures for Field crops grown with residual moisture under rainfed condition

2 (A) Optimal residual moisture

2A1. Land type- UPLAND

- a) Cropping system- Maize/ Groundnut-/Finger Millet –Vegetable/Toria/Mustard / Linseed
- b) Crop name- Toria, Mustard,Linseed, Vegetable- Tomato, Brinjal, Round melon (Tinda),Radish (Under limited Source of water)
- c) Sowing Window- Toria- 3rd week of September – 4th week of September, Linseed- 1st week of October- 3rd week of October, Vegetables (Tomato, Brinjal, Radish)- 1st week of October- 4th week of November
- d) Variety- Toria-PT 203, Panchali; Mustard- Pusa Mahek, Pusa Mustard 25, NRCHB 101, Bharat Sarson 1, Pusa 28, 30; Linseed- Sharda, Priyam, Divya

e) Agronomic management practices

- Rain water harvesting and recycling.
- Deeping of water storing structure(Shallow and deep) in April and May month
- Deep summer ploughing in April and May month.
- Strengthing and raising of field bunds in April and May months
- Sowing in defined window for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better crop stand (Plant population)
- Application of Lime or Dolomite (3-5 q/ha) in soil
- Soil application of Sulphur (20 kg/ha) and boron (1kg/ha) in oilseed, pulses and vegetables.
- Foliar spray of Urea (2 %) at flower initiation and pod formation stage in oilseed and pulses
- Follow seed priming (warm water for 4-6 hrs.) before sowing
- Follow seed treatment with fungicide-insecticide-rhizobium
- Irrigate only at critical stages
- Pre and post emergence weedicide application
- Follow hoeing after manual weeding
- Follow RDF, INM and IPM



- For Water use efficiency use antitranspirant, reflectant and mulches
- Regular monitoring of field for disease and insect attack
- Use pheromone trap and attractant
- Promote protected vegetable cultivation under naturally ventilated polyhouse and net house.
- Timely sowing for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population)
- Proper water management

Take care of Aphid, white rust in Mustard, Early, late blight and leaf curling in potato

Mustard/Toria – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Linseed - Follow seed treatment, Irrigate only at critical stages, Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Precautions for pod borer, bud fly insect and powdery mildew disease management.

2A2. Land type- MEDIUM LAND

- a) Cropping system- Rice –Wheat, Rice –Chickpea, Rice –Lentil, Rice-Linseed, Rice-Mustard, Soybean-Vegetable, Maize-Sunflower/ Safflower ,Rice-Lathyrus, Blackgram-Barley, Rice-Fodder
- b) Crop name- Irrigated-Wheat (Zero tillage), Mustard, Potato, Sunflower, Vegetables (Tomato, Chili, Brinjal, capsicum, Cabbage, Cauliflower, Broccoli, vegetable pea, French bean, Radish, Coriander, Amaranthus, Spinach); Rainfed (Zero tillage)- Barley, Safflower, Chickpea, Lentil, Linseed, Lathyrus; Paira cropping- Linseed, Lathyrus; Fodder crop-Oat, Maize
- c) Sowing Window- Wheat (Timely)- 3rd week of October – 4th week of November, Wheat (Late)-1st week of December – 2nd week of December, Mustard- 2nd week of October – 1st week of November, Sunflower-1st week of October-2nd week of October, Linseed- 2nd week of October – 2nd week of November, Chickpea – 2nd week of October – 1st week of November, Lentil- 3rd week opf October- 2nd week of November, Vegetables- 1st week of October – 4th week of November, Lathyrus- 2nd week of October – 4th week of November, Oat- 2nd week of October – 4th week of October, Maize- 2nd week of October – 4th week of October
- d) Variety- Irrigated Wheat- Timely sown (120-125)- HD2967, WH 1105, K307, HD2733; Late sown (105-110)- HD 3059, DBW 14, HI 1563 (seed rate 25 % more than timely sown); Mustard- Pusa mahek,Pusa mustard 25, NRCHB 101, Bharat sarson 1, Pusa 28, 30; Potato- Kufri Surya, Kufri Arun, Kufri Sutlej, Kufri Laukar, Kufri Lalima; Sunflower – DRSP 108 (OP), KBSH 44; Safflower- A 300, Akshagiri 59-2-1, A 1; Chickpea- Jaki 9218, Kak 2, Birsa Chana 3; Lentil – HUL 57,WBL 77, KLS 218 Linseed-Sharda, Priyam, Divya; Lathyrus-Maha Teora, Oat- Kent, Maize- J 1006, Pratap Makka Chari 6

e) Agronomic management practices

- Seed tretment with Azotobacter and Azospirillum and also soil application
- Timely sowing for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population)
- Follow deep summer ploughing
- Proper water management
- Follow seed treatment
- Irrigate only at critical stages
- Pre emergence weedicide application
- Follow RDF, INM and IPM
- Take care of Painted Bud, Aphid, white rust in Mustard; Early, late blight and leaf curling and grub in potato; Collar rot, Dry root rot, Pod borer in Chickpea; Wilt in Lentil.

Wheat-Seed tretment with Azotobacter and Azospirillum and also soil application.Timely sowing for better establishment. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Follow deep summer ploughing. Proper water management. Irrigate only at critical stages(3-6). Pre emergence weedicide application. Follow RDF, INM and IPM. Take case of Loose Smut Disease. 1st irrigation should be after CRI stage i.e at 30-35 DAS

Mustard – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Potato- Seed treatment. Proper spacing. Frequent irrigation. Take care for leaf curling, Early, late blight and grub infestation. Irrigate during cold day and night to get relief from frost attack. Produce smoke during cooler day and night. Earthing up in potato at 30-35 DAS or crop attended the height of 20-25 cm

Sunflower- Seed treatment, Irrigate the field at the time of final land preparation and follow sowing after two days of irrigation for good germination and crop establishment. Follow proper crop geometry for good crop stand. Other than RDF of NPK also needed to apply Sulphur. Follow hoeing at 20-25 DAS, 35-40 DAS. Keep weed free upto 60 days. Use post emergence weedicide. Follow earthing up at Knee height. Irrigate at Critical stages (20-25 DAS, 55-60 DAS, 70-75 DAS. For summer season- 4-10 light irrigation is required

Chickpea - Seed treatment with Rhizobium culture and Phosphate solubilizing bacteria (PSB) and Trichoderma. Management for Collar rot during temperature fall and dry root rot during temperature increment. Pre emergence weedicide application. Irrigate at Critical stages. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray

Lentil - Foliar spray of Sulphur and Boron is necessary. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Follow deep summer ploughing. Proper water management. Follow seed treatment with Rhizobium culture and PSB. Irrigate only at critical stages. Pre emergence weedicide application. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray.

Linseed - Follow seed treatment, Irrigate only at critical stages, Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Precautions for pod borer, bud fly insect and powdery mildew disease management.

Management for wilt disease. One hand weeding followed by two hoeing for management of weeds (HW-20-25 DAS and Hoeing 30-32 and 40-42 DAS).

Lathyrus- Proper seed rate and Follow RDF for potential yield. Follow two nipping in between 25-45 DAS. Irrigate after every nipping. Take care of diseases

Forage-Oat- Proper seed rate for better crop stand. 1st and 2nd cutting at 30 and 45 DAS and 3rd before flowering. Berseem- 1st at 50 DAS and follow 2nd, 3rd and 4th cutting every at an interval of 30-40 days. Lucern- Same as Berseem. Japani Mustard- 1st at 50 DAS during fruiting and rest cutting every at an interval of 30 days. Follow RDF. For Lucern other than N P K use Lime , Boron and Molybdenum micro nutrients for better yield.

2A3. Land type- LOW LAND

- a) Cropping system-Rice –Wheat, Rice- Linseed (Paira), Rice-Vegetable
- b) Crop name- Linseed/ lathyrus (Paira cropping), Wheat, Vegetables (Onion, Fodder Crop- Oat, Maize, lathyrus
- c) Sowing Window- Linseed- 4th week of October – 2nd week of November, Wheat- Timely- 1st week of November- 3rd week of December, Late Sown - 1st week of December- 4th week of December, Lathyrus- 4th week of October – 2nd week of November, Fodder-2nd week of November – 2nd week of December
- d) Variety- Linseed- Sharda, Priyam, Divya, Wheat-Timely), K 8027, HD 2967, K 1006, K 307, HDR 77, HD 2733; Late sown wheat- HD 3059, PBW 373, DBW 14, 39, HI 1563; Lathyrus-Maha Teora; Fodder- Oat— Kent, Maize- Pratap Makka(Chari 6), J 1006, Berseem-Vardhan

e) Agronomic management practices

Linseed - Follow seed treatment. Irrigate only at critical stages. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Follow RDF, INM and IPM. Be cautious for pod borer, bud fly insect and powdery mildew disease management.

Wheat – For surface seeding increase seed rate and Nitrogenous fertilizer by 25 per cent. Remove excess water by making deep furrow around their fields. Planking should be done after seed placement for better germination and crop stands. Follow RDF, INM and IPM. Pre emergence weedicide application

Lathyrus- Proper seed rate and Follow RDF for potential yield. Follow two nipping in between 25-45 DAS. Irrigate after every nipping. Take care of diseases



Forage-Oat- Proper seed rate for better crop stand. 1st and 2nd cutting at 30 and 45 DAS and 3rd before flowering, Berseem- 1st at 50 DAS and follow 2nd, 3rd and 4th cutting every at an interval of 30-40 days, Lucern-Same as Berseem, Japani Mustard- 1st at 50 DAS during fruiting and rest cutting every at an interval of 30 days. Follow RDF. For Lucern other than N P K use Lime , Boron and Molybdenum micro nutrients for better yield.

2 (B) Less than optimum moisture i.e., 25% less than normal, which can happen due to insufficient rainfall during September/October months. Deficit of 20-40% rainfall

2B1 Land type- UP LAND

a) Cropping system- Maize-Toria, Maize-Linseed

b) Crop name – Toria, Linseed

c) Sowing Window- Toria- 3rd week of September – 4th week of September, Linseed- 1st week of October- 3rd week of October

d) Variety- Toria- PT 203, Panchali; Linseed- Sharda, Priyam, Divya

e) Agronomic management practices

- Rain water harvesting and recycling.
- Deeping of water storing structure(Shallow and deep) in April and May month
- Deep summer ploughing in April and May month.
- Strengthening and raising of field bunds in April and May months
- Sowing in defined window for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better crop stand (Plant population)
- Application of Lime or Dolomite (3-5 q/ha) in soil
- Soil application of Sulphur (20 kg/ha) and boron (1kg/ha) in oilseed, pulses and vegetables.
- Foliar spray of Urea (2 %) at flower initiation and pod formation stage in oilseed and pulses
- Follow seed priming (warm water for 4-6 hrs.) before sowing
- Follow seed treatment with fungicide-insecticide-rhizobium
- Follow deep summer ploughing
- Irrigate only at critical stages
- Pre and post emergence weedicide application
- Follow hoeing after hand weeding
- Follow RDF, INM and IPM
- For Water use efficiency use antitranspirant, reflectant and mulches
- Regular monitoring of field for disease and insect attack
- Use pheromone trap and attractant
- Promote protected vegetable cultivation under naturally ventilated polyhouse and net house.
- Zero Tillage for seed placement at proper depth for better germination
- One hand weeding followed by one hoeing for management of germinating weeds
- For Water use efficiency use antitranspirant, reflectant and mulches

Toria—Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Linseed - Follow seed treatment, Irrigate only at critical stages, Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Precautions for pod borer, bud fly insect and powdery mildew disease management.

2B2 Land type- MEDIUM LAND

a) Cropping system- Rice –Barley, Rice –Chickpea, Rice –Lentil, Rice-Linseed, Rice-Fodder

b) Crop name – Rainfed (Zero tillage)- Barley, Chickpea, Lentil, Linseed, Fodder Crop- Oat, Maize, lathyrus

c) Sowing Window- Barley– 3rd week of October – 2nd week of November, Chickpea – 2nd week of October – 2nd week of November, Lentil- 3rd week of October- 2nd week of November, Linseed- 1st week of October – 4th week of October, Fodder-2nd week of October – 2nd week of November



d) Variety- Barley- Ratna, Azad, Vijaya, Amber, Jyoti, Kailash, BG 108; Chickpea- KWR 108, HK 94134, Jaki 9218, Birsa Chana 3; Lentil –WBL 77, KLS 218; Linseed- Sharda, Priyam, Divya, Fodder Crop-Oat— Kent, Maize- Pratap Makka(Cheri 6), J 1006, Lathyrus- Maha Teora

e) Agronomic management practices

Barley- Proper seed rate and spacing for better crop standard. Pre emergence weedicide application. Irrigate at critical stages (2 irrigation at 30-35 DAS and 55-60 DAS). Two weeding in between 25-45 DAS. Follow RDF, INM and IPM. Take care of covered and loose Smut disease and manage for termite attack. Mustard – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Chickpea - Seed treatment with Rhizobium culture and Phosphate solublizing bacteria (PSB) and Trichoderma. Management for Collar rot during temperature fall and dry root rot during temperature increment in. Pre emergence weedicide application. Irrigate at Critical stages. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray

Lentil - Foliar spray of Sulphur and Boron is necessary. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Follow deep summer ploughing. Proper water management. Follow seed treatment with Rhizobium culture and PSB. Irrigate only at critical stages. Pre emergence weedicide application. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray. Management for wilt disease. One hand weeding followed by two hoeing for management of weeds (HW-20-25 DAS and Hoeing 30-32 and 40-42 DAS)

Linseed - Follow seed treatment. Irrigate only at critical stages. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Follow RDF, INM and IPM. Precaution for pod borer, bud fly insect and powdery mildew disease management

Forage- Oat- Proper seed rate for better crop stand. 1st and 2nd cutting at 30 and 45 DAS and 3rd before flowering. Berseem- 1st at 50 DAS and follow 2nd, 3rd and 4th cutting every at an interval of 30-40 days. Lucern-Same as Berseem. Japani Mustard- 1st at 50 DAS during fruiting and rest cutting every at an interval of 30 days. Follow RDF. For Lucern other than N P K use Lime , Boron and Molybdenum micro nutrients for better yield.

2B3 Land type- LOW LAND

- a) Cropping system- Rice- Linseed/Lathyrus (Paira), Rice-Vegetable(Using harvested water), Rice-Fodder
- b) Crop name- Linseed/ lathyrus (Paira cropping), Vegetables (Tomato, Coriander, Radish, vegetable Pea, Spinach, Fodder Crop-Oat, Maize, lathyrus
- c) Sowing Window- Linseed- 4th week of October – 2nd week of November; Lathyrus- 4th week of October – 2nd week of November; Vegetable- 3rd week of November- 4th week of December; Fodder-2nd week of November – 2nd week of December
- c) Variety- Linseed- Sharda, Priyam, Divya, Lathyrus-Maha Teora; Fodder- Oat— Kent, Maize- Pratap Makka (Cheri 6), J 1006

e) Agronomic management practices

Linseed - Follow seed treatment. Irrigate only at critical stages. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Follow RDF, INM and IPM. Precaution for pod borer, bud flies insect and powdery mildew disease management.

Lathyrus- Proper seed rate and Follow RDF for potential yield. Follow two nipping in between 25-45 DAS. Irrigate after every nipping. Take care of diseases

Forage- Oat- Proper seed rate for better crop stand. 1st and 2nd cutting at 30 and 45 DAS and 3rd before flowering. Berseem- 1st at 50 DAS and follow 2nd, 3rd and 4th cutting every at an interval of 30-40 days. Lucern-Same as Berseem. Japani Mustard- 1st at 50 DAS during fruiting and rest cutting every at an interval of 30 days. Follow RDF. For Lucern other than N P K use Lime , Boron and Molybdenum micro nutrients for better yield.



CONTINGENT STRATEGIES FOR LIVESTOCK, POULTRY & FISHERIES

1 Livestock

Suggested contingency measures under DROUGHT event

a) Before the event

Feed and fodder availability

Preservation of surplus fodder, encourage fodder cultivation and tree plantation and also encourage supply of molasses to cattle feed plants

- Preservation of surplus fodder

Green grass is a good source of vitamin A which is present in the form of Carotene. One kg of green grass provides 50mg of vitamin A and 15 to 20g protein to the animal. Cowpea, beans, subabul leaves etc. give 30 to 40g of protein. From grass fodder herbivorous animals get the carbohydrates (energy source), proteins ("building material" of the body) and vitamins (especially carotene), which are the main drives of sustainable operation of the body.

Two methods are available for preserving or conserving the seasonal excess of green fodder, viz. hay making and silage making. Each method has its own limitations and advantageous. Ensiling is preferred on the basis of fodder quality.

Hay making

Hay -refers to cereals, grasses or legumes that are harvested at appropriate stage, dried and stored

Ensilage / Silage making

Silage may be defined as the green succulent roughage preserved under controlled anaerobic fermentation in the absence of oxygen by compacting green chops in air and watertight receptacles.

- Complete Feed Blocks

Supply enriched complete feed blocks containing dry roughage, concentrates/ unconventional supplements 50:50 ratio. Complete feed blocks may be sourced from different commercial sources.

Feeding practices for livestock in India at present separate feeding of roughage and concentrate

❖ Chopped roughage and soaked concentrate mixed together

❖ Chopped roughage mechanically mixed with concentrate as mash

❖ Chopped roughage and concentrate ingredients mixed and densified as Complete Feed Block

Concept of densified complete feeds with fibrous crop residues is a noble way to increase the intake and improve the nutrients utilization. A complete feed block has been defined as a system of feeding all ingredients including roughages, processed and mixed uniformly, to be made available ad lib to the animals.

- Urea molasses mineral block licks

Urea-molasses mineral block lick can sustain the animals by providing protein, energy and essential minerals. It is cost effective, easy to handle and transport and available commercially through milk cooperatives. Therefore, it is required that urea molasses blocks stocks (UMBS) are made available in the rain-deficient areas.

- Methods used for improving nutritive quality of straws and other crop residues like urea treatment

Spray dry roughages such as paddy and wheat straw with about 10% molasses and 2% urea for maintenance of animals in fodder deficit areas.

Preparation of 100 kg roughage-based enriched feed containing 88.8 kg wheat straw or any other straw/stover, 10 kg molasses, 1 kg urea and 0.5 kg mineral mixture will cost about Rs. 375-450 per quintal.

- Utilization of forest byproducts for feeding of livestock

Use of dry and fallen tree leafs like Pipal, Neem, Mango and Kathal etc.

- Making Leaf meal

- Use of conventional and non conventional feeds

- Rice Mills

The main by-products of rice are rice straw, rice husk or hull, and rice bran. Rice straw is produced when harvesting paddy. Rice husks generated during the first stage of rice milling, when rough rice or paddy rice is husked.

- Aquatic plants

❖ One kg DM/100 kg BW

❖ Water hyacinth, aquatic spinach, Stalks & leaves of lotus plant, Hydrilla, Pistia etc.

- Encourage supply of molasses to cattle feed plants

Molasses and Bagasse are the byproducts from sugarcane industry and are available in abundance. They can be used as cattle feed after supplementation with urea. Such a ration is a ready feed during drought and scarcity conditions when nothing else is available for feeding to animals.

- Crop Residue Enrichment & Densification

Crop residues can be fortified with feed ingredients like cakes, barns, grains, molasses, hay, minerals and then densified into blocks or pellets to save on storage and transport costs. Also balanced ration in the form of complete diet or total mixed ration as per need of animals can be supplied for improved productivity.



• Demonstration of Re-vegetation of Common Grazing Land

The grazing lands play an important role in the lives of rural people who are getting fodder, fuel, drinking water from commons. However, such lands are being continuously degraded due to overgrazing and overexploitation by locals. Re-vegetation of such lands on scientific lines suiting to agro-climatic conditions is to be demonstrated through strengthening institutional arrangement at village level. Fodder production from such lands can be enhanced substantially by introducing high yielding cultivated fodder crops, grasses and pasture legumes. An integrated approach of growing cultivated crops, grasses, trees and shrubs under silvi-pastoral/ horti - silvopasture system will improve overall productivity of such land.

Drinking water

Repairs of tube wells, clear off the sludge in the canals and local water catchments and clean the water tanks, large ponds and lakes

Health and Hygiene

Tick damage and tick-borne diseases

- Tick damage - Vaccinate the cattle against tick-borne diseases
- Tick-borne diseases- Vaccination is best done in calves under 6 months of age and one dose is sufficient
- Babesiosis (Red water)- Treatment involves keeping the cattle calm. They should not be driven over long distances and should be injected with Berenil or Imizol. The dose for Berenil is 5 ml of made up solution (1 packet mixed with 12.5 ml of sterile water) for each 100 kg (for example, 20 ml for a 400 kg animal)
- Sarcoptic Mange in pigs- Not applicable before event

Diseases caused by biting insects

- Trypanosomiasis- Fly control is important for prevention of the disease.
- Three-day stiff sickness- Prevention is by vaccination
- Lumpy-skin disease- Prevention is by vaccination

Diet related Disease problems

- Eating plastic bags and wire(Pica)- Feed cattle well, especially in winter. Clear wires and plastic bags from the grazing area. Watch cattle closely when they are grazing. Mineral mixture supplement should be given to the animal
- Poisonous plants- Not applicable before event
- Botulism- Prevention involves vaccination and good nutrition. Burn or bury all carcasses, bones or decaying material

Deficiency diseases

Cattle grazing on drought-stricken pastures can suffer serious depletion of reserves of minerals and vitamins.

- Copper and Cobalt- Not applicable before event
- Calcium, Phosphorous & Vit. D- Not applicable before event
- Vitamin A- Not applicable before event

Infectious Diseases

- Foot and Mouth Disease (FMD)- Vaccination at the age 4 months and above. Booster should be given 1 month after first dose then every six monthly
- Haemorrhagic Septicaemia (HS)- Vaccination at the age 6 months and above. Annually in endemic areas. Vaccinate the animal before onset of monsoon every year preferably in the month of May and June.
- Black Quarter (BQ)- Vaccination at the age 6 months and above. Annually in endemic areas. Vaccinate the animal before onset of monsoon every year preferably in the month of May and June
- Anthrax- Vaccination at the age 4 months and above. Annually in endemic areas. Vaccinate the animal before onset of monsoon every year preferably in the month of May and June.
- Rabies (Post bite therapy only)- Not applicable
- Enterotoxaemia (pulpy kidney)- Vaccinate the animal at the age of 3-4 months, repeat after 15 days and then annually.
- Pneumonia- Not applicable

Non-Infectious Diseases

- Ruminant tympany (Bloat)- Not applicable
- Rumen acidosis- Not applicable
- Intussusception- Deforming should be give
- Pregnancy toxemia (Ketosis)- Fed the pregnant animal with balanced ration.

Poisoning

- Organochlorine compounds- Not applicable
- Organophosphorus compounds- This group consists of malathion, darathion, chlorathion, carbophenothion, demton, dasnon, dimethylparathion, trichlorphon, dioxalthion etc. Symptoms of toxicity are profuse salivation, muscle stiffness, dyspnoea with open mouth breathing, tremors. Treatment consists of administering antidote, usually atropine sulphate.
- Snake bite- Not applicable



b) During the event

Feed and fodder availability

- Lactating and pregnant animals need to be provided enriched feed to meet the requirements and rest of animals be provided the maintenance diet. In case of acute shortage, lactating animals be provided feed meeting 50% of the requirements to maintain minimum level of production.
- Drought tolerant fodder crops (like sorghum PC 6 and MP chari, cowpea - BL 1 and 2) and fodder grasses (like stylo, *cenchrus ciliaris*, *athropogon*, etc.) should be cultivated. Under the mini kit programme, the developmental department need to provide fodder crop seeds in the drought-affected areas.
- Provide salt dose daily through feed (40-50 g of salt per adult animal and 10-20 g for small ruminants and calves).

Issue

- Large scale migration -Creating additional resources in drought prone area
- Grazing of poisonous plants/toxicity problems -Inventory of anti nutritional/toxic factors. Creating awareness in farmer for avoiding nitrate/nitrite HCN poisoning.
- Transport of fodder from normal DPA-Establishing feed and fodder banks. Effective mechanism for distribution of fodder/feed to productive animals. Densification/baling/briquette technologies

Drinking water

Harnessing water through the existing reservoirs and exploitation of groundwater.

Health and Hygiene

Tick damage and tick-borne diseases

- Tick damage - If disease occurs Treat the cattle against tick-borne diseases. Consult Veterinarian.
- Tick-borne diseases- Prevention is by tick control, treatment of diseased animal and vaccination. Consult Veterinarian.
- Babesiosis (Red water)- Treatment involves keeping the cattle calm. They should not be driven over long distances and should be injected with Berenil or Imizol. The dose for Berenil is 5 ml of made up solution (1 packet mixed with 12,5 ml of sterile water) for each 100 kg (for example, 20 ml for a 400 kg animal). Consult Veterinarian.
- Sarcoptic Mange in pigs- Itching; dermatitis; rubbing; scratching; reduced growth rate. Miticidal sprays; pour-ones injection and in-feed premix. Consult Veterinarian.

Diseases caused by biting insects

- Trypanosomiasis- Treated with SURAMIN through intramuscular injection or intravenous infusion if sufficient observation is possible. Consult Veterinarian.
- Three-day stiff sickness- It is important that the animal is given food and water if it is unable to stand
- Animal should be treated by Veterinarian
- Lumpy-skin disease- If your cattle get this disease, you should speak to your state veterinarian

Diet related Disease problems

- Eating plastic bags and wire (Pica) - Mostly occurring in those animals which are having shortage of feeds and fodder and deficiency of Phosphorus. Prevention involves the following: - Feed cattle well, especially in winter. Clear wires and plastic bags from the grazing area. Watch cattle closely when they are Grazing. Mineral mixture supplement should be given to the animal. Once the cow has eaten plastic bags or wire, the only effective treatment is an operation. Consult Veterinarian.
- Poisonous plants- Due to scarcity of feed s and fodder animals used to consume poisonous plans and are more likely to get toxicity. Poisoning can also happen when owner or animal handler move cattle to new paddocks where toxic plants occur. Consult Veterinarian.
- Botulism- Botulism can occur when cattle eat carcass and bone material when there is a lack of feed during drought or if they have a phosphorus deficiency
- Treatment is only possible in the early stages and requires an antitoxin. Consult Veterinarian.

Deficiency diseases

- Cattle grazing on drought-stricken pastures can suffer serious depletion of reserves of minerals and vitamins.
- Copper and Cobalt- Characterized by anorexia and wasting. Deficiency affects growth and fertility of the cattle. Anemia, diarrhoea and unthriftiness occur in extreme cases. Copper or cobalt sulphate in the form of mineral mixture supplement causes rapid disappearance of the symptoms
 - Calcium, Phosphorous & Vit. D- Deficiency may result in rickets in calves and osteomalacia in adults. Mineral supplementation in diet is essential to prevent this deficiency.
 - Vitamin A- Vit. A deficiency occurs in cattle on dry countryside during periods of drought. Symptoms include night blindness, corneal keratinization, ptyriasis, hoof defects, loss of weight and infertility. Animals should have access to green pasture and should be supplied with Vit. A in feed to prevent deficiency.



Infectious Diseases

- Foot and Mouth Disease (FMD)- If outbreak occurs then the animal should be treated. Foot lesion should be washed with soap / detergent and apply Povidon iodine lotion while in mouth lesion boroglyserine should be applied. Consult Veterinarian.
- Haemorrhagic Septicaemia (HS)- If disease occurs animal should be treated with Broad Spectrum Antibiotic like penicillin at higher dose and other supportive medicine as per the symptoms. Consult local Veterinarian.
- Black Quarter (BQ)- If disease occurs animal should be treated with Broad Spectrum Antibiotic like penicillin at higher dose and other supportive medicine as per the symptoms. Consult local Veterinarian.
- Anthrax- If disease occurs animal should be treated with Broad Spectrum Antibiotic like penicillin at higher dose and other supportive medicine as per the symptoms. Consult local Veterinarian.
- Rabies (Post bite therapy only)- Vaccinate the animal immediately after suspected bite. Booster should be given on 3, 7, 14, 28 and 90 (optional) days after first dose.
- Enterotoxaemia (pulpy kidney)- Not applicable
- Pneumonia- Not applicable

Non-Infectious Diseases

- Rumenal tympany (Bloat)- Not applicable
- Rumen acidosis- Ingestion of large amounts of highly fermentable carbohydrate feeds causes an acute illness due to excess production of lactic acid in the rumen. Clinically, the disease is manifested by dehydration, blindness, recumbency, complete rumen stasis and a high mortality rate. Normal saline, sodium bicarbonate and antihistaminic are advised.
- Intussusceptions- It occurs commonly due to nodular worms, change in feed and local intestinal problems. The animal is dull, off-feed, kicking at the belly with no rise of temperature, frequent straining with no defecation, colic symptoms, and at later stages, recumbency. Emergency surgery is the only rational treatment.
- Pregnancy toxæmia (Ketosis)- It is a highly fatal disease caused due to a decline in the plane of nutrition and short periods of starvation (40 hrs) during the last two months of pregnancy. Treatment comprises intravenous administration of 50% glucose. Supply of molasses in the ration and concentrate in the last two months of pregnancy helps in preventing the condition.

Poisoning

- Organochlorine compounds- Not applicable
- Organophosphorous compounds- This group consists of malathion, darathion, chlorathion, carbophenothion, demton, dasnon, dimethylparathion, trichlorphon, dioxalthion etc. Symptoms of toxicity are profuse salivation, muscle stiffness, dyspnoea with open mouth breathing, tremors. Treatment consists of administering antidote, usually atropine sulphate.
- Snake bite- Usually bitten on the scrotum or udder. The presence of hair may obscure the typical fang marks. Prolonged pain, muscular weakness, impaired vision, nausea and paralysis are generally exhibited along with symptoms of shock. If anti-venin is not available and the bite is located in an area where a tourniquet cannot be applied, excision of an area of skin and sub-cutaneous tissue can be life-saving

c) After the event

Feed and fodder availability

Promotion of fodder seed production, cultivation and storage, establishment of fodder block making machines in fodder surplus areas

Post flood feeding management

- Animal should not be allowed to graze in water logged area
- Feeds to be protected from fungal contamination & wet feeds to be dried & fed
- Provides clean drinking water to animals
- Provide ready to eat feed blocks particularly the pregnant and lactating animals
- Requirement of energy may be met providing crude molasses
- Top feeds/ tree leaves available in the area to be provided to meet the DM requirement

Specific contingencies can be adopted for livestock feeding depending upon availability as under in different regions during drought situation

Neem seed kernel cake (NSKC), Saw dust, Paper waste, Agave (Ketki), Cactus, Tree leaves and vegetable leaves, Cher leaves and fruits, Straw and gotars, Sugarcane bagasse as animal feed and Use of damaged grains as feed

Feed and fodder availability

Promotion of fodder seed production, cultivation and storage, establishment of fodder block making machines in fodder surplus areas

Post flood feeding management

- Animal should not be allowed to graze in water logged area



- Feeds to be protected from fungal contamination & wet feeds to be dried & fed
- Provides clean drinking water to animals
- Provide ready to eat feed blocks particularly the pregnant and lactating animals
- Requirement of energy may be met providing crude molasses
- Top feeds/ tree leaves available in the area to be provided to meet the DM requirement

Specific contingencies can be adopted for livestock feeding depending upon availability as under in different regions during drought situation

Neem seed kernel cake (NSKC), Saw dust, Paper waste, Agave (Ketki), Cactus, Tree leaves and vegetable leaves, Cher leaves and fruits, Straw and gotars, Sugarcane bagasse as animal feed and Use of damaged grains as feed

Drinking water

To strengthen reservoirs by promoting recharging of water and rain water harvesting during rainy season.

Health and Hygiene

Tick damage and tick-borne diseases

- Tick damage - Treat the cattle against tick-borne diseases. Consult Veterinarian.
- Tick-borne diseases- Prevention is by tick control, treatment of diseased animal and vaccination. Consult Veterinarian.
- Babesiosis (Red water)- Treatment involves keeping the cattle calm. They should not be driven over longdistances and should be injected with Berenil or Imizol. The dose for Berenil is 5 ml of made up solution (1packet mixed with 12.5 ml of sterile water) for each 100 kg (for example, 20 ml for a 400 kg animal).Consult Veterinarian.
- Sarcoptic Mange in pigs- Not applicable after event

Diseases caused by biting insects

- Trypanosomiasis- Treated with SURAMIN through intramuscular injection or intravenous infusion if sufficient observation is possible. Consult Veterinarian
- Three-day stiff sickness- It is important that the animal is given food and water if it is unable to stand.
- Animal should be treated by Veterinarian
- Lumpy-skin disease- If your cattle get this disease, you should speak to your state veterinarian

Diet related Disease problems

- Eating plastic bags and wire (Pica)- Feed cattle well, especially in winter. Clear wires and plastic bags from the grazing area. Watch cattle closely when they are grazing. Mineral mixture supplement should be given to the animal
- Poisonous plants- Not applicable
- Botulism- Prevention involves vaccination and good nutrition. Burn or bury all carcasses, bones or decaying material

Deficiency diseases

Cattle grazing on drought-stricken pastures can suffer serious depletion of reserves of minerals and vitamins.

- Copper and Cobalt- Not applicable
- Calcium, Phosphorous & Vit. D- Not applicable
- Vitamin A- Not applicable

Infectious Diseases

- Foot and Mouth Disease (FMD)- If outbreak occurs then the animal should be treated. Foot lesion should be washed with soap / detergent the apply Povidon iodine lotion while in mouth lesion boroglyserine should be applied. Consult Veterinarian.
- Haemorrhagic Septicaemia (HS)- Not applicable

- Black Quarter (BQ)- Not applicable
- Anthrax- Not applicable
- Rabies (Post bite therapy only)- Not applicable
- Enterotoxaemia (pulpy kidney) - It affects the animals in a high state of nutrition on a lush feed, grass or grain. Morbidity rates seldom exceed 10% but mortality rate approximates 100%. Under certain conditions, the organism proliferated rapidly in the intestines and produces lethal quantity of toxin. Suphadimidine with other supportive medicine may be effective for treatment
- Pneumonia- It is one of the most common and important pathological conditions. It is characterized clinically by increased respiration, coughing and abdominal breathing. Treatment with broad spectrum antibiotic, nabulization and other supportive drugs is effective.

Non-Infectious Diseases

- Ruminal tympany (Bloat)- It is the over-distension of the left flank either due to free gas or froth. This is generally encountered in "greedy feeders" when lush green pasture is available. Oral administration of sweet oil with turpentine oil or at times with formalin is advised.
- Rumen acidosis- Not applicable
- Intussusceptions- Not applicable
- Pregnancy toxæmia (Ketosis)- Not applicable

Poisoning

- Organochlorine compounds- This group includes DDT, BHC, lindane, aldrin, dieldrin, chlordane, toxaphane, methoxychlor etc. which are used as pesticides on crops. Toxicity symptoms include increased excitability and irritability followed by muscle tremors, weakness, paralysis etc. Treatment consists of administering antidote, usually short-acting barbiturates.
- Organophosphorus compounds- This group consists of malathion, darathion, chlorathion, carbophenothion, demeton, dasnon, dimethylparathion, trichlorphon, dioxalthion etc. Symptoms of toxicity are profuse salivation, muscle stiffness, dyspnoea with open mouth breathing, tremors. Treatment consists of administering antidote, usually atropine sulphate.
- Snake bite- Usually bitten on the scrotum or udder. The presence of hair may obscure the typical fang marks. Prolonged pain, muscular weakness, impaired vision, nausea and paralysis are generally exhibited along with symptoms of shock. If anti-venin is not available and the bite is located in an area where a tourniquet cannot be applied, excision of an area of skin and sub-cutaneous tissue can be life-saving

2 Poultry

Suggested contingency measures under DROUGHT event

a) Before the event

Shelter management

Optimum space should be provided. Orientation of shed (Long axis) should be in North – South. Plantation of tree around shed to provide cool environment. Provision of ad lib. Fresh water

Shortage of feed ingredients

Storage of feed

Drinking water

Manage clean drinking water. Storage facility should be made. Water quality should be checked before drinking to animal

Health and disease management

- Newcastle Disease- regular vaccination - Broiler birds should be with RD vaccine (Lasota 'F' strain) at the age of 4-7 days through Intra-nasal or Intra-ocular route. Layer birds should be vaccinated with NDV vaccine at the age of 9-14 day, 4 weeks, 13-14 weeks in drinking water/eye drop. Then at the age of 17 week with NDV vaccine through Intra-muscular (IM) route
- Marek's disease Marek's disease- Birds should be vaccinated with Herpes virus turkey vaccine at the age of 1 day through Subcutaneous route.
- Fowl pox- Chick embryo adopted fowl pox vaccine at the age 6-8 weeks. It important for the layer and broiler birds.
- Drop in Egg Production or Quality- Not applicable
- Nervous Signs and Lameness- Not applicable
- Diarrhoea- Not applicable
- Upper Respiratory Diseases- Vaccination against the some of the viral diseases like Newcastle disease, influenza, infectious bronchitis, infectious laryngotracheitis which are also responsible for the respiratory symptoms can prevent this syndrome. Antifungal and antiparasitic drugs should be given.

Heat Wave

Plantation of tree around shed to provide cooler environment. Proper ventilation should be provided. Optimum space should be provided. Orientation of shed (Long axis) should be in East- West. Plantation of tree around shed to provide cool environment. Provision of ad lib. Fresh water. Manage green fodder and silage preparation. Height of roof should be minimum 220 – 240 cm. Roof of shed should be painted with white.

Cold Wave

Provide ad lib fresh water. Proper ventilation should be provided. Optimum space should be provided. Orientation of shed (Long axis) should be in North – South. Plantation of tree around shed to provide break cold wave. Provision of ad lib. Fresh water. Manage green fodder and silage preparation. Height of roof should be minimum 220 – 240 cm Roof of shed should be painted with Black Floor of shed should be Dry

b) During the event

Shelter management

Optimum space should be provided, there must not be overcrowded. Protect the animal from direct sun light. Try to provide them cool water. Proper ventilation should be maintained. Allow for grazing during night/early morning. Provision of ad lib. Fresh water

Shortage of feed ingredients

Provide non conventional feed, supplement anti oxidant and anti stress



Drinking water

Provide clean fresh and cold drinking water all the time. Water availability may increase by 20-50% depending upon feed quality and environmental temperature. Soft drinking water should be preferred. Add vit-C and other anti stress ingredients with water

Health and disease management

- Newcastle Disease- Vaccination and treatment of diseased one. Newcastle disease is the most important disease for poultry farmers around the world. This disease causes a large number of deaths in chickens and huge losses to farmers and the industry. Diseased birds should be slaughtered immediately. Consult Veterinarian.
- Marek's disease Marek's disease- It is one of the important diseases of poultry caused by virus. Mortality is very high and causes economic losses to the farmer and poultry industry.
- Fowl pox- It is a viral infection of chickens and turkeys characterized by proliferative lesions in the skin (Cutaneous form), it also affect the GI tract and respiratory tract (Diphtheritic form)
- Drop in Egg Production or Quality- There are many different types of organisms that can cause a drop in egg production or quality. These include: Bacteria (E. coli, Salmonella), Mycoplasma, Viruses (Newcastle disease, influenza, infectious bronchitis, infectious laryngotracheitis, avian encephalomyelitis, egg drop syndrome). The Parasites, lack of Nutrition and Stress factor also support the onset of this condition. Adding vitamins and minerals to the water or feed may help. Consult Veterinarian
- Nervous Signs and Lameness- Chickens lie down because they cannot stand up. They also walk with a limp or are reluctant to move. Nervous signs may include staring into the sky, pulling the head and neck over their backs, paralysis. Sores on the breast muscles from lying down
- Diarrhoea- The stool or droppings of the chickens are not firm but very loose, watery, not of the normal colour and may contain blood. This may cause the feathers of the vent to be soiled and caked together, Depression, reluctance to eat, drink and move about, poor growth and death. Use an antibiotic or coccidiostatic drug in the water that was recommended by the animal health technician or veterinarian in the water for 3 to 5 days. Stress preparations that contain electrolytes, vitamins and minerals can be added to the water
- Upper Respiratory Diseases- Not applicable

Heat Wave

Water sprinkling to animal. Prevent the animal from direct sunlight. Optimum space should be provided, there must not be overcrowded. Fan should be provided to make the body cool. Try to provide them cool drinking water all time

Proper ventilation should be maintained. Allow for grazing during night/early morning. Try to provide green fodder and silage. Stocking density should be less. Roof should be covered with tiles, paddy, dry leaves to protect from direct sun light

Cold Wave

Luke worm water should be provided at least 4-6 times a day. Prevent the animal from direct cold wave by closing the windows and doors. Optimum space should be provided, there must not be overcrowded. Proper ventilation should be maintained. Allow for grazing during sunny day time. Try to provide green fodder and silage. During extreme cold condition electric heater of wood fire heat should be provided. Try to make the environment inside and outside the shed dry. Gunny bags or blanket may be used to cover the body. Bedding material like paddy straw, Gunny Bag, Bhusa should be provided specially to young one shed.

c) After the event

Shelter management

Optimum space should be provided. Take care and fulfill the requirement of water along with proper nutrients. Take care of proper feeding as per requirement. Provision of ad lib. Fresh water

Shortage of feed ingredients

Not applicable

Drinking water

Provide adlib. Drinking water

Health and disease management

- Newcastle Disease- Disposal of dead birds
- Marek's disease Marek's disease- Disposal of dead birds
- Fowl pox- Disposal of dead birds
- Drop in Egg Production or Quality-Not applicable
- Nervous Signs and Lameness- A complete hygiene and disinfection programme should be planned together with the animal health technician or veterinarian. Antibiotics will only be effective against bacteria and can be used as recommended. If it is a viral disease, such as Newcastle disease, urgent steps have to be taken to prevent possible spread because it causes serious production losses
- Diarrhoea- Disposal of dead birds



- Upper Respiratory Diseases- There is many different types of organisms that can cause disease in the upper respiratory tract. These include: Mycoplasma Bacteria (E. coli, Pasteurella, Haemophilus), Viruses (Newcastle disease, influenza, infectious bronchitis, infectious laryngotracheitis), Parasites (mites and worms) And Fungi (Aspergillus). Cold stress is also one of the predisposing factors for the occurrence of respiratory problems. Use an antibiotic drug that was recommended by your animal health technician or veterinarian in the water for 3 to 5 days
- Stress preparations that contain electrolytes, vitamins and minerals can be added to the water

Heat Wave

Optimum space should be provided. Take care and fulfill the requirement of water along with proper nutrients. Take care of proper feeding as per requirement. Provision of ad lib. Fresh water

Cold Wave

Provide ad lib. Normal drinking water. Optimum space should be provided. Take care and fulfill the requirement of water along with proper nutrients. Take care of proper feeding as per requirement. Provision of ad lib. Fresh water

3 Fisheries

Suggested contingency measures under DROUGHT event

a) Before the event

Aquaculture

- Shallow water in ponds due to insufficient rains/inflow- Increase depth of pond, Repair dyke, outlet and inlet of pond; Prepare duck/pig house & stock pig @ 50-60,duck @ 450-500 no/ha if farmer involve in Integrated fish farming, Allow manure and urine directly in pond, Remove unwanted, predatory & old fishes and for this apply Mahua oil cake @ 2500kg/ha. Fixed net in outlet & inlet to prevent escaping of fish, Plough the pond and apply lime @ 250 kg/ha, Check the natural feed (plankton)@ 1.0 1.5 ml/50 lt. water; otherwise apply organic manure, Stock yearling (stunted grow fish) @ 6,000-8,000 no/ha
- Impact of salt load build up in ponds / change in water quality- Prevent entry of polluted water or apply lime at inlet.

Heat wave and cold wave

- Changes in pond environment (water quality) - Increase depth of pond. Reduce application of organic manure and supplementary feeds
- Health and Disease management- Apply lime @ 50 kg/ha

b) During the event

Aquaculture

- Shallow water in ponds due to insufficient rains/inflow- Reduce the stocking density from 25000 fry (1inches size) to 10000-15000/ha, fingerling 6,000-8,000 no/ha. Check the availability of natural food, if it is not sufficient provide supplementary feed at fixed place, time, amount and ratio & if it is more greenish stop supplementary feed & manure, store manure in separate place for agricultural purpose. Check the growth & health status by regular netting, Apply lime @ 50kg/ha.
- Impact of salt load build up in ponds / change in water quality- Apply lime @ 50 kg/ha on every 15-30 days. Aerate the water as per need

Heat wave and cold wave

- Changes in pond environment (water quality) - Stop or reduce supplementary feed and manure, Remove bigger size fishes. Reduce/stop application of feed and fertilizer.
- Health and Disease management- Apply lime/salt as per need

c) After the event

Aquaculture


- Shallow water in ponds due to insufficient rains/inflow- Remove the bigger size fishes (0.5kg). In winter season fish reduces feed consumption so reduce supplementary feed, duck start egg laying so they should not allow before 9'oclock otherwise loss of egg is possible, pig may attain 50 – 60 kg so that can be sell out and again stock same no of piglets. Apply bleaching powder @ 10kg/ha at place of litter deposition.
- Impact of salt load build up in ponds / change in water quality- Apply lime as per need @ 50 kg/ha

Heat wave and cold wave

- Changes in pond environment (water quality) - Stop or reduce supplementary feed and manure, Remove bigger size fishes. Harvest the bigger fishes, Reduce/stop application of supplementary feed, Apply lime @ 50 kg/ha and potassium per magnet in perforated plastic ball- 5-10g in each ball
- Health and Disease management- Apply lime/salt as per need

SARAIKELA - KHARSAWAN DISTRICT

Sl. No.	CONTENTS	Page No.
1.	District Agriculture profile and land use pattern	1
	PART-I CONTINGENCY PLAN FOR KHARIF	
2.	Contingency plan for 2 weeks delay in monsoon arrival (onset in 4th week of June) A1, A4, A7. Upland A2, A5, A8. Midland A3, A6. Lowland Contingency plan for 4 weeks delay in monsoon arrival (onset in 2nd week of July) B1, B4, B7. Upland B2, B5, B8. Midland B3, B6, B9. Lowland Contingency plan for 6 weeks delay in monsoon arrival (onset in 6th week of July) C1, C6. Upland C2, C4, C7. Midland C3, C5, C8. Lowland	2-12 12-23 24-32
	PART-II	
3.	A. Contingency Plan for normal monsoon onset followed by 15-20 days dry spell A1. Upland A2. Midland A3. Lowland	33-34
4.	B. Contingency plan for mid season drought Upland B1. At vegetative phase B2. At Flowering/Fruiting stage Midland B3. At vegetative phase B4. At Flowering/Fruiting stage Lowland B5. At vegetative phase B6. At Flowering/Fruiting stage	34-36
5.	C. Contingency plan for Late season drought/Terminal drought (Early withdrawal of monsoon) At fruiting/pre physiological maturity stage C1.1. Upland C1.2. Midland C1.3. Lowland	37
	PART-III	
6.	A. Unusual rains : Continuous high rainfall in a short span leading to water logging Crop management Pest and disease management	38-40
7.	B. Extreme weather events (Hail storm, Heat wave, Cold wave, Frost) CONTINGENCY PLAN FOR RABI	40-41
8.	1. Sowing window information 2. Contingency measures for field crops grown with residual moisture under rainfed condition 2(A) Optimal residual moisture 2A.1 Upland 2A.2 Midland 2A.3 Lowland 2 (B) Less than optimal soil moisture (25 % less than normal-Deficit of 20-40 % rainfall) 2B.1 Upland 2B.2 Midland 2B.3 Lowland	42 42-46
	CONTINGENCY STRATEGIES FOR LIVESTOCK, POULTRY AND FISHERIES	
9.	1. Livestock a) Before the event b) During the event c) After the event 2. Poultry a) Before the event b) During the event c) After the event 3. Fisheries a) Before the event b) During the event c) After the event	47-54





District Agriculture profile

Agro-Climatic Zone	AZ - 59		
Agro Ecological Sub Region (ICAR)	Eastern plateau (chotanagpur) And Eastern Ghats, Hot Subhumid Eco-Region (12.3)		
Agro-Climatic Zone (Planning Commission)	Eastern Plateau And Hills Region (VII)		
Agro Climatic Zone (NARP)	South Eastern Plateau Sub Zone - VI		
Meteorological Subdivision	8 th		
List all the districts falling under the NARP Zone (>50% area falling in the zone)	East Singhbhum, Saraikela-Kharsawan, West Singhbhum		
Geographic coordinates of district headquarters	Latitude		Altitude
	22° 29'02" N - 23°09' 39" N		85° 30' 47"E - 86°14'56" E 388m
Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/ RRTTS	Zonal Research Station (ZRS), Darisai, Birsa Agricultural University, Ranchi		
Mention the KVK located in the district with address	Krishi Vignan Kendra, Darisai, Vill-Kharia, PO. Barakhurshi Distt. East Singhbhum-832 304		
Name and address of the nearest Agromet Field Unit (AMFU, IMD) for agro-advisories in the Zone	ZRS, Darisai, Eas Singhbhum-832 304		

Land use pattern of the district (area: '000 ha)

Geographical area	Cultivable area	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable wasteland	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
237.231	95.046	64.972	21.559	3.557	11.091	3.259	16.477	31.292	21.27



CONTINGENCY PLAN FOR KHARIF

PART-I

A Monsoon/Weather Situation: 2 Weeks Delay (Onset: 4th Week of June) – Early Season Drought

A1. Major Farming Situation/Land Situation: Upland High rainfall, High rainfall, shallow red sandysoil	
Normal Crop/cropping system	Upland Rice , Finger millet, Pigeonpea + Sorghum, Pigeonpea + Maize
Suggested Contingency measures	
a) Change in crop/cropping system	
<p><u>Discard rice Crop</u> <u>Sole crop</u> Pigeonpea, Finger millet, Groundnut, Soybean, Maize, Blackgram, Greengram Sorghum, Cowpea, lady's Finger <u>Intercrop :</u> Pigeonpea + Lady's finger (1:2), Pigeonpea + Sorghum (1:1), Pigeonpea + Maize (1:1), Pigeonpea + Groundnut (1:2), Maize + lady's finger (1:2), Maize + Cowpea (1:2)</p>	
<p><u>Horticulture Crop</u> Vegetable : Tomato/ Brinjal/French bean/ Cucumber_ <u>Variety</u> Pigeonpea- Birsa Arhar (200-220), Narendra Arhar 1, ICPH 2671 (200) Finger millet- BM 2, BM 3 (BBM 10), VL 149 Groundnut- Birsa mungfali 3, 4, Girnar 3 Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335 Birsa safed soybean 2 (105-110), RKS 18, RAUS 5 Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80) Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain) Greengram- HUM 16, IPM-02-03-60-65, SML 668 (summer) Sorghum- CSV 20-110-20, MP cheri, CSV 1616 Cowpea-rainy – Birsa sweta(80-90), Swarn sweta(80-90), Swarn harit (80-90) Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika <u>Vegetable crop</u> Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Hybrid- Swarn sampada, Swarn samridh, Pusa hybrid 1 Suraksha Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6 French bean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata Cucumber-Japani long green, Pusa sanyog, Balamkhira, Puna khira, Swarn ageti</p>	
b) Agronomic measures	
<ul style="list-style-type: none"> • Summer deep ploughing with Mouldboard or disc • Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations. • RD Spacing • Zero tillage practices • Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing • RDF and in case of Intercropping reduce 1/3rd dose for intercrop • Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables • Bund construction for unbunded upland • Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables • Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables • Inter-cropping to meet the consequences of occasional Drought. • Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed. • In case of phosphogypsum for soil application apply @ 120 kg/ha • Lime or dolomite application for pulses and oilseed @ 3-5 q/ha in furrow at the time of sowing. • In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha • Follow recommended seed rate • Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha • Foliar application of Urea 2% solution + lime in lady's finger • Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure 	

c) Remarks on Implementation

- Linkage with RKVY , ATMAs, and NFSM
 - Vermicomposting through KVKS ATMAs and NHM
 - Goatry and poultry rearing through KVKS, ATMAs and Veterinary Dept of. Govt. and BAU for livelihood support.
 - Awareness about balanced use of fertilizers to increase their fertility, productivity and sustainability
 - A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology.
 - Awareness for more and more use of organic manures, biopesticides for organic cultivation with IFS (eight components linkages)
 - Upland- 15-20 % upland area should be covered with orchard
1. Mango based orchard-

Variety- Amrapali (30 June-5 July), Mallika (15020 JuneRegular bearer), Sunder langra(15-20 May)
 Spacing- 5 m X 5m

 - i) Recommended package of Practices- Intercrops
 - a) Mango + Papaya (Filler crop for two years) + Blackgram (rainy)/ Chickpea
 - b) Mango + Custard apple (for 10 years and renovate or remove after 10 years) + Blackgram/Chickpea
 - Variety- Langra (15 June)/Bombay green(15 May)/ Himsagar (20-25 May irregular bearer),
 Spacing- 10 m X 10m
 - ii) Recommended package of practices
 - a) Mango + Guava(Upto 10 years as filler) + Papaya (Less than 3 years) + Blackgram-Chickpea/Lentil
 - b) Mango + Lemon + Papaya + Rabi pulses/vegetables
 - c) Mango + Custard apple + Papaya + Blackgram – Pea/Chickpea/Lentil/ Vegetables
 2. Guava base orchard-

Variety- Arka Mridula, Pant Prabhat, Allahabad safeda, L 49
 Spacing- 5m X 5m

 - Recommended package of practices- Intercrops
 - a) Guava + Papaya (For 3 years) + Blackgram-Chickpea
 - b) Guava + Custard apple + Blackgram/Soybean- Pea/Vegetables
 3. Ber Based Orchard –

Variety- Banarsi, Karakka, Gola, Apple ber
 Spacing- 5m X 5m

 - Recommended package of practices Intercrops

Ber + Custard apple + Sesame/Blackgram- Toria/Linseed/Safflower
- N.B.-** Cucurbits, beans or any creeper or climber vegetable should be avoided
- Field crops having height more than one meter should be avoided such as Pigenpea, Maize, Sorghum
 - After 3-5 years when shading effects started shade loving crops like ginger, Turmeric, Ol or leafy vegetables should be grown
 - In citrus leaf minor and aphid susceptible crops should be avoided
 - Aphid should be managed of mustard /atoria taken in citrus orchard
4. Cashewnut based orchard for Kolhan region
 5. Cassava should be grown for the requirement as feed for pig animals
 6. Moringa should also be grown as fodder or vegetable purpose on upland main field bunds as shelter belt/ wind break. Every year pruning and thinning should be followed for bushy look.

A2. Major Farming Situation/Land Situation: Midland High rainfall, slightly deep yellow loamy to sandy loam soil

Normal Crop/cropping system	Rice, Finger millet, Pigeonpea, Sorghum, Maize, Pigeonpea + Sorghum,Pigeonpea + Maize
-----------------------------	---

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807

Don 3

Ridge and Furrow method or raise bed broad furrow: Replace Rice with cereal/ Pulse/ Vegetable
 Cereal – Maize/ Sorghum

Pulse- Pigeonpea + Lady's Finger / Groundnut/ Blackgram/ Finger millet

Vegetable- Tomato,/Brinjal, Cucurbits/ Chili/ Lady's finger/ Cowpea/ Dolichos bean

Variety

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)



Sorghum- CSV 20-110-20, MP cheri, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)

Groundnut- Birsa mungfali 3, 4, Girnar 3

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80) small grain

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149

Vegetable crop

Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Hybrid- Swarn sampada, Swarn samridhi, Pusa hybrid 1 Suraksha Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6

Cucurbits

Bitter gourd- Arka harit, Pusa domausami,

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, Local

Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white

Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,

Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi

Chili- Spices- AndhraJyoti, Pusasadabahar, NP 46, Jwala, KA 2, California wonder, Chinese giant, Yellow wonder, Bharat

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Creeper- Birsa sweta, Swarna sweta, Swarn harit

Dolichos bean-Swarna utkrist, Swarna rituwar

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INP
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P_2O_5 + 1 kg K_2O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P_2O_5 : K_2O (Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P_2O_5 + 40 K_2O /ha ((Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K_2O
- DSR-Use plastic drum seeder rice tools
- Follow RDF and
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/l.t. water, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04%, Blast- Beam or Tricyclazole @ 0.6 gm /l.t water, Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/t. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/l.t. water, Mosaic- Methyl Demeton @ 1.5 ml/l.t. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Tricoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/l.t. water

c) Remarks on Implementation

- A campaign through RKVY, ATMAs, NFSM, KVKS, NHM and other State Govt. line departments are needed to be launched through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.



A3. Major Farming Situation/Land Situation: Lowland High rainfall, medium depth, grayish heavy textured clay loam soil.

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2</u> in <u>Don 1</u> DSR(Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days) Transplanting (Hybrid rice varieties) Var.-Arize 6444 (Gold), Uday 111, PHB 71, 26P52, 25P25	
b) Agronomic Measures	
<ul style="list-style-type: none"> • Staggered Nursery raising by MAT/ DAPOG method • Follow community based nursery raising • Follow RDF,INPM • Use Post emergence weedicide • Use early to mid early duration of rice variety. • Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² • Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice • Topdressing above mentioned dose 10-15 days after sowing • In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling • Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm • Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering. • DSR-Use plastic drum seeder rice tools • Use of post weedicide • Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lt. water, Gundhi bug, leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /lt water 	
c) Remarks on Implementation	
<ul style="list-style-type: none"> • Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme • Supply of Plastic drum seeder through line departments • Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept. • Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon. • Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates • Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation. 	

A4. Major Farming Situation/Land Situation: Upland High rainfall, shallow sandy soil

Normal Crop/cropping system	Upland Rice, Pigeonpea, Maize, Pigeonpea + Maize, Pigeonpea + Sorghum
Suggested Contingency measures	
a) Change in crop/cropping system	
<p><u>Discard rice Crop</u> <u>Sole crop</u> Pigeonpea, Finger millet, Groundnut, Soybean, Maize, Blackgram, Greengram Sorghum, Cowpea, lady's finger <u>Intercrop</u> Pigeonpea + Lady's finge (1:2), Pigeonpea + Sorghum (1:1), Pigeonpea + Maize(1:1), Pigeonpea + Groundnut (1:2), Maize + lady's finger (1:2), Maize + Cowpea (1:2)</p>	
<p><u>Horticulture Crop</u> Vegetable : Tomato/ Brinjal/French bean/ Cucumber <u>Variety</u> Pigeonpea- Birsa Arhar (200-220), Narendra Arhar 1, ICPH 2671 (200) Finger millet- BM 2, BM 3 (BBM 10), VL 149 Groundnut- Birsa mungfali 3, 4, Girnar 3 Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335 Birsa safed soybean 2 (105-110), RKS 18, RAUS 5</p>	

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain)

Greengram- HUM 16, IPM-02-03-60-65, SML 668 (summer)

Sorghum- CSV 20-110-20, MP cheri, CSV 1616

Cowpea-rainy – Birsa sweta(80-90), Swarn sweta(80-90), Swarn harit (80-90)

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Vegetable crop

Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Hybrid- Swarn sampada, Swarn samridh, Pusa hybrid 1 Suraksha Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6

French bean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata

Cucumber-Japani long green, Pusa sanyog, Balamkhira, Puna khira, Swarn ageti

b) Agronomic measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin@ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbundled upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite application for pulses and oilseed @ 3-5 q/ha in furrow at the time of sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Dry seeding with 15% to 20% higher seed rate. Seed treatment with Rhizobium in pulses. Maximum use of organic manure

c) Remarks on Implementation

- Linkage with RKVY , ATMAS, and NFSM
- Vermicomposting through KVKS ATMAS and NHM
- Goatry and poultry rearing through KVKS, ATMAS and Veterinary Dept of. Govt. and BAU for livelihood support.
- Awareness about balanced use of fertilizers to increase their fertility, productivity and sustainability
- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology.
- Awareness for more and more use of organic manures, biopesticides for organic cultivation with IFS (eight components linkages)
- Upland- 15-20 % upland area should be covered with orchard

1. Mango based orchard-

Variety- Amrapali (30 June-5 July), Mallika (15-20 June regular bearer), Sunder langra(15-20 May)

Spacing- 5 m X 5m

i) Recommended package of Practices- Intercrops

a) Mango + Papaya (Filler crop for two years) + Blackgram (rainy)/ Chickpea

b) Mango + Custard apple (for 10 years and renovate or remove after 10 years) + Blackgram/Chickpea



Variety- Langra (15 June)/Bombay green(15 May)/ Himsagar (20-25 May irregular bearer), Spacing- 10 m X 10m						
ii) Recommended package of practices a).Mango + Guava(Up to 10 years as filler) + Papaya (Less than 3 years) + Blackgram-Chickpea/Lentil b).Mango + Lemon + Papaya + Rabi pulses/vegetables c).Mango + Custard apple + Papaya + Blackgram – Pea/Chickpea/Lentil/ Vegetables						
2. Guava base orchard- Variety- Arka Mridula, Pant Prabhat, Allahabad safeda, L 49 Spacing- 5m X 5m Recommended package of practices- Intercrops a) Guava + Papaya (For 3 years) + Blackgram-Chickpea b) Guava + Custard apple + Blackgram/Soybean- Pea/Vegetables						
3. Ber Based Orchard – Variety- Banarsi, Karakka, Gola, Apple ber Spacing- 5m X 5m Recommended package of practices Intercrops Ber + Custard apple + Sesame/Blackgram- Toria/Linseed/Safflower						
N.B.- Cucurbits, beans or any creeper or climber vegetable should be avoided <ul style="list-style-type: none">• Field crops having height more than one meter should be avoided such as Pigenpea, Maize, Sorghum• After 3-5 years when shading effects started shade loving crops like ginger, Turmeric, OI or leafy vegetables should be grown• In citrus leaf minor and aphid susceptible crops should be avoided• Aphid should be managed of mustard /atoria taken in citrus orchard 4. Cashewnut based orchard for Kolhan region 5. Cassava should be grown for the requirement as feed for pig animals 6. Moringa should also be grown as fodder or vegetable purpose on upland main field bunds as shelter belt/wind break. Every year pruning and thinning should be followed for bushy look.						
A5. Major Farming Situation/Land Situation: Midland High rainfall, shallow sandy soil <table border="1"><tr><td>Normal Crop/cropping system</td><td>Rice, Pigeonpea, Maize, Pigeonpea + Maize , Pigeonpea + Sorghum</td></tr><tr><td align="center" colspan="2">Suggested Contingency measures</td></tr><tr><td align="center" colspan="2">a) Change in crop/cropping system</td></tr></table> <p><u>Don 2</u> DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807</p> <p><u>Don 3</u> Ridge and Furrow method or raise bed broad furrow: Replace Rice with cereal/ Pulse/ Vegetable Cereal – Maize/ Sorghum Pulse- Pigeonpea + Lady's Finger / Groundnut/ Soybean/ Finger millet/ Blackgram Vegetable-Radish/ Lady's finger/ Cowpea/ Cucurbits <u>Variety</u> Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80) Sorghum- CSV 20-110-20, MP cheri, CSV 1616 Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200) Groundnut- Birsa mungfali 3, 4, Girnar 3 Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149 <u>Vegetable crop</u> Radish- Pusa chetki (summer), Pusa deshi, Kashi hansi, Jaunpur/ Pusa himani, Japanese white, Pusa roshni Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika Cucurbits Bitter gourd- Arka hait, Pusa domausami, Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, local, Arka harit Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia, Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukh</p>	Normal Crop/cropping system	Rice, Pigeonpea, Maize, Pigeonpea + Maize , Pigeonpea + Sorghum	Suggested Contingency measures		a) Change in crop/cropping system	
Normal Crop/cropping system	Rice, Pigeonpea, Maize, Pigeonpea + Maize , Pigeonpea + Sorghum					
Suggested Contingency measures						
a) Change in crop/cropping system						



b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INP
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O
- DSR-Use plastic drum seeder rice tools
- Follow RDF and
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1 ml/l.t. water, Gundhi bug,leaf folder and BPH -Quinolophos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04%, Blast- Beam or Tricyclazole @ 0.6 gm /l.t water, Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/t. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/l.t. water, Mosaic- Methyl Demeton @ 1.5 ml/l.t. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/l.t. water

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKS, NHM and other State Govt. line departments are needed to be launched through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

A6. Major Farming Situation/Land Situation: Lowland High rainfall, medium depth, yellow to gray, heavy textured, loamy to clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2</u> in <u>Don 1</u>	

DSR(Improved Rice variety) Var.- Shabhagi, IR 64-Drt 1, Abhishek (120 days), MTU-1001, MTU 1010, Transplanting (Hybrid rice varieties) Var.- Arize 6444 (Gold), Uday 111, PHB 71, 26P52, 25P25, 27P31, 27P36

b) Agronomic Measures

- Staggered Nursery raising by MAT/ DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use Post emergence weedicide
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm



- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water, Gundhi bug, leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /ltr water

c) Remarks on Implementation

- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation.

A7. Major Farming Situation/Land Situation: Upland Less rainfall, shallow red light textured sandy & acidic soil

Normal Crop/cropping system	Upland Rice , Pigeonpea, Maize, Pigeonpea + Maize
Suggested Contingency measures	
a) Change in crop/cropping system	
<u>Discard rice Crop</u> <u>Sole crop</u> Pigeonpea, Finger millet, Gundli, Kodo, Sawan,Groundnut, Soybean, Maize, Blackgram, Greengram, Sorghum, Cowpea, lady;s Finger <u>Intercrop</u> Pigeonpea + Lady's finger (1:2), Pigeonpea + Sorghum (1:1), Pigeonpea + Maize (1:1) Pigeonpea + Groundnut(1:2), Maize + lady's finger (1:2), Maize + Cowpea (1:2) <u>Horticulture Crop</u> Vegetable : Tomato/ Brinjal/French bean/ Cucumber/ Radish <u>Variety</u> Pigeonpea- Birsa Arhar (200-220), Narendra Arhar 1, ICPH 2671 (200) Finger millet- BM 2, BM 3 (BBM 10), VL 149 Groundnut- Birsa mungfali 3, 4, Girnar 3 Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335 Birsa safed soybean 2 (105-110), RKS 18, RAUS 5 Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80) Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain) Greengram- HUM 16, IPM-02-03-60-65, SML 668 (summer) Sorghum- CSV 20-110-20, MP cheri, CSV 1616 Cowpea-rainy – Birsa sweta(80-90), Swarn sweta(80-90), Swarn harit (80-90) Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika <u>Vegetable crop</u> Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Hybrid- Swarn sampada, Swarn samridhi, Pusa hybrid 1 Suraksha Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6 Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata Cucumber-Japani long green, Pusa sanyog, Balamkhira, Puna khira, Swarn ageti Radish- Pusa chetki (summer), Pusa deshi, Kashi hansi, Jaunpur/ Pusa himani, Japanese white, Pusa roshni	
b) Agronomic measures	
<ul style="list-style-type: none"> Summer deep ploughing with Mouldboard or disc Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations. 	

- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbundled upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite application for pulses and oilseed @ 3-5 q/ha in furrow at the time of sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Dry seeding with 15% to 20% higher seed rate. Seed treatment with Rhizobium in pulses. Maximum use of organic manure

c) Remarks on Implementation

- Linkage with RKVY, ATMAS, and NFSM
- Vermicomposting through KVKS ATMAS and NHM
- Goatry and poultry rearing through KVKS, ATMAS and Veterinary Dept of. Govt. and BAU for livelihood support.
- Awareness about balanced use of fertilizers to increase their fertility, productivity and sustainability
- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology.
- Awareness for more and more use of organic manures, biopesticides for organic cultivation with IFS (eight components linkages)
- Upland- 15-20 % upland area should be covered with orchard

1. Mango based orchard-

Variety- Amrapali (30 June-5 July), Mallika (15-20 June regular bearer), Sunder langra(15-20 May)

Spacing- 5 m X 5m

i) Recommended package of Practices- Intercrops

- Mango + Papaya (Filler crop for two years) + Blackgram (rainy)/ Chickpea
- Mango + Custard apple (for 10 years and renovate or remove after 10 years) + Blackgram/Chickpea

Variety- Langra (15 June)/Bombay green(15 May)/ Himsagar (20-25 May irregular bearer),

Spacing- 10 m X 10m

ii) Recommended package of practices

- Mango + Guava(Up to 10 years as filler) + Papaya (Less than 3 years) + Blackgram-Chickpea/Lentil
- Mango + Lemon + Papaya + Rabi pulses/vegetables
- Mango + Custard apple + Papaya + Blackgram – Pea/Chickpea/Lentil/ Vegetables

2. Guava base orchard-

Variety- Arka Mridula, Pant Prabhat, Allahabad safeda, L 49

Spacing- 5m X 5m

Recommended package of practices- Intercrops

- Guava + Papaya (For 3 years) + Blackgram-Chickpea
- Guava + Custard apple + Blackgram/Soybean- Pea/Vegetables

3. Ber Based Orchard –

Variety- Banarsi, Karakka, Gola, Apple ber

Spacing- 5m X 5m

Recommended package of practices Intercrops

Ber + Custard apple + Sesame/Blackgram- Toria/Linseed/Safflower

N.B.- Cucurbits, beans or any creeper or climber vegetable should be avoided

- Field crops having height more than one meter should be avoided such as Pigenpea, Maize, Sorghum
- After 3-5 years when shading effects started shade loving crops like ginger, Turmeric, OI or leafy vegetables should be grown
- In citrus leaf minor and aphid susceptible crops should be avoided



- Aphid should be managed of mustard /toria taken in citrus orchard
- 4. Cashewnut based orchard for Kolhan region
- 5. Cassava should be grown for the requirement as feed for pig animals
- 6. Moringa should also be grown as fodder or vegetable purpose on upland main field bunds as shelter belt/wind break. Every year pruning and thinning should be followed for bushy look.

A8. Major Farming Situation/Land Situation: Midland Less rainfall, shallow red light textured sandy & acidic soil

Normal Crop/cropping system	Rice, Pigeonpea, Maize, Pigeonpea + Maize , Pigeonpea + Sorghum
Suggested Contingency measures	
a) Change in crop/cropping system	
<u>Don 2</u>	
DSR (Improved rice variety) Var -Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, MTU 1001Rice +Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR	
Transplanting (Hybrid Rice varieties) Var. –Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 27P31, DRRH 2	
<u>Don 3</u>	
Ridge and Furrow method or raise bed broad furrow:_Replace Rice with cereal/ Pulse/ Vegetable	
Cereal – Maize/ Sorghum/ Gundli/ Kodo/ Sawan,	
Pulse- Pigeonpea + Lady's finger (1:1) / Groundnut (1:2)/ Soybean (1:2)/ Finger millet (1:1)	
Vegetable-Radish/ Lady's Finger/ Cowpea	
<u>Variety</u>	
Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)	
Sorghum- CSV 20-110-20, MP cheri, CSV 1616	
Gundli- Birsa gundli 1	
Pigeonpea- Birsa Arhar (200-220), Asha (200-220), ICPH 2671 (200)	
Groundnut- Birsa mungfali 3, 4, Girnar 3	
Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain	
Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149	
<u>Vegetable crop</u>	
Radish- Pusa chetki (summer), Pusa deshi, Kashi hansi, Jaunpur/ Pusa himani, Japanese white, Pusa roshni	
Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika	
Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Creeper- Birsa sweta, Swarna sweta, Swarn harit	

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INP
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P_2O_5 + 1 kg K_2O for 100 m^2
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m^2 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P_2O_5 : K_2O (Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P_2O_5 + 40 K_2O /ha ((Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K_2O
- DSR-Use plastic drum seeder rice tools
- Follow RDF and
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/l. water, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /lt water, Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/l. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/l. water, Mosaic- Methyl Demeton @ 1.5 ml/l. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/ m^2 before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/l. water

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKs, NHM and other State Govt. line departments are needed to be launched through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

B. Monsoon/Weather Situation: 4 Weeks Delay (Onset: 2nd Week of July) – Early Season Drought

B1. Major Farming Situation/Land Situation: Upland High rainfall, shallow red sandy soil	
Normal Crop/cropping system	
Suggested Contingency measures	
a) Change in crop/cropping system	
<p><u>Discard Rice crop</u></p> <p><u>Sole crop</u></p> <p>Cowpea, Sorghum, Blackgram Finger millet, Soybean, Maize</p> <p><u>Intercrop</u></p> <p>Pigeonpea + Cowpea (1:2), Pigeonpea + Sorghum (1:1), Pigeonpea + Maize (1:1)</p> <p>Pigeonpea + Lady's finger (1:2), Maize + lady's finger (1:2), Maize + Cowpea (1:2)</p> <p><u>Horticulture Crop</u></p> <p>Vegetable- Tomato/ Brinjal/ Chili/ Amaranthus leaf/ Lobia/ French bean/ Cucumber</p> <p><u>Fodder Crop</u></p> <p>Cowpea/Sorghum/ Maize/ Blackgram</p> <p><u>Variety</u></p> <p>Cowpea-rainy – Birsa sweta(80-90), Swarn sweta(80-90), Swarn harit (80-90)</p> <p>Sorghum- CSV 20-110-20, MP cheri, CSV 1616</p> <p>Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain)</p> <p>Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149</p> <p>Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335</p> <p>Birsa safed soybean 2 (105-110), RKS 18, RAUS 5</p> <p>Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)</p> <p>Groundnut- Birsa mungfali 3, 4, Girnar 3</p> <p><u>Vegetable crops</u></p> <p>Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Hybrid- Swarn sampada, Swarn samridh, Pusa hybrid 1 Suraksha</p> <p>Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6</p> <p>Chili- Spices- Andhrajyoti, Pusasadabahar, NP 46, Jwala, KA 2, California wonder, Chinese giant, Yellow wonder, Bharat</p> <p>Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Creeper- Birsa sweta, Swarna sweta, Swarn harit</p> <p>Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata</p> <p>Cucumber-Japani long green, Pusa sanyog, Balamkhira, Puna khira, Swarn ageti</p> <p>Potato- early- Kufri ashoka (80-90), Kufri pukhraj, Mid early- Kufri kanchan , Kufri puskar, Kufri lalima (90-110), Late- Kufri sinduri (more than 110 days)</p> <p><u>Fodder crop</u></p> <p>Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4</p> <p>Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid</p> <p>Maize- African tall, JS-1006 and Vijaya composite.</p> <p>Blackgram- PU 19/31/35 (70-75), WBU 109 (70-75)</p>	
b) Agronomic Measures	
<ul style="list-style-type: none"> • Summer deep ploughing with Mouldboard or disc • Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations. 	

- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demoton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demoton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water, Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/lit. water or Cartap hydrochloride @ 2 gm/lit. water, hairy caterpiller -Quinolphos 1.5ml /lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- Linkage with RKVY , ATMAs and NFSM
- Vermicomposting Awareness through KVKS, ATMAs and NHM
- Backyard Goatry and poultry rearing Awareness campaign through KVKS, ATMAs and Veterinary Dept of. Govt. and BAU for livelihood support.
- A special programme is needed to be launched in such areas on priority basis to motivate the farmers to adopt improved technology for stress management through ATMAs, KVKS, Govt Dept., NGOs
- Campaign for Awareness of crop-weather insurance to meet the losses due to drought/cyclone like weather vagaries.

B2. Major Farming Situation/Land Situation: Midland High rainfall, shallow sandy soil

Normal Crop/cropping system	Rice, Finger millet, Pigeonpea, Sorghum, Maize, Pigeonpea + Sorghum, Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures



a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Rice + Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807

Don 3

Ridge and Furrow method or raise bed broad furrow: Replace rice with Cereal/Pulses/ vegetable/ Fodder crop
Cereal – Maize/Sorghum

Pulses- Pigeonpea+ Lady's Finger/Blackgram/Soybean/

Vegetables- Radish/ Coriander leaf/ Cucurbits/ Lady's Finger/ Cowpea/French bean/ Dolichos bean

Fodder Crop

Rice bean (Moth bean)/ Maize / Cowpea/ Maize/ Sorghum

Variety

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)

Sorghum- CSV 20-110-20, MP cheri, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Asha (200-220), ICPH 2671 (200)

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain

Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Vegetable crop

Radish- Pusa chetki (summer), Pusa deshi, Kashi hansi, Jaunpur/ Pusa himani, Japanese white, Pusa roshni

Coriander- Pant haritima, Rajendra swati

Cucurbits

Bitter gourd- Arka harit, Pusa domausami,

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, Local

Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white

Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,

Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Creeper- Birsa sweta, Swarna sweta, Swarn harit

French bean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata

Dolichos bean-Swarna utkrust, Swarna rituwar

Fodder crop

Maize- Aprican tall

Cowpea- UPC-5286, GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin@ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate



- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility.
- Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water, Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water.
- Rice pest and disease management -Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /lt water. Termite- Methyl parathion dust @ 25 kg/ha

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be awared through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

B3. Major Farming Situation/Land Situation: Lowland High rainfall, medium depth, heavy textured clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of Don 2 in Don 1 DSR(Improved Rice variety) var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days) Transplanting (Hybrid rice varieties) Var.-Arize 6444 (Gold), Uday 111, PHB 71, 26P52, 25P25	
b) Agronomic Measures	
<ul style="list-style-type: none">• Staggered Nursery raising by MAT/ DAPOG method• Follow community based nursery raising• Follow RDF,INPM• Use Post emergence weedicide• Use early to mid early duration of rice variety.• Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²• Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice• Topdressing above mentioned dose 10-15 days after sowing• In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling• Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm	



- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /ltr water

c) Remarks on Implementation

- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation

B4. Major Farming Situation/Land Situation: Upland High rainfall, shallow sandy soil

Normal Crop/cropping system	Upland Ric, Pigeonpea, Maize, Pigeonpea + Maize (Mixed)
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Rice crop	
<u>Sole crop</u>	
Pigeonpea, Sorghum, Finger millet, Soybean, Blackgram, Greengram, Lady's finger	
<u>Intercrop</u>	
Pigeonpea + Maize (1:1), Pigeonpea + Lady's finger (1:2)/Greengram (1:2), Maize+ creeper vegetables (Sponge Gourd) (1:2), Pigeonpea + Sorghum (1:1)	
<u>Horticulture Crop</u>	
Vegetable :Radish/Early Cabbage/ Cauliflower/ French bean / Cowpea	
<u>Variety</u>	
Pigeonpea- Birsa Arhar (200-220), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)	
Finger millet- BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149	
Soybean- JS 335	
Birsa safed soybean 2 (105-110), RKS 18, RAUS 5	
Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75)	
Greengram- HUM 16, IPM-02-03-60-65, SML 668	
<u>Vegetable crops</u>	
Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika	
Cucurbits	
Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white	
Radish- Pusa chetki, Pusa deshi, Japanese white, Pusa roshni	
Cabbage- early- Golden acer, Early drumfead, Pride of India, Shri ganesh cabbage 8	
Cauliflower- - Early kuwari, Early synthetic, Pusa him jyoti, Pant subhra	
Frenchbean- Bushy- Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata	
Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Creeper- Birsa sweta, Swarna sweta, Swarn harit	
b) Agronomic Measures	
<ul style="list-style-type: none"> Summer deep ploughing with Mouldboard or disc Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations. RD Spacing Zero tillage practices Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing RDF and in case of Intercropping reduce 1/3rd dose for intercrop Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathypor or Pendimithilin@ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables Bund construction for unbunded upland 	



- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/l.t. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/t. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/l.t. water, Mosaic- Methyl Demeton @ 1.5 ml/l.t. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/l.t. water, Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuran 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/l.t. water

c) Remarks on Implementation

- Linkage with RKVY, ATMAs and NFSM
- Vermicomposting Awareness through KVks, ATMAs and NHM
- Backyard Goatry and poultry rearing Awareness campaign through KVks, ATMAs and Veterinary Dept of Govt. and BAU for livelihood support.
- A special programme is needed to be launched in such areas on priority basis to motivate the farmers to adopt improved technology for stress management through ATMAs, KVks, Govt Dept., NGOs
- Campaign for Awareness of crop-weather insurance to meet the losses due to drought/cyclone like weather vagaries.

B5. Major Farming Situation/Land Situation: Midland shallow red light textured sandy & acidic soil

Normal Crop/cropping system	Rice, Pigeonpea, Maize, Pigeonpea + Maize , Pigeonpea + Sorghum
-----------------------------	---

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111,Rice + Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 27P31, DRRH 2
Don 3

Ridge and Furrow method or raise bed broad furrow along the slope: Replace rice with Cereal/Pulses/ vegetable/ Fodder crop

Cereal – Maize/Sorghum/ Gundli/Kodo/Sawan/

Pulses- Pigeonpea+ Lady's Finger/Blackgram/Soybean/ Groundnut/Maize

Vegetables- Lady's Finger/ Cowpea/ Dolichos bean/

Fodder Crop

Rice bean (Moth bean)/ Maize / Cowpea/ Sweet Sorghum

Variety

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)



Sorghum- CSV 20-110-20, MP chari, CSV 1616

Gundli- Birsa gundli 1

Pigeonpea- Birsa Arhar (200-220), Asha (200-220), ICPH 2671 (200)

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain

Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Groundnut- Birsa mungfali 3, 4, Girnar 3

Vegetable crops

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Creeper- Birsa sweta, Swarna sweta, Swarn harit

Dolichos bean-Swarna utkrist, Swarna rituwar

Fodder crop

Maize- African tall, JS-1006 and Vijaya composite.

Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin@ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility.
- Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/ltr. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/ltr. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/ltr. water, Mosaic- Methyl Demeton @ 1.5 ml/ltr. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/ltr. water, Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/ltr. water or Cartap hydrochloride @ 2 gm/ltr. water, hairy caterpillar –Quinolophos 1.5ml /ltr. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 ltr. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/ltr. water.



- Rice pest and disease management -Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /lt water. Termite- Methyl parathion dust @ 25 kg/ha

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be awared through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

B6. Major Farming Situation/Land Situation: Lowland High rainfall, medium depth, heavy textured clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2</u> in <u>Don 1</u> DSR(Improved Rice variety) var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days)	
Transplanting (Hybrid rice varieties) Var.-Arize 6444 (Gold), Uday 111, PHB 71, 26P52, 25P25	
b) Agronomic Measures	
<ul style="list-style-type: none"> Staggered Nursery raising by MAT/ DAPOG method Follow community based nursery raising Follow RDF,INPM Use Post emergence weedicide Use early to mid early duration of rice variety. Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice Topdressing above mentioned dose 10-15 days after sowing In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering. DSR-Use plastic drum seeder rice tools Use of post weedicide Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lt, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /lt water 	

c) Remarks on Implementation

- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation

B7. Major Farming Situation/Land Situation: Upland Less rainfall, shallow red light textured sandy & acidic soil

Normal Crop/cropping system	Upland Ric, Pigeonpea, Maize, Pigeonpea + Maize (Mixed)
Suggested Contingency measures	



a) Change in crop/cropping system

Discard rice Crop

Sole crop

Finger millet, Sorghum, Blackgram, Soybean, Maize, Gundli, Kodo, Sawan,

Intercrop :

Pigeonpea + Cowpea (1:2), Pigeonpea + Sorghum (1:1), Pigeonpea + Maize (1:1), Pigeonpea + Lady's finger (1:2)

Maize + lady's Finger (1:2), Maize + Cowpea (!:2)

Horticulture Crop

Vegetable- Radish/Early Cabbage/ Cauliflower/ French bean / Cowpea (Lobia)

Fodder Crop

Cowpea/Sorghum/ Maize/ Blackgram

Variety

Finger millet- BM 2, BM 3 (BBM 10), VL 149

Sorghum- CSV 20-110-20, MP cheri, CSV 1616

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain)

Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335, Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)

Gundli- Birsa gundli 1

Pigeonpea- Birsa Arhar (200-220), Narendra Arhar 1, ICPH 2671 (200)

Cowpea-rainy – Birsa sweta(80-90), Swarn sweta(80-90), Swarn harit (80-90)

Vegetable crops

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Radish- Pusa chetki (summer), Pusa deshi, Kashi hansh, Jaunpur/ Pusa himani, Japanese white, Pusa roshni

Cabbage- early- Golden acer, Early drumfead, Pride of India Late- Late drunhead, sabyay cabbage, 7 Ganga, Jamuna, Kaveri, Shri ganesh cabbage 8

Cauliflower-Summer- Early kuwari, early- Kuwari, Pusa katki, Pusadipali, Early synthetic, Mid early- Pusa ketaki, Pusadipali, Pusa him jyoti, Pant subhra, Late- Maghi, Snowbowl 16, dania, Pusa Snowbowl, K Pusa Snowbowl, Hybrid- Himani, Swati, Endum early Pusa hybrid 1

Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Creeper- Birsa sweta, Swarna sweta, Swarn harit

Fodder crop

Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4.

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid.

Maize- African tall, JS-1006 and Vijaya composite

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain)

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.



- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/t. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- Linkage with RKVY , ATMAs and NFSM
- Vermicomposting Awareness through KVks, ATMAs and NHM
- Backyard Goatry and poultry rearing Awareness campaign through KVks, ATMAs and Veterinary Dept of. Govt. and BAU for livelihood support.
- A special programme is needed to be launched in such areas on priority basis to motivate the farmers to adopt improved technology for stress management through ATMAs, KVks, Govt Dept., NGOs
- Campaign for Awareness of crop-weather insurance to meet the losses due to drought/cyclone like weather vagaries.

B8. Major Farming Situation/Land Situation: Midland Less rainfall, shallow depth , light textured sandy and acidic soil

Normal Crop/cropping system	Rice, Pigeonpea, Maize Pigeonpea + Maize, Pigeonpea + Sorghum, Rice-vegetable
-----------------------------	---

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 27P31, DRRH 2
Don 3

Ridge and Furrow method or raise bed broad furrow along the slope :_Replace rice with Cereal/Pulses/ vegetable/ Fodder crop

Cereal – Maize/Sorghum

Pulses- Pigeonpea+ Lady's Finger/Blackgram/Soybean/ Groundnut/Maize

Vegetables- Lady's Finger/ Cowpea/ Dolichos bean/

Fodder Crop -

Rice bean (Moth bean)/ Maize / Cowpea/ Sweet Sorghum

Variety

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 – Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Blackgram- Birsa urd 1 (75-80), PU 31 (70-75), WBU 109 (70-75)
Soybean- Birsa soybean 1 black(120-125), JS 335
Birsa safed soybean 2 (105-110), RKS 18, RAUS 5
Groundnut- Birsa mungfali 3, 4, Girnar 3
Maize- Birsa makka (Vikash) 2 (75-80), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)
Vegetable crops
Lady's finger- Pusa A 4, Sarika
Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati, Swarn harit
Dolichos bean-Swarna utkrist, Swarna rituwar
Fodder crop
Maize- African tall, JS-1006 and Vijaya composite.
Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4
Sorghum-PC-1, PC-6, PC-23, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin@ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility.
- Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like: Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water, Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/lit. water or Cartap hydrochloride @ 2 gm/lit. water, hairy caterpillar –Quinolphos 1.5ml /lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuran 3G @ 3 gm/m² before 10 days of transplanting followed by application of Tricoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water.



- Rice pest and disease management -Gundhi bug, leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /lt water. Termite- Methyl parathion dust @ 25 kg/ha

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be awared through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

B9. Major Farming Situation/Land Situation: Lowland Less rainfall, medium gray, heavy textured loamy & clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2</u> in <u>Don 1</u> DSR(Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), MTU-1001, MTU 1010, Transplanting (Hybrid rice varieties) Var.–Arize 6444 (Gold), Arize Tej (Gold), Akshay Dhan, PHB 71, 26P52, 25P25	
Vegetable	
Cucurbits-	
Bitter gourd- Arka hait, Pusa domausami,	
Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, Local, Arka harit	
Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white	
Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,	
Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi	

b) Agronomic Measures

- Staggered Nursery raising by MAT/ DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use Post emergence weedicide
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lt. water, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /lt water

c) Remarks on Implementation

Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme

Supply of Plastic drum seeder through line departments

Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.

Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.

Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates

Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation



C. Monsoon/Weather Situation: 6 Weeks Delay (Onset: 6th Week of July) – Early Season Drought

C1. Major Farming Situation/Land Situation: Upland High rainfall, shallow red sandy soils

Normal Crop/cropping system	Upland Rice, Finger millet, Pigeonpea + Sorghum, Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Discard Rice Crop

Sole crop

Pigeonpea (early maturity) Soybean, Gundli, Kodo, Sawan, Finger millet, Blackgram, Greengram, Sorghum,

Intercrop

Pigeonpea + Lady's Finger (1:2), Lady's finger + Blackgram (1:2), Maize + Blackgram (1:2), Maize + Cowpea (1:2), Maize + lady's finger (1:2), Pigeonpea + Maize (1:1), Pigeonpea + Sorghum (1:1)

Horticulture crop

Vegetable –Lady's Finger, Ridge Gourd, Bottle Gourd, Cowpea

Fodder Crop

Chara badam/ Anjan grass/ Hybrid napier/ Cactus/ Sweet Sorghum

Variety

Pigeonpea- Birsa Arhar (200-220), Asha (200-220), ICPH 2671 (200)

Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149

Greengram- HUM 16, IPM-02-03-60-65, SML 668 (summer)

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain)

Sorghum- CSV 20-110-20, MP cheri, CSV 1616

Vegetable crops

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Cucurbits

Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, local, Arka harit

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Creeper- Birsa sweta, Swarna sweta, Swarn harit

Fodder crop

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid.

b) Agronomic Measures

- Top dressing of urea and DAP after receipt of the rain for all crops
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10-15 kg/ha
- Replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Soybean, Sesame, Horsegram , Niger, Cowpea, Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour
- Follow mulch after cultural operations to control the weeds in vegetables.
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges
- Foliar application of 2 % DAP or 0.5 to 1 % potassium chloride (KCl) +0.3 % Boric acid or 2% urea at pre-flowering and flowering stage in pulses and vegetables
- 2 % DAP spray for pulses.
- Use antitranspirants :
- Stomatal closure (Growth hormones like ABA, Ethrel, TIBA, succinic acid, ascorbic acid and Cycocel (CCC)
- Reflectant (Calcium bicarbonate, Lime water) Thin film (Hexadecanol (Higher alcohols) Cetyl alcohol, Methanol
- Acidic soils should be reclaimed by application of soil ameliorants.
- Follow integrated pest management.
- Weed control by applying pre-emergence 5-6 DAS (Pendimethalin) or Post-emergence 18-28 DAS (Bispyribac)
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/ltr. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/ltr. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/ltr. water, Mosaic- Methyl Demeton @ 1.5 ml/ltr water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/ltr. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 ltr. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/ltr. water

c) Remarks on Implementation

- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology for stress management through ATMAs, KVks, Govt Dept., NGOs and others. Soybean and fodder crops may be promoted.
- Promote Knowingness about climate resilient agriculture at district, block, panchayat and village level through involvement of KV's, ATMAs, DAO, NGO's and other State Agril. Govt line departments.
- Awareness of mechanization and Supply of Mouldboard and disc chisel/harrow through govt. scheme on subsidized way.
- Promote for double their income by curtailing cost of cultivation by introduction of early duration crops variety.
- Campaign for Awareness programme about crop-weather insurance

C2. Major Farming Situation/Land Situation: Midland High rainfall, slightly deep yellow loamy to sandy loam soil

Normal Crop/cropping system	Rice, Finger millet, Pigeonpea, Sorghum, Maize, Pigeonpea + Sorghum, Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 110, 111, MTU 1001, Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 25P25, 27P36

Don 3

Ridge and Furrow method or raise bed broad furrow :Replace Rice with pulses and cereals/ Vegetable/ Fodder crop Pulses and cereal-Pigeonpea/ Soybean/ Maize/ Sorghum

Vegetable- Lady's finger/Tomato,/ Brinjal, Cucurbits/Chili

Fodder Crop

Blackgram/ Greengram/ Cowpea/ Sorghum/ Maize

Variety

Pigeonpea- Birsa Arhar (200-220), Asha (200-220), ICPH 2671 (200)

Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Vegetable crop

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Hybrid- Swarn sampada, Swarn samridhi, Pusa hybrid 1 Suraksha

Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6

Cucurbits

Bitter gourd- Arka harit, Pusa domausami,

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, local

Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white

Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,

Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi

Chili- Spices- AndhraJyoti, Pusasadabahar, NP 46, Jwala, KA 2, California wonder, Chinese giant, Yellow wonder, Bharat

Fodder crop

Blackgram- WBU 109 (70-75), Uttara (75-80)

Greengram- HUM 16, IPM-02-03-60-65

Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

Maize- African tall, JS-1006 and Vijaya composite.

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²



- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water; Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /lit water; Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- Campaign for awareness improved technology through RKVY, ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be at different district, block, panchayat and village level
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency crops through Lamps within one months.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

C3. Major Farming Situation/Land Situation: Lowland High rainfall, medium gray, heavy clay loam soil.

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of Don 2 in Don 1 DSR(Improved Rice variety) Var- Shabhagi, IR 64-Drt 1, Abhishek (120 days), MTU-1001, MTU 1010, Transplanting (Hybrid rice varieties) Var.-PAC 801,25P25, 27P31 Arize 6444 (Gold), PHB 71, 26P52, 27P36 Fodder crop - In case of fallow (Late heavy rainfall) Dallis grass/ Para grass/ Arundino Grass	
b) Agronomic Measures	
<ul style="list-style-type: none"> • Staggered Nursery raising by MAT/ DAPOG method • Follow community based nursery raising • Follow RDF,INPM • Use Post emergence weedicide • Use early to mid early duration of rice variety. • Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² • Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice • Topdressing 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² at 10-15 days after sowing • In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling • Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm • Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering. • DSR-Use plastic drum seeder rice tools • Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water; Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /lit water • In DSR- Plastic drum seeder with increased seed rate by 20-25 Per cent 	



- Maximum use of organic manure
- Post emergence weedicide application in soil
- Follow RDF, INM and IPM
- Proper spacing for the use of conoweeder
- Use Azolla or BGA

c) Remarks on Implementation

- Awareness programme of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments in case of DSR
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency mid early rice varieties through Lamps within one month Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation
- Contingency technology Awareness programme through KVKs, ATMAs, NGOs and DAOs
- Achieve maximum fallow area in case of late drought and suggest to go for cultivation of early duration rabi and fodder crops.

C4. Major Farming Situation/Land Situation: Midland High rainfall, shallow sandy soil

Normal Crop/cropping system	Rice, Pigeonpea, Maize, Pigeonpea + Maize, Pigeonpea + Sorghum
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 110, 111, MTU 1001, Transplanting (Hybrid Rice Varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 25P25, 27P36

Don 3

Ridge and Furrow method or raise bed broad furrow: Replace Rice with pulses and cereals/ Vegetable/ Fodder crop Pulses and cereal-Pigeonpea/ Maize/ Sorghum

Vegetable- Lady's finger/Tomato,/ Brinjal, Cucurbits/Chili

Fodder Crop

Blackgram/ Greengram/ Cowpea/ Sorghum/ Maize

Vegetable crop

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Hybrid- Swarn sampada, Swarn samridhi, Pusa hybrid 1 Suraksha Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6

Cucurbits-

Bitter gourd- Arka harit, Pusa domausami,

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, local

Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green, Long white

Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,

Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi

Chili- Spices- Andhrayoti, Pusasadabahar, NP 46, Jwala, KA 2, California wonder, Chinese giant, Yellow wonder, Bharat

Fodder crop

Blackgram- WBU 109 (70-75), Uttara (75-80)

Greengram- HUM 16, IPM-02-03-60-65

Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

Maize- African tall, JS-1006 and Vijaya composite.

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm

- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water; Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /lit water; Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- Campaign for awareness improved technology through RKVY, ATMAS, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be at different district, block, panchayat and village level
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency crops through Lamps within one months.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

C5. Major Farming Situation/Land Situation: Lowland High rainfall, medium gray, heavy textured loamy & clay loam soils.

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna, BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2 in Don 1</u> DSR (Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), MTU-1001, MTU 1010, Transplanting (Hybrid rice varieties) Var.-PAC 801,25P25, 27P31 Arize 6444 (Gold), PHB 71, 26P52, 27P36, KRH 2, DRRH 2 Fodder crop- In case of fallow (Late heavy rainfall) Dallis grass/ Para grass/ Arundino Grass	
b) Agronomic Measures	
<ul style="list-style-type: none"> Staggered Nursery raising by MAT/ DAPOG method Follow community based nursery raising Follow RDF,INPM Use Post emergence weedicide Use early to mid early duration of rice variety. Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice Topdressing 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² at 10-15 days after sowing In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm Fertilizer dose- 80:40:20 kg/ha N : P₂O₅: K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering. DSR-Use plastic drum seeder rice tools Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit; Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /lit water In DSR- Plastic drum seeder with increased seed rate by 20-25 Per cent Maximum use of organic manure Post emergence weedicide application in soil Follow RDF, INM and IPM Proper spacing for the use of conoweede Use Azolla or BGA 	

c) Remarks on Implementation

- Awareness programme of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments in case of DSR
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency mid early rice varieties through Lamps within one month Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation
- Contingency technology Awareness programme through KVK's, ATMAS, NGO's and DAO's
- Achieve maximum fallow area in case of late drought and suggest to go for cultivation of early duration rabi and fodder crops.

C6. Major Farming Situation/Land Situation: Upland Less rainfall, shallow depth , light textured sandy and acidic soil.

Normal Crop/cropping system	Upland Rice, Pigeonpea + Sorghum, Pigeonpea, Maize
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Rice Crop	
<u>Sole crop</u>	
Raised bed or ridge and furrow method: Niger, Horsegram, Sorghum, Blackgram, Soybean, Finger millet, Pigeonpea, Lady's finger, Maize	
<u>Intercrop</u>	
Pigeonpea + Horsegram (1:2), Pigeonpea + Maize (1:1)	
<u>Horticulture crop</u>	
Vegetable –Lady's Finger, Ridge Gourd, Bottle Gourd, Cowpea	
<u>Fodder Crop</u>	
Deenanath grass/ Guinea grass/ Sadabahar/ Stylo hemata/ Chara badam/ Anjan grass/ Hybrid napier/ Cactus/ Sweet Sorghum	
<u>Variety</u>	
Niger- Birsa niger 1, 2 and 3 (95-105)	
Horsegram- Birsa kulthi1 (90-95)	
Sorghum- CSV 20-110-20, MP chari,	
Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75)	
Soybean- R 518 (110), JS 9752 (100), RKS 18, RAUS 5	
Finger millet- A 404, BM 2, BM 3 (BBM 10) Pigeonpea- Birsa Arhar (200-220), Asha (200-220)	
Lady's finger- Varsa uphar, Hybrid- Sonal, Sarika	
Maize- Birsa makka (Vikash) 2 (75-80), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85)	
<u>Fodder crop</u>	
Sorghum-HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid	

b) Agronomic Measures

- Top dressing of urea and DAP after receipt of the rain for all crops
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10-15 kg/ha
- Replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Soybean, Sesame, Horsegram , Niger, Cowpea, Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour
- Follow mulch after cultural operations to control the weeds in vegetables.
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges
- Foliar application of 2 % DAP or 0.5 to 1 % potassium chloride (KCl) +0.3 % Boric acid or 2% urea at pre-flowering and flowering stage in pulses and vegetables
- 2 % DAP spray for pulses.
- Use antitranspirants : Stomatal closure (Growth hormones like ABA, Ethrel, TIBA, succinic acid, ascorbic acid and Cycocel (CCC), Reflectant (Calcium bicarbonate, Lime water) Thin film (Hexadecanol (Higher alcohols) Cetyl alcohol, Methanol
- Acidic soils should be reclaimed by application of soil ameliorants.
- Follow integrated pest management.
- Weed control by applying pre-emergence 5-6 DAS (Pendimethalin) or Post-emergence 18-28 DAS (Bispyribac)



- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/ltr. water; Pigeonpea-leaf folder-Methyl demeton @ 1.5 ml/ltr. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/ltr. water, Mosaic- Methyl Demeton @ 1.5 ml/ltr. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/ltr. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 ltr. water; vegetables-Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/ltr. water

c) Remarks on Implementation

- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology for stress management through ATMAS, KVKs, Govt Dept., NGOs and others. Soybean and fodder crops may be promoted.
- Promote Knowingness about climate resilient agriculture at district, block, panchayat and village level through involvement of KVK's, ATMAS, DAO, NGO's and other State Agril. Govt line departments.
- Awareness of mechanization and Supply of Mouldboard and disc chisel/harrow through govt. scheme on subsidized way.
- Promote for double their income by curtailing cost of cultivation by introduction of early duration crops variety.
- Campaign for Awareness programme about crop-weather insurance

C7. Major Farming Situation/Land Situation: Midland Less rainfall, shallow red light textured sandy & acidic soil

Normal Crop/cropping system	Rice, Pigeonpea, Maize, Pigeonpea + Maize, Pigeonpea + Sorghum
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 110, 111, Vardhan
Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 27P31, 25P31, CR Dhan 40, Akshya Dhan

Don 3

Ridge and Furrow method or raise bed broad furrow along the slope:_Replace Rice with pulse/ vegetable/ Fodder crop Pulse & Millet- Pigeonpea + Blackgram (1:2)/Sesame (1:2)/Maize (1:1)/ Lady's finger (1:1)/ Niger (1:2)/ Horsegram (1:2)/ Gundli (1:2)/Kodo/Sawan/

Vegetable

Capsicum/ French bean/ Coriander leaf/ Amaranthus leaf/Lady's finger/Tomato,/ Brinjal/Chili/
Fodder Crop

Cowpea/ Sorghum/ Maize/ Rice bean (Moth bean)/ Sudan grass(SC)/Thin Napier, Late August-September-Berseem (MC)/ Oat (MC)/ Rye grass

Variety

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)

Sesame- RT 346 (90), Kanke safed (95-100), Krishna (95-100)

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 –Yuvral gold (80-85), VMH 4106 (Sweet corn hybrid), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Niger- Birsa niger 1, 2 and 3 (95-105), Puja 1 (90), VLG 19

Horsegram- Birsa kulthi1 (90-95)

Gundli- Birsa gundli 1

Vegetable crops

Capsicum-California wonder, Yellow wonder, Arka mohini, Arka gaurav, Hybrid- Bharat, Indra

Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata

Coriander- Pant haritima, Rajendra swati

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Tomato- Swarn lalima, BT 12, Swarn vaibhaw, Samrat, Hybrid- Swarn sampada, Swarn samridhi, Pusa hybrid 1 Suraksha

Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant, Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6

Chili- Spices- Andhra jyoti, Pusasadabahar, NP 46, Jwala, KA 2, California wonder, Chinese giant, Yellow wonder, Bharat

**Fodder crop-**

Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4.

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid Maize- African tall, JS-1006 and Vijaya composite.

Berseem (MC)-Vardan

Oat (MC)

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water; Gundhi bug,leaf folder and BPH -Quinolophos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /lit water; Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/t. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit., Mosaic- Methyl Demeton @ 1.5 ml/lit. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- Campaign for awareness improved technology through RKVY, ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be at different district, block, panchayat and village level
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency crops through Lamps within one months.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

C8. Major Farming Situation/Land Situation: Lowland Less rainfall, medium gray, heavy textured loamy & clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2 in Don 1</u>	
DSR (Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), MTU 1010, BVD 203, BVS 1	
Transplanting (Hybrid rice varieties) Var.-PAC 801,25P25, 27P31 Arize 6444 (Gold), PHB 71, 26P52, 27P36, KRH 2, DRRH 2	
Fodder crop -In case of fallow (Late heavy rainfall) Para grass	



b) Agronomic Measures

- Staggered Nursery raising by MAT/ DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use Post emergence weedicide
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² at 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water; Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /ltr water
- In DSR- Plastic drum seeder with increased seed rate by 20-25 Per cent
- Maximum use of organic manure
- Post emergence weedicide application in soil
- Follow RDF, INM and IPM
- Proper spacing for the use of conoweede
- Use Azolla or BGA

c) Remarks on Implementation

- Awareness programme of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments in case of DSR
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency mid early rice varieties through Lamps within one month Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation
- Contingency technology Awareness programme through KVKS, ATMAS, NGO's and DAO's
- Achieve maximum fallow area in case of late drought and suggest to go for cultivation of early duration rabi and fodder crops.

PART-II

A. Monsoon/Weather Situation: Normal onset followed by 15-20 days dry spell after sowing (Early Season Drought-Normal onset)

A1. Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils	
Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
Suggested Contingency measures	
a) Change management Cultivate drought tolerant promising non paddy crops like pigeonpea, blackgram, greengram, rice bean, finger millet, guar, sesame, soyabean, sorghum, pearl millet, sweet potato, castor and vegetables like radish, tomato,, brinjal, creeper bean, chili, lady's finger wherever possible in place of upland rice	
<ul style="list-style-type: none"> • Maximum use of organic manures for early seedling vigour along with RDF (N:P₂O₅:K₂O) • Recommend to resow with subsequent rains for better plant stand. • When damage is Less than 30 per cent then go for Gap filling in all upland crops • When damage is More than 50 per cent then go resowing in all upland crops • Removing excess plants where are over crowded,to reduce crop stand to conserve soil moisture • Water spraying during evening and early morning 	
b) Soil nutrient & moisture conservation measures <ul style="list-style-type: none"> • Avoid top dressing of Urea during dry spell and wait till downpour • Go for in-situ moisture conservation • One hand weeding followed by hoeing and simultaneous eartingup after 20 DAS is highly recommended in all upland crops. 	
c) Remarks on Implementation Awareness for Construction of rain water harvesting structures for recycling of water during dry spell like DOVAS through SHG or on subsidized basis through State Govt.schemes.	
A2. Major Farming Situation/Land Situation: MID LAND Sandy loam soils	
Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change management <u>Don 2</u> <ul style="list-style-type: none"> • If possible, go for staggered raising of nursery in rice crop • If possible, raise community nursery of rice at a reliable water source to save time for further delay. • In case, if rice population is less than 40-50 percent, gap filled by retransplanting the rice crop and for more than 50 per cent mortality use fresh seeding for fresh transplanting. • Follow gap filling by removing seedlings from profuse tillers to have a uniform distribution of same aged plants • For termite and disease management in nursery spray Indofil M 45 and Chlorpyriphos @ 0.2 per cent • life saving irrigation • DSR on receipt of rain by using Paddy drum seeder or • High yielding varieties- follow transplanting while, Improved varieties – follow DSR • In case of DSR- Use sprouted seeds in plastic drum seeder with increased seed rate by 20-25 per cent for good crop stand • Late transplanted rice during early season drought results in the occurrence of sheath rot and grain discoloration diseases. • Follow pre emergence and post emergence weedicide to disturb/check the crop-weed competition for nutrient • Provide life saving and protective irrigation to over aged seedling in nursery through dovas (harvested rain water). Also, take care of blast disease in nursery and avoid using urea in nursery. • Strengthen the bunds to check the drainage holes and seepage loss in transplanted and direct sown medium land rice regularly <u>Don3</u> <ul style="list-style-type: none"> • Follow raised bed broad furrow or Ridge and furrow method for Maize/ Pigeonpea/ Lady's finger/ Blackgram/ Soybean • Adopt surface mulching with crop residue or tree loppings of <i>Glyricidia</i> wherever possible.If farm waste is not available, use blade to form a thin layer of soil mulch to avoid cracks • Life saving irrigation • In case of transplanting of over aged seedling (35-45 days), increase number of seedling per hill (5-6 seedling/hill) 	

b) Soil nutrient & moisture conservation measures

- Dry seeding of rice with application of pre and post emergence weedicide in over aged seedlings (>25 DOS)
- Split application of Urea fertilizer
- Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells

c) Remarks on Implementation

Awareness for Construction of rain water harvesting structures for recycling of water during dry spell like DOVAS through SHG or on subsidized basis through State Govt.schemes.

A3. Major Farming Situation/Land Situation: LOW LAND Sandy clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
<p>a) Change management</p> <ul style="list-style-type: none"> • If possible, go for staggered nursery raising in rice crop • If possible, raise community nursery of rice at a reliable water source to save time for further delay. • In case, if rice population is less than 40-50 percent, gap filled by retransplanting the rice crop and for more than 50 per cent mortality use fresh seeding for fresh transplanting. • Follow gap filling by removing seedlings from profuse tillers to have a uniform distribution of same aged plants • Prefer mid early rice variety instead of late variety • Use pre and post emergence weedicide • Over aged seedling should be top cut and treat the seedlings root by Dursban/Chlorpyriphos @ 5 ml per lt water and transplant immediately after treated seedlings with 2 per cent Urea solution • In case of transplanting over aged seedling (35-45 days), increase number of seedling per hill (5-6 seedling/hill) • In fallow land go for cultivation of mid early duration rice variety through DSR @ 70-80 Kg/ha 	

b) Soil nutrient & moisture conservation measures

- Split application of Urea fertilizer
- Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells

c) Remarks on Implementation

Awareness for Construction of Ponds, check dam through water shed management & MNREGA scheme through SHG or on subsidized basis through State Govt.schemes.

B. Monsoon/Weather Situation: Mid season drought (long dry spell, consecutive 2 weeks rainless (<2.5 mm) period

B1. At vegetative phase

B1.1. Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils

Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
Suggested Contingency measures	

a) Change management

- Use organic mulches such as tree leaves, straw and other available crop residue to conserve soil moisture
- Avoid top dressing of fertilizers till sufficient moisture is available in soil
- Use reflectant or antitranspirant like Kaolin @ 3-5 kg/100 lt or
- In pulses, at weekly interval foliar spray of KCl @ 0.5- 1 % + 100 ppm Boric acid followed by foliar spraying of 2 percent urea during evening time
- Spray wax emulser
- Manual weeding followed by hoeing for germinating weeds.
- For termite and leaf folder control spraying or drenching of Chlorpyriphos @ 2ml/lt water and for all pulses and cereals.
- For leaf folder control in Maize (Stem borer) and Pigeonpea apply Carbofuran 3 G @ 12 Kg/acre or Phorate 10 G @ 4 kg/acre or Quinolphos @ 1 ml/lt water in Maize for leaf folder
- Also, spray @ 20/40/60 ppm CaCl₂ in pulses
- Vegetables- Foliar spray of water with 2 per cent KCl + 100 ppm Boron
- Tomato- Foliar spray of CaCl₂ @ 20/40/60 ppm
- Gap filling may be done with pigeonpea to maintain adequate plant stand.
- For termites in pigeonpea, maize and other standing cereal crops which can be controlled by soil drenching with chlorpyriphos 20 EC @ 2 ml/lt water or by adding Chlorpyriphos 1.5% dust @ 8– 10 kg/ha or Carbofuran 3G @ 12 kg or Phorate 10 G @ 4 kg.acre before final land preparation and also control Gallmidge
- In green and blackgram, cowpea, bean and lady's finger the spread of YMV by insect vector may increase. Hence, to control insect vectors spray Dimethoate @ 1ml/ lt or Imidacloprid 4 ml/10 lt twice at 10 days interval



- In groundnut crop termites and white grub incidence is expected to be more. Methods suggested in rice may be followed to reduce the pest infestation.
- Incidence of leaf miner in groundnut may increase which can be managed by spraying Monocrotophos 36 SL or Triazophos 40 EC @ 1 ml/lit. water twice at fortnight intervals.
- Under dry condition incidence of mites is expected to be more in vegetable crops which can be brought down by spraying of dicofol @ 2 ml/lit water.
- Early and mid season drought favours disease like brown spot of rice, bacterial wilt of brinjal and other vegetables

b) Soil nutrient & moisture conservation measures

- Foliar spraying of DAP @ 2 per cent along with Boric acid @ 0.3 per cent. Also, spray Urea @ 1 per cent
- Provide micro- irrigation with drip for wide spaced crops such as chillies and vegetables and Sprinklers for groundnut, maize and vegetables wherever ground/ surface water is available.
- Go for life saving and protective irrigation from constructed dovas.

c) Remarks on Implementation

Promote construction of Rain water harvesting structure watershed programme and MNREGA

B2. At flowering/ fruiting stage

B2.1 Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils

Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
-----------------------------	--

Suggested Contingency measures

a) Change management

- Maize- Harvest it for fodder use
- Pulses- and vegetables- At 2-3 days interval spraying of water followed by 2 per cent KCl + 100 ppm Boron during evening time is recommended.
- In case of groundnut maturing in the month of September which can be harvested after providing light irrigation through dohas to loose the soil.

b) Soil nutrient & moisture conservation measures

Go for life saving and protective irrigation from constructed DOVAS.

c) Remarks on Implementation

Promote for the construction of Rain water harvesting structure watershed programme and MNREGA

B3. At vegetative phase

B3.1. Major Farming Situation/Land Situation: MID LAND Sandy loam soils

Normal Crop/cropping system	Rice
-----------------------------	------

Suggested Contingency measures

a) Crop management

Don 2

- Manual weeding followed by hoeing for germinating weeds
- Take care of mealy bug and termite attack which are more prevalent in dry weather .
- Top dressing should be followed only after receipt of rain .
- No urea should be top dressed until receipt of rainfall in rice crop.
- For BPH, dusting field bunds and around with Carbaryl (Savin)4% or malathion 5% @ 10 – 12 kg/acre

Don 3

- One manual weeding for germinating weeds
- Apply 4 Kg N/acre in sorghum and oilseed crops soon after receipt of rains.
- In pigeonpea, if the drought affected plants to recoup with the revival of the rains, spray 2 to 3% urea after the foliage is wetted with the rains.
- Foliar application of Sulphur @ 1ppm to mitigate the stress condition in oilseed is necessary after receipt of rainfall
- Apply post emergence weedicide for controlling weeds in oilseed (Groundnut) to undisturb the pegging process.
- During 40-45 DAS, if there is a severe moisture stress, thinning may be done in kharif sorghum and pearl millet.

b) Soil nutrient & moisture conservation measures

- Foliar spray of KCl or ZNSO₄ @ 2 per cent
- Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells
- Life saving irrigation through dovas,well, ponds, check dams and bora bandh

c) Remarks on Implementation

Promote for the construction of Rain water harvesting structure watershed programme and MNREGA



B4. At flowering/ fruiting stage

B4.1. Major Farming Situation/Land Situation: MID LAND Sandy loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<p><u>Don 2 and Don 3</u></p> <ul style="list-style-type: none"> • Life saving irrigation with harvested water • Spray of urea @ 1-2 percent • Drought condition during the month of August-September onwards shall result in severe incidence of foliar blast and brown spot diseases in rice. It is advised to spray Tricyclazole (Tilt) @ 6 g/ 10 lt or Casugamycin @ or Kasu B @ 2 ml/lt. water twice at 10 days intervals during drought period. 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Foliar spray of KCl or ZNSO₄ @ 2 per cent • Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells • Life saving irrigation through dovas, well, ponds, check dams and bora bandh 	
c) Remarks on Implementation	
Promote for the construction of Rain water harvesting structure watershed programme and MNREGA	

B5. At vegetative phase

B5.1. Major Farming Situation/Land Situation: LOW LAND Sandy clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<ul style="list-style-type: none"> • Foliar spray of 2 per cent KCL followed by 1-2 per cent Urea. • Weeding should be done • Drought makes the crop vulnerable to sheath rot and sheath blight diseases. Maintenance of field sanitation followed by twice spraying at 10 days interval with validamycin 2-3 ml/lt water or Tricyclazole @ 6g/10 lt or carbendazim @ 2 g/lt water are advised. • Life saving irrigation 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Foliar spray of Foliar spray of Urea @ 2 per cent • Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells • Life saving irrigation through dovas, well, ponds, check dams and bora bandh 	
c) Remarks on Implementation	
Awareness for Construction of Ponds, check dam through water shed management & MNREGA scheme through SHG or on subsidized basis through State Govt.schemes.	

B6. At flowering/ fruiting stage

B6.1. Major Farming Situation/Land Situation: LOW LAND Sandy clayloam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<ul style="list-style-type: none"> • Drought condition during flowering and fruiting and onwards shall result in severe incidence of foliar blast and brown spot diseases in rice. It is advised to spray Tricyclazole (Tilt) @ 6 g/ 10 lt or Casugamycin @ or Kasu B @ 2 ml/lt. water twice at 10 days intervals during drought period. • Life saving irrigation • During drought, attack of gundhi bug shall be more more. Apply Quinolphos or Monocrotophos @ 1-2 ml per lt. water. 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Weeding and foliar spray of urea @ 2 per cent • Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells • Life saving irrigation through dovas, well, ponds, check dams and bora bandh 	
c) Remarks on Implementation	
Promote for the construction of Rain water harvesting structure watershed programme and MNREGA	



C. Monsoon/Weather Situation: Terminal drought (Early withdrawal of monsoon)

C1. At fruiting/pre physiological maturity stage	
C1.1. Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils	
Normal Crop/cropping system	Upland Rice , Finger millet, Pigeonpea + Sorghum, Pigeonpea + Maize
Suggested Contingency measures	
<p>a) Change management</p> <ul style="list-style-type: none"> • Life saving irrigation to vegetables through stored moisture from constructed DOVA • If not possible to make survival harvest it for fodder use 	
<p>b) Rabi Crop planning</p> <ul style="list-style-type: none"> • Cultivation of Niger, Horsegram, Toria, linseed as relay/paira cropping • In case of availability of irrigation, go for cultivation of early Potato and pea (early Arkel group) • Prepare kachha check dam or Bora Bandh for Water conservation • Mid early variety of radish cultivation is recommended 	
<p>c) Remarks on Implementation</p> <p>Promote for the construction of Farm ponds through watershed management programme and MNREGA</p>	
C1.2. Major Farming Situation/Land Situation: MID LAND Sandy loam soils	
Normal Crop/cropping system	Rice
Suggested Contingency measures	
<p>a) Crop management</p> <p><u>Don 2</u></p> <ul style="list-style-type: none"> • At milking , soft and dough stage spray KCL @ 2 per cent • In case of gundhi bug attack found more than ETL(>2 gundhibug /m²), spray Chlorpyriphos dust or Monocrotophos @ 1 ml/lt • If possible go for life saving irrigation • Late season drought generally results in outbreak of foliar, node, collar or neck blast of rice depending on the stage of crop. <p><u>Don 3</u></p> <p>Instead of grain purpose crops like sorghum, pearlmillet, maize, cowpea, black and Greengram that can be harvested for fodder use</p>	
<p>b) Rabi crop planning</p> <ul style="list-style-type: none"> • Ensure for all inputs required for rabi season in advance. • In case of failure of kharif crops prefer sowing of pre rabi catch crops like, Toria, Niger, Horsegram, blackgram, sesame linseed in uplands to medium lands 	
<p>c) Remarks on Implementation</p> <p>Promote construction of Rain water harvesting structure watershed programme and MNREGA</p>	
C1.3. Major Farming Situation/Land Situation: LOW LAND Sandy loam soils	
Normal Crop/cropping system	Rice
Suggested Contingency measures	
<p>a) Crop management</p> <ul style="list-style-type: none"> • Life saving irrigation. • The land should be tilled properly in case <i>kharif</i> crop fails sow <i>rabi</i> crops like safflower, pigeonpea in sept-Oct (Short duration) • Spray KCL @ 2 per cent followed by Urea @ 2 per cent • Mid early rice crop may be harvested at Physiological maturity • Cultivate vegetables like Tomato, Brinjal, Capsicum, Shimla mirch, Broccoli, Cabbage and Cauliflower, green pea and potato as per suitability near and around tributaries 	
<p>b) Rabi crop planning</p> <p>Prefere early sowing of wheat, Mustard, Chickpea,linseed and lentil as sole or intercrop Wheat + Chickpea (4:2) Wheat+ Mustard (4:3)</p>	
<p>c) Remarks on Implementation</p> <p>Promote construction of Rain water harvesting structure watershed programme and MNREGA</p>	

PART-III

A. Unusual rains: Continuous high rainfall in a short span leading to water logging

Suggested Contingency measures
a) Crop management
Pigeonpea /Sorghum/Pearl millet
Vegetative stage- Prefer ridge and furrow method of sowing. Ensure for proper drainage through channel. Collect runoff water in Dovas for further use.
Flowering stage- Ensure for proper drainage through channel. Collect runoff water in Dovas for further use.
Crop maturity stage- No such situation at the time of maturity
Post harvest- After Sun drying follow grading and storing
Blackgram and other pulses/Oilseeds
Vegetative stage- Follow Ridge and furrow sowing
Ensure for proper drainage through channel
Collect runoff water in Dovas for further use
Avoid application of fertilizer
Flowering stage- Ensure for proper drainage through channel
Collect runoff water in Dovas for further use
Avoid application of fertilizer
Prophylactic measure for jassid and YMV
Crop maturity stage-
Post harvest-
Rice
Vegetative stage- Safe disposal of excess water from rice field. Bund repairing and strengthen. Application of insecticides in the afternoon hours is preferred seeing the weather condition or after spraying weather should remain rain free for at least 4-5 hrs. Retransplant to maintain plant population in case of mortality more than 50 %
In partially damaged crop, allow to withstand upright. Flood occurs due to heavy storm in mid and lowland which when recedes probability of occurrence of swarming caterpillar on field bunds and around of rice crop is more. So, when it crosses the Economic Threshold Limit (ETL) i.e., one larva / hill then spray the crop with Chlorpyriphos/ Triazophos/ Profenophos @ 2 ml/lt water or dust the crop with Quinalphos @ 1.5% D @ 10kg/ acre. To prevent migration of larvae from one field to other, bunds should be heavily dusted with the dust formulation mentioned above. In partially ponded field, rice caseworm and in general leaf folder attack is expected. If 1-2 cases or folded leaves/hill is seen spray the crop with Monocrotophos / Chlorpyriphos @ 1 ml/lt water or with Cartap Hydrochloride 50 SP / Fipronil 5 SP @ 200 g/acre. Rain storms during <i>kharif</i> may result in severe occurrence of bacterial leaf streak and bacterial blight in rice. It is advised to spray the crop immediately after every rainspell with streptocycline @ 1g/10 lt water or plantomycin @ 1g/lt water or bacterinol @ 2g/lt. water. Control snail occurrence by Acaricide
Flowering stage- Safe disposal of excess water from rice field. Bund repairing and strengthen. Avoid application of fertilizer. Flood occurs due to heavy storm in mid and lowland which when recedes probability of occurrence of swarming caterpillar, BPH and cut worm on field bunds and around of rice crop is more. So, when it crosses the Economic Threshold Limit (ETL) i.e., one larva / hill then spray Chlorpyriphos/ Triazophos/ Profenophos @ 2 ml/lt water or dust the crop with Quinalphos @ 1.5% D @ 10kg/ acre. To prevent migration of larvae from one field to other, bunds should be heavily dusted with the dust formulation mentioned above. In partially ponded field, rice caseworm and in general leaf folder attack is expected. If 1-2 cases or folded leaves/hill is seen spray the crop with Monocrotophos / Chlorpyriphos @ 1 ml/lt water or with Cartap Hydrochloride 50 SP / Fipronil 5 SP @ 200 g/acre. Unusual and heavy rain during <i>kharif</i> may result in severe occurrence of bacterial leaf streak and bacterial blight in rice. It is advised to spray the crop immediately after every rain spell with streptocycline @ 1g/10 lt water or plantomycin @ 1g/lt water or bacterinol @ 2g/lt. water. Control snail occurrence by Acaricide.
Crop maturity stage- Provide drainage for fast removal of water from the field to favour harvesting
Post harvest- Protect the grain from rain and store it after sun drying for 2-3 days
Maize
Vegetative stage- Prefer ridge and furrow method of sowing. Ensure for proper drainage through channel. Earthingup after downpour. At Knee stage apply thimate 10 G @ 4-6 grains in whirl
Flowering stage- Ensure for proper drainage through channel. At flowering and silking stage for ant attack apply dust on silks @ 0.5 g / cob
Crop maturity stage- Provide drainage for fast removal of water from the field to favour harvesting
Post harvest- Protect grains from rain and store it after sun drying for 2-3 days



Horticulture

Vegetative stage- Prefer ridge and furrow method for sowing and proper drainage. Ensure for proper drainage through water ways. Collect runoff water in Dovas for further use. Soil drenching Carbofuran 3G @ 3 g/lt water against insects. In case of web formation with leaves apply (Nuvan)DDVP @ 1 ml/lit water as a fumigant
Flowering stage- Apply hormone to prevent flower drop. Ensure for proper drainage. Take precaution against wilting and fruit rot. In Tomato and Brinjal-drenching Bavisting @ 2 ml/lit. water + Streptocycline @ 1-2 g/lit water. In Cauliflower -In case of Incidence of collar rot -Spraying of Saaf (Metalaxyl + Mancozeb) @ 2 g/lit water solution. Drainage of excess water. In Lady's finger- YVMV- Spray insecticide followed by fungicide. Soil drenching Carbofuran 3G @ 3 g/lit water against insects. In case of web formation with leaves apply (Nuvan) DDVP @ 1 ml/lit water as a fumigant

Crop maturity stage- Take precaution against wilting and fruit rot. For wilting– Soil drenching with Bavistin @ 2 ml/lit + Streptocycline @ 1-2 g/lit water. In YMVM- Insecticide followed by fungicid

Post harvest- Immediate harvest and safe disposal of produce

Vegetables- (Cucurbits,/ Tomato/ Brinjal/ Cauliflower/ Cabbage/ Lady's finger/Dolichos bean/Amaranthus leaf/ Cariander leaf/Radish)

Vegetative stage- Sowing on ridge and drainage through furrow. Prophylactic measures against pest and diseases. Damaged twigs and leaves may be removed and follow fungicide spraying and stacking

Flowering stage- Apply hormone to prevent flower drop. Ensure for proper drainage. Take precaution againsts wilting and fruit rot. In Tomato and Brinjal-drenching Bavisting @ 2 ml/lit + Streptocycline @ 1-2 g/lit water. In Cauliflower -In case of Incidence of collar rot -Spraying of Saaf (Metalaxyl + Mancozeb) @ 2 g/lit water solution. Drainage of excess water. In Lady's finger- YVMV- Spray insecticide followed by fungicide. Provide support through stacking

Crop maturity stage- Take precaution against wilting and fruit rot. In Wilting– Soil drenching with Bavistin @ 2 ml/lit. water + Streptocycline @ 1-2 g/lit water. In YMVM- Insecticide followed by fungicide

Provide support through stacking.
Post harvest- Immediate harvest and sell produce safely in the market

b) Pest and disease management

Rice

Vegetative stage- Sheath blight- Hexaconazole @ 1ml/lit water. Blast- Tricyclazole @ 6 g/10 lt water

Flowering stage- Sheath blight- Hexaconazole @ 1ml/lit water. Blast- Tricyclazole @ 6 g/10 lt water. Falsesmut-Nativo @ 4g/10 lt water

Crop maturity stage- False Smut - Control- Nativo @ 4g/10 lt water or Propiconazole + Tricyclazole 52.5 SE @ 1ml/lit water. In case of grain discolourness (Grain blast). Spray Tricyclazole @ 6 ml / 10 liter water

Post harvest- Store grains after proper sun drying to minimize the incidence of stored grain pest

Maize

Vegetative stage- Stem borer Control- Carbofuran 3 G @ 12 Kg/acre or Phorate 10G@ 4 kg/acre

Flowering stage- Sheath blight Control- Hexaconazole1-2 ml/lit water

Vegetables- (Cucurbits,/ Tomato/ Brinjal/ cauliflower/ cabbage/ lady's finger/Dolichos bean/Amaranthus leaf/ Cariander leaf/Radish)

Vegetative stage- Before sowing apply in soil, Carbofuran 3 G @2-3 g/m². Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt streptocycline or 2-3 g/lit plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lit water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lit. water against downy mildew diseases of cucurbits.

Flowering stage- Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt. water streptocycline or 2-3 g/lit. water plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lit water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lit. water against downy mildew diseases of cucurbits. YVM Control- Carbofuran 3G @ 3 or Phorate 10 G @ 1 g/m² followed by any fungicide

Crop maturity stage- Stop spraying 1 week before harvesting

Post harvest- Harvest and sell produce in the market

French bean-

Vegetative stage- Rust disease Control- Mancozeb 2g/ lt water. Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt. water streptocycline or 2-3 g/lit. water plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lit water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lit. water against downy mildew diseases of cucurbits.



Flowering stage- Take care of pod borer and aphid attack. Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt streptocycline or 2-3 g/lt plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lt water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lt. water against downy mildew diseases of cucurbits.
Crop maturity stage- Stop spraying 1 week before harvesting
Post harvest- Harvest and sell produce in the market

B. Extreme Weather Events

Suggested Contingency measures

Hail storm

Seedling / nursery stage- Vegetable nursery should be raised in poly house or make proper arrangement of low height Polly tunnels in open area or cover with plastic sheet or thatching should be done

Vegetative stage- In vegetables-Remove damaged parts immediately and apply insecticide followed by fungicide as prophylactic measures. Follow fertilization through foliar as well as broadcasting

Reproductive stage- n vegetables- Remove damaged parts immediately and apply insecticide followed by fungicide as prophylactic measures. Follow fertilization through foliar as well as broadcasting for proper fruiting

At harvest- Safely sell in the market after grading for immediate returns

Heat Wave

Wheat Chickpea/pea

Seedling / nursery stage- For protection from heat and cold wave there is intervention to sow the rabi crops in between 2nd week of October to 2nd week of November to protect their vegetative phase from ground/radiation frost results from cold wave/wind chill injury and reproductive phase from terminal heat stress on Mustard, Chickpea, Wheat, Lentil, Linseed and pea crops. Life saving irrigation

Vegetative stage- Timely sown crop never face heat stress while very late sown(January) crop face heat stress hence only one option is to provide life saving irrigation and water spray during evening time frequently at 2-3 days intervals. Take care of termite attack by spraying Chlorpyriphos @ 1 ml/lt. water and drenching @ 3-5 ml/lt water

In Chickpea because of high soil and ambient temperature (> 35 °C) favours the dry root rot disease starts during flowering/reproductive stage (spraying Captan or thiram or carbendazim or ridomil MZ or Saaf @ 1,5-2 g/lt water)

Reproductive stage- To minimize the terminal heat stress during the month of March and April the only and only way is to provide frequent protective irrigation irrespective of theirs stages (Life saving irrigation). Take care of termite attack by spraying Chlorpyriphos @ 1 ml/lt. water and drenching @ 3-5 ml/lt. water. In Chickpea because of high soil and ambient temperature (> 35 °C) favours the dry root rot disease starts during flowering/reproductive stage (spraying Captan or thiram or carbendazim or ridomil MZ or Saaf @ 1,5-2 g/lt water)

At harvest- Frequent irrigation should be provided to meet the evaporative losses.

Tomato/Brinjal/ lady's finger/Cucurbits

Seedling / nursery stage- Due to heat stress wilting and mortality is more hence frequent irrigation and cover the nursery with mulch(Straw/leaves)

Vegetative stage- Due to heat stress wilting and mortality is more hence frequent irrigation and cover the nursery with mulch(Straw/leaves)

Reproductive stage- Drying of flower- Spray PCOA. Follow mulching after irrigation

At harvest- Immediate harvest after irrigation and shift it to safer place

Cold wave

Wheat

Seedling / nursery stage- Cold environment during tillering or branching stage favours more number of tillers in wheat and more branching in mustard, chickpea, lentil and linseed crops which prospects for high yield but it is detrimental for potato, tomato, brinjal, pea, creeper vegetables and fruits. Irrigation. Balanced fertilizer application.

Foliar spray of nutrients

Vegetative stage- Light irrigation. Mulching with crop residue \ weeds. Fertilizer application

Reproductive stage- Irrigation, fertilizer application

At harvest- N/A

Pigeonpea/Mustard/Linseed/Chickpea/pea

Seedling / nursery stage- In Mustard because of cool weather aphid insects attack is more prominent (spraying Rogor (Dimethoate) @ 2 ml or or Monocrotophos 36 EC @ 1 ml /lt water during evening time is advised).

In linseed Alternaria blight (For blight spray Double dose (Iprodione 25 % WP + Carbendazim 25 % WP) @ 2 g per lt. water) and powdery mildew (prophylactic spraying of Sulfex @ 3 g or Kairathen 1 ml per lt water twice at weekly interval during evening time) disease are more common. For powdery mildew in pea (spraying Calixin (Tridemorph 80 % EC @ 5 ml per 10 lt water twice are highly recommended).



In Chickpea-Cold and wet environment (High humidity) during seedling stage cause collar rot, black root rot, wet rot, Pythium root and seed rot in Chickpea, while in potato, pea and tomato favours late blight (spraying of Krilaksil or Ridomil MZ chemical @ 1.5 g per liter water), powdery mildew (spraying newly emerged fungicide Double dose (Iprodione 25 % WP + Carbendazim 25 % WP) 2 g per Lt water twice at weekly interval) and bacterial wilt, leaf spot and canker (spraying carbendazim @ 2g/Lt water and streptocycline @ 1g/10 Lt. water at 10 DAP, 25 DAP and 40 DAP) diseases in respective vegetable crops. Anthracnose in cucurbitaceous species. Vegetative stage- Provide light irrigation. Follow mulching with crop residue \ weeds\straw\leaves. **In Mustard** because of cool weather aphid insects attack is more prominent (spraying Rogor (Dimethoate) @ 2 ml or or Monocrotophos 36 EC @ 1 ml /Lt water during evening time is advised)

Reproductive stage- Pigeonpea- During flowering and pod formation stage attack of Pod borer/sucking bug, mites, blister beetle insects as well as sterility disease may occur more (spraying Profenophos 50 EC, methomyl 40 SP or monocrotophos 36 SL kill the larvae but as the webs protect them from contact insecticides hence along with contact insecticides, mixing of fumigant insecticide such as DDVP @ 0.5 ml/Lt. water is required to make the larvae come out from the web. For Mites and Aphids, Dimethoate 30 EC @ 2ml/Lt. water and acaricides such as Dicofol 18.5 EC @ 2.5 ml/l water , for Blister beetle synthetic pyrethroids such as Cypermethrin 10 EC @ 1.0 ml/Lt. waterw or Lamda cyhalothrin 5 EC @ 1.0 ml/l water; for sterility mosaic Dicofol 18.5 EC 2.5 ml or Oxydemeton methyl 25 EC or Dimethoate 30 EC 2.0 ml or ml/l water on alternate row twice at an interval of 10 days are recommended).

Vegetables

Seedling / nursery stage- Raising seedling in Poly house, re sowing if damage is more. Provide shelter belt (Wind break) at appropriate spacing with Sisham, Ghamhar. Provide irrigation and mulching with straw and leaves

Vegetative stage- Provide light irrigation. Follow mulching with crop residue \ weeds\straw\leaves. Disease and pest control, care for chilling injury or replanting

Reproductive stage- Drying of flower- Spray PCOA. Follow mulching after irrigation

At harvest- Grading and safely dispose produce in the marketing

Frost

Wheat

Seedling / nursery stage- N/A

Vegetative stage- Provide light irrigation. Follow mulching with crop residue \ weeds\straw\leaves

Pigeonpea

Seedling / nursery stage- Exposure of crop to smoke by burning waste material during night time

Vegetative stage- Exposure of crop to smoke by burning waste material during night time, Light sprinkler irrigation

Reproductive stage- Exposure of crop to smoke by burning waste material during night time, Light sprinkler irrigation

At harvest- Exposure of crop to smoke by burning waste material during night time, Light sprinkler irrigation

Tomato & Potato and Horticultural crops (fruit)

Seedling / nursery stage- Create smoke around the field by using waste materials or set afire with used mobile oil in north-west or west-north direction towards incoming cold waves. Use polythene or bamboo hoogli in small horticultural /nursery/cash vegetable crops during morning hour and remove it during daytime. In Perennial or Horticulture crop (Fruit) also frequent irrigation followed by mulching, thatching , creating smoke screen s and lighting of fire should be practiced in availability of irrigation facility

Vegetative stage- Earthing up, Irrigation and create smoke around the field by using waste materials or set a fire with used mobile oil in north-west or west-north direction towards incoming cold waves. Use polythene or bamboo hoogli in small horticultural /nursery/cash vegetable crops during morning hour and remove it during daytime. In Perennial or Horticulture crop (Fruit) also frequent irrigation followed by mulching, thatching , creating smoke screen s and lighting of fire should be practiced

Reproductive stage- Immediate harvesting and disposal

At harvest- Harvest in dry weather

Cyclone- Not applicable



CONTINGENCY PLANS FOR RABI

1. Sowing window information

Land type	Cropping system	Crop name	Optimum sowing window (Please mention along with week)
1.Upland	Maize/ Groundnut-/Finger Millet –Vegetable/ Toria Linseed/ Safflower	Toria Linseed, Safflower Vegetable,- Cauliflower, Tomato, cabbage,	Toria-1 st week of September – 4 th week of September Linseed- 4 th week of October- 2 nd week of November Safflower- 1 st week of October- 1 st week of November Vegetables- 2 nd week of September – 4 th week of October
2. Mid Land	Rice –Wheat Rice –Chickpea Rice –Lentil Rice-Linseed Rice-Mustard Soybean- Vegetable Maize-Sunflower/ Safflower Rice-Lathyrus Blackgram-Barley Rice-Fodder	Irrigated-Wheat (Zero tillage), Mustard, Potato, Sunflower, Vegetables (Tomato, Chili, Brinjal, capsicum, Cabbage, Cauliflower, Broccoli, vegetable pea, french bean, Radish, Coriander, Amaranthus, Spinach) Rainfed (Zero tillage)- Barley, Safflower, Chickpea, Lentil, Linseed, Lathyrus Paira cropping-Linseed, Lathyrus Fodder crop- Oat, Maize,	Wheat (Timely)- 3 rd week of October – 4 th week of November Wheat (Late)-1 st week of December – 2 nd week of December Mustard- 2 nd week of October – 1 st week of November Sunflower-1 st week of October-2 nd week of October Linseed- 2 nd week of October – 2 nd week of November Chickpea – 2 nd week of October – 1 st week of November Lentil- 3 rd week opf October- 2 nd week of November Vegetables- 1 st week of October – 4 th week of November Lathyrus- 2 nd week of October – 4 th week of November Oat- 2 nd week of October – 4 th week of October Maize- 2 nd week of October – 4 th week of October
3. Low Land	Rice-Linseed (Paira) Rice-Vegetable (Stream Line based)	Linseed (Paira cropping), Vegetable- Cauliflower, Tomato, cabbage, Fodder crop- Oat, Maize, Lathyrus	Linseed- 4 th week of October – 2 nd week of November Oat- 3 rd week of October – 1 st week of November Maize- 3 rd week of October – 1 st week of November Lathyrus- 4 th week of October – 4 th week of November Vegetable – 1 st week of November -4 th week of December

2. Contingency measures for Field crops grown with residual moisture under rainfed condition

2 (A) Optimal residual moisture

2A1.Land type- UPLAND

a) Cropping system- Maize/ Groundnut-/Finger Millet –Vegetable/Toria/Mustard / Linseed

b) Crop name- Toria Linseed, Safflower Vegetable,- Cauliflower, Tomato, cabbage

c)Sowing Window- Toria- 3rd week of September – 4th week of September, Linseed- 1st week of October- 3rd week of October; Safflower- 1st week of October- 1st week of November; Vegetables - Vegetables- 2nd week of September – 4th week of October

d)Variety- Toria-PT 203, Panchali; Linseed- (rainfed)-Skekhar, Subra, Sweta, T397; Safflower-A 300, Akshagiri 59-2-1, A 1

e) Agronomic management practices

- Rain water harvesting and recycling.
- Deeping of water storing structure(Shallow and deep) in April and May month
- Deep summer ploughing in April and May month.
- Strengthing and raising of field bunds in April and May months
- Sowing in defined window for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better crop stand (Plant population)
- Application of Lime or Dolomite (3-5 q/ha) in soil
- Soil application of Sulphur (20 kg/ha) and boron (1kg/ha) in oilseed, pulses and vegetables.
- Foliar spray of Urea (2 %) at flower initiation and pod formation stage in oilseed and pulses
- Follow seed priming (warm water for 4-6 hrs.) before sowing
- Follow seed treatment with fungicide-insecticide-rhizobium
- Irrigate only at critical stages
- Pre and post emergence weedicide application
- Follow hoeing after manual weeding
- Follow RDF, INM and IPM
- For Water use efficiency use antitranspirant, reflectant and mulches
- Regular monitoring of field for disease and insect attack
- Use pheromone trap and attractant
- Promote protected vegetable cultivation under naturally ventilated polyhouse and net house.
- Timely sowing for better establishment



- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population)
- Lime or Dolomite application in soil
- Foliar spray of Sulphur and boron
- Proper water management

Take care of Aphid, white rust in Mustard, Early, late blight and leaf curling in potato

Toria – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Linseed - Follow seed treatment, Irrigate only at critical stages, Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Precautions for pod borer, bud fly insect and powdery mildew disease management.

2A2. Land type- MEDIUM LAND

a) Cropping system- Rice –Wheat, Rice –Chickpea, Rice –Lentil, Rice-Linseed, Rice-Mustard, Soybean-Vegetable, Maize-Sunflower/ Safflower, Rice-Lathyrus, Blackgram-Barley, Rice-Fodder

b) Crop name- Irrigated-Wheat (Zero tillage), Mustard, Potato, Sunflower, Vegetables (Tomato, Chili, Brinjal, capsicum, Cabbage, Cauliflower, Broccoli, vegetable pea, french bean, Radish, Coriander, Amaranthus, Spinach); Rainfed (Zero tillage)- Barley, Safflower, Chickpea, Lentil, Linseed, Lathyrus; Paira cropping- Linseed, Lathyrus; Fodder crop-Oat, Maize

c) Sowing Window- Wheat (Timely)- 3rd week of October – 4th week of November, Wheat (Late)-1st week of December – 2nd week of December, Mustard- 2nd week of October – 1st week of November, Sunflower-1st week of October-2nd week of October, Linseed- 2nd week of October – 2nd week of November, Chickpea – 2nd week of October – 1st week of November, Lentil- 3rd week of October- 2nd week of November, Vegetables- 1st week of October – 4th week of November, Lathyrus- 2nd week of October – 4th week of November, Oat- 2nd week of October – 4th week of October, Maize- 2nd week of October – 4th week of October

d) Variety- Irrigated Wheat- Timely sown (120-125)- HD2967, WH 1105, K307, HD2733; Late sown (105-110)- HD 3059, DBW 14, HI 1563 (seed rate 25 % more than timely sown); Mustard- Pusa mahek,Pusa mustard 25, NRCHB 101, Bharat sarson 1, Pusa 28, 30; Potato- Kufri Surya, Kufri Arun, Kufri Sutlej, Kufri Laukar, Kufri Lalima; Sunflower – DRSF 108 (OP), KBSH 44; Safflower- A 300, Akshagiri 59-2-1, A 1; Chickpea- Jaki 9218, Kak 2, Birsa Chana 3; Lentil – HUL 57,WBL 77, KLS 218 Linseed-Sharda, Priyam, Divya; Lathyrus-Maha Teora, Oat- Kent, Maize- J 1006, Pratap Makka Chari 6

e) Agronomic management practices

- Seed treatment with Azotobacter and Azospirillum and also soil application
- Timely sowing for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population)
- Follow deep summer ploughing
- Proper water management
- Follow seed treatment
- Irrigate only at critical stages
- Pre emergence weedicide application
- Follow RDF, INM and IPM
- Take care of Painted Bud, Aphid, white rust in Mustard; Early, late blight and leaf curling and grub in potato; Collar rot, Dry root rot, Pod borer in Chickpea; Wilt in Lentil.

Wheat-Seed tretment with Azotobacter and Azospirillum and also soil application.Timely sowing for better establishment. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Follow deep summer ploughing. Proper water management. Irrigate only at critical stages(3-6). Pre emergence weedicide application. Follow RDF, INM and IPM. Take case of Loose Smut Disease. 1st irrigation should be after CRI stage i.e at 30-35 DAS

Mustard – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Potato- Seed treatment. Proper spacing. Frequent irrigation. Take care for leaf curling, Early, late blight and grub infestation. Irrigate during cold day and night to get relief from frost attack. Produce smoke during cooler day and night. Earthing up in potato at 30-35 DAS or crop attended the height of 20-25 cm

Sunflower- Seed treatment, Irrigate the field at the time of final land preparation and follow sowing after two days of irrigation for good germination and crop establishment. Follow proper crop geometry for good crop stand. Other than RDF of NPK also needed to apply Sulphur. Follow hoeing at 20-25 DAS, 35-40 DAS. Keep weed free upto 60 days. Use post emergence weedicide. Follow earthing up at Knee height. Irrigate at Critical stage (20-25 DAS, 55-60 DAS, 70-75 DAS. For summer season- 4-10 light irrigation is required

Chickpea - Seed treatment with Rhizobium culture and Phosphate solubilizing bacteria (PSB) and Trichoderma. Management for Collar rot during temperature fall and dry root rot during temperature increment. Pre emergence weedicide application. Irrigate at Critical stages. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray

Lentil - Foliar spray of Sulphur and Boron is necessary. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Follow deep summer ploughing. Proper water management. Follow seed treatment with Rhizobium culture and PSB. Irrigate only at critical stages. Pre emergence weedicide application. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray.

Linseed - Follow seed treatment, Irrigate only at critical stages, Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Precautions for pod borer, bud fly insect and powdery mildew disease management. Management for wilt disease. One hand weeding followed by two hoeing for management of weeds (HW-20-25 DAS and Hoeing 30-32 and 40-42 DAS).

Lathyrus- Proper seed rate and Follow RDF for potential yield. Follow two nipping in between 25-45 DAS. Irrigate after every nipping. Take care of diseases

Forage-Oat- Proper seed rate for better crop stand. 1st and 2nd cutting at 30 and 45 DAS and 3rd before flowering. Berseem- 1st at 50 DAS and follow 2nd, 3rd and 4th cutting every at an interval of 30-40 days. Lucern-Same as Berseem. Japani Mustard- 1st at 50 DAS during fruiting and rest cutting every at an interval of 30 days. Follow RDF. For Lucern other than N P K use Lime , Boron and Molybdenum micro nutrients for better yield.

2A3 Land type- LOW LAND

- a) Cropping system-Rice- Linseed (Paira), Rice-Vegetable (stream kine baesd)
- b) Crop name- Linseed/ lathyrus (Paira cropping), Vegetables (Cauliflower, Tomato, Cabbage), Fodder Crop- Oat, Maize, lathyrus
- c) Sowing Window - Linseed- 4th week of October – 2nd week of November, Oat- 3rd week of October – 1st week of November, Maize- 3rd week of October – 1st week of November, Lathyrus- 4th week of October – 4th week of November, Vegetable – 1st week of November -4th week of December
- d) Variety- Linseed- Sharda, Priyam, Divya, Lathyrus-Maha Teora; Fodder- Oat— Kent, Maize- Pratap Makka(Chari 6), J 1006, Berseem-Vardhan

e) Agronomic management practices

Linseed - Follow seed treatment. Irrigate only at critical stages. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Follow RDF, INM and IPM. Be cautious for pod borer, bud fly insect and powdery mildew disease management.

Lathyrus- Proper seed rate and Follow RDF for potential yield. Follow two nipping in between 25-45 DAS. Irrigate after every nipping. Take care of diseases

Forage-Oat- Proper seed rate for better crop stand. 1st and 2nd cutting at 30 and 45 DAS and 3rd before flowering, Berseem- 1st at 50 DAS and follow 2nd, 3rd and 4th cutting every at an interval of 30-40 days, Lucern-Same as Berseem, Japani Mustard- 1st at 50 DAS during fruiting and rest cutting every at an interval of 30 days. Follow RDF. For Lucern other than N P K use Lime , Boron and Molybdenum micro nutrients for better yield.



2 (B) Less than optimum moisture i.e., 25% less than normal, which can happen due to insufficient rainfall during September/October months. Deficit of 20-40% rainfall

2B1 Land type- UP LAND

- a) Cropping system- Maize/Groundnut/Finger millet –Toria/Linseed/Safflower
- b) Crop name – Toria, Linseed, Safflower
- c) Sowing Window- Toria- 3rd week of September – 4th week of September, Linseed- 1st week of October- 3rd week of October, Safflower- 1st week of October- 1st week of November
- d) Variety- Toria- PT 203, Panchali; Linseed- Sharda, Priyam, Divya,; Safflower-A 300, Akshagiri 59-2-1, A 1

e) Agronomic management practices

- Rain water harvesting and recycling.
- Deeping of water storing structure(Shallow and deep) in April and May month
- Deep summer ploughing in April and May month.
- Strengthening and raising of field bunds in April and May months
- Sowing in defined window for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better crop stand (Plant population)
- Application of Lime or Dolomite (3-5 q/ha) in soil
- Soil application of Sulphur (20 kg/ha) and boron (1kg/ha) in oilseed, pulses and vegetables.
- Foliar spray of Urea (2 %) at flower initiation and pod formation stage in oilseed and pulses
- Follow seed priming (warm water for 4-6 hrs.) before sowing
- Follow seed treatment with fungicide-insecticide-rhizobium
- Follow deep summer ploughing
- Irrigate only at critical stages
- Pre and post emergence weedicide application
- Follow hoeing after hand weeding
- Follow RDF, INM and IPM
- For Water use efficiency use antitranspirant, reflectant and mulches
- Regular monitoring of field for disease and insect attack
- Use pheromone trap and attractant
- Promote protected vegetable cultivation under naturally ventilated polyhouse and net house.
- Zero Tillage for seed placement at proper depth for better germination
- One hand weeding followed by one hoeing for management of germinating weeds

For Water use efficiency use antitranspirant, reflectant and mulches

Toria – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Linseed - Follow seed treatment, Irrigate only at critical stages, Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Precautions for pod borer, bud fly insect and powdery mildew disease management.

Safflower- Seed treatment, Irrigate the field at the time of final land preparation and follow sowing after two days of irrigation for good germination and crop establishment. Follow proper crp geometry for good crop stand. Other than RDF of NPK also needed to apply Sulphur. Follow hoeing at 20-25 DAS, 35-40 DAS. Keep weed free upto 60 days. Use post emergence weedicide. Follow earthing up at Knee height. Irrigate at Critical stage (20-25 DAS, 55-60 DAS, 70-75 DAS. For summer season- 4-10 light irrigation is required

2B2 Land type- MEDIUM LAND

- a) Cropping system- Rice –Chickpea, Rice –Lentil, Rice-Linseed, Rice-Mustard, Rice-Safflower, Soybean-Vegetable (Early) Maize/ Rice-Lathyrus, Blackgram-Barley, Rice-Fodder
- b) Crop name – Mustard, Vegetables (Tomato, vegetable pea, Radish, Coriander, Amaranthus, Spinach), Rainfed (Zero tillage)- Barley, Safflower, Chickpea, Lentil, Linseed, Lathyrus, Pairs cropping-Linseed, Lathyrus,Fodder crop- Oat, Maize
- c) Sowing Window- Mustard- 2nd week of October – 1st week of November, Safflower-2nd week of October-2nd week of November, Linseed- 2nd week of October – 2nd week of November, Chickpea – 2nd week of October – 1st week of November, Lentil- 3rd week opf October- 2nd week of November, Vegetables- 1st week of October – 4th week of November, Lathyrus- 2nd week of October – 4th week of November, Fodder Crops- Oat- 2nd week of October – 4th week of October



d) Variety - Mustard-Pusa mahek, Pusa mustard 25, NRCHB 101, Bharat sarson 1 Pusa 28, 30; Safflower- A 300, Akshagiri 59-2-1, A 1, Chickpea- Jaki 9218, Kak 2, Birsa Chana 3; Lentil – HUL 57,WBL 77, KLS 218 Linseed- Sharda, Priyam, Divya; Lathyrus-Maha Teora, Oat- Kent, Maize- J 1006, Pratap Makka Cheri 6

e) Agronomic management practices

Mustard – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Sunflower- Seed treatment, Irrigate the field at the time of final land preparation and follow sowing after two days of irrigation for good germination and crop establishment. Follow proper crop geometry for good crop stand. Other than RDF of NPK also needed to apply Sulphur. Follow hoeing at 20-25 DAS, 35-40 DAS. Keep weed free upto 60 days. Use post emergence weedicide. Follow earthing up at Knee height. Irrigate at Critical stages (20-25 DAS, 55-60 DAS, 70-75 DAS. For summer season- 4-10 light irrigation is required

Chickpea - Seed treatment with Rhizobium culture and Phosphate solubilizing bacteria (PSB) and Trichoderma. Management for Collar rot during temperature fall and dry root rot during temperature increment in. Pre emergence weedicide application. Irrigate at Critical stages. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray

Lentil - Foliar spray of Sulphur and Boron is necessary. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Follow deep summer ploughing. Proper water management. Follow seed treatment with Rhizobium culture and PSB. Irrigate only at critical stages. Pre emergence weedicide application. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray. Management for wilt disease. One hand weeding followed by two hoeing for management of weeds (HW-20-25 DAS and Hoeing 30-32 and 40-42 DAS)

Linseed - Follow seed treatment. Irrigate only at critical stages. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Follow RDF, INM and IPM. Precaution for pod borer, bud fly insect and powdery mildew disease management

Forage- Oat- Proper seed rate for better crop stand. 1st and 2nd cutting at 30 and 45 DAS and 3rd before flowering. Berseem- 1st at 50 DAS and follow 2nd , 3rd and 4th cutting every at an interval of 30-40 days. Lucern- Same as Berseem. Japani Mustard- 1st at 50 DAS during fruiting and rest cutting every at an interval of 30 days. Follow RDF. For Lucern other than N P K use Lime , Boron and Molybdenum micro nutrients for better yield.

2B3 Land type- LOW LAND

a) Cropping system- Rice- Linseed/Lathyrus (Paira), Rice-Vegetable(Using harvested water)

b) Crop name- Linseed/ lathyrus (Paira cropping), Vegetables (Cauliflower,Tomato, Cabbage) Spinach, Fodder Crop-boat, Maize, lathyrus

c) Sowing Window- Linseed- 4th week of October – 2nd week of November; Lathyrus- 4th week of October – 2nd week of November; Vegetable- 3rd week of November- 4th week of December; Fodder-2nd week of November – 2nd week of December, Fodder Crops-Oat- 3rd week of October – 1st week of November , Maize- 3rd week of October – 1st week of November

d) Variety- Linseed- Sharda, Priyam, Divya, Lathyrus-Maha Teora; Fodder- Oat— Kent, Maize- Pratap Makka(Cheri 6), J 1006

e) Agronomic management practices

Linseed - Follow seed treatment. Irrigate only at critical stages. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Follow RDF, INM and IPM. Precaution for pod borer, bud fly insect and powdery mildew disease management.

Forage- Oat- Proper seed rate for better crop stand. 1st and 2nd cutting at 30 and 45 DAS and 3rd before flowering. Berseem- 1st at 50 DAS and follow 2nd , 3rd and 4th cutting every at an interval of 30-40 days. Lucern- Same as Berseem. Japani Mustard- 1st at 50 DAS during fruiting and rest cutting every at an interval of 30 days. Follow RDF. For Lucern other than N P K use Lime , Boron and Molybdenum micro nutrients for better yield.



CONTINGENT STRATEGIES FOR LIVESTOCK, POULTRY & FISHERIES

1 Livestock

Suggested contingency measures under DROUGHT event

a) Before the event

Feed and fodder availability

Preservation of surplus fodder, encourage fodder cultivation and tree plantation and also encourage supply of molasses to cattle feed plants

- Preservation of surplus fodder

Green grass is a good source of vitamin A which is present in the form of Carotene. One kg of green grass provides 50mg of vitamin A and 15 to 20g protein to the animal. Cowpea, beans, subabul leaves etc. give 30 to 40g of protein. From grass fodder herbivorous animals get the carbohydrates (energy source), proteins ("building material" of the body) and vitamins (especially carotene), which are the main drives of sustainable operation of the body.

Two methods are available for preserving or conserving the seasonal excess of green fodder, viz. hay making and silage making. Each method has its own limitations and advantageous. Ensiling is preferred on the basis of fodder quality.

Hay making

Hay -refers to cereals, grasses or legumes that are harvested at appropriate stage, dried and stored

Ensilage / Silage making

Silage may be defined as the green succulent roughage preserved under controlled anaerobic fermentation in the absence of oxygen by compacting green chops in air and watertight receptacles.

- Complete Feed Blocks

Supply enriched complete feed blocks containing dry roughage, concentrates/ unconventional supplements 50:50 ratio. Complete feed blocks may be sourced from different commercial sources.

Feeding practices for livestock in India at present separate feeding of roughage and concentrate

- ❖ Chopped roughage and soaked concentrate mixed together

- ❖ Chopped roughage mechanically mixed with concentrate as mash

- ❖ Chopped roughage and concentrate ingredients mixed and densified as Complete Feed Block

Concept of densified complete feeds with fibrous crop residues is a noble way to increase the intake and improve the nutrients utilization. A complete feed block has been defined as a system of feeding all ingredients including roughages, processed and mixed uniformly, to be made available ad lib to the animals.

- Urea molasses mineral block licks

Urea-molasses mineral block lick can sustain the animals by providing protein, energy and essential minerals. It is cost effective, easy to handle and transport and available commercially through milk cooperatives. Therefore, it is required that urea molasses blocks stocks (UMBS) are made available in the rain-deficient areas.

- Methods used for improving nutritive quality of straws and other crop residues like urea treatment

Spray dry roughages such as paddy and wheat straw with about 10% molasses and 2% urea for maintenance of animals in fodder deficit areas.

Preparation of 100 kg roughage-based enriched feed containing 88.8 kg wheat straw or any other straw/stover, 10 kg molasses, 1 kg urea and 0.5 kg mineral mixture will cost about Rs. 375-450 per quintal.

- Utilization of forest byproducts for feeding of livestock

Use of dry and fallen tree leafs like Pipal, Neem, Mango and Kathal etc.

- Making Leaf meal

- Use of conventional and non conventional feeds

- Rice Mills

The main by-products of rice are rice straw, rice husk or hull, and rice bran. Rice straw is produced when harvesting paddy. Rice husks generated during the first stage of rice milling, when rough rice or paddy rice is husked.

- Aquatic plants

- ❖ One kg DM/100 kg BW

- ❖ Water hyacinth, aquatic spinach, Stalks & leaves of lotus plant, Hydrilla, Pistia etc.

- Encourage supply of molasses to cattle feed plants

Molasses and Bagasse are the byproducts from sugarcane industry and are available in abundance. They can be used as cattle feed after supplementation with urea. Such a ration is a ready feed during drought and scarcity conditions when nothing else is available for feeding to animals.

- Crop Residue Enrichment & Densification

Crop residues can be fortified with feed ingredients like cakes, barns, grains, molasses, hay, minerals and then densified into blocks or pellets to save on storage and transport costs. Also balanced ration in the form of complete diet or total mixed ration as per need of animals can be supplied for improved productivity.

- Demonstration of Re-vegetation of Common Grazing Land

The grazing lands play an important role in the lives of rural people who are getting fodder, fuel, drinking water from commons. However, such lands are being continuously degraded due to overgrazing and overexploitation by locals. Re-vegetation of such lands on scientific lines suiting to agro-climatic conditions is to be demonstrated through strengthening institutional arrangement at village level. Fodder production from such lands can be enhanced substantially by introducing high yielding cultivated fodder crops, grasses and pasture legumes. An integrated approach of growing cultivated crops, grasses, trees and shrubs under silvi-pastoral/ horti - silvipasture system will improve overall productivity of such land.

Drinking water

Repairs of tube wells, clear off the sludge in the canals and local water catchments and clean the water tanks, large ponds and lakes

Health and Hygiene

Tick damage and tick-borne diseases

- Tick damage - Vaccinate the cattle against tick-borne diseases
- Tick-borne diseases- Vaccination is best done in calves under 6 months of age and one dose is sufficient
- Babesiosis (Red water)- Treatment involves keeping the cattle calm. They should not be driven over long distances and should be injected with Berenil or Imizol. The dose for Berenil is 5 ml of made up solution (1 packet mixed with 12.5 ml of sterile water) for each 100 kg (for example, 20 ml for a 400 kg animal)
- Sarcoptic Mange in pigs- Not applicable before event

Diseases caused by biting insects

- Trypanosomiasis- Fly control is important for prevention of the disease.
- Three-day stiff sickness- Prevention is by vaccination
- Lumpy-skin disease- Prevention is by vaccination

Diet related Disease problems

- Eating plastic bags and wire(Pica)- Feed cattle well, especially in winter. Clear wires and plastic bags from the grazing area. Watch cattle closely when they are grazing. Mineral mixture supplement should be given to the animal
- Poisonous plants- Not applicable before event
- Botulism- Prevention involves vaccination and good nutrition. Burn or bury all carcasses, bones or decaying material

Deficiency diseases

Cattle grazing on drought-stricken pastures can suffer serious depletion of reserves of minerals and vitamins.

- Copper and Cobalt- Not applicable before event
- Calcium, Phosphorous & Vit. D- Not applicable before event
- Vitamin A- Not applicable before event

Infectious Diseases

- Foot and Mouth Disease (FMD)- Vaccination at the age 4 months and above. Booster should be given 1 month after first dose then every six monthly
- Haemorrhagic Septicaemia (HS)- Vaccination at the age 6 months and above. Annually in endemic areas. Vaccinate the animal before onset of monsoon every year preferably in the month of May and June.
- Black Quarter (BQ)- Vaccination at the age 6 months and above. Annually in endemic areas. Vaccinate the animal before onset of monsoon every year preferably in the month of May and June
- Anthrax- Vaccination at the age 4 months and above. Annually in endemic areas. Vaccinate the animal before onset of monsoon every year preferably in the month of May and June.
- Rabies (Post bite therapy only)- Not applicable
- Enterotoxaemia (pulpy kidney)- Vaccinate the animal at the age of 3-4 months, repeat after 15 days and then annually.
- Pneumonia- Not applicable

Non-Infectious Diseases

- Ruminant tympany (Bloat)- Not applicable
- Rumen acidosis- Not applicable
- Intussusception- Deforming should be given
- Pregnancy toxemia (Ketosis)- Fed the pregnant animal with balanced ration.

Poisoning

- Organochlorine compounds- Not applicable
- Organophosphorus compounds- This group consists of malathion, darathion, chlorathion, carbophenothion, demeton, dasnon, dimethylparathion, trichlorphon, dioxalthion etc. Symptoms of toxicity are profuse salivation, muscle stiffness, dyspnoea with open mouth breathing, tremors. Treatment consists of administering antidote, usually atropine sulphate.
- Snake bite- Not applicable

b) During the event

Feed and fodder availability

- Lactating and pregnant animals need to be provided enriched feed to meet the requirements and rest of animals be provided the maintenance diet. In case of acute shortage, lactating animals be provided feed meeting 50% of the requirements to maintain minimum level of production.
- Drought tolerant fodder crops (like sorghum PC 6 and MP chari, cowpea - BL 1 and 2) and fodder grasses (like stylo, *cenchrus ciliaris*, *athropogon*, etc.) should be cultivated. Under the mini kit programme, the developmental department need to provide fodder crop seeds in the drought-affected areas.
- Provide salt dose daily through feed (40-50 g of salt per adult animal and 10-20 g for small ruminants and calves).

Issue

- Large scale migration -Creating additional resources in drought prone area
- Grazing of poisonous plants/toxicity problems -Inventory of anti nutritional/toxic factors. Creating awareness in farmer for avoiding nitrate/nitrite HCN poisoning.
- Transport of fodder from normal DPA-Establishing feed and fodder banks. Effective mechanism for distribution of fodder/feed to productive animals. Densification/baling/briquette technologies

Drinking water

Harnessing water through the existing reservoirs and exploitation of groundwater.

Health and Hygiene

Tick damage and tick-borne diseases

- Tick damage - If disease occurs Treat the cattle against tick-borne diseases. Consult Veterinarian.
- Tick-borne diseases- Prevention is by tick control, treatment of diseased animal and vaccination. Consult Veterinarian.
- Babesiosis (Red water)- Treatment involves keeping the cattle calm. They should not be driven over long distances and should be injected with Berenil or Imizol. The dose for Berenil is 5 ml of made up solution (1 packet mixed with 12,5 ml of sterile water) for each 100 kg (for example, 20 ml for a 400 kg animal). Consult Veterinarian.
- Sarcoptic Mange in pigs- Itching; dermatitis; rubbing; scratching; reduced growth rate. Miticidal sprays; pour-ones injection and in-feed premix. Consult Veterinarian.

Diseases caused by biting insects

- Trypanosomiasis- Treated with SURAMIN through intramuscular injection or intravenous infusion if sufficient observation is possible. Consult Veterinarian.
- Three-day stiff sickness- It is important that the animal is given food and water if it is unable to stand
- Animal should be treated by Veterinarian
- Lumpy-skin disease- If your cattle get this disease, you should speak to your state veterinarian

Diet related Disease problems

- Eating plastic bags and wire (Pica) - Mostly occurring in those animals which are having shortage of feeds and fodder and deficiency of Phosphorus. Prevention involves the following: - Feed cattle well, especially in winter. Clear wires and plastic bags from the grazing area. Watch cattle closely when they are Grazing. Mineral mixture supplement should be given to the animal. Once the cow has eaten plastic bags or wire, the only effective treatment is an operation. Consult Veterinarian.
- Poisonous plants- Due to scarcity of feed s and fodder animals used to consume poisonous plans and are more likely to get toxicity. Poisoning can also happen when owner or animal handler move cattle to new paddocks where toxic plants occur. Consult Veterinarian.
- Botulism- Botulism can occur when cattle eat carcass and bone material when there is a lack of feed during drought or if they have a phosphorus deficiency
- Treatment is only possible in the early stages and requires an antitoxin. Consult Veterinarian.

Deficiency diseases

Cattle grazing on drought-stricken pastures can suffer serious depletion of reserves of minerals and vitamins.

- Copper and Cobalt- Characterized by anorexia and wasting. Deficiency affects growth and fertility of the cattle. Anemia, diarrhoea and unthriftiness occur in extreme cases. Copper or cobalt sulphate in the form of mineral mixture supplement causes rapid disappearance of the symptoms
- Calcium, Phosphorous & Vit. D- Deficiency may result in rickets in calves and osteomalacia in adults. Mineral supplementation in diet is essential to prevent this deficiency.
- Vitamin A- Vit. A deficiency occurs in cattle on dry countryside during periods of drought. Symptoms include night blindness, corneal keratinization, ptyriasis, hoof defects, loss of weight and infertility. Animals should have access to green pasture and should be supplied with Vit. A in feed to prevent deficiency.

Infectious Diseases

- Foot and Mouth Disease (FMD)- If outbreak occurs then the animal should be treated. Foot lesion should be washed with soap / detergent and apply Povidon iodine lotion while in mouth lesion boroglyserine should be applied. Consult Veterinarian.
- Haemorrhagic Septicaemia (HS)- If disease occurs animal should be treated with Broad Spectrum Antibiotic like penicillin at higher dose and other supportive medicine as per the symptoms. Consult local Veterinarian.
- Black Quarter (BQ)- If disease occurs animal should be treated with Broad Spectrum Antibiotic like penicillin at higher dose and other supportive medicine as per the symptoms. Consult local Veterinarian.
- Anthrax- If disease occurs animal should be treated with Broad Spectrum Antibiotic like penicillin at higher dose and other supportive medicine as per the symptoms. Consult local Veterinarian.
- Rabies (Post bite therapy only)- Vaccinate the animal immediately after suspected bite. Booster should be given on 3, 7, 14, 28 and 90 (optional) days after first dose.
- Enterotoxaemia (pulpy kidney)- Not applicable
- Pneumonia- Not applicable

Non-Infectious Diseases

- Ruminant tympany (Bloat)- Not applicable
- Rumen acidosis- Ingestion of large amounts of highly fermentable carbohydrate feeds causes an acute illness due to excess production of lactic acid in the rumen. Clinically, the disease is manifested by dehydration, blindness, recumbency, complete rumen stasis and a high mortality rate. Normal saline, sodium bicarbonate and antihistaminic are advised.
- Intussusceptions- It occurs commonly due to nodular worms, change in feed and local intestinal problems. The animal is dull, off-feed, kicking at the belly with no rise of temperature, frequent straining with no defecation, colic symptoms, and at later stages, recumbency. Emergency surgery is the only rational treatment.
- Pregnancy toxæmia (Ketosis)- It is a highly fatal disease caused due to a decline in the plane of nutrition and short periods of starvation (40 hrs) during the last two months of pregnancy. Treatment comprises intravenous administration of 50% glucose. Supply of molasses in the ration and concentrate in the last two months of pregnancy helps in preventing the condition.

Poisoning

- Organochlorine compounds- Not applicable
- Organophosphorus compounds- This group consists of malathion, darathion, chlorathion, carbophenothion, demton, dasnon, dimethylparathion, trichlorphon, dioxalthion etc. Symptoms of toxicity are profuse salivation, muscle stiffness, dyspnoea with open mouth breathing, tremors. Treatment consists of administering antidote, usually atropine sulphate.
- Snake bite- Usually bitten on the scrotum or udder. The presence of hair may obscure the typical fang marks. Prolonged pain, muscular weakness, impaired vision, nausea and paralysis are generally exhibited along with symptoms of shock. If anti-venin is not available and the bite is located in an area where a tourniquet cannot be applied, excision of an area of skin and sub-cutaneous tissue can be life-saving

c) After the event

Feed and fodder availability

Promotion of fodder seed production, cultivation and storage, establishment of fodder block making machines in fodder surplus areas

Post flood feeding management

- Animal should not be allowed to graze in water logged area
- Feeds to be protected from fungal contamination & wet feeds to be dried & fed
- Provides clean drinking water to animals
- Provide ready to eat feed blocks particularly the pregnant and lactating animals
- Requirement of energy may be met providing crude molasses
- Top feeds/ tree leaves available in the area to be provided to meet the DM requirement

Specific contingencies can be adopted for livestock feeding depending upon availability as under in different regions during drought situation

Neem seed kernel cake (NSKC), Saw dust, Paper waste, Agave (Ketki), Cactus, Tree leaves and vegetable leaves, Cher leaves and fruits, Straw and gotars, Sugarcane bagasse as animal feed and Use of damaged grains as feed

Drinking water

To strengthen reservoirs by promoting recharging of water and rain water harvesting during rainy season.

Health and Hygiene

Tick damage and tick-borne diseases

- Tick damage - Treat the cattle against tick-borne diseases. Consult Veterinarian.
- Tick-borne diseases- Prevention is by tick control, treatment of diseased animal and vaccination. Consult Veterinarian.
- Babesiosis (Red water)- Treatment involves keeping the cattle calm. They should not be driven over longdistances and should be injected with Berenil or Imizol. The dose for Berenil is 5 ml of made up solution (1packet mixed with 12.5 ml of sterile water) for each 100 kg (for example, 20 ml for a 400 kg animal). Consult Veterinarian.
- Sarcoptic Mange in pigs- Not applicable after event

Diseases caused by biting insects

- Trypanosomiasis- Treated with SURAMIN through intramuscular injection or intravenous infusion if sufficient observation is possible. Consult Veterinarian
- Three-day stiff sickness- It is important that the animal is given food and water if it is unable to stand.
- Animal should be treated by Veterinarian
- Lumpy-skin disease- If your cattle get this disease, you should speak to your state veterinarian

Diet related Disease problems

- Eating plastic bags and wire (Pica)- Feed cattle well, especially in winter. Clear wires and plastic bags from the grazing area. Watch cattle closely when they are grazing. Mineral mixture supplement should be given to the animal
- Poisonous plants- Not applicable
- Botulism- Prevention involves vaccination and good nutrition. Burn or bury all carcasses, bones or decaying material

Deficiency diseases

Cattle grazing on drought-stricken pastures can suffer serious depletion of reserves of minerals and vitamins.

- Copper and Cobalt- Not applicable
- Calcium, Phosphorous & Vit. D- Not applicable
- Vitamin A- Not applicable

Infectious Diseases

- Foot and Mouth Disease (FMD)- If outbreak occurs then the animal should be treated. Foot lesion should be washed with soap / detergent the apply Povidon iodine lotion while in mouth lesion boroglyserine should be applied. Consult Veterinarian.
- Haemorrhagic Septicaemia (HS)- Not applicable
- Black Quarter (BQ)- Not applicable
- Anthrax- Not applicable
- Rabies (Post bite therapy only)- Not applicable
- Enterotoxaemia (pulpy kidney) - It affects the animals in a high state of nutrition on a lush feed, grass or grain. Morbidity rates seldom exceed 10% but mortality rate approximates 100%. Under certain conditions, the organism proliferated rapidly in the intestines and produces lethal quantity of toxin. Suphadimidine with other supportive medicine may be effective for treatment
- Pneumonia- It is one of the most common and important pathological conditions. It is characterized clinically by increased respiration, coughing and abdominal breathing. Treatment with broad spectrum antibiotic, nabulization and other supportive drugs is effective.

Non-Infectious Diseases

- Ruminal tympany (Bloat)- It is the over-distension of the left flank either due to free gas or froth. This is generally encountered in "greedy feeders" when lush green pasture is available. Oral administration of sweet oil with turpentine oil or at times with formalin is advised.
- Rumen acidosis- Not applicable
- Intussusceptions- Not applicable
- Pregnancy toxæmia (Ketosis)- Not applicable

Poisoning

- Organochlorine compounds- This group includes DDT, BHC, lindane, aldrin, dieldrin, chlordane, toxaphane, methocyclor etc. which are used as pesticides on crops. Toxicity symptoms include increased excitability and irritability followed by muscle tremors, weakness, paralysis etc. Treatment consists of administering antidote, usually short-acting barbiturates.
- Organophosphorous compounds- This group consists of malathion, darathion, chlorathion, carbophenothion, demton, dasnon, dimethylparathion, trichlorphon, dioxalthion etc. Symptoms of toxicity are profuse salivation, muscle stiffness, dyspnoea with open mouth breathing, tremors. Treatment consists of administering antidote, usually atropine sulphate.



- Snake bite- Usually bitten on the scrotum or udder. The presence of hair may obscure the typical fang marks. Prolonged pain, muscular weakness, impaired vision, nausea and paralysis are generally exhibited along with symptoms of shock. If anti-venin is not available and the bite is located in an area where a tourniquet cannot be applied, excision of an area of skin and sub-cutaneous tissue can be life-saving

2 Poultry

Suggested contingency measures under DROUGHT event

a) Before the event

Shelter management

Optimum space should be provided. Orientation of shed (Long axis) should be in North – South. Plantation of tree around shed to provide cool environment. Provision of ad lib. Fresh water

Shortage of feed ingredients

Storage of feed

Drinking water

Manage clean drinking water. Storage facility should be made. Water quality should be checked before drinking to animal

Health and disease management

- Newcastle Disease- Regular vaccination - Broiler birds should be with RD vaccine (Lasota 'F' strain) at the age of 4-7 days through Intra-nasal or Intra-ocular route. Layer birds should be vaccinated with NDV vaccine at the age of 9-14 day, 4 weeks, 13-14 weeks in drinking water/eye drop. Then at the age of 17 week with NDV vaccine through Intra-muscular (IM) route
- Marek's disease Marek's disease- Birds should be vaccinated with Herpes virus turkey vaccine at the age of 1 day through Subcutaneous route.
- Fowl pox- Chick embryo adopted fowl pox vaccine at the age 6-8 weeks. It important for the layer and broiler birds.
- Drop in Egg Production or Quality- Not applicable
- Nervous Signs and Lameness- Not applicable
- Diarrhoea- Not applicable
- Upper Respiratory Diseases- Vaccination against the some of the viral diseases like Newcastle disease, influenza, infectious bronchitis, infectious laryngotracheitis which are also responsible for the respiratory symptoms can prevent this syndrome. Antifungal and antiparasitic drugs should be given.

Heat Wave

Plantation of tree around shed to provide cooler environment. Proper ventilation should be provided. Optimum space should be provided. Orientation of shed (Long axis) should be in East- West. Plantation of tree around shed to provide cool environment. Provision of ad lib. Fresh water. Manage green fodder and silage preparation. Height of roof should be minimum 220 – 240 cm. Roof of shed should be painted with white.

Cold Wave

Provide ad lib fresh water. Proper ventilation should be provided. Optimum space should be provided. Orientation of shed (Long axis) should be in North – South. Plantation of tree around shed to provide break cold wave. Provision of ad lib. Fresh water. Manage green fodder and silage preparation. Height of roof should be minimum 220 – 240 cm Roof of shed should be painted with Black Floor of shed should be Dry

b) During the event

Shelter management

Optimum space should be provided, there must not be overcrowded. Protect the animal from direct sun light. Try to provide them cool water. Proper ventilation should be maintained. Allow for grazing during night/early morning. Provision of ad lib. Fresh water

Shortage of feed ingredients

Provide non conventional feed, supplement anti oxidant and anti stress

Drinking water

Provide clean fresh and cold drinking water all the time. Water availability may increase by 20-50% depending upon feed quality and environmental temperature. Soft drinking water should be preferred. Add vit-C and other anti stress ingredients with water

Health and disease management

- Newcastle Disease- Vaccination and treatment of diseased one. Newcastle disease is the most important disease for poultry farmers around the world. This disease causes a large number of deaths in chickens and huge losses to farmers and the industry. Diseased birds should be slaughtered immediately. Consult Veterinarian.
- Marek's disease Marek's disease- It is one of the important diseases of poultry caused by virus. Mortality is very high and causes economic losses to the farmer and poultry industry.

- Fowl pox- It is a viral infection of chickens and turkeys characterized by proliferative lesions in the skin (Cutaneous form), it also affect the GI tract and respiratory tract (Diphtheritic form)
- Drop in Egg Production or Quality- There are many different types of organisms that can cause a drop in egg production or quality. These include: Bacteria (E. coli, Salmonella), Mycoplasma, Viruses (Newcastle disease, influenza, infectious bronchitis, infectious laryngotracheitis, avian encephalomyelitis, egg drop syndrome). The Parasites, lack of Nutrition and Stress factor also support the onset of this condition. Adding vitamins and minerals to the water or feed may help. Consult Veterinarian
- Nervous Signs and Lameness- Chickens lie down because they cannot stand up. They also walk with a limp or are reluctant to move. Nervous signs may include staring into the sky, pulling the head and neck over their backs, paralysis. Sores on the breast muscles from lying down
- Diarrhoea- The stool or droppings of the chickens are not firm but very loose, watery, not of the normal colour and may contain blood. This may cause the feathers of the vent to be soiled and caked together, Depression, reluctance to eat, drink and move about, poor growth and death. Use an antibiotic or coccidiostatic drug in the water that was recommended by the animal health technician or veterinarian in the water for 3 to 5 days. Stress preparations that contain electrolytes, vitamins and minerals can be added to the water
- Upper Respiratory Diseases- Not applicable

Heat Wave

Water sprinkling to animal. Prevent the animal from direct sunlight. Optimum space should be provided, there must not be overcrowded. Fan should be provided to make the body cool. Try to provide them cool drinking water all time

Proper ventilation should be maintained. Allow for grazing during night/early morning. Try to provide green fodder and silage. Stocking density should be less. Roof should be covered with tiles, paddy, dry leaves to protect from direct sun light

Cold Wave

Luke worm water should be provided at least 4-6 times a day. Prevent the animal from direct cold wave by closing the windows and doors. Optimum space should be provided, there must not be overcrowded. Proper ventilation should be maintained. Allow for grazing during sunny day time. Try to provide green fodder and silage. During extreme cold condition electric heater of wood fire heat should be provided. Try to make the environment inside and outside the shed dry. Gunny bags or blanket may be used to cover the body. Bedding material like paddy straw, Gunny Bag, Bhusa should be provided specially to young one shed.

c) After the event

Shelter management

Optimum space should be provided. Take care and fulfill the requirement of water along with proper nutrients. Take care of proper feeding as per requirement. Provision of ad lib. Fresh water

Shortage of feed ingredients

Not applicable

Drinking water

Provide adlib. Drinking water

Health and disease management

- Newcastle Disease- Disposal of dead birds
- Marek's disease Marek's disease- Disposal of dead birds
- Fowl pox- Disposal of dead birds
- Drop in Egg Production or Quality-Not applicable
- Nervous Signs and Lameness- A complete hygiene and disinfection programme should be planned together with the animal health technician or veterinarian. Antibiotics will only be effective against bacteria and can be used as recommended. If it is a viral disease, such as Newcastle disease, urgent steps have to be taken to prevent possible spread because it causes serious production losses
- Diarrhoea- Disposal of dead birds
- Upper Respiratory Diseases- There is many different types of organisms that can cause disease in the upper respiratory tract. These include: Mycoplasma Bacteria (E. coli, Pasteurella, Haemophilus), Viruses (Newcastle disease, influenza, infectious bronchitis, infectious laryngotracheitis), Parasites (mites and worms) And Fungi (Aspergillus). Cold stress is also one of the predisposing factors for the occurrence of respiratory problems. Use an antibiotic drug that was recommended by your animal health technician or veterinarian in the water for 3 to 5 days
- Stress preparations that contain electrolytes, vitamins and minerals can be added to the water

Heat Wave

Optimum space should be provided. Take care and fulfill the requirement of water along with proper nutrients. Take care of proper feeding as per requirement. Provision of ad lib. Fresh water

Cold Wave

Provide ad lib. Normal drinking water. Optimum space should be provided. Take care and fulfill the requirement of water along with proper nutrients. Take care of proper feeding as per requirement. Provision of ad lib. Fresh water

3 Fisheries

Suggested contingency measures under DROUGHT event

a) Before the event

Aquaculture

- Shallow water in ponds due to insufficient rains/inflow- Increase depth of pond, Repair dyke, outlet and inlet of pond; Prepare duck/pig house & stock pig @ 50-60,duck @ 450-500 no/ha if farmer involve in Integrated fish farming, Allow manure and urine directly in pond, Remove unwanted, predatory & old fishes and for this apply Mahua oil cake @ 2500kg/ha. Fixed net in outlet & inlet to prevent escaping of fish, Plough the pond and apply lime @ 250 kg/ha, Check the natural feed (plankton)@ 1.0 1.5 ml/50 ltr of water; otherwise apply organic manure, Stock yearling (stunted grow fish) @ 6,000-8,000 no/ha
- Impact of salt load build up in ponds / change in water quality- Prevent entry of polluted water or apply lime at inlet.

Heat wave and cold wave

- Changes in pond environment (water quality) - Increase depth of pond. Reduce application of organic manure and supplementary feeds
- Health and Disease management- Apply lime @ 50 kg/ha

b) During the event

Aquaculture

- Shallow water in ponds due to insufficient rains/inflow- Reduce the stocking density from 25000 fry (1inches size) to 10000-15000/ha, fingerling 6,000-8,000 no/ha. Check the availability of natural food, if it is not sufficient provide supplementary feed at fixed place, time, amount and ratio & if it is more greenish stop supplementary feed & manure, store manure in separate place for agricultural purpose. Check the growth & health status by regular netting, Apply lime @ 50kg/ha.
- Impact of salt load build up in ponds / change in water quality- Apply lime @ 50 kg/ha on every 15-30 days. Aerate the water as per need

Heat wave and cold wave

- Changes in pond environment (water quality) - Stop or reduce supplementary feed and manure, Remove bigger size fishes. Reduce/stop application of feed and fertilizer.
- Health and Disease management- Apply lime/salt as per need

c) After the event

Aquaculture


- Shallow water in ponds due to insufficient rains/inflow- Remove the bigger size fishes (0.5kg). In winter season fish reduces feed consumption so reduce supplementary feed, duck start egg laying so they should not allow before 9'oclock otherwise loss of egg is possible, pig may attain 50 – 60 kg so that can be sell out and again stock same no of piglets. Apply bleaching powder @ 10kg/ha at place of litter deposition.
- Impact of salt load build up in ponds / change in water quality- Apply lime as per need @ 50 kg/ha

Heat wave and cold wave

- Changes in pond environment (water quality) - Stop or reduce supplementary feed and manure, Remove bigger size fishes. Harvest the bigger fishes, Reduce/stop application of supplementary feed, Apply lime @ 50 kg/ha and potassium per magnet in perforated plastic ball- 5-10g in each ball
- Health and Disease management- Apply lime/salt as per need

WEST SINGHBHUM DISTRICT

Sl. No.	CONTENTS	Page No.
1.	District Agriculture profile and land use pattern	1
PART-I		
CONTINGENCY PLAN FOR KHARIF		
2.	Contingency plan for 2 weeks delay in monsoon arrival (onset in 4th week of June)	2-8
A1, A4. Upland A2, A5. Midland A3, A6. Lowland		
Contingency plan for 4 weeks delay in monsoon arrival (onset in 2nd week of July)		
B1, B4. Upland B2, B5. Midland B3, B6. Lowland		
Contingency plan for 6 weeks delay in monsoon arrival (onset in 6th week of July)		
C1, C4. Upland C2, C5. Midland C3, C6. Lowland		
PART-II		
3.	A. Contingency Plan for normal monsoon onset followed by 15-20 days dry spell	23-24
A1. Upland A2. Midland A3. Lowland		
4.	B. Contingency plan for mid season drought	24-26
Upland		
B1. At vegetative phase B2. At Flowering/Fruiting stage		
Midland		
B3. At vegetative phase B4. At Flowering/Fruiting stage		
Lowland		
B5. At vegetative phase B6. At Flowering/Fruiting stage		
5.	C. Contingency plan for Late season drought/Terminal drought (Early withdrawal of monsoon)	27
At fruiting/pre physiological maturity stage		
C1.1. Upland C1.2. Midland C1.3. Lowland		
PART-III		
6.	A. Unusual rains : Continuous high rainfall in a short span leading to water logging	28-30
Crop management Pest and disease management		
7.	B. Extreme weather events (Hail storm, Heat wave, Cold wave, Frost)	30-31
CONTINGENCY PLAN FOR RABI		
8.	1. Sowing window information	32
2. Contingency measures for field crops grown with residual moisture under rainfed condition		
2(A) Optimal residual moisture		
2A.1 Upland 2A.2 Midland 2A.3 Lowland		
2 (B) Less than optimal soil moisture (25 % less than normal-Deficit of 20-40 % rainfall)		
2B.1 Upland 2B.2 Midland 2B.3 Lowland		
CONTINGENCY STRATEGIES FOR LIVESTOCK, POULTRY AND FISHERIES		
9.	1. Livestock	36-42
a) Before the event b) During the event c) After the event		
2. Poultry		
a) Before the event b) During the event c) After the event		
3. Fisheries		
a) Before the event b) During the event c) After the event		





District Agriculture profile

Agro-Climatic Zone	AZ - 59		
Agro Ecological Sub Region (ICAR)	Eastern plateau (chotanagpur) And Eastern Ghats, Hot Subhumid Eco-Region (12.3)		
Agro-Climatic Zone (Planning Commission)	Eastern Plateau And Hills Region (VII)		
Agro Climatic Zone (NARP)	South Eastern Plateau Sub Zone - VI		
Meteorological Subdivision	8 th		
List all the districts falling under the NARP Zone (>50% area falling in the zone)	East Singhbhum, Saraikela-Kharsawan, West Singhbhum		
Geographic coordinates of district headquarters	Latitude		Altitude
	23° 24'36" N - 23°59' 12" N		85° 34' 51"E - 86°29'31" E 388m
Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/ RRTTS	Zonal Research Station (ZRS), Darisai, Birsa Agricultural University, Ranchi		
Mention the KVK located in the district with address	Krishi Vignan Kendra, Darisai, Vill-Kharia, PO. Barakhurshi Distt. East Singhbhum-832 304		
Name and address of the nearest Agromet Field Unit (AMFU, IMD) for agro-advisories in the Zone	ZRS, Darisai, Eas Singhbhum-832 304		

Land use pattern of the district (area: '000 ha)

Geographical area	Cultivable area	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable wasteland	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
567.769	241.194	142.336	4.998	3.712	37.502	4.998	55.335	87.783	77.694



CONTINGENCY PLAN FOR KHARIF

PART-I

A Monsoon/Weather Situation: 2 Weeks Delay (Onset: 4th Week of June) – Early Season Drought

A1. Major Farming Situation/Land Situation: Upland High rainfall, shallow light textured sandy soils

Normal Crop/cropping system	Upland Rice , Finger millet, Pigeonpea + Sorghum, Pigeonpea + Maize
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Rice crop	
<u>Sole crop</u>	
Pigeonpea, Finger millet, Sorghum, Soybean, Blackgram, Greengram, Sweet potato, lady's Finger	
Intercrop :	
Pigeonpea + Maize (1:1), Pigeonpea + Lady's finger (1:2)/Greengram (1:2), Maize+ creeper vegetables (Sponge Gourd) (1:2), Pigeonpea + Groundnut (1:2), Pigeonpea + Sorghum (1:1)	
<u>Horticulture Crop</u>	
Vegetable : Amaranthus leaf/ Coriander leaf/ Early Cabbage/ Cauliflower/ French bean/ Creepers_ Variety	
Pigeonpea- Birsa Arhar (200-220), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)	
Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149	
Sorghum- CSV 20-110-20, MP chari, CSV 1616	
Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335	
Birsa safed soybean 2 (105-110), RKS 18, RAUS 5	
Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain)	
Greengram- HUM 16, IPM-02-03-60-65, SML 668 (summer)	
Sweet potato-Shribhadra (80-90), Kalinga, Birsa sakarkand 1, Gauri	
Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika	
<u>Vegetable crops</u>	
Coriander- Pant haritima, Rajendra swati	
Cabbage- early- Golden acer, Early drumfead, Pride of India Late- Late drunhead, sabyay cabbage, 7 Ganga, Jamuna, Kaveri, Shri ganesh cabbage 8	
Cauliflower-Summer- Early kuwari, early- Kuwari, Pusa katki, Pusadipali, Early synthetic, Mid early- Pusa ketaki, Pusadipali, Pusa him jyoti, Pant subhra, Late- Maghi, Snowbowl 16, dania, Pusa Snowbowl, K Pusa Snowbowl, Hybrid- Himani, Swati, Endum early Pusa hybrid 1	
Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata	
Cucurbits-	
Bitter gourd- Arka harit, Pusa domausami,	
Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, Local	
Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white	
Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,	
Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi	
b) Agronomic measures	
<ul style="list-style-type: none">Summer deep ploughing with Mouldboard or discDobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.RD SpacingZero tillage practicesSeed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacingRDF and in case of Intercropping reduce 1/3rd dose for intercropWeed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flocloralin or Basalin and also for vegetablesBund construction for unbunded uplandBroadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetablesRidge and furrow method for moisture Conservation in cereals, pulses and vegetablesInter-cropping to meet the consequences of occasional Drought.Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.In case of phosphogypsum for soil application apply @ 120 kg/ha	



- Lime or dolomite application for pulses and oilseed @ 3-5 q/ha in furrow at the time of sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure

c) Remarks on Implementation

- Linkage with RKVY , ATMAs, and NFSM
- Vermicomposting through KVKS ATMAs and NHM
- Goatry and poultry rearing through KVKS, ATMAs and Veterinary Dept of. Govt. and BAU for livelihood support.
- Awareness about balanced use of fertilizers to increase their fertility, productivity and sustainability
- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology.
- Awareness for more and more use of organic manures, biopesticides for organic cultivation with IFS (eight components linkages)
- Upland- 15-20 % upland area should be covered with orchard

1. Mango based orchard-

Variety- Amrapali (30 June-5 July), Mallika (15-20 June regular bearer), Sunder langra(15-20 May)
Spacing- 5 m X 5m

i) Recommended package of Practices- Intercrops

- Mango + Papaya (Filler crop for two years) + Blackgram (rainy)/ Chickpea
- Mango + Custard apple (for 10 years and renovate or remove after 10 years) + Blackgram/Chickpea

Variety- Langra (15 June)/Bombay green(15 May)/ Himsagar (20-25 May irregular bearer),

Spacing- 10 m X 10m

ii) Recommended package of practices

- Mango + Guava(Up to 10 years as filler) + Papaya (Less than 3 years) + Blackgram-Chickpea/Lentil
- Mango + Lemon + Papaya + Rabi pulses/vegetables
- Mango + Custard apple + Papaya + Blackgram – Pea/Chickpea/Lentil/ Vegetables

2. Guava base orchard-

Variety- Arka Mridula, Pant Prabhat, Allahabad safeda, L 49

Spacing- 5m X 5m

Recommended package of practices- Intercrops

- Guava + Papaya (For 3 years) + Blackgram-Chickpea
- Guava + Custard apple + Blackgram/Soybean- Pea/Vegetables

3. Ber Based Orchard –

Variety- Banarsi, Karakka, Gola, Apple ber

Spacing- 5m X 5m

Recommended package of practices Intercrops

Ber + Custard apple + Sesame/Blackgram- Toria/Linseed/Safflower

N.B.- Cucurbits, beans or any creeper or climber vegetable should be avoided

- Field crops having height more than one meter should be avoided such as Pigenpea, Maize, Sorghum
- After 3-5 years when shading effects started shade loving crops like ginger, Turmeric, OI or leafy vegetables should be grown
- In citrus leaf minor and aphid susceptible crops should be avoided
- Aphid should be managed of mustard /toria taken in citrus orchard

4. Cashewnut based orchard for Kolhan region

5. Cassava should be grown for the requirement as feed for pig animals

- 6. Moringa should also be grown as fodder or vegetable purpose on upland main field bunds as shelter belt/ wind break. Every year pruning and thinning should be followed for bushy look.

A2. Major Farming Situation/Land Situation: Midland High rainfall, slightly deep sandy loam soils

Normal Crop/cropping system	Rice, Finger millet, Pigeonpea, Sorghum, Maize, Pigeonpea + Sorghum, Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures



a) Change in crop/cropping system

Don 2

DSR (Improved variety) Var.-IR 64 Drt 1, Shabhagi Dhan, BVD 111,Rice + Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var.-PAC 801, 807, PHB 71, Arize Tej (Gold)

Don3

Ridge and Furrow method or raise bed broad furrow: Replace Rice with cereal/ Pulse/ Vegetable

Cereal – Maize/ Sorghum/

Pulse- Pigeonpea + Lady's finger (1:2) / Soybean (1:2)/ Finger millet (1:1)

Vegetable-Amaranthus leaf/Coriander leaf/ Lady's Finger/ Cowpea/ Dolichos bean

Variety

Maize- Birsa makka (Vikash) 2 (75-80), Shaktiman 1(105-1010), KDMH, LG 32-81 –Yuvral gold (80-85), Malvia makka 2 (90), Kanchan(K 25) 100-110

Sorghum- CSV 20-110-20, MP cheri, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Asha (200-220), ICPH 2671 (200

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149

Vegetable crops

Coriander- Pant haritima, Rajendra swati

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Creeper- Birsa sweta, Swarna sweta, Swarn harit Dolichos bean-Swarna utkrist, Swarna rituwar

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INP
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P_2O_5 + 1 kg K_2O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P_2O_5 : K_2O (Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P_2O_5 + 40 K_2O /ha ((Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K_2O
- DSR-Use plastic drum seeder rice tools
- Follow RDF and
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/l.t. water, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /l.t water, Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/l.t. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/l.t. water, Mosaic- Methyl Demeton @ 1.5 ml/l.t. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/l.t. water

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKS, NHM and other State Govt. line departments are needed to be launched through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.



A3. Major Farming Situation/Land Situation: Lowland High rainfall, medium depth, heavy textured clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of Don 2 in Don 1 DSR(Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), Vardhan, MTU 1010, MTU 1001 Transplanting (Hybrid rice varieties) var.-Arize 6444 (Gold), Arize Tej (Gold), Akshay Dhan, PHB 71, 26P52, 25P25	

b) Agronomic Measures

- Staggered Nursery raising by MAT/ DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use Post emergence weedicide
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water Gundhi bug, leaf folder and BPH -Quinolhos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /ltr water

c) Remarks on Implementation

- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation.

A4. Major Farming Situation/Land Situation: Upland Less rainfall, shallow depth red light textured sandy soils

Normal Crop/cropping system	Upland Rice , Pigeonpea, Maize, Pigeonpea + Maize,
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Rice crop	
<u>Sole crop</u>	
Pigeonpea, Sorghum, Finger millet, Soybean, Blackgram, Greengram, Lady's finger	
<u>Intercrop</u>	
Pigeonpea + Maize (1:1), Pigeonpea + Lady's finger (1:2)/Greengram (1:2), Maize+ creeper vegetables (Sponge gourd) (1:2), Pigeonpea + Sorghum (1:1)	
<u>Horticulture Crop</u>	
Vegetable :Radish/Early Cabbage/ Cauliflower/ French bean / Cowpea	
<u>Variety</u>	
Pigeonpea- Birsa Arhar (200-220), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200) Sorghum- CSV 20-110-20, MP chari, CSV 1616	
Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149	
Soybean- R 518 (110), JS 9752 (100), Birsa soybean 1 black(120-125), JS 335	
Birsa safed soybean 2 (105-110), RKS 18, RAUS 5	
Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain)	
Greengram- HUM 16, IPM-02-03-60-65, SML 668 (summer)	
Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika	



Vegetable crops

Radish- Pusa chetki, Pusa deshi, Jaunpur/ Pusa himani, Pusa roshni

Cabbage- early- Golden acer, Early drumfead, Pride of India

Cauliflower-Early- Kuwari, Pusa katki, Pusa him jyoti, Pant subhra, Hybrid- Himani, Swati, Endum early Pusa hybrid 1

Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati

b) Agronomic measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbundled upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite application for pulses and oilseed @ 3-5 q/ha in furrow at the time of sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Dry seeding with 15% to 20% higher seed rate. Seed treatment with Rhizobium in pulses. Maximum use of organic manure

c) Remarks on Implementation

- Linkage with RKVY , ATMAS, and NFSM
- Vermi composting through KVKS ATMAS and NHM
- Goatry and poultry rearing through KVKS, ATMAS and Veterinary Dept of. Govt. and BAU for livelihood support.
- Awareness about balanced use of fertilizers to increase their fertility, productivity and sustainability
- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology.
- Awareness for more and more use of organic manures, biopesticides for organic cultivation with IFS (eight components linkages)
- Upland- 15-20 % upland area should be covered with orchard

1. Mango based orchard-

Variety- Amrapali (30 June-5 July), Mallika (15-20 June regular bearer), Sunder langra(15-20 May)
Spacing- 5 m X 5m

i) Recommended package of Practices- Intercrops

- a) Mango + Papaya (Filler crop for two years) + Blackgram (rainy)/ Chickpea
- b) Mango + Custard apple (for 10 years and renovate or remove after 10 years) + Blackgram/Chickpea

Variety- Langra (15 June)/Bombay green(15 May)/ Himsagar (20-25 May irregular bearer),

Spacing- 10 m X 10m

ii) Recommended package of practices

- a) Mango + Guava(Up to 10 years as filler) + Papaya (Less than 3 years) + Blackgram-Chickpea/Lentil
- b) Mango + Lemon + Papaya + Rabi pulses/vegetables
- c) Mango + Custard apple + Papaya + Blackgram – Pea/Chickpea/Lentil/ Vegetables

2. Guava base orchard-

Variety- Arka Mridula, Pant Prabhat, Allahabad safeda, L 49
Spacing- 5m X 5m

Recommended package of practices- Intercrops



- a) Guava + Papaya (For 3 years) + Blackgram-Chickpea
 b) Guava + Custard apple + Blackgram/Soybean- Pea/Vegetables
3. Ber Based Orchard –
 Variety- Banarsi, Karakka, Gola, Apple ber
 Spacing- 5m X 5m
 Recommended package of practices Intercrops
 Ber + Custard apple + Sesame/Blackgram- Toria/Linseed/Safflower
- N.B.- Cucurbits, beans or any creeper or climber vegetable should be avoided
- Field crops having height more than one meter should be avoided such as Pigenpea, Maize, Sorghum
 - After 3-5 years when shading effects started shade loving crops like ginger, Turmeric, Ol or leafy vegetables should be grown
 - In citrus leaf minor and aphid susceptible crops should be avoided
 - Aphid should be managed of mustard /toria taken in citrus orchard
4. Cashewnut based orchard for Kolhan region
 5. Cassava should be grown for the requirement as feed for pig animals
 6. Moringa should also be grown as fodder or vegetable purpose on upland main field bunds as shelter belt/wind break. Every year pruning and thinning should be followed for bushy look.

A5. Major Farming Situation/Land Situation: Midland Less rainfall, medium depth, light to medium textured sandy loam soils.

Normal Crop/cropping system	Rice, Pigeonpea, Maize, Pigeonpea + Maize , Pigeonpea + Sorghum
Suggested Contingency measures	
a) Change in crop/cropping system	
<u>Don 2</u>	
DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Rice + Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR	
<i>Transplanting (Hybrid Rice varieties)</i> Var.-PAC 801, 807, PHB 71, Arize Tej (Gold)	
<u>Don 3</u>	
Ridge and Furrow method or raise bed broad furrow:_Replace Rice with cereal/ Pulse/ Vegetable	
Cereal – Maize/ Sorghum	
Pulse- Pigeonpea + Lady's finger (1:2) /Soybean (1:2)/ Finger millet (1:1)	
Vegetable-Radish/ Lady's Finger/ Cowpea/ Dolichos bean	
<u>Variety</u>	
Maize- Birsa makka (Vikash) 2 (75-80), Shaktiman 1(105-1010), KDMH, LG 32-81 –Yuvral gold (80-85), Malvia makka 2 (90), Kanchan(K 25) 100-110	
Sorghum- CSV 20-110-20, MP cheri, CSV 1616	
Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Asha (200-220), ICPH 2671 (200	
Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149	
<u>Vegetable crops</u>	
Radish- Pusa deshi, Kashi hansi, Jaunpur/ Pusa himani, Japanese white,	
Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika	
Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati	
Dolichos bean-Swarna utkrist, Swarna rituwar	

b) Agronomic measures

- Follow RDF,INP
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P_2O_5 + 1 kg K_2O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P_2O_5 : K_2O (Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P_2O_5 + 40 K_2O /ha ((Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K_2O
- DSR-Use plastic drum seeder rice tools
- Follow RDF and
- INPM
- Use of post weedicide



- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /lt water, Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKS, NHM and other State Govt. line departments are needed to be launched through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

A6. Major Farming Situation/Land Situation: Lowland medium depth, heavy textured clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of Don 2 in Don 1 DSR (Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), Vardhan, MTU 1010, MTU 1001 Transplanting (Hybrid rice varieties) var.-Arize 6444 (Gold), Arize Tej (Gold), Akshay Dhan, PHB 71, 26P52, 25P25	
b) Agronomic Measures	
<ul style="list-style-type: none"> Staggered Nursery raising by MAT/ DAPOG method Follow community based nursery raising Follow RDF,INPM Use Post emergence weedicide Use early to mid early duration of rice variety. Nursery management- 1 kg N + 1 kg P₂O₅ + 1 kg K₂O for 100 m² Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice Topdressing above mentioned dose 10-15 days after sowing In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering. DSR-Use plastic drum seeder rice tools Use of post weedicide Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /lt water 	

c) Remarks on Implementation

- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation.



B. Monsoon/Weather Situation: 4 Weeks Delay (Onset: 2nd Week of July) – Early Season Drought

B1. Major Farming Situation/Land Situation: Upland High rainfall, shallow light textured sandy soils	
Normal Crop/cropping system	Upland Rice , Finger millet , Pigeonpea + Sorghum, Pigeonpea + Maize
Suggested Contingency measures	
a) Change in crop/cropping system	
<p>Discard Rice Crop</p> <p>Sole crop</p> <p>Pigeonpea, Sorghum, Finger millet, Soybean, Blackgram, Greengram, Sweet potato, lady's Finger</p> <p>Intercrop :</p> <p>Pigeonpea/Maize + Lady's finger (1:2), Pigeonpea + Maize (1:1) , Pigeonpea + Blackgram (1:2)/ Greengram (1:2)</p> <p>Pigeonpea + Sorghum (1:1),</p> <p>Horticulture Crop-</p> <p>Vegetable- Tomato/ Brinjal/ Chili/ Amaranthus leaf/ Lobia/ French bean/ Creepers</p> <p>Fodder Crop</p> <p>Sorghum, Maize, Blackgram, Greengram</p> <p>Verity</p> <p>Pigeonpea- Birsa Arhar (200-220), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)</p> <p>Finger millet- BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149</p> <p>Soybean- JS 335</p> <p>Birsa safed soybean 2 (105-110), RKS 18, RAUS 5</p> <p>Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75)</p> <p>Greengram- HUM 16, IPM-02-03-60-65, SML 668</p> <p>Sweet potato-Shribhadra (80-90), Kalinga, Birsa sakarkand 1, Gauri</p> <p>Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika</p> <p>Vegetable crops</p> <p>Tomato- Hybrid- Swarn sampada, Swarn samridhi, Pusa hybrid 1 Suraksha</p> <p>Brinjal- Pusa purple long, Pusa purple round, Pusa purple cluster, Mukta keshi, Banaras giant,</p> <p>Chili- Spices- Andhra jyoti, Pusasadabahar, Yellow wonder, Bharat</p> <p>Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati</p> <p>French bean- Bushy- Pant anupma, Swarna priya, Arka Komal,</p> <p>Cucurbits</p> <p>Bitter gourd- Arka harit, Pusa domausami,</p> <p>Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, local</p> <p>Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white</p> <p>Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,</p> <p>Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi</p> <p>Fodder crop</p> <p>Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid.</p> <p>Maize- African tall, JS-1006 and Vijaya composite.</p> <p>Blackgram- PU 19/31/35 (70-75)</p> <p>Greengram- HUM 16, IPM-02-03-60-65</p>	
b) Agronomic Measures	
<ul style="list-style-type: none"> • Summer deep ploughing with Mouldboard or disc • Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations. • RD Spacing • Zero tillage practices • Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing • RDF and in case of Intercropping reduce 1/3rd dose for intercrop • Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables • Bund construction for unbunded upland • Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables • Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables • Inter-cropping to meet the consequences of occasional Drought. • Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed. • In case of phosphogypsum for soil application apply @ 120 kg/ha • Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing. 	



- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lit. water; vegetables- Nursery management- Application of carbofuran 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- Linkage with RKVY, ATMAS and NFSM
- Vermicomposting Awareness through KVKS, ATMAS and NHM
- Backyard Goatry and poultry rearing Awareness campaign through KVKS, ATMAS and Veterinary Dept of Govt. and BAU for livelihood support.
- A special programme is needed to be launched in such areas on priority basis to motivate the farmers to adopt improved technology for stress management through ATMAS, KVKS, Govt Dept., NGOs
- Campaign for Awareness of crop-weather insurance to meet the losses due to drought/cyclone like weather vagaries.

B2. Major Farming Situation/Land Situation: Midland High rainfall, slightly deep sandy loam soils

Normal Crop/cropping system	Rice, Finger millet, Pigeonpea, Sorghum, Maize, Pigeonpea + Sorghum, Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807

Don 3

Ridge and Furrow method or raise bed broad furrow: Replace rice with Cereal/Pulses/ vegetable/ Fodder crop
Cereal – Maize/Sorghum

Pulses- Pigeonpea+ Lady's finger(1:1)/Blackgram (1:2)/Soybean (1:2)

Vegetables- Lady's Finger/ Cowpea/ Dolichos bean/

Fodder Crop

Rice bean (Moth bean)/ Maize / Cowpea

Variety

Maize- Birsa makka (Vikash) 2 (75-80), Shaktiman 1(105-1010), KDMH, LG 32-81 –Yuvral gold (80-85), Malvia makka 2 (90), Kanchan(K 25) 100-110

Sorghum- CSV 20-110-20, MP cheri, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Asha (200-220), ICPH 2671 (200

Blackgram- Birsa urd 1 (75-80), WBU 109 (70-75), Uttara (75-80)

Soybean- R 518 (110), JS 9752 (100), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5



Vegetable crops

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati

Dolichos bean-Swarna utkrist, Swarna rituwar

Fodder crop

Maize- African tall, JS-1006 and Vijaya composite.

Cowpea-EC-4216, UPC-287

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility.
- Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/ltr. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/ltr. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/ltr. water, Mosaic- Methyl Demeton @ 1.5 ml/ltr. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/ltr. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 ltr. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Tricoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/ltr. water.
- Rice pest and disease management -Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /ltr water. Termite- Methyl parathion dust @ 25 kg/ha



c) Remarks on Implementation

- A campaign through RKVY , ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be awarded through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

B3. Major Farming Situation/Land Situation: Lowland High rainfall, medium depth, heavy textured clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2</u> in <u>Don 1 DSR(Improved Rice variety)</u> Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), Vardhan, MTU 1001 Transplanting (Hybrid rice Varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), Akshay Dhan, PHB 71, 26P52, 25P25	
b) Agronomic Measures	
<ul style="list-style-type: none"> • Staggered Nursery raising by MAT/ DAPOG method • Follow community based nursery raising • Follow RDF,INPM • Use Post emergence weedicide • Use early to mid early duration of rice variety. • Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² • Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice • Topdressing above mentioned dose 10-15 days after sowing • In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling • Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm • Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering. • DSR-Use plastic drum seeder rice tools • Use of post weedicide • Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water, Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /lit water 	
c) Remarks on Implementation	
<ul style="list-style-type: none"> • Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme • Supply of Plastic drum seeder through line departments • Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept. • Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon. • Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates • Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation 	

B4. Major Farming Situation/Land Situation: Upland Less rainfall, shallow depth, light textured sandy and acidic soil.

Normal Crop/cropping system	Upland Ric, Pigeonpea, Maize, Pigeonpea + Maize (Mixed)
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Rice crop <u>Solecrop</u> Pigeonpea, Blackgram, Greengram, Soybean, Groundnut, <u>Intercrop</u> Pigeonpea + Groundnut(1:2), Pigeonpea + Maize 1:1), Pigeonpea + Sorghum (1:1) <u>Horticulture Crop</u> Vegetable- Radish/Early Cabbage/ Cauliflower/ French bean / Cowpea (Lobia)	



Fodder Crop

Cowpea/Sorghum/ Maize/ Blackgram

Variety

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), ICPH 2671 (200)

Blackgram- Birsa urd 1 (75-80), PU 35 (70-75), Uttara (75-80 small grain)

Greengram- HUM 16, IPM-02-03-60-65, SML 668

Soybean- JS 9752 (100), JS 335, Birsa safed soybean 2 (105-110), RAUS 5

Groundnut- Birsa mungfali 3, 4, Girnar 3

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Shaktiman 1(105-1010), LG 32-81 –Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Vegetable crops

Radish- Pusa chetki (summer), Pusa deshi, Kashi hansi, Jaunpur/ Pusa himani, Japanese white, Pusa roshni

Coriander- Pant haritima, Rajendra swati

Cabbage- Golden acer, Early drumhead

Cauliflower-Pusa katki, Pusadipali, Early synthetic, Pusa him jyoti, Pant subhra, Endum early Pusa hybrid 1

Frenchbean- Bushy- Pant anupma, Swarna priya, Arka Komal, Stringless, Creeper- Kentucky wonder, Birsa priya, Swarna lata

Cowpea- bushy- CP 4, Arka garima, Pusa komal

Fodder Crop

Cowpea- UPC-5286, GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, HC-136, HC-171, PSC-1, Sorghum Sudan hybrid

Maize- African tall, JS-1006 and Vijaya composite

Blackgram- Uttara (75-80)

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing
- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathypor or Pendimithilin@ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbundled upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofurran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages



- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/ltr. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/ltr. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/ltr. water, Mosaic- Methyl Demeton @ 1.5 ml/ltr. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/ltr. water, Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/ltr. water or Cartap hydrochloride @ 2 gm/ltr. water, hairy caterpillar –Quinolphos 1.5ml / ltr. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 ltr; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/ltr. water

c) Remarks on Implementation

- Linkage with RKVY , ATMAs and NFSM
- Vermicomposting Awareness through KVKS, ATMAs and NHM
- Backyard Goatry and poultry rearing Awareness campaign through KVKS, ATMAs and Veterinary Dept of. Govt. and BAU for livelihood support.
- A special programme is needed to be launched in such areas on priority basis to motivate the farmers to adopt improved technology for stress management through ATMAs, KVKS, Govt Dept., NGOs
- Campaign for Awareness of crop-weather insurance to meet the losses due to drought/cyclone like weather vagaries.

B5. Major Farming Situation/Land Situation: Midland Less rainfall, shallow depth , light textured sandy and acidic soil

Normal Crop/cropping system	Rice, Pigeonpea, Maize Pigeonpea + Maize, Pigeonpea + Sorghum, Rice-vegetable
-----------------------------	---

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 111, Dhaincha/Sunhemp (Green manuring/ Brown manuring) in DSR

Transplanting (Hybrid Rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), PAC 801, 807, 27P31, DRRH 2

Don 3

Ridge and Furrow method or raised broad bed furrow along the slope :_Replace rice with Cereal/Pulses/ vegetable/ Fodder crop

Cereal – Maize/Sorghum

Pulses- Pigeonpea+ Lady's Finger/Blackgram/Soybean/ Groundnut/Maize

Vegetables- Lady's Finger/ Cowpea/ Dolichos bean/

Fodder Crop -

Rice bean (Moth bean)/ Maize / Cowpea/ Sweet Sorghum

Variety

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Pusa HM 9(AQH 9), KDMH, P3544, LG 32-81 – Yuvral gold (80-85), Malvia makka 2 (90), Vivek hybrid 9 (80)

Sorghum- CSV 20-110-20, MP chari, CSV 1616

Pigeonpea- Birsa Arhar (200-220), Malvia 13 (240-250), Narendra Arhar 1 and 2 (240-250), Asha (200-220), ICPH 2671 (200)

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Blackgram- Birsa urd 1 (75-80), PU 31 (70-75), WBU 109 (70-75)

Soybean- Birsa soybean 1 black(120-125), JS 335

Birsa safed soybean 2 (105-110), RKS 18, RAUS 5

Groundnut- Birsa mungfali 3, 4, Girnar 3

Maize- Birsa makka (Vikash) 2 (75-80), Malvia makka 2 (90), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)

Vegetable crops

Lady's finger- Pusa A 4, Sarika

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati, Swarn harit

Dolichos bean-Swarna utkrist, Swarna rituwar

Fodder crop

Maize- African tall, JS-1006 and Vijaya composite.

Cowpea-EC-4216, UPC-287, UPC-5286, GFC-1, GFC-2 and GFC-4

Sorghum-PC-1, PC-6, PC-23, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

b) Agronomic Measures

- Summer deep ploughing with Mouldboard or disc
- Dobha construction for In-situ rain water conservation Line sowing in upland rice areas through suitable seeding devices is required to be made popularized for desired plant population. This will facilitate to control weeds and also to carry out intercultural operations.
- RD Spacing



- Zero tillage practices
- Seed rate – Sole- full quantity and in case of Intercropping reduce seed rate by 30-40 % according to spacing
- RDF and in case of Intercropping reduce 1/3rd dose for intercrop
- Weed control (Maize- Atrazine as pre-emergence, Pulses- pre-emergence Imizathyper or Pendimithilin @ 1 kg a.i./ha, Soybean- Flucloralin or Basalin and also for vegetables
- Bund construction for unbunded upland
- Broadcast Well rotten FYM along with 1/4th N + Full basal application of P, K of recommended dose for all crops and for vigorous seedling growth of vegetables
- Ridge and furrow method for moisture Conservation in cereals, pulses and vegetables
- Inter-cropping to meet the consequences of occasional Drought.
- Follow RDF for all upland crops and add Sulphur @ 20kg/ha soil application for pulses and oilseed.
- In case of phosphogypsum for soil application apply @ 120 kg/ha
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- In vegetable nursery apply carbofuran @ 3gm/m² or phorate 10 G @ 1gm/ m² or neem cake @ 50 kg/ha
- Follow recommended seed rate
- Treat the leguminous seed in the sequence of FIR (Bavistin @ 2gm/kg, Imidacloprid@ 3 ml or Chlorpyriphos @ 5ml/kg, Rhizobium 500 gm/ha , PSB @ 500 gm/ha and for non leguminous treat seed with Fungicide + Insecticide + soil application Azotobacter @ 2kg /ha
- Foliar application of Urea 2% solution + lime in lady's finger
- Application of required fungicide and insecticide in case of population count more than the ETL or as prophylactic measure
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility.
- Apply Borax @ 10 kg/ha
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges and furrows
- Cultivate vegetables like Brinjal Tomato, Cucurbits, Lady's finger, Chili, Coriander leaf, Amaranthus leaf, Oel, Arvi, Dolichos bean, Cole crop, French bean Cowpea etc.
- Gap filling and resowing should be done If mortality is more than 50 per cent and even if necessary replace the crops with short duration high yielding low water requiring crops like: Greengram, Blackgram, Horsegram, Niger, Cowpea Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour.
- Weed control by applying pre-emergence 5-6 DAS (Pendimithilin) or Post-emergence 18-28 DAS (Bispyribac).
- Irrigate only at critical stages
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit. water; Pigeonpea-leaf folder- Methyl demoton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water water, Mosaic- Methyl Demoton @ 1.5 ml/lit. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water, Groundnut- Tikka and leaf minor- Hexaconazole(Cartap) @ 1ml/lit. water or Cartap hydrochloride @ 2 gm/lit. water, hairy caterpiller -Quinolphos 1.5ml /lit. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water.
- Rice pest and disease management -Gundhi bug, leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha. Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %. Blast- Beam or Tricyclazole @ 0.6 gm /lt water. Termite- Methyl parathion dust @ 25 kg/ha

c) Remarks on Implementation

- A campaign through RKVY , ATMAS, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be awared through different district, block, panchayat and village level programme.
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

B6. Major Farming Situation/Land Situation: Lowland Less rainfall, medium depth, heavy textured clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	

a) Change in crop/cropping system

Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of Don 2 in Don 1 DSR(Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), MTU-1001, MTU 1010, Transplanting (Hybrid rice varieties) Var.–Arize 6444 (Gold), Arize Tej (Gold), Akshay Dhan, PHB 71, 26P52, 25P25 Vegetable

Cucurbits-

Bitter gourd- Arka harit, Pusa domausami,

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, Local

Sponge gourd- Pusa chikni, Pusa supriya, Rajendra nema, Long green,Long white

Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,

Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi

b) Agronomic Measures

- Staggered Nursery raising by MAT/ DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use Post emergence weedicide
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water, Gundhi bug,leaf folder and BPH -Quinolophos 25 EC(Ekalux) dust @ 25 kg/ha, Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %, Blast- Beam or Tricyclazole @ 0.6 gm /lt water

c) Remarks on Implementation

- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds through Lamps one month prior to the arrival of monsoon.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation

C. Monsoon/Weather Situation: 6 Weeks Delay (Onset: 6th Week of July) – Early Season Drought

C1. Major Farming Situation/Land Situation: Upland High rainfall, shallow light textured sandy soils

Normal Crop/cropping system	Upland Rice,Finger millet, Pigeonpea+ Sorghum, Pigeonpea + Maize
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Discard Rice Crop

Sole crop

Ridge and Furrow method)-Niger, Horsegram, Cowpea, Finger millet, Blackgram, Greengram

Intercrop

Ridge and Furrow method:Pigeonpea + Lady's finger (1:2), Pigeonpea + Maize (1:1)

Pigeonpea + Blackgram(1:2)/Greengram (1:2), Maize + lady's finger (1:2)

Horticulture crop

Ridge and Furrow method:Vegetable –Lady's Finger, Ridge Gourd, Bottle Gourd, Cowpea

Fodder Crop

Sadabahar/Chara badam/ Anjan grass/ Hybrid napier/ Cactus/ Sweet Sorghum



Variety

Niger- Birsa niger 1, 2 and 3 (95-105), Puja 1 (90), VLG 19

Horsegram- Birsa kulthi1 (90-95)

Cowpea-rainy – Birsa sweta(80-90), Swarn sweta(80-90), Swarn harit (80-90)

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), WBU 109 (70-75), Uttara (75-80 small grain)

Greengram- HUM 16, IPM-02-03-60-65, SML 668 (summer)

Maize- Birsa makka (Vikash) 2 (75-80), Pusa HM 9(AQH 9), KDMH, P3544, Malvia makka 2 (90), Vivek hybrid 9 (80)

Vegetable crops

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Cucurbits

Ridge gourd- Swarn manjari, Swarn uphar, Swarn baha, Pusa nasdar, Satputia,

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtur long green, local

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Birsa sweta, Swarna sweta, Swarn harit

b) Agronomic Measures

- Top dressing of urea and DAP after receipt of the rain for all crops
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10-15 kg/ha
- Replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Soybean, Sesame, Horsegram , Niger, Cowpea, Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour
- Follow mulch after cultural operations to control the weeds in vegetables.
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges
- Foliar application of 2 % DAP or 0.5 to 1 % potassium chloride (KCl) +0.3 % Boric acid or 2% urea at pre-flowering and flowering stage in pulses and vegetables
- 2 % DAP spray for pulses.
- Use antitranspirants :
 - Stomatal closure (Growth hormones like ABA, Ethrel, TIBA, succinic acid, ascorbic acid and Cycocel (CCC) Reflectant (Calcium bicarbonate, Lime water) Thin film (Hexadecanol (Higher alcohols) Cetyl alcohol, Methanol
- Acidic soils should be reclaimed by application of soil ameliorants.
- Follow integrated pest management.
- Weed control by applying pre-emergence 5-6 DAS (Pendimethalin) or Post-emergence 18-28 DAS (Bispyribac)
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/lit; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/lit. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/lit. water, Mosaic- Methyl Demeton @ 1.5 ml/lit; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/lit. water; Finger millet- Leaf/ finger/neck and collar blast- Tricyclazole @ 6 gm/10 lt. water; vegetables- Nursery management- Application of carbofuran 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/lit. water

c) Remarks on Implementation

- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology for stress management through ATMAS, KVKS, Govt Dept., NGOs and others. Soybean and fodder crops may be promoted.
- Promote Knowingness about climate resilient agriculture at district, block, panchayat and village level through involvement of KVKS, ATMAS, DAO, NGOs and other State Agril. Govt line departments.
- Awareness of mechanization and Supply of Mouldboard and disc chisel/harrow through govt. scheme on subsidized way.
- Promote for double their income by curtailing cost of cultivation by introduction of early duration crops variety.
- Campaign for Awareness programme about crop-weather insurance

C2. Major Farming Situation/Land Situation: Midland High rainfall, slightly deep sandy loam soils

Normal Crop/cropping system	Rice, Finger millet, Pigeonpea, Sorghum, Maize, Pigeonpea + Sorghum,Pigeonpea + Maize
-----------------------------	---

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 110, 111 Vardhan

Transplanting (Hybrid Rice varieties) var.-CR Dhan 40, DRRH 2, Arize Tej (Gold), 27P31, PAC 801, PHB 71

Don3

Ridge and Furrow method or raised broad bed furrow: Replace Rice with Pulses and cereals/ Vegetable/ Fodder crop

Pulse and cereal - Blackgram, Greengram, Maize, Finger millet,

Vegetable- Lady's finger/Tomato,/ Brinjal, creepers/Chili

Fodder Crop

Blackgram/ Greengram/ Cowpea/ Sorghum/ Maize

Variety

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), Uttara (75-80)

Greengram-IPM-02-03-60-65, SML 668

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Pusa HM 9(AQH 9), LG 32-81 –Yuvral gold (80-85), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149

Vegetable crops

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Tomato-Hybrid- Swarn sampada, Swarn samridhi, Pusa hybrid 1 Suraksha

Brinjal- Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6

Cucurbits

Bitter gourd- Arka harit, Pusa domausami,

Bottle gourd- Arka bahar, Pusa samar, Coimbatore long green

Sponge gourd- Rajendra nema, Long green, Long white

Ridge gourd- Swarn uphar, Swarn baha, Pusa nasdar,

Red Pumpkin- CO 1, CO 2, Arka chandan, Arka suryamukhi

Chili- Spices- Andhra Jayoti, Pusa Sadabahar, NP 46, Jwala, KA 2, California wonder, Chinese giant, Yellow wonder, Bharat

Fodder crop

Blackgram- WBU 109 (70-75), Uttara (75-80)

Greengram- HUM 16, IPM-02-03-60-65,

Cowpea- EC-4216, UPC-287

Maize- African tall, JS-1006 and Vijaya composite

Sorghum- PC-1, PC-6, HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF, INPM
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water; Gundhi bug, leaf folder and BPH -Quinalphos 25 EC(Ekalux) dust @ 25 kg/ha; False smut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /ltr water; Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/ltr. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/ltr. water, Mosaic- Methyl Demeton @ 1.5 ml/ltr. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/ltr. water

c) Remarks on Implementation

- Campaign for awareness improved technology through RKVY , ATMAS, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be at different district, block, panchayat and village level
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme



- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency crops through Lamps within one months.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

C3. Major Farming Situation/Land Situation: Lowland High rainfall, medium depth, heavy textured clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of Don 2 in Don 1 DSR(Improved Rice variety) Var.- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), Vardhan, MTU 1001, MTU 1010 Transplanting (Hybrid rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), Akshay Dhan, PHB 71, 26P52, 25P25, DRRH 2, KRH 2	
Fodder crop- In case of fallow (Late heavy rainfall) Dallis grass/ Para grass/ Arundino Grass	
b) Agronomic Measures	
<ul style="list-style-type: none"> • Staggered Nursery raising by MAT/ DAPOG method • Follow community based nursery raising • Follow RDF,INPM • Use Post emergence weedicide • Use early to mid early duration of rice variety. • Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² • Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice • Topdressing 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² at 10-15 days after sowing • In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling • Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm • Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering. • DSR-Use plastic drum seeder rice tools • Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water; Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /ltr water 	
c) Remarks on Implementation	
<ul style="list-style-type: none"> • Awareness programme of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme • Supply of Plastic drum seeder through line departments in case of DSR • Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept. • Supply of improved and hybrid seeds of contingency mid early rice varieties through Lamps within one month • Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates • Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation • Contingency technology Awareness programme through KVKS, ATMAS, NGOs and DAOs • Achieve maximum fallow area in case of late drought and suggest to go for cultivation of early duration rabi and fodder crops. 	

C4. Major Farming Situation/Land Situation: Upland Less rainfall, shallow depth red light textured sandy soils

Normal Crop/cropping system	Upland Rice, Pigeonpea, Maize, Pigeonpea+ Maize
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard rice crop	
<u>Sole crop</u>	
Raised bed or ridge and furrow method-Niger, Horsegram, Sorghum, Pigeonpea, Blackgram, Greengram, Lady's Finger Intercrop	
Raised bed or ridge and furrow method- Pigeonpea+ Lady's finger (1:2), Lady's finger + Blackgram (1:2)	
Maize + Lady's finger (1:2), Pigeonpea + Maize (1:1), Pigeonpea + Greengram (1:2)	
Maize+ Blackgram (1:2)/Greengram (1:2)	



Horticulture crop

Vegetable –Lady's Finger, Bottle Gourd, Cowpea (lobia)

Fodder Crop

Maize/Chara badam/ Thin napier/ Cactus/ Sweet Sorghum

Variety

Niger- Birsa niger 1, 2 and 3 (95-105), Puja 1 (90), VLG 19

Horsegram- Birsa kulthi1 (90-95

Cowpea-rainy – Birsa sweta(80-90), Swarn sweta(80-90), Swarn harit (80-90)

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149

Blackgram- Birsa urd 1 (75-80), PU

Greengram- HUM 16, IPM-02-03-60-65, SML 668 (summer)

Maize- Birsa makka (Vikash) 2 (75-80), Pusa HM 9(AQH 9), KDMH, P3544, Malvia makka 2 (90), Vivek hybrid 9 (80)

Vegetable crops

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Bottle gourd- Arka bahar, Pusa samar, Pusa Naveen, PusaMeghdoot, Coimbtor long green, Local

Cowpea- bushy- CP 4, Arka garima, Pusa komal, Pusa barsati Birsa sweta, Swarna sweta, Swarn harit

Fodder crop

Maize- African tall, JS-1006 and Vijaya composite

Sorghum-HC-171, PSC-1, Pant Chari-5, Pant Chari-6 and Sorghum Sudan hybrid

b) Agronomic Measures

- Top dressing of urea and DAP after receipt of the rain for all crops
- Lime or dolomite (3-5 q/ha) sulphur and phosphogypsum (30 Kg/ha) with compost application few days before sowing.
- Leguminous/pulse crops may be included in the cropping system in order to improve the soil fertility. Apply Borax @ 10-15 kg/ha
- Replace the crops with short duration high yielding low water requiring crops like : Greengram, Blackgram, Soybean, Sesame, Horsegram , Niger, Cowpea, Fodder maize, Fodder cowpea, Fodder sorghum, Fodder pearl millet, Sweet potato, Gundli, Guarfalli after receiving the downpour
- Follow mulch after cultural operations to control the weeds in vegetables.
- For in-situ moisture conservation in vegetables, 15-20 DAS follow intercultural operations by making ridges
- Foliar application of 2 % DAP or 0.5 to 1 % potassium chloride (KCl) +0.3 % Boric acid or 2% urea at pre-flowering and flowering stage in pulses and vegetables
- 2 % DAP spray for pulses.
- Use antitranspirants :
Stomatal closure (Growth hormones like ABA, Ethrel, TIBA, succinic acid, ascorbic acid and Cycocel (CCC)
Reflectant (Calcium bicarbonate, Lime water) Thin film (Hexadecanol (Higher alcohols) Cetyl alcohol, Methanol
- Acidic soils should be reclaimed by application of soil ameliorants.
- Follow integrated pest management.
- Weed control by applying pre-emergence 5-6 DAS (Pendimethalin) or Post-emergence 18-28 DAS (Bispyribac)
- Pest and disease management- Maize- Stem borer Monocrotophos @ 1ml/ltr. water; Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/ltr. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/ltr., Mosaic- Methyl Demeton @ 1.5 ml/ltr. water; Soybean- Cercospora leaf spot- Indofil M 45 1 ml/ltr. water; Finger millet- Leaf/finger/neck and collar blast- Tricyclazole @ 6 gm/10 ltr. water; vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/ltr. water

c) Remarks on Implementation

- A special programme is needed to be launched in such areas to motivate the farmers to adopt improved technology for stress management through ATMAS, KVKS, Govt Dept., NGOs and others. Soybean and fodder crops may be promoted.
- Promote Knowingness about climate resilient agriculture at district, block, panchayat and village level through involvement of KVKS, ATMAS, DAO, NGOs and other State Agril. Govt line departments.
- Awareness of mechanization and Supply of Mouldboard and disc chisel/harrow through govt. scheme on subsidized way.
- Promote for double their income by curtailing cost of cultivation by introduction of early duration crops variety.
- Campaign for Awareness programme about crop-weather insurance



C5. Major Farming Situation/Land Situation: Midland Less rainfall, medium depth, light to medium textured sandy loam soils.

Normal Crop/cropping system	Rice, Pigeonpea, Maize, Pigeonpea + Maize, Pigeonpea + Sorghum
-----------------------------	--

Suggested Contingency measures

a) Change in crop/cropping system

Don 2

DSR (Improved rice variety) Var.-Shabhagi Dhan, Abhishek, IR 64-Drt 1, BVD 110, 111
Transplanting (Hybrid Rice varieties) Var.-Arize Tej (Gold), PHB 71, 27P52, 27P31, 25P25

Don 3

Ridge and Furrow method or raise bed broad furrow along the slope :_Replace Rice with pulse/ vegetable/
Fodder crop

Pulse and cereal/ Blackgram/ Greengram/ Maize/ Finger millet

Vegetable

Lady's finger/Tomato,/ Brinjal/Chili/

Fodder Crop

Cowpea/ Sorghum/ Maize/ Rice bean (Moth bean)/ Sudan grass(SC)/Thin Napier Late August-September-Berseem (MC)/ Oat (MC)/ Rye grass

Variety

Blackgram- Birsa urd 1 (75-80), PU 19/31/35 (70-75), Uttara (75-80)

Greengram-IPM-02-03-60-65, SML 668

Maize- Birsa makka (Vikash) 2 (75-80), HQPM 1 (90-100), Pusa HM 9(AQH 9), LG 32-81 –Yuvral gold (80-85), Kanchan(K 25) 100-110 , Vivek hybrid 9 (80)

Finger millet- A 404, BM 2, BM 3 (BBM 10), GPU 28, 67, VL 149

Vegetable crops

Lady's finger- Pusa A 4, Arka anamika, Varsa uphar, Hybrid- Sonal, Sarika

Tomato-Hybrid- Swarn sampada, Swarn samridhi, Pusa hybrid 1 Suraksha

Brinjal- Swarn pratibha, Swarn mani, Swarn shayamali, hybrid-Swarn shakti , Vijay, Swarna sampada 6

Chili- Spices- Andhra jyoti, Pusasadabahar, NP 46, Jwala, KA 2, California wonder, Chinese giant, Yellow wonder, Bharat

Fodder crop

Cowpea-EC-4216, UPC-287

Sorghum-PC-1, PC-6, HC-171, PSC-1, Pant Chari-5

Maize- African tall, JS-1006 and Vijaya composite

b) Agronomic Measures

- Staggered Nursery raising by MAT/DAPOG method
- Follow community based nursery raising
- Follow RDF,INPM
- Use early to mid early duration of rice variety.
- Nursery management- 1 kg N + 1kg P_2O_5 + 1 kg K_2O for 100 m²
- Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice
- Topdressing above mentioned dose 10-15 days after sowing
- In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling
- Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm
- Fertilizer dose- 80:40:20 kg/ha N : P_2O_5 : K_2O (Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P_2O_5 + 40 K_2O /ha ((Basal 1/2 N + full dose P_2O_5 + 2/3rd K_2O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K_2O at the time of flowering.
- DSR-Use plastic drum seeder rice tools
- INPM
- Use of post weedicide
- Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/ltr. water; Gundhi bug,leaf folder and BPH -Quinolophos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /ltr water; Termite- Methyl parathion dust @ 25 kg/ha
- Pest and disease management- Pigeonpea-leaf folder- Methyl demeton @ 1.5 ml/ltr. water; Blackgram and Greengram- Leaf minor- Monocrotophos @ 1ml/ltr. water, Mosaic- Methyl Demeton @ 1.5 ml/ltr. water; S vegetables- Nursery management- Application of carbofuron 3G @ 3 gm/m² before 10 days of transplanting followed by application of Trichoderma along with half rotten cow dung @ 1 kg Trichoderma in 100 kg cow dung (20 days staying period required for cow dung treated with Trichoderma), rainy potato-Ridomyl MZ @ 1-2 gm/ltr. water



c) Remarks on Implementation

- Campaign for awareness improved technology through RKVY , ATMAs, NFSM, KVKS, NHM programme and other State Govt. line departments are needed to be at different district, block, panchayat and village level
- Awareness of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments
- Awareness about climate smart agriculture through Birsa Agricultural University and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency crops through Lamps within one months.
- Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone situation.

C6. Major Farming Situation/Land Situation: Lowland Less rainfall, medium depth, heavy textured clay loam soil

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change in crop/cropping system	
Discard Long duration variety (Swarna , BPT 5204 and Rajshree) with Medium duration rice variety of <u>Don 2 in Don 1</u> DSR(Improved Rice variety) Var.-- Shabhagi Dhan, IR 64-Drt 1, Abhishek (120 days), MTU 1010, BVD 203, BVS 1	
Transplanting (Hybrid rice varieties) Var.-Arize 6444 (Gold), Arize Tej (Gold), Akshay Dhan, PHB 71, 26P52, 25P25, DRRH 2, KRH 2	
Fodder crop -In case of fallow (Late heavy rainfall)-Para grass/Dallis grass	
b) Agronomic Measures	
<ul style="list-style-type: none"> • Staggered Nursery raising by MAT/ DAPOG method • Follow community based nursery raising • Follow RDF,INPM • Use Post emergence weedicide • Use early to mid early duration of rice variety. • Nursery management- 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² • Seed rate 80-100 kg/ha for improved rice variety and 15 kg/ha for hybrid rice • Topdressing 1 kg N + 1kg P₂O₅ + 1 kg K₂O for 100 m² at 10-15 days after sowing • In nursery- Carbofuron 3G @300 gm/100 m² 10 days before uprooting of seedling • Spacing DSR- 20 cm row for PDS and for transplanting 20-25 X 15-25 cm • Fertilizer dose- 80:40:20 kg/ha N : P₂O₅ : K₂O (Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O and rest before flowering and for hybrid 120-150 kg N + 60 Kg P₂O₅ + 40 K₂O/ha ((Basal 1/2 N + full dose P₂O₅ + 2/3rd K₂O ; 1/4th N at 20-25 DAS; 1/4th N at 45 DAS ; 1/3rd K₂O at the time of flowering. • DSR-Use plastic drum seeder rice tools • Rice pest and disease management- Stem borer- Carbofuron 3 G 12 kg/acre , Gall midge- Monocrotophos @ 1ml/lit. water; Gundhi bug,leaf folder and BPH -Quinolphos 25 EC(Ekalux) dust @ 25 kg/ha; Falsesmut- 1st spraying at time of flowering and 2nd 10 days after 1st spraying of Propiconazole @ 0.1 % or Nativo @ 0.04 %; Blast- Beam or Tricyclazole @ 0.6 gm /lt water 	

c) Remarks on Implementation

- Awareness programme of MAT/DAPOG method of raising nursery and nursery management through different district, block, panchayat and village level programme
- Supply of Plastic drum seeder through line departments in case of DSR
- Awareness about climate smart agriculture through Birsa Agricultural university and state Govt. Ag. Dept.
- Supply of improved and hybrid seeds of contingency mid early rice varieties through Lamps within one month Linkages of farmers through different line departments and BAU for providing alertness, warning and weather vagaries updates
- Campaign for more and more crop-weather insurance to meet losses in case of drought/cyclone upside-down situation
- Contingency technology Awareness programme through KVKS's, ATMAs, NGO's and DAO's
- Achieve maximum fallow area in case of late drought and suggest to go for cultivation of early duration rabi and fodder crops.

PART-II

A. Monsoon/Weather Situation: Normal onset followed by 15-20 days dry spell after sowing
(Early Season Drought-Normal onset)

A1. Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils	
Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
Suggested Contingency measures	
a) Change management Cultivate drought tolerant promising non paddy crops like pigeonpea, blackgram, greengram, rice bean, finger millet, guar, sesame, soyabean, sorghum, pearl millet, sweet potato, castor and vegetables like radish, tomato,, brinjal, creeper bean, chili, lady's finger wherever possible in place of upland rice	
<ul style="list-style-type: none"> • Maximum use of organic manures for early seedling vigour along with RDF (N:P₂O₅:K₂O) • Recommend to resow with subsequent rains for better plant stand. • When damage is Less than 30 per cent then go for Gap filling in all upland crops • When damage is More than 50 per cent then go resowing in all upland crops • Removing excess plants where are over crowded,to reduce crop stand to conserve soil moisture • Water spraying during evening and early morning 	
b) Soil nutrient & moisture conservation measures <ul style="list-style-type: none"> • Avoid top dressing of Urea during dry spell and wait till downpour • Go for in-situ moisture conservation • One hand weeding followed by hoeing and simultaneous eartingup after 20 DAS is highly recommended in all upland crops. 	
c) Remarks on Implementation Awareness for Construction of rain water harvesting structures for recycling of water during dry spell like DOVAS through SHG or on subsidized basis through State Govt.schemes.	
A2. Major Farming Situation/Land Situation: MID LAND Sandy loam soils	
Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change management <u>Don 2</u> <ul style="list-style-type: none"> • If possible, go for staggered raising of nursery in rice crop • If possible, raise community nursery of rice at a reliable water source to save time for further delay. • In case, if rice population is less than 40-50 percent, gap filled by retransplanting the rice crop and for more than 50 per cent mortality use fresh seeding for fresh transplanting. • Follow gap filling by removing seedlings from profuse tillers to have a uniform distribution of same aged plants • For termite and disease management in nursery spray Indofil M 45 and Chlorpyriphos @ 0.2 per cent • life saving irrigation • DSR on receipt of rain by using Paddy drum seeder or • High yielding varieties- follow transplanting while, Improved varieties – follow DSR • In case of DSR- Use sprouted seeds in plastic drum seeder with increased seed rate by 20-25 per cent for good crop stand • Late transplanted rice during early season drought results in the occurrence of sheath rot and grain discoloration diseases. • Follow pre emergence and post emergence weedicide to disturb/check the crop-weed competition for nutrient • Provide life saving and protective irrigation to over aged seedling in nursery through dovas (harvested rain water). Also, take care of blast disease in nursery and avoid using urea in nursery. • Strengthen the bunds to check the drainage holes and seepage loss in transplanted and direct sown medium land rice regularly 	
<u>Don3</u> <ul style="list-style-type: none"> • Follow raised broad bed furrow or Ridge and furrow method for Maize/ Pigeonpea/ Lady's finger/ Blackgram/ Soybean • Adopt surface mulching with crop residue or tree loppings of <i>Glyricidia</i> wherever possible. If farm waste is not available, use blade to form a thin layer of soil mulch to avoid cracks • Life saving irrigation • In case of transplanting of over aged seedling (35-45 days), increase number of seedling per hill (5-6 seedling/hill) 	



b) Soil nutrient & moisture conservation measures

- Dry seeding of rice with application of pre and post emergence weedicide in over aged seedlings (>25 DOS)
- Split application of Urea fertilizer
- Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells

c) Remarks on Implementation

Awareness for Construction of rain water harvesting structures for recycling of water during dry spell like DOVAS through SHG or on subsidized basis through State Govt.schemes.

A3. Major Farming Situation/Land Situation: LOW LAND Sandy clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Change management	
<ul style="list-style-type: none"> • If possible, go for staggered nursery raising in rice crop • If possible, raise community nursery of rice at a reliable water source to save time for further delay. • In case, if rice population is less than 40-50 percent, gap filled by retransplanting the rice crop and for more than 50 per cent mortality use fresh seeding for fresh transplanting. • Follow gap filling by removing seedlings from profuse tillers to have a uniform distribution of same aged plants • Prefer mid early rice variety instead of late variety • Use pre and post emergence weedicide • Over aged seedling should be top cut and treat the seedlings root by Dursban/Chlorpyriphos @ 5 ml per lt water and transplant immediately after treated seedlings with 2 per cent Urea solution • In case of transplanting over aged seedling (35-45 days), increase number of seedling per hill (5-6 seedling/hill) • In fallow land go for cultivation of mid early duration rice variety through DSR @ 70-80 Kg/ha 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Split application of Urea fertilizer • Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells 	
c) Remarks on Implementation	
Awareness for Construction of Ponds, check dam through water shed management & MNREGA scheme through SHG or on subsidized basis through State Govt.schemes.	

B. Monsoon/Weather Situation: Mid season drought (long dry spell, consecutive 2 weeks rainless (<2.5 mm) period

B1. At vegetative phase

B1.1 Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils

Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
Suggested Contingency measures	
a) Change management	
<ul style="list-style-type: none"> • Use organic mulches such as tree leaves, straw and other available crop residue to conserve soil moisture • Avoid top dressing of fertilizers till sufficient moisture is available in soil • Use reflectant or antitranspirant like Kaolin @ 3-5 kg/100 lt or • In pulses, at weekly interval foliar spray of KCl @ 0.5- 1 % + 100 ppm Boric acid followed by foliar spraying of 2 percent urea during evening time • Spray wax emulser • Manual weeding followed by hoeing for germinating weeds. • For termite and leaf folder control spraying or drenching of Chlorpyriphos @ 2ml/lt water and for all pulses and cereals. • For leaf folder control in Maize (Stem borer) and Pigeonpea apply Carbofuran 3 G @ 12 Kg/acre or Phorate 10 G @ 4 kg/acre or Quinolphos @ 1 ml/lt water in Maize for leaf folder • Also, spray @ 20/40/60 ppm CaCl₂ in pulses • Vegetables- Foliar spray of water with 2 per cent KCl + 100 ppm Boron • Tomato- Foliar spray of CaCl₂ @ 20/40/60 ppm • Gap filling may be done with pigeonpea to maintain adequate plant stand. • For termites in pigeonpea, maize and other standing cereal crops which can be controlled by soil drenching with chlorpyriphos 20 EC @ 2 ml/lt water or by adding Chlorpyriphos 1.5% dust @ 8– 10 kg/ha or Carbofuran 3G @ 12 kg or Phorate 10 G @ 4 kg.acre before final land preparation and also control Gallmidge 	



- In green and blackgram, cowpea, bean and lady's finger the spread of YMV by insect vector may increase. Hence, to control insect vectors spray Dimethoate @ 1ml/ lt or Imidacloprid 4 ml/10 lt twice at 10 days interval
- In groundnut crop termites and white grub incidence is expected to be more. Methods suggested in rice may be followed to reduce the pest infestation.
- Incidence of leaf miner in groundnut may increase which can be managed by spraying Monocrotophos 36 SL or Triazophos 40 EC @ 1 ml/lt. water twice at fortnight intervals.
- Under dry condition incidence of mites is expected to be more in vegetable crops which can be brought down by spraying of dicofol @ 2 ml/lt water.
- Early and mid season drought favours disease like brown spot of rice, bacterial wilt of brinjal and other vegetables

b) Soil nutrient & moisture conservation measures

- Foliar spraying of DAP @ 2 per cent along with Boric acid @ 0.3 per cent. Also, spray urea @ 1 per cent
- Provide micro- irrigation with drip for wide spaced crops such as chillies and vegetables and Sprinklers for groundnut, maize and vegetables wherever ground/ surface water is available.
- Go for life saving and protective irrigation from constructed dovas.

c) Remarks on Implementation

Promote construction of Rain water harvesting structure watershed programme and MNREGA

B2. At flowering/ fruiting stage

B2.1. Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils

Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
-----------------------------	--

Suggested Contingency measures

a) Change management

- Maize- Harvest it for fodder use
- Pulses- and vegetables- At 2-3 days interval spraying of water followed by 2 per cent KCl + 100 ppm Boron during evening time is recommended.
- In case of groundnut maturing in the month of September which can be harvested after providing light irrigation through dohas to loose the soil.

b) Soil nutrient & moisture conservation measures

Go for life saving and protective irrigation from constructed DOVAS.

c) Remarks on Implementation

Promote for the construction of Rain water harvesting structure watershed programme and MNREGA

B3 At vegetative phase

B3.1. Major Farming Situation/Land Situation: MID LAND Sandy loam soils

Normal Crop/cropping system	Rice
-----------------------------	------

Suggested Contingency measures

a) Crop management

Don 2

- Manual weeding followed by hoeing for germinating weeds
- Take care of mealy bug and termite attack which are more prevalent in dry weather .
- Top dressing should be followed only after receipt of rain .
- No urea should be top dressed until receipt of rainfall in rice crop.
- For BPH, dusting field bunds and around with Carbaryl (Savin)4% or malathion 5% @ 10 – 12 kg/acre

Don 3

- One manual weeding for germinating weeds
- Apply 4 Kg N/acre in sorghum and oilseed crops soon after receipt of rains.
- In pigeonpea, if the drought affected plants to recoup with the revival of the rains, spray 2 to 3% urea after the foliage is wetted with the rains.
- Foliar application of Sulphur @ 1ppm to mitigate the stress condition in oilseed is necessary after receipt of rainfall
- Apply post emergence weedicide for controlling weeds in oilseed (Groundnut) to undisturb the pegging process.
- During 40-45 DAS, if there is a severe moisture stress, thinning may be done in kharif sorghum and pearl millet.

b) Soil nutrient & moisture conservation measures

- Foliar spray of KCl or ZNSO₄ @ 2 per cent
- Foliar spray of 2% KNO₃ or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21- 21 to supplement nutrition during dry spells
- Life saving irrigation through dovas,well, ponds, check dams and bora bandh

c) Remarks on Implementation

Promote for the construction of Rain water harvesting structure watershed programme and MNREGA



B4. At flowering/ fruiting stage

B4.1 Major Farming Situation/Land Situation: MID LAND Sandy loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<p>Don 2 and Don 3</p> <ul style="list-style-type: none"> • Life saving irrigation with harvested water • Spray of urea @ 1-2 percent • Drought condition during the month of August-September onwards shall result in severe incidence of foliar blast and brown spot diseases in rice. It is advised to spray Tricyclazole (Tilt) @ 6 g/ 10 lt. water or Casugamycin @ or Kasu B @ 2 ml/lt. water twice at 10 days intervals during drought period. 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Foliar spray of KCl or $ZNSO_4$ @ 2 per cent • Foliar spray of 2% KNO_3 or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells • Life saving irrigation through dovas, well, ponds, check dams and bora bandh 	
c) Remarks on Implementation	
Promote for the construction of Rain water harvesting structure watershed programme and MNREGA	

B5. At vegetative phase

B5.1 Major Farming Situation/Land Situation: LOW LAND Sandy clay loam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<ul style="list-style-type: none"> • Foliar spray of 2 per cent KCL followed by 1-2 per cent Urea. • Weeding should be done • Drought makes the crop vulnerable to sheath rot and sheath blight diseases. Maintenance of field sanitation followed by twice spraying at 10 days interval with validamycin 2-3 ml/lt water or Tricyclazole @ 6g/10 lt or carbendazim @ 2 g/lt water are advised. • Life saving irrigation 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Foliar spray of Foliar spray of Urea @ 2 per cent • Foliar spray of 2% KNO_3 or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells • Life saving irrigation through dovas, well, ponds, check dams and bora bandh 	
c) Remarks on Implementation	
Awareness for Construction of Ponds, check dam through water shed management & MNREGA scheme through SHG or on subsidized basis through State Govt.schemes.	

B6. At flowering/ fruiting stage

B6.1. Major Farming Situation/Land Situation: LOW LAND Sandy clayloam soils

Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management	
<ul style="list-style-type: none"> • Drought condition during flowering and fruiting and onwards shall result in severe incidence of foliar blast and brown spot diseases in rice. It is advised to spray Tricyclazole (Tilt) @ 6 g/ 10 lt or Casugamycin @ or Kasu B @ 2 ml/lt. water twice at 10 days intervals during drought period. • Life saving irrigation • During drought, attack of gundhi bug shall be more. Apply Quinolphos or Monocrotophos @ 1-2 ml per lt. water. 	
b) Soil nutrient & moisture conservation measures	
<ul style="list-style-type: none"> • Weeding and foliar spray of urea @ 2 per cent • Foliar spray of 2% KNO_3 or 2% urea solution or 1% water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 to supplement nutrition during dry spells • Life saving irrigation through dovas, well, ponds, check dams and bora bandh 	
c) Remarks on Implementation	
Promote for the construction of Rain water harvesting structure watershed programme and MNREGA	



Monsoon/Weather Situation: Terminal drought (Early withdrawal of monsoon)

C1. At fruiting/pre physiological maturity stage	
C1.1 Major Farming Situation/Land Situation: UP LAND Sandy red lateritic soils	
Normal Crop/cropping system	Upland rice, Maize, Vegetables, Cowpea, Pigeonpea + Groundnut, Maize + Pigeonpea, Bhindi + Maize
Suggested Contingency measures	
a) Change management <ul style="list-style-type: none"> Life saving irrigation to vegetables through stored moisture from constructed DOVA If not possible to make survival harvest it for fodder use 	
b) Rabi Crop planning <ul style="list-style-type: none"> Cultivation of Niger, Horsegram, Toria, linseed as relay/paira cropping In case of availability of irrigation, go for cultivation of early Potato and pea (early Arkel group) Prepare kachha check dam or Bora Bandh for Water conservation Mid early variety of radish cultivation is recommended 	
(c) Remarks on Implementation Promote for the construction of Farm ponds through watershed management programme and MNREGA	
C1.2. Major Farming Situation/Land Situation: MID LAND Sandy loam soils	
Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management <u>Don 2</u> <ul style="list-style-type: none"> At milking , soft and dough stage spray KCL @ 2 per cent In case of gundhi bug attack found more than ETL(>2 gundhibug /m²), spray Chlorpyriphos dust or Monocrotophos @ 1 ml/lit. water If possible go for life saving irrigation Late season drought generally results in outbreak of foliar, node, collar or neck blast of rice depending on the stage of crop. 	
<u>Don 3</u> Instead of grain purpose crops like sorghum, pearmillet, maize, cowpea, black and greengram that can be harvested for fodder use	
b) Rabi crop planning <ul style="list-style-type: none"> Ensure for all inputs required for rabi season in advance. In case of failure of kharif crops prefer sowing of pre rabi catch crops like, Toria, Niger, Horsegram, blackgram, sesame linseed in uplands to medium lands 	
c) Remarks on Implementation Promote construction of Rain water harvesting structure watershed programme and MNREGA	
C1.3. Major Farming Situation/Land Situation: LOW LAND Sandy loam soils	
Normal Crop/cropping system	Rice
Suggested Contingency measures	
a) Crop management <ul style="list-style-type: none"> Life saving irrigation. The land should be tilled properly in case <i>kharif</i> crop fails sow <i>rabi</i> crops like safflower, pigeonpea in sept-Oct (Short duration) Spray KCL @ 2 per cent followed by Urea @ 2 per cent Mid early rice crop may be harvested at Physiological maturity Cultivate vegetables like Tomato, Brinjal, Capsicum, Shimla mirch, Broccoli, Cabbage and Cauliflower, green pea and potato as per suitability near and around tributaries 	
b) Rabi crop planning Prefere early sowing of wheat, Mustard, Chickpea,linseed and lentil as sole or intercrop Wheat + Chickpea (4:2) Wheat+ Mustard (4:3)	
c) Remarks on Implementation Promote construction of Rain water harvesting structure watershed programme and MNREGA	

PART-III

A. Unusual rains: Continuous high rainfall in a short span leading to water logging

Suggested Contingency measures
a) Crop management
Pigeonpea /Sorghum/Pearl millet
Vegetative stage- Prefer ridge and furrow method of sowing. Ensure for proper drainage through channel. Collect runoff water in Dovas for further use.
Flowering stage- Ensure for proper drainage through channel. Collect runoff water in Dovas for further use.
Crop maturity stage- No such situation at the time of maturity
Post harvest- After Sun drying follow grading and storing
Blackgram and other pulses/Oilseeds
Vegetative stage- Follow Ridge and furrow sowing
Ensure for proper drainage through channel
Collect runoff water in Dovas for further use
Avoid application of fertilizer
Flowering stage- Ensure for proper drainage through channel
Collect runoff water in Dovas for further use
Avoid application of fertilizer
Prophylactic measure for jassid and YMV
Crop maturity stage-
Post harvest-
Rice
Vegetative stage- Safe disposal of excess water from rice field. Bund repairing and strengthen. Application of insecticides in the afternoon hours is preferred seeing the weather condition or after spraying weather should remain rain free for at least 4-5 hrs. Retransplant to maintain plant population in case of mortality more than 50 %
In partially damaged crop, allow to withstand upright. Flood occurs due to heavy storm in mid and lowland which when recedes probability of occurrence of swarming caterpillar on field bunds and around of rice crop is more. So, when it crosses the Economic Threshold Limit (ETL) i.e., one larva / hill then spray the crop with Chlorpyriphos/ Triazophos/ Profenophos @ 2 ml/lt water or dust the crop with Quinalphos @ 1.5% D @ 10kg/ acre. To prevent migration of larvae from one field to other, bunds should be heavily dusted with the dust formulation mentioned above. In partially ponded field, rice caseworm and in general leaf folder attack is expected. If 1-2 cases or folded leaves/hill is seen spray the crop with Monocrotophos / Chlorpyriphos @ 1 ml/lt water or with Cartap Hydrochloride 50 SP / Fipronil 5 SP @ 200 g/acre. Rain storms during <i>kharif</i> may result in severe occurrence of bacterial leaf streak and bacterial blight in rice. It is advised to spray the crop immediately after every rain spell with streptocycline @ 1g/10 lt water or plantomycin @ 1g/lt water or bacterinol @ 2g/lt. water. Control snail occurrence by Acaricide
Flowering stage- Safe disposal of excess water from rice field. Bund repairing and strengthen. Avoid application of fertilizer. Flood occurs due to heavy storm in mid and lowland which when recedes probability of occurrence of swarming caterpillar, BPH and cut worm on field bunds and around of rice crop is more. So, when it crosses the Economic Threshold Limit (ETL) i.e., one larva / hill then spray Chlorpyriphos/ Triazophos/ Profenophos @ 2 ml/lt water or dust the crop with Quinalphos @ 1.5% D @ 10kg/ acre. To prevent migration of larvae from one field to other, bunds should be heavily dusted with the dust formulation mentioned above. In partially ponded field, rice caseworm and in general leaf folder attack is expected. If 1-2 cases or folded leaves/hill is seen spray the crop with Monocrotophos / Chlorpyriphos @ 1 ml/lt water or with Cartap Hydrochloride 50 SP / Fipronil 5 SP @ 200 g/acre. Unusual and heavy rain during <i>kharif</i> may result in severe occurrence of bacterial leaf streak and bacterial blight in rice. It is advised to spray the crop immediately after every rain spell with streptocycline @ 1g/10 lt water or plantomycin @ 1g/lt water or bacterinol @ 2g/lt. water. Control snail occurrence by Acaricide.
Crop maturity stage- Provide drainage for fast removal of water from the field to favour harvesting
Post harvest- Protect the grain from rain and store it after sun drying for 2-3 days
Maize
Vegetative stage- Prefer ridge and furrow method of sowing. Ensure for proper drainage through channel. Earthingup after downpour. At Knee stage apply thimate 10 G @ 4-6 grains in whirl
Flowering stage- Ensure for proper drainage through channel. At flowering and silking stage for ant attack apply dust on silks @ 0.5 g / cob
Crop maturity stage- Provide drainage for fast removal of water from the field to favour harvesting
Post harvest- Protect grains from rain and store it after sun drying for 2-3 days



Horticulture

Vegetative stage- Prefer ridge and furrow method for sowing and proper drainage. Ensure for proper drainage through water ways. Collect runoff water in Dovas for further use. Soil drenching Carbofuran 3G @ 3 g/lt water against insects. In case of web formation with leaves apply (Nuvan)DDVP @ 1 ml/lit water as a fumigant
Flowering stage- Apply hormone to prevent flower drop. Ensure for proper drainage. Take precaution against wilting and fruit rot. In Tomato and Brinjal-drenching Bavisting @ 2 ml/lit. water + Streptocycline @ 1-2 g/lit water. In Cauliflower -In case of Incidence of collar rot -Spraying of Saaf (Metalaxyl + Mancozeb) @ 2 g/lit water solution. Drainage of excess water. In Lady's finger- YVMV- Spray insecticide followed by fungicide. Soil drenching Carbofuran 3G @ 3 g/lit water against insects. In case of web formation with leaves apply (Nuvan) DDVP @ 1 ml/lit water as a fumigant

Crop maturity stage- Take precaution against wilting and fruit rot. For wilting– Soil drenching with Bavistin @ 2 ml/lit. water + Streptocycline @ 1-2 g/lit water. In YMVM- Insecticide followed by fungicid

Post harvest- Immediate harvest and safe disposal of produce

Vegetables- (Cucurbits,/ Tomato/ Brinjal/ Cauliflower/ Cabbage/ Lady's finger/Dolichos bean/Amaranthus leaf/ Cariander leaf/Radish)

Vegetative stage- Sowing on ridge and drainage through furrow. Prophylactic measures against pest and diseases. Damaged twigs and leaves may be removed and follow fungicide spraying and stacking

Flowering stage- Apply hormone to prevent flower drop. Ensure for proper drainage. Take precaution againsts wilting and fruit rot. In Tomato and Brinjal-drenching Bavisting @ 2 ml/lit. water

+ Streptocycline @ 1-2 g/lit water. In Cauliflower -In case of Incidence of collar rot -Spraying of Saaf (Metalaxyl + Mancozeb) @ 2 g/lit water solution. Drainage of excess water. In Lady's finger- YVMV- Spray insecticide followed by fungicide. Provide support through stacking

Crop maturity stage- Take precaution against wilting and fruit rot. In Wilting– Soil drenching with Bavistin @ 2 ml/lit + Streptocycline @ 1-2 g/lit water. In YMVM- Insecticide followed by fungicide

Provide support through stacking.

Post harvest- Immediate harvest and sell produce safely in the market

b) Pest and disease management

Rice

Vegetative stage- Sheath blight- Hexaconazole @ 1ml/lit wate. Blast- Tricyclazole @ 6 g/10 lt water

Flowering stage- Sheath blight- Hexaconazole @ 1ml/lit water. Blast- Tricyclazole @ 6 g/10 lt water. Falsesmut-Nativo @ 4g/10 lt water

Crop maturity stage- False Smut - Control- Nativo @ 4g/10 lt water or Propiconazole + Tricyclazole 52.5 SE @ 1ml/lit water. In case of grain discolourness (Grain blast). Spray Tricyclazole @ 6 ml / 10 liter water

Post harvest- Store grains after proper sun drying to minimize the incidence of stored grain pest

Maize

Vegetative stage- Stem borer Control- Carbofuron 3 G @ 12 Kg/acre or Phorate 10G@ 4 kg/acre

Flowering stage- Sheath blight Control- Hexaconazole1-2 ml/lit water

Vegetables- (Cucurbits,/ Tomato/ Brinjal/ Cauliflower/ Cabbage/ Lady's finger/Dolichos bean/Amaranthus leaf/ Cariander leaf/Radish)

Vegetative stage- Before sowing apply in soil, Carbofuran 3 G @2-3 g/m². Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt streptocycline or 2-3 g/lit plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lit water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lit. water against downy mildew diseases of cucurbits.

Flowering stage- Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt streptocycline or 2-3 g/lit plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lit water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lit. water against downy mildew diseases of cucurbits. YVM Control- Carbofuran 3G @ 3 or Phorate 10 G @ 1 g/m² followed by any fungicide

Crop maturity stage- Stop spraying 1 week before harvesting

Post harvest- Harvest and sell produce in the market

French bean-

Vegetative stage- Rust disease Control- Mancozeb 2g/ lt water. Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt. water streptocycline or 2-3 g/lit. water plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lit water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lit. water against downy mildew diseases of cucurbits.



Flowering stage- Take care of pod borer and aphid attack. Post heavy rainfall affected areas where vegetable is grown are likely to be affected by bacterial wilt, leaf spot and canker disease. Therefore, total package of practices starting from planting need to be followed as given below. Seedling root dip for 30 minutes in 1g/10 lt. water streptocycline or 2-3 g/lt. water plantomycin. Perform soil drenching to the base of the plants with a solution of carbendazim @ 2g/lt water and streptocycline @ 1g/10 lt. water at 10 DAP, 25 DAP and 40 DAP coinciding with intercultural operations. Spray Ridomil MZ @ 1.5 ml/lt. water against downy mildew diseases of cucurbits.
Crop maturity stage- Stop spraying 1 week before harvesting
Post harvest- Harvest and sell produce in the market

B. Extreme Weather Events

Suggested Contingency measures

Hail storm

Seedling / nursery stage- Vegetable nursery should be raised in poly house or make proper arrangement of low height Polly tunnels in open area or cover with plastic sheet or thatching should be done

Vegetative stage- In vegetables-Remove damaged parts immediately and apply insecticide followed by fungicide as prophylactic measures. Follow fertilization through foliar as well as broadcasting

Reproductive stage- n vegetables- Remove damaged parts immediately and apply insecticide followed by fungicide as prophylactic measures. Follow fertilization through foliar as well as broadcasting for proper fruiting

At harvest- Safely sell in the market after grading for immediate returns

Heat Wave

Wheat Chickpea/pea

Seedling / nursery stage- For protection from heat and cold wave there is intervention to sow the rabi crops in between 2nd week of October to 2nd week of November to protect theirs vegetative phase from ground/radiation frost results from cold wave/wind chill injury and reproductive phase from terminal heat stress on Mustard, Chickpea, Wheat, Lentil, Linseed and pea crops. Life saving irrigation

Vegetative stage- Timely sown crop never face heat stress while very late sown(January) crop face heat stress hence only one option is to provide life saving irrigation and water spray during evening time frequently at 2-3 days intervals. Take care of termite attack by spraying Chlorpyriphos @ 1 ml/lt and drenching @ 3-5 ml/lt water

In Chickpea because of high soil and ambient temperature (> 35 °C) favours the dry root rot disease starts during flowering/reproductive stage (spraying Captan or thiram or carbendazim or ridomil MZ or Saaf @ 1,5-2 g/lt water)

Reproductive stage- To minimize the terminal heat stress during the month of March and April the only and only way is to provide frequent protective irrigation irrespective of theirs stages (Life saving irrigation). Take care of termite attack by spraying Chlorpyriphos @ 1 ml/lt and drenching @ 3-5 ml/lt water. In Chickpea because of high soil and ambient temperature (> 35 °C) favours the dry root rot disease starts during flowering/reproductive stage (spraying Captan or thiram or carbendazim or ridomil MZ or Saaf @ 1,5-2 g/lt water)

At harvest- Frequent irrigation should be provided to meet the evaporative losses.

Tomato/Brinjal/ Lady's finger/Cucurbits

Seedling / nursery stage- Due to heat stress wilting and mortality is more hence frequent irrigation and cover the nursery with mulch(Straw/leaves)

Vegetative stage- Due to heat stress wilting and mortality is more hence frequent irrigation and cover the nursery with mulch(Straw/leaves)

Reproductive stage- Drying of flower- Spray PCOA. Follow mulching after irrigation

At harvest- Immediate harvest after irrigation and shift it to safer place

Cold wave

Wheat

Seedling / nursery stage- Cold environment during tillering or branching stage favours more number of tillers in wheat and more branching in mustard, chickpea, lentil and linseed crops which prospects for high yield but it is detrimental for potato, tomato, brinjal, pea, creeper vegetables and fruits. Irrigation. Balanced fertilizer application.

Foliar spray of nutrients

Vegetative stage- Light irrigation. Mulching with crop residue \ weeds. Fertilizer application

Reproductive stage- Irrigation, fertilizer application

At harvest- N/A

Pigeonpea/Mustard/Linseed/Chickpea/pea

Seedling / nursery stage- In Mustard because of cool weather aphid insects attack is more prominent (spraying Rogor (Dimethoate) @ 2 ml or Monocrotophos 36 EC @ 1 ml /lt water during evening time is advised).

In linseed Alternaria blight (For blight spray Double dose (Iprodione 25 % WP + Carbendazim 25 % WP) @ 2 g per lt. water) and powdery mildew (prophylactic spraying of Sulfex @ 3 g or Kairathen 1 ml per lt water twice at weekly interval during evening time) disease are more common. For powdery mildew in pea (spraying Calixin (Tridemorph 80 % EC @ 5 ml per 10 lt water twice are highly recommended).



In Chickpea-Cold and wet environment (High humidity) during seedling stage cause collar rot, black root rot, wet rot, Pythium root and seed rot in Chickpea, while in potato, pea and tomato favours late blight (spraying of Krilaksil or Ridomil MZ chemical @ 1.5 g per liter water), powdery mildew (spraying newly emerged fungicide Double dose (Iprodione 25 % WP + Carbendazim 25 % WP) 2 g per Lt water twice at weekly interval) and bacterial wilt, leaf spot and canker (spraying carbendazim @ 2g/Lt water and streptocycline @ 1g/10 Lt. water at 10 DAP, 25 DAP and 40 DAP) diseases in respective vegetable crops. Anthracnose in cucurbitaceous species. Vegetative stage- Provide light irrigation. Follow mulching with crop residue \ weeds\straw\leaves. In Mustard because of cool weather aphid insects attack is more prominent (spraying Rogor (Dimethoate) @ 2 ml or Monocrotophos 36 EC @ 1 ml /Lt water during evening time is advised)

Reproductive stage- Pigeonpea- During flowering and pod formation stage attack of Pod borer/sucking bug, mites, blister beetle insects as well as sterility disease may occur more (spraying Profenophos 50 EC, methomyl 40 SP or monocrotophos 36 SL kill the larvae but as the webs protect them from contact insecticides hence along with contact insecticides, mixing of fumigant insecticide such as DDVP @ 0.5 ml/l is required to make the larvae come out from the web. For Mites and Aphids, Dimethoate 30 EC @ 2ml/l and acaricides such as Dicofol 18.5 EC @ 2.5 ml/l water, for Blister beetle synthetic pyrethroids such as Cypermethrin 10 EC @ 1.0 ml/l or Lamda cyhalothrin 5 EC @ 1.0 ml/l water; for sterility mosaic Dicofol 18.5 EC 2.5 ml or Oxydemeton methyl 25 EC or Dimethoate 30 EC 2.0 ml or ml/l water on alternate row twice at an interval of 10 days are recommended).

Vegetables

Seedling / nursery stage- Raising seedling in Poly house, re sowing if damage is more. Provide shelter belt (Wind break) at appropriate spacing with Sisham, Ghamhar. Provide irrigation and mulching with straw and leaves

Vegetative stage- Provide light irrigation. Follow mulching with crop residue \ weeds\straw\leaves. Disease and pest control, care for chilling injury or replanting

Reproductive stage- Drying of flower- Spray PCOA. Follow mulching after irrigation

At harvest- Grading and safely dispose produce in the marketing

Frost

Wheat

Seedling / nursery stage- N/A

Vegetative stage- Provide light irrigation. Follow mulching with crop residue \ weeds\straw\leaves

Pigeonpea

Seedling / nursery stage- Exposure of crop to smoke by burning waste material during night time

Vegetative stage- Exposure of crop to smoke by burning waste material during night time, Light sprinkler irrigation

Reproductive stage- Exposure of crop to smoke by burning waste material during night time, Light sprinkler irrigation

At harvest- Exposure of crop to smoke by burning waste material during night time, Light sprinkler irrigation

Tomato & Potato and Horticultural crops (fruit)

Seedling / nursery stage- Create smoke around the field by using waste materials or set afire with used mobile oil in north-west or west-north direction towards incoming cold waves. Use polythene or bamboo hoogli in small horticultural /nursery/cash vegetable crops during morning hour and remove it during daytime. In Perennial or Horticulture crop (Fruit) also frequent irrigation followed by mulching, thatching , creating smoke screen s and lighting of fire should be practiced in availability of irrigation facility

Vegetative stage- Earthing up, Irrigation and create smoke around the field by using waste materials or set a fire with used mobile oil in north-west or west-north direction towards incoming cold waves. Use polythene or bamboo hoogli in small horticultural /nursery/cash vegetable crops during morning hour and remove it during daytime. In Perennial or Horticulture crop (Fruit) also frequent irrigation followed by mulching, thatching , creating smoke screen s and lighting of fire should be practiced

Reproductive stage- Immediate harvesting and disposal

At harvest- Harvest in dry weather

Cyclone- Not applicable



CONTINGENCY PLANS FOR RABI

1. Sowing window information

Land type	Cropping system	Crop name	Optimum sowing window (Please mention along with week)
1. Upland	Maize/ Groundnut/- Finger Millet – Vegetable/Toria/ Mustard / Linseed/ Safflower	Toria, Mustard, Linseed, Safflower, Vegetable- (Cauliflower, Tomato, Cabbage, Pea, Radish, Coriander, Amaranthus, French bean, Carrot, Spinach (with harvested rain water)	Toria- 1 st week of September – 4 th week of September Mustard- 1 st week of October– 4 th week of October Linseed- 4 th week of October- 2 nd week of November Safflower- 1 st week of October- 3 rd week of October Vegetables- 2 nd week of September – 4 th week of October
2. Mid Land	Rice –Wheat Rice –Chickpea Rice –Lentil Rice-Linseed Rice-Potato Rice-Vegetable	Irrigated-Wheat (Zero tillage), Mustard, Rabi Maize, Potato, Vegetables (Tomato, Chili, Brinjal, capsicum, Cabbage, Cauliflower, Broccoli, Vegetable pea, French bean, Radish, Coriander, Amaranthus, Spinach) Rainfed (Zero tillage)- Barley, Safflower, Chickpea, Lentil, Linseed, Lathyrus, Paira cropping Linseed, Lathyrus, Fodder crop-Oat, Maize,	Wheat (Timely)– 3 rd week of October – 4 th week of November Wheat (Late)-1 st week of December – 2 nd week of December Mustard- 2 nd week of October – 1 st week of November Rabi Maize- 2 nd week of October – 2 nd week of November Potato- 4 th week of October -2 nd Week of November Linseed- 2 nd week of October – 2 nd week of November Chickpea – 2 nd week of October – 1 st week of November Lentil- 3 rd week of October- 2 nd week of November Vegetables- 1 st week of October – 4 th week of November Lathyrus- 2 nd week of October – 4 th week of November Oat- 2 nd week of October – 4 th week of October Maize- 2 nd week of October – 4 th week of October
3. Low Land	Rice- Linseed (Paira) Rice-Vegetable	Paira cropping Linseed, Lathyrus Fodder crop- Oat,maize	Linseed- 4 th week of October – 4 th week of November Lathyrus- 4 th week of October – 4 th week of November Oat- 3 rd week of October – 1 st week of November Maize- 3 rd week of October – 1 st week of November

2. Contingency measures for Field crops grown with residual moisture under rainfed condition

2 (A) Optimal residual moisture

2A1 Land type- UPLAND

- a) Cropping system- Maize, Finger Millet, Toria, Mustard, Linseed, Vegetable
- b) Crop name- Toria, Mustard, Linseed, Vegetable- Cauliflower, Tomato, Cabbage, Pea, radish, Cariander, Amaranthus, French bean, Carrot, Spinach (with harvested rain water)
- c) Sowing Window- Toria-1st week of September – 4th week of September, Mustard-1st week of October– 4th week of October, Linseed- 4th week of October- 2nd week of November, Vegetables- 2nd week of September – 4th week of October
- d) Variety- Toria-PT 203, Panchali; Mustard- Pusa Mahek, Pusa Mustard 25, NRCHB 101, Bharat Sarson 1, Pusa 28, 30; Linseed- Sharda, Priyam, Divya

e) Agronomic management practices

- Rain water harvesting and recycling.
- Deeping of water storing structure(Shallow and deep) in April and May month
- Deep summer ploughing in April and May month.
- Strengthing and raising of field bunds in April and May months
- Sowing in defined window for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better crop stand (Plant population)
- Application of Lime or Dolomite (3-5 q/ha) in soil
- Soil application of Sulphur (20 kg/ha) and boron (1kg/ha) in oilseed, pulses and vegetables.
- Foliar spray of Urea (2 %) at flower initiation and pod formation stage in oilseed and pulses
- Follow seed priming (warm water for 4-6 hrs.) before sowing
- Follow seed treatment with fungicide-insecticide-rhizobium
- Irrigate only at critical stages
- Pre and post emergence weedicide application
- Follow hoeing after manual weeding
- Follow RDF, INM and IPM
- For Water use efficiency use antitranspirant, reflectant and mulches
- Regular monitoring of field for disease and insect attack
- Use pheromone trap and attractant
- Promote protected vegetable cultivation under naturally ventilated polyhouse and net house.
- Timely sowing for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population)



- Lime or Dolomite application in soil
- Foliar spray of Sulphur and boron
- Proper water management
- Take care of Aphid, white rust in Mustard, Early, late blight and leaf curling in potato

Mustard/Toria – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Linseed - Follow seed treatment, Irrigate only at critical stages, Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Precautions for pod borer, bud fly insect and powdery mildew disease management.

2A2 Land type- MEDIUM LAND

- a) Cropping system- Rice –Wheat, Rice –Chickpea, Rice –Lentil, Rice-Linseed, Rice-Potato, Rice-Vegetable
- b) Crop name- Irrigated-Wheat(Zero tillage), Mustard, Rabi Maize, Potato, Vegetables (Tomato, Chili, Brinjal, capsicum, Cabbage, Cauliflower, Broccoli, vegetable pea, french bean, Radish, Coriander, Amaranthus, Spinach); Rainfed (Zero tillage)- Chickpea, Lentil, Linseed, Lathyrus; Paira cropping; Linseed, Lathyrus
- c) Sowing Window- Wheat (Timely)- 3rd week of October – 4th week of November; Wheat (Late)-1st week of December – 2nd week of December, Mustard- 2nd week of October – 1st week of November; Rabi Maize- 2nd week of October – 2nd week of November; Potato- 4th week of October -2nd Week of November; Linseed- 2nd week of October – 2nd week of November; Chickpea – 2nd week of October – 1st week of November; Lentil- 3rd week opf October- 2nd week of November; Vegetables- 1st week of October – 4th week of November; Lathyrus- 2nd week of October – 4th week of November
- d) Variety- Wheat-(Timely), K 8027, HD 2967, K 1006, K 307, HDR 77, HD 2733; Late sown wheat- HD 3059, PBW 373, DBW 14, 39, HI 1563; Mustard- Pusa mahek, Pusa mustard 25, NRCHB 101, Bharat sarson 1, Pusa 28, 30; Rabi Maize- BVM 2, Malvia Makka 2, HQPM 5; Potato- Kufri Surya, Kufri Arun, Kufri Sutlej, Kufri Laukar, Kufri Lalima; Linseed- (rainfed)- Sharda, Priyam, Divya,; Chickpea- Jaki 9218, Kak 2, Birsa Chana 3; Lentil – HUL 57,WBL 77, KLS 218; Lathyrus- Maha Teora

e) Agronomic management practices

- Seed treatment with Azotobacter and Azospirillum and also soil application
- Timely sowing for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population)
- Follow deep summer ploughing
- Proper water management
- Follow seed treatment
- Irrigate only at critical stages
- Pre emergence weedicide application
- Follow RDF, INM and IPM
- Take care of Painted Bud, Aphid, white rust in Mustard; Early, late blight and leaf curling and grub in potato; Collar rot, Dry root rot, Pod borer in Chickpea; Wilt in Lentil.

Wheat-Seed treatment with Azotobacter and Azospirillum and also soil application. Timely sowing for better establishment Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population), Follow deep summer ploughing. Proper water management. Irrigate only at critical stages(3-6). Pre emergence weedicide application

Follow RDF, INM and IPM. Take case of Loose Smut Disease. 1st irrigation should be after CRI stage i.e at 30-35 DAS

Mustard – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew.

Potato- Seed treatment. Proper spacing. Frequent irrigation. Take care for leaf curling,Early, late blight and grub infestation Irrigate during cold day and night to get relief from frost attack. Produce smoke during cooler day and night

Linseed - Follow seed treatment, Irrigate only at critical stages, Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Precautions for pod borer, bud fly insect and powdery mildew disease management.



Chickpea - Seed treatment with Rhizobium culture and Phosphate solubilizing bacteria (PSB) and Trichoderma. Management for Collar rot during temperature fall and dry root rot during temperature increment. Pre emergence weedicide application. Irrigate at Critical stages. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray

Potato- Seed treatment. Proper spacing. Frequent irrigation. Take care for leaf curling, Early, late blight and grub infestation. Irrigate during cold day and night to get relief from frost attack. Produce smoke during cooler day and night. Earthing up in potato at 30-35 DAS or crop attended the height of 20-25 cm

Lentil - Foliar spray of Sulphur and Boron is necessary. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Follow deep summer ploughing. Proper water management. Follow seed treatment with Rhizobium culture and PSB. Irrigate only at critical stages. Pre emergence weedicide application. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray. Management for wilt disease. One hand weeding followed by two hoeing for management of weeds (HW-20-25 DAS and Hoeing 30-32 and 40-42 DAS).

2A3 Land type- LOW LAND

a) Cropping system-Rice- Linseed (Paira), Rice-Vegetable

b) Crop name- Linseed/ lathyrus (Paira cropping), Vegetables (Onion, Fodder Crop- Oat, Maize, lathyrus

c) Sowing Window Linseed- 4th week of October – 2nd week of Lathyrus- 4th week of October – 4th week of November

d) Variety- Linseed- Sharda, Priyam, Divya.; Lathyrus-Maha Teora

e) Agronomic management practices

Linseed - Follow seed treatment. Irrigate only at critical stages. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Follow RDF, INM and IPM. Be cautious for pod borer, bud fly insect and powdery mildew disease management.

2 (B) Less than optimum moisture i.e., 25% less than normal, which can happen due to insufficient rainfall during September/October months. Deficit of 20-40% rainfall

2B1 Land type- UP LAND

a) Cropping system- Maize-Toria, Finger millet-Linseed

b) Crop name – Toria, Linseed

c) Sowing Window- Toria- 3rd week of September – 4th week of September, Linseed- 1st week of October- 3rd week of October

d) Variety- Toria- PT 203, Panchali; Linseed- Sharda, Priyam, Divya

e) Agronomic management practices

- Rain water harvesting and recycling.
- Deepening of water storing structure(Shallow and deep) in April and May month
- Deep summer ploughing in April and May month.
- Strengthening and raising of field bunds in April and May months
- Sowing in defined window for better establishment
- Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better crop stand (Plant population)
- Application of Lime or Dolomite (3-5 q/ha) in soil
- Soil application of Sulphur (20 kg/ha) and boron (1kg/ha) in oilseed, pulses and vegetables.
- Foliar spray of Urea (2 %) at flower initiation and pod formation stage in oilseed and pulses
- Follow seed priming (warm water for 4-6 hrs.) before sowing
- Follow seed treatment with fungicide-insecticide-rhizobium
- Follow deep summer ploughing
- Irrigate only at critical stages
- Pre and post emergence weedicide application
- Follow hoeing after hand weeding
- Follow RDF, INM and IPM
- For Water use efficiency use antitranspirant, reflectant and mulches
- Regular monitoring of field for disease and insect attack
- Use pheromone trap and attractant
- Promote protected vegetable cultivation under naturally ventilated polyhouse and net house.
- Zero Tillage for seed placement at proper depth for better germination
- One hand weeding followed by one hoeing for management of germinating weeds
- For Water use efficiency use antitranspirant, reflectant and mulches



Toria – Seed treatment and proper seed rate for optimum crop stand. Thinning, weeding at 30-35 DAS followed by irrigation. Follow RDF, INM and IPM. Irrigate at three critical stages 30-35 DAS, before flowering and capsule/pod formation. Apply second dose of Urea before flowering. Management for painted bug, aphid and Powdery mildew

Linseed - Follow seed treatment, Irrigate only at critical stages, Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Precautions for pod borer, bud fly insect and powdery mildew disease management.

2B.2 Land type- MEDIUM LAND

- a) Cropping system- Rice –Chickpea, Rice –Lentil, Rice-Linseed, Rice-Fodder
- b) Crop name – Chickpea, Lentil,Paira cropping- Linseed, lathyrus
- c) Sowing Window- Linseed- Sharda, Priyam, Divya; Chickpea – 2nd week of October – 2nd week of November, Lentil- 3rd week of October- 2nd week of November, Lathyrus- 2nd week of October – 4th week of November
- d) Variety- Linseed- Sharda, Priyam, Divya; Chickpea- KWR 108, HK 94134, Jaki 9218, Birsa Chana 3; Lentil –WBL 77, KLS 218; Lathyrus- Maha Teora

e) Agronomic management practices

Linseed - Follow seed treatment. Irrigate only at critical stages. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Follow RDF, INM and IPM. Precaution for pod borer, bud fly insect and powdery mildew disease management

Chickpea - Seed treatment with Rhizobium culture and Phosphate solublizing bacteria (PSB) and Trichoderma. Management for Collar rot during temperature fall and dry root rot during temperature increment in. Pre emergence weedicide application. Irrigate a critical stages. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray

Lentil - Foliar spray of Sulphur and Boron is necessary. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Follow deep summer ploughing. Proper water management. Follow seed treatment with Rhizobium culture and PSB. Irrigate only at critical stages. Pre emergence weedicide application. Follow RDF, INM and IPM also drench FYM treated with Trichoderma and Foliar spray. Management for wilt disease. One hand weeding followed by two hoeing for management of weeds (HW-20-25 DAS and Hoeing 30-32 and 40-42 DAS)

Lathyrus- Proper seed rate and Follow RDF for potential yield. Follow two nipping in between 25-45 DAS. Irrigate after every nipping. Take care of diseases.

2B3 Land type- LOW LAND

- a) Cropping system- Rice- Linseed/Lathyrus (Paira), Rice-Fodder
- b) Crop name- Linseed/ lathyrus (Paira cropping), Fodder Crop-Oat, Maize
- c) Sowing Window- Linseed- 4th week of October – 4th week of November; Lathyrus- 4th week of October – 4th week of November; Fodder Crop- Maize- 4th week of November – 2nd week of December; Oat- 4th week of November – 2nd week of December
- d) Variety- Linseed- Sharda, Priyam, Divya, Lathyrus-Maha Teora; Fodder- Oat— Kent, Maize- Pratap Makka(Chari 6), J 1006

e)Agronomic management practices

Linseed - Follow seed treatment. Irrigate only at critical stages. Seed placement at proper depth for better germination and optimum seed rate with right crop geometry for better establishment (Plant population). Soil as well as seed treatment for the management of disease and termite attack. Pre and post emergence weedicide application. One hand weeding followed by one hoeing for management of weeds between 20 and 45 DAS. Irrigate at critical stages (before flowering and at pod formation). For dual purpose (seed and flax) increase N fertilizer dose by 25 percent. Follow RDF, INM and IPM. Follow deep summer ploughing. Proper water management. Pre emergence weedicide application. Follow RDF, INM and IPM. Precaution for pod borer, bud flies insect and powdery mildew disease management.

Forage- Oat- Proper seed rate for better crop stand. 1st and 2nd cutting at 30 and 45 DAS and 3rd before flowering. Berseem- 1st at 50 DAS and follow 2nd, 3rd and 4th cutting every at an interval of 30-40 days. Lucern- Same as Berseem. Japani Mustard- 1st at 50 DAS during fruiting and rest cutting every at an interval of 30 days. Follow RDF. For Lucern other than N P K use Lime , Boron and Molybdenum micro nutrients for better yield.



CONTINGENT STRATEGIES FOR LIVESTOCK, POULTRY & FISHERIES

1 Livestock

Suggested contingency measures under DROUGHT event

a) Before the event

Feed and fodder availability

Preservation of surplus fodder, encourage fodder cultivation and tree plantation and also encourage supply of molasses to cattle feed plants

- Preservation of surplus fodder

Green grass is a good source of vitamin A which is present in the form of Carotene. One kg of green grass provides 50mg of vitamin A and 15 to 20g protein to the animal. Cowpea, beans, subabul leaves etc. give 30 to 40g of protein. From grass fodder herbivorous animals get the carbohydrates (energy source), proteins ("building material" of the body) and vitamins (especially carotene), which are the main drives of sustainable operation of the body.

Two methods are available for preserving or conserving the seasonal excess of green fodder, viz. hay making and silage making. Each method has its own limitations and advantageous. Ensiling is preferred on the basis of fodder quality.

Hay making

Hay -refers to cereals, grasses or legumes that are harvested at appropriate stage, dried and stored

Ensilage / Silage making

Silage may be defined as the green succulent roughage preserved under controlled anaerobic fermentation in the absence of oxygen by compacting green chops in air and watertight receptacles.

- Complete Feed Blocks

Supply enriched complete feed blocks containing dry roughage, concentrates/ unconventional supplements 50:50 ratio. Complete feed blocks may be sourced from different commercial sources.

Feeding practices for livestock in India at present separate feeding of roughage and concentrate

- ❖ Chopped roughage and soaked concentrate mixed together
- ❖ Chopped roughage mechanically mixed with concentrate as mash
- ❖ Chopped roughage and concentrate ingredients mixed and densified as Complete Feed Block

Concept of densified complete feeds with fibrous crop residues is a noble way to increase the intake and improve the nutrients utilization. A complete feed block has been defined as a system of feeding all ingredients including roughages, processed and mixed uniformly, to be made available ad lib to the animals.

- Urea molasses mineral block licks

Urea-molasses mineral block lick can sustain the animals by providing protein, energy and essential minerals. It is cost effective, easy to handle and transport and available commercially through milk cooperatives. Therefore, it is required that urea molasses blocks stocks (UMBS) are made available in the rain-deficient areas.

- Methods used for improving nutritive quality of straws and other crop residues like urea treatment

Spray dry roughages such as paddy and wheat straw with about 10% molasses and 2% urea for maintenance of animals in fodder deficit areas.

Preparation of 100 kg roughage-based enriched feed containing 88.8 kg wheat straw or any other straw/stover, 10 kg molasses, 1 kg urea and 0.5 kg mineral mixture will cost about Rs. 375-450 per quintal.

- Utilization of forest byproducts for feeding of livestock

Use of dry and fallen tree leafs like Pipal, Neem, Mango and Kathal etc.

- Making Leaf meal

- Use of conventional and non conventional feeds

- Rice Mills

The main by-products of rice are rice straw, rice husk or hull, and rice bran. Rice straw is produced when harvesting paddy. Rice husks generated during the first stage of rice milling, when rough rice or paddy rice is husked.

- Aquatic plants

- ❖ One kg DM/100 kg BW
- ❖ Water hyacinth, aquatic spinach, Stalks & leaves of lotus plant, Hydrilla, Pistia etc.

- Encourage supply of molasses to cattle feed plants

Molasses and Bagasse are the byproducts from sugarcane industry and are available in abundance. They can be used as cattle feed after supplementation with urea. Such a ration is a ready feed during drought and scarcity conditions when nothing else is available for feeding to animals.

- Crop Residue Enrichment & Densification

Crop residues can be fortified with feed ingredients like cakes, barns, grains, molasses, hay, minerals and then densified into blocks or pellets to save on storage and transport costs. Also balanced ration in the form of complete diet or total mixed ration as per need of animals can be supplied for improved productivity.

- Demonstration of Re-vegetation of Common Grazing Land

The grazing lands play an important role in the lives of rural people who are getting fodder, fuel, drinking water from commons. However, such lands are being continuously degraded due to overgrazing and overexploitation by locals. Re-vegetation of such lands on scientific lines suiting to agro-climatic conditions is to be demonstrated through strengthening institutional arrangement at village level. Fodder production from such lands can be enhanced substantially by introducing high yielding cultivated fodder crops, grasses and pasture legumes. An integrated approach of growing cultivated crops, grasses, trees and shrubs under silvi-pastoral/ horti - silvipasture system will improve overall productivity of such land.



Drinking water

Repairs of tube wells, clear off the sludge in the canals and local water catchments and clean the water tanks, large ponds and lakes

Health and Hygiene

Tick damage and tick-borne diseases

- Tick damage - Vaccinate the cattle against tick-borne diseases
- Tick-borne diseases- Vaccination is best done in calves under 6 months of age and one dose is sufficient
- Babesiosis (Red water)- Treatment involves keeping the cattle calm. They should not be driven over long distances and should be injected with Berenil or Imizol. The dose for Berenil is 5 ml of made up solution (1 packet mixed with 12.5 ml of sterile water) for each 100 kg (for example, 20 ml for a 400 kg animal)
- Sarcoptic Mange in pigs- Not applicable before event

Diseases caused by biting insects

- Trypanosomiasis- Fly control is important for prevention of the disease.
- Three-day stiff sickness- Prevention is by vaccination
- Lumpy-skin disease- Prevention is by vaccination

Diet related Disease problems

- Eating plastic bags and wire(Pica)- Feed cattle well, especially in winter. Clear wires and plastic bags from the grazing area. Watch cattle closely when they are grazing. Mineral mixture supplement should be given to the animal
- Poisonous plants- Not applicable before event
- Botulism- Prevention involves vaccination and good nutrition. Burn or bury all carcasses, bones or decaying material

Deficiency diseases

Cattle grazing on drought-stricken pastures can suffer serious depletion of reserves of minerals and vitamins.

- Copper and Cobalt- Not applicable before event
- Calcium, Phosphorous & Vit. D- Not applicable before event
- Vitamin A- Not applicable before event

Infectious Diseases

- Foot and Mouth Disease (FMD)- Vaccination at the age 4 months and above. Booster should be given 1 month after first dose then every six monthly
- Haemorrhagic Septicaemia (HS)- Vaccination at the age 6 months and above. Annually in endemic areas. Vaccinate the animal before onset of monsoon every year preferably in the month of May and June.
- Black Quarter (BQ)- Vaccination at the age 6 months and above. Annually in endemic areas. Vaccinate the animal before onset of monsoon every year preferably in the month of May and June
- Anthrax- Vaccination at the age 4 months and above. Annually in endemic areas. Vaccinate the animal before onset of monsoon every year preferably in the month of May and June.
- Rabies (Post bite therapy only)- Not applicable
- Enterotoxaemia (pulpy kidney)- Vaccinate the animal at the age of 3-4 months, repeat after 15 days and then annually.
- Pneumonia- Not applicable

Non-Infectious Diseases

- Ruminal tympany (Bloat)- Not applicable
- Rumen acidosis- Not applicable
- Intussusception- Deforming should be give
- Pregnancy toxemia (Ketosis)- Fed the pregnant animal with balanced ration.

Poisoning

- Organochlorine compounds- Not applicable
- Organophosphorus compounds- This group consists of malathion, darathion, chlorathion, carbophenothion, demton, dasnon, dimethylparathion, trichlorphon, dioxalthion etc. Symptoms of toxicity are profuse salivation, muscle stiffness, dyspnoea with open mouth breathing, tremors. Treatment consists of administering antidote, usually atropine sulphate.
- Snake bite- Usually bitten on the scrotum or udder. The presence of hair may obscure the typical fang marks. Prolonged pain, muscular weakness, impaired vision, nausea and paralysis are generally exhibited along with symptoms of shock. If anti-venin is not available and the bite is located in an area where a tourniquet cannot be applied, excision of an area of skin and sub-cutaneous tissue can be life-saving

b) During the event

Feed and fodder availability

- Lactating and pregnant animals need to be provided enriched feed to meet the requirements and rest of animals be provided the maintenance diet. In case of acute shortage, lactating animals be provided feed meeting 50% of the requirements to maintain minimum level of production.
- Drought tolerant fodder crops (like sorghum PC 6 and MP chari, cowpea - BL 1 and 2) and fodder grasses (like stylo, *cenchrus ciliaris*, *athropogon*, etc.) should be cultivated. Under the mini kit programme, the developmental department need to provide fodder crop seeds in the drought-affected areas.
- Provide salt dose daily through feed (40-50 g of salt per adult animal and 10-20 g for small ruminants and calves).

Issue

- Large scale migration -Creating additional resources in drought prone area
- Grazing of poisonous plants/toxicity problems -Inventory of anti nutritional/toxic factors. Creating awareness in farmer for avoiding nitrate/nitrite HCN poisoning.
- Transport of fodder from normal DPA-Establishing feed and fodder banks. Effective mechanism for distribution of fodder/feed to productive animals. Densification/baling/briquette technologies

Drinking water

- Harnessing water through the existing reservoirs and exploitation of groundwater.

Health and Hygiene

Tick damage and tick-borne diseases

- Tick damage - If disease occurs Treat the cattle against tick-borne diseases. Consult Veterinarian.
- Tick-borne diseases- Prevention is by tick control, treatment of diseased animal and vaccination. Consult Veterinarian.
- Babesiosis (Red water)- Treatment involves keeping the cattle calm. They should not be driven over long distances and should be injected with Berenil or Imizol. The dose for Berenil is 5 ml of made up solution (1 packet mixed with 12.5 ml of sterile water) for each 100 kg (for example, 20 ml for a 400 kg animal). Consult Veterinarian.
- Sarcoptic Mange in pigs- Itching; dermatitis; rubbing; scratching; reduced growth rate. Miticidal sprays; pour-ones injection and in-feed premix. Consult Veterinarian.

Diseases caused by biting insects

- Trypanosomiasis- Treated with SURAMIN through intramuscular injection or intravenous infusion if sufficient observation is possible. Consult Veterinarian.
- Three-day stiff sickness- It is important that the animal is given food and water if it is unable to stand
- Animal should be treated by Veterinarian
- Lumpy-skin disease- If your cattle get this disease, you should speak to your state veterinarian

Diet related Disease problems

- Eating plastic bags and wire (Pica) - Mostly occurring in those animals which are having shortage of feeds and fodder and deficiency of Phosphorus. Prevention involves the following: - Feed cattle well, especially in winter. Clear wires and plastic bags from the grazing area. Watch cattle closely when they are Grazing. Mineral mixture supplement should be given to the animal. Once the cow has eaten plastic bags or wire, the only effective treatment is an operation. Consult Veterinarian.
- Poisonous plants- Due to scarcity of feed s and fodder animals used to consume poisonous plans and are more likely to get toxicity. Poisoning can also happen when owner or animal handler move cattle to new paddocks where toxic plants occur. Consult Veterinarian.
- Botulism- Botulism can occur when cattle eat carcass and bone material when there is a lack of feed during drought or if they have a phosphorus deficiency
- Treatment is only possible in the early stages and requires an antitoxin. Consult Veterinarian.

Deficiency diseases

- Cattle grazing on drought-stricken pastures can suffer serious depletion of reserves of minerals and vitamins.
- Copper and Cobalt- Characterized by anorexia and wasting. Deficiency affects growth and fertility of the cattle. Anemia, diarrhoea and unthriftiness occur in extreme cases. Copper or cobalt sulphate in the form of mineral mixture supplement causes rapid disappearance of the symptoms
 - Calcium, Phosphorous & Vit. D- Deficiency may result in rickets in calves and osteomalacia in adults. Mineral supplementation in diet is essential to prevent this deficiency.
 - Vitamin A- Vit. A deficiency occurs in cattle on dry countryside during periods of drought. Symptoms include night blindness, corneal keratinization, ptyriasis, hoof defects, loss of weight and infertility. Animals should have access to green pasture and should be supplied with Vit. A in feed to prevent deficiency.

Infectious Diseases

- Foot and Mouth Disease (FMD)- If outbreak occurs then the animal should be treated. Foot lesion should be washed with soap / detergent the apply Povidon iodine lotion while in mouth lesion boroglyserine should be applied. Consult Veterinarian.
- Haemorrhagic Septicaemia (HS)- If disease occurs animal should be treated with Broad Spectrum Antibiotic like penicillin at higher dose and other supportive medicine as per the symptoms. Consult local Veterinarian.
- Black Quarter (BQ)- If disease occurs animal should be treated with Broad Spectrum Antibiotic like penicillin at higher dose and other supportive medicine as per the symptoms. Consult local Veterinarian.
- Anthrax- If disease occurs animal should be treated with Broad Spectrum Antibiotic like penicillin at higher dose and other supportive medicine as per the symptoms. Consult local Veterinarian.
- Rabies (Post bite therapy only)- Vaccinate the animal immediately after suspected bite. Booster should be given on 3, 7, 14, 28 and 90 (optional) days after first dose.
- Enterotoxaemia (pulpy kidney)- Not applicable
- Pneumonia- Not applicable

Non-Infectious Diseases

- Ruminal tympany (Bloat)- Not applicable



- Rumen acidosis- Ingestion of large amounts of highly fermentable carbohydrate feeds causes an acute illness due to excess production of lactic acid in the rumen. Clinically, the disease is manifested by dehydration, blindness, recency, complete rumen stasis and a high mortality rate. Normal saline, sodium bicarbonate and antihistaminic are advised.
- Intussusceptions- It occurs commonly due to nodular worms, change in feed and local intestinal problems. The animal is dull, off-feed, kicking at the belly with no rise of temperature, frequent straining with no defecation, colic symptoms, and at later stages, recency. Emergency surgery is the only rational treatment.
- Pregnancy toxæmia (Ketosis)- It is a highly fatal disease caused due to a decline in the plane of nutrition and short periods of starvation (40 hrs) during the last two months of pregnancy. Treatment comprises intravenous administration of 50% glucose. Supply of molasses in the ration and concentrate in the last two months of pregnancy helps in preventing the condition.

Poisoning

- Organochlorine compounds- Not applicable
- Organophosphorous compounds- This group consists of malathion, darathion, chlorathion, carbophenothon, demton, dasnon, dimethylparathion, trichlorphon, dioxalthion etc. Symptoms of toxicity are profuse salivation, muscle stiffness, dyspnoea with open mouth breathing, tremors. Treatment consists of administering antidote, usually atropine sulphate.
- Snake bite- Usually bitten on the scrotum or udder. The presence of hair may obscure the typical fang marks. Prolonged pain, muscular weakness, impaired vision, nausea and paralysis are generally exhibited along with symptoms of shock. If anti-venin is not available and the bite is located in an area where a tourniquet cannot be applied, excision of an area of skin and sub-cutaneous tissue can be life-saving

c) After the event

Feed and fodder availability

Promotion of fodder seed production, cultivation and storage, establishment of fodder block making machines in fodder surplus areas

Post flood feeding management

- Animal should not be allowed to graze in water logged area
- Feeds to be protected from fungal contamination & wet feeds to be dried & fed
- Provides clean drinking water to animals
- Provide ready to eat feed blocks particularly the pregnant and lactating animals
- Requirement of energy may be met providing crude molasses
- Top feeds/ tree leaves available in the area to be provided to meet the DM requirement

Specific contingencies can be adopted for livestock feeding depending upon availability as under in different regions during drought situation

Neem seed kernel cake (NSKC), Saw dust, Paper waste, Agave (Ketki), Cactus, Tree leaves and vegetable leaves, Cher leaves and fruits, Straw and gotars, Sugarcane bagasse as animal feed and Use of damaged grains as feed

Drinking water

To strengthen reservoirs by promoting recharging of water and rain water harvesting during rainy season.

Health and Hygiene

Tick damage and tick-borne diseases

- Tick damage - Treat the cattle against tick-borne diseases. Consult Veterinarian.
- Tick-borne diseases- Prevention is by tick control, treatment of diseased animal and vaccination. Consult Veterinarian.
- Babesiosis (Red water)- Treatment involves keeping the cattle calm. They should not be driven over longdistances and should be injected with Berenil or Imizol. The dose for Berenil is 5 ml of made up solution (1packet mixed with 12.5 ml of sterile water) for each 100 kg (for example, 20 ml for a 400 kg animal).Consult Veterinarian.
- Sarcoptic Mange in pigs- Not applicable after event

Diseases caused by biting insects

- Trypanosomiasis- Treated with SURAMIN through intramuscular injection or intravenous infusion if sufficient observation is possible. Consult Veterinarian
- Three-day stiff sickness- It is important that the animal is given food and water if it is unable to stand.
- Animal should be treated by Veterinarian
- Lumpy-skin disease- If your cattle get this disease, you should speak to your state veterinarian

Diet related Disease problems

- Eating plastic bags and wire (Pica)- Feed cattle well, especially in winter. Clear wires and plastic bags from the grazing area. Watch cattle closely when they are grazing. Mineral mixture supplement should be given to the animal
- Poisonous plants- Not applicable
- Botulism- Prevention involves vaccination and good nutrition. Burn or bury all carcasses, bones or decaying material

Deficiency diseases

Cattle grazing on drought-stricken pastures can suffer serious depletion of reserves of minerals and vitamins.

- Copper and Cobalt- Not applicable
- Calcium, Phosphorous & Vit. D- Not applicable
- Vitamin A- Not applicable

Infectious Diseases

- Foot and Mouth Disease (FMD)- If outbreak occurs then the animal should be treated. Foot lesion should be washed with soap / detergent the apply Povidon iodine lotion while in mouth lesion boroglyserine should be applied. Consult Veterinarian.
- Haemorrhagic Septicaemia (HS)- Not applicable
- Black Quarter (BQ)- Not applicable
- Anthrax- Not applicable
- Rabies (Post bite therapy only)- Not applicable
- Enterotoxaemia (pulpy kidney) - It affects the animals in a high state of nutrition on a lush feed, grass or grain. Morbidity rates seldom exceed 10% but mortality rate approximates 100%. Under certain conditions, the organism proliferated rapidly in the intestines and produces lethal quantity of toxin. Suphadimidine with other supportive medicine may be effective for treatment
- Pneumonia- It is one of the most common and important pathological conditions. It is characterized clinically by increased respiration, coughing and abdominal breathing. Treatment with broad spectrum antibiotic, nabulization and other supportive drugs is effective.

Non-Infectious Diseases

- Ruminant tympany (Bloat)- It is the over-distension of the left flank either due to free gas or froth. This is generally encountered in "greedy feeders" when lush green pasture is available. Oral administration of sweet oil with turpentine oil or at times with formalin is advised.
- Rumen acidosis- Not applicable
- Intussusceptions- Not applicable
- Pregnancy toxæmia (Ketosis)- Not applicable

Poisoning

- Organochlorine compounds- This group includes DDT, BHC, lindane, aldrin, dieldrin, chlordane, toxaphane, methocyclor etc. which are used as pesticides on crops. Toxicity symptoms include increased excitability and irritability followed by muscle tremors, weakness, paralysis etc. Treatment consists of administering antidote, usually short-acting barbiturates.
- Organophosphorus compounds- This group consists of malathion, darathion, chlorathion, carbophenothion, demton, dasnon, dimethylparathion, trichlorphon, dioxathion etc. Symptoms of toxicity are profuse salivation, muscle stiffness, dyspnoea with open mouth breathing, tremors. Treatment consists of administering antidote, usually atropine sulphate.
- Snake bite-Usually bitten on the scrotum or udder. The presence of hair may obscure the typical fang marks. Prolonged pain, muscular weakness, impaired vision, nausea and paralysis are generally exhibited along with symptoms of shock. If anti-venin is not available and the bite is located in an area where a tourniquet cannot be applied, excision of an area of skin and sub-cutaneous tissue can be life-saving

2 Poultry

Suggested contingency measures under DROUGHT event

a) Before the event

Shelter management

Optimum space should be provided. Orientation of shed (Long axis) should be in North – South. Plantation of tree around shed to provide cool environment. Provision of ad lib. Fresh water

Shortage of feed ingredients

Storage of feed

Drinking water

Manage clean drinking water. Storage facility should be made. Water quality should be checked before drinking to animal

Health and disease management

- Newcastle Disease- regular vaccination - Broiler birds should be with RD vaccine (Lasota 'F' strain) at the age of 4-7 days through Intra-nasal or Intra-ocular route. Layer birds should be vaccinated with NDV vaccine at the age of 9-14 day, 4 weeks, 13-14 weeks in drinking water/eye drop. Then at the age of 17 week with NDV vaccine through Intramuscular (IM) route
- Marek's disease Marek's disease- Birds should be vaccinated with Herpes virus turkey vaccine at the age of 1 day through Subcutaneous route.
- Fowl pox-Chick embryo adopted fowl pox vaccine at the age 6-8 weeks. It important for the layer and broiler birds.
- Drop in Egg Production or Quality- Not applicable
- Nervous Signs and Lameness- Not applicable
- Diarrhoea- Not applicable
- Upper Respiratory Diseases- Vaccination against the some of the viral diseases like Newcastle disease, influenza, infectious bronchitis, infectious laryngotracheitis which are also responsible for the respiratory symptoms can prevent this syndrome. Antifungal and antiparasitic drugs should be given.

Heat Wave

Plantation of tree around shed to provide cooler environment. Proper ventilation should be provided. Optimum space should be provided. Orientation of shed (Long axis) should be in East- West. Plantation of tree around shed to provide cool environment. Provision of ad lib. Fresh water. Manage green fodder and silage preparation. Height of roof should be minimum 220 – 240 cm. Roof of shed should be painted with white.



Cold Wave

Provide ad lib fresh water. Proper ventilation should be provided. Optimum space should be provided. Orientation of shed (Long axis) should be in North – South. Plantation of tree around shed to provide break cold wave. Provision of ad lib. Fresh water. Manage green fodder and silage preparation. Height of roof should be minimum 220 – 240 cm
Roof of shed should be painted with Black Floor of shed should be Dry

b) During the event

Shelter management

Optimum space should be provided, there must not be overcrowded. Protect the animal from direct sun light. Try to provide them cool water. Proper ventilation should be maintained. Allow for grazing during night/early morning. Provision of ad lib. Fresh water

Shortage of feed ingredients

Provide non conventional feed, supplement anti oxidant and anti stress

Drinking water

Provide clean fresh and cold drinking water all the time. Water availability may increase by 20-50% depending upon feed quality and environmental temperature. Soft drinking water should be preferred. Add vit-C and other anti stress ingredients with water

Health and disease management

- Newcastle Disease- Vaccination and treatment of diseased one. Newcastle disease is the most important disease for poultry farmers around the world. This disease causes a large number of deaths in chickens and huge losses to farmers and the industry. Diseased birds should be slaughtered immediately. Consult Veterinarian.
- Marek's disease Marek's disease- It is one of the important diseases of poultry caused by virus. Mortality is very high and causes economic losses to the farmer and poultry industry.
- Fowl pox- It is a viral infection of chickens and turkeys characterized by proliferative lesions in the skin (Cutaneous form), it also affect the GI tract and respiratory tract (Diphtheritic form)
- Drop in Egg Production or Quality- There are many different types of organisms that can cause a drop in egg production or quality. These include: Bacteria (E. coli, Salmonella), Mycoplasma, Viruses (Newcastle disease, influenza, infectious bronchitis, infectious laryngotracheitis, avian encephalomyelitis, egg drop syndrome). The Parasites, lack of Nutrition and Stress factor also support the onset of this condition. Adding vitamins and minerals to the water or feed may help. Consult Veterinarian
- Nervous Signs and Lameness- Chickens lie down because they cannot stand up. They also walk with a limp or are reluctant to move. Nervous signs may include staring into the sky, pulling the head and neck over their backs, paralysis. Sores on the breast muscles from lying down
- Diarrhoea- The stool or droppings of the chickens are not firm but very loose, watery, not of the normal colour and may contain blood. This may cause the feathers of the vent to be soiled and caked together, Depression, reluctance to eat, drink and move about, poor growth and death. Use an antibiotic or coccidiostatic drug in the water that was recommended by the animal health technician or veterinarian in the water for 3 to 5 days. Stress preparations that contain electrolytes, vitamins and minerals can be added to the water
- Upper Respiratory Diseases- Not applicable

Heat Wave

Water sprinkling to animal. Prevent the animal from direct sunlight. Optimum space should be provided, there must not be overcrowded. Fan should be provided to make the body cool. Try to provide them cool drinking water all time Proper ventilation should be maintained. Allow for grazing during night/early morning. Try to provide green fodder and silage. Stocking density should be less. Roof should be covered with tiles, paddy, dry leaves to protect from direct sun light

Cold Wave

Luke worm water should be provided at least 4-6 times a day. Prevent the animal from direct cold wave by closing the windows and doors. Optimum space should be provided, there must not be overcrowded. Proper ventilation should be maintained. Allow for grazing during sunny day time. Try to provide green fodder and silage. During extreme cold condition electric heater or wood fire heat should be provided. Try to make the environment inside and outside the shed dry. Gunny bags or blanket may be used to cover the body. Bedding material like paddy straw, Gunny Bag, Bhusa should be provided specially to young one shed.

c) After the event

Shelter management

Optimum space should be provided. Take care and fulfill the requirement of water along with proper nutrients. Take care of proper feeding as per requirement. Provision of ad lib. Fresh water

Shortage of feed ingredients

Not applicable

Drinking water

Provide adlib. Drinking water

Health and disease management

- Newcastle Disease- Disposal of dead birds
- Marek's disease Marek's disease- Disposal of dead birds
- Fowl pox- Disposal of dead birds

- Drop in Egg Production or Quality-Not applicable
- Nervous Signs and Lameness- A complete hygiene and disinfection programme should be planned together with the animal health technician or veterinarian. Antibiotics will only be effective against bacteria and can be used as recommended. If it is a viral disease, such as Newcastle disease, urgent steps have to be taken to prevent possible spread because it causes serious production losses
- Diarrhoea- Disposal of dead birds
- Upper Respiratory Diseases- There are many different types of organisms that can cause disease in the upper respiratory tract. These include: Mycoplasma Bacteria (E. coli, Pasteurella, Haemophilus), Viruses (Newcastle disease, influenza, infectious bronchitis, infectious laryngotracheitis), Parasites (mites and worms) And Fungi (Aspergillus). Cold stress is also one of the predisposing factors for the occurrence of respiratory problems. Use an antibiotic drug that was recommended by your animal health technician or veterinarian in the water for 3 to 5 days
- Stress preparations that contain electrolytes, vitamins and minerals can be added to the water

Heat Wave

Optimum space should be provided. Take care and fulfill the requirement of water along with proper nutrients. Take care of proper feeding as per requirement. Provision of ad lib. Fresh water

Cold Wave

Provide ad lib. Normal drinking water. Optimum space should be provided. Take care and fulfill the requirement of water along with proper nutrients. Take care of proper feeding as per requirement. Provision of ad lib. Fresh water

3 Fisheries

Suggested contingency measures under DROUGHT event

a) Before the event

Aquaculture

- Shallow water in ponds due to insufficient rains/inflow- Increase depth of pond, Repair dyke, outlet and inlet of pond; Prepare duck/pig house & stock pig @ 50-60, duck @ 450-500 no/ha if farmer involve in Integrated fish farming, Allow manure and urine directly in pond, Remove unwanted, predatory & old fishes and for this apply Mahua oil cake @ 2500kg/ha. Fixed net in outlet & inlet to prevent escaping of fish, Plough the pond and apply lime @ 250 kg/ha, Check the natural feed (plankton) @ 1.0 1.5 ml/50 lt. water; otherwise apply organic manure, Stock yearling (stunted grow fish) @ 6,000-8,000 no/ha
- Impact of salt load build up in ponds / change in water quality- Prevent entry of polluted water or apply lime at inlet.

Heat wave and cold wave

- Changes in pond environment (water quality) - Increase depth of pond. Reduce application of organic manure and supplementary feeds
- Health and Disease management- Apply lime @ 50 kg/ha

b) During the event

Aquaculture

- Shallow water in ponds due to insufficient rains/inflow- Reduce the stocking density from 25000 fry (1inches size) to 10000-15000/ha, fingerling 6,000-8,000 no/ha. Check the availability of natural food, if it is not sufficient provide supplementary feed at fixed place, time, amount and ratio & if it is more greenish stop supplementary feed & manure, store manure in separate place for agricultural purpose. Check the growth & health status by regular netting, Apply lime @ 50kg/ha.
- Impact of salt load build up in ponds / change in water quality- Apply lime @ 50 kg/ha on every 15-30 days. Aerate the water as per need

Heat wave and cold wave

- Changes in pond environment (water quality) - Stop or reduce supplementary feed and manure, Remove bigger size fishes. Reduce/stop application of feed and fertilizer.
- Health and Disease management- Apply lime/salt as per need

c) After the event

Aquaculture

- Shallow water in ponds due to insufficient rains/inflow- Remove the bigger size fishes (0.5kg). In winter season fish reduces feed consumption so reduce supplementary feed, duck start egg laying so they should not allow before 9'oclock otherwise loss of egg is possible, pig may attain 50 – 60 kg so that can be sell out and again stock same no of piglets. Apply bleaching powder @ 10kg/ha at place of litter deposition.
- Impact of salt load build up in ponds / change in water quality- Apply lime as per need @ 50 kg/ha

Heat wave and cold wave

- Changes in pond environment (water quality) - Stop or reduce supplementary feed and manure, Remove bigger size fishes. Harvest the bigger fishes, Reduce/stop application of supplementary feed, Apply lime @ 50 kg/ha and potassium per magnet in perforated plastic ball- 5-10g in each ball
- Health and Disease management- Apply lime/salt as per need

Contributors of DACPs



- **Dr. A. Wadood** (Chairman AMES and Nodal Officer, BAU Contingency Cell) 9431371693
- **Dr. D. N. Singh**(Director Research & Chairman Contingency Cell, BAU) 9430362061
- **Dr. Ramesh Kumar** (University Professor, AMES) 9431353240
- **Dr. Pragyan Kumari** (Assistant Professor, AMES) 9431389204
- **Sri Binod Kumar** (Research Associate, GKMS, Darisai) 9334729740
- **Sri Sanjiv Kumar** (Research Associate, GKMS, Ranchi) 9431354072
- **Dr. BK Agarwal** (University Professor, Department of Soil Science & Agriculture Chemistry)... 9431222937
- **Dr. PB Saha** (ZRS Dumka) 9934525212
- **Dr. MK Barnwal** (ZRS Darisai)..... 9431358373
- **Dr. Sudhir Kumar Jha** (KVK Garhwa) 9431381183
- **Dr. Shankar Kumar Singh** (KVK Lohardagga)..... 9431100641
- **Dr. Ranjay Kumar Singh** (KVK Chatra) 9431339380
- **Dr. Pramod Kumar** (KVK Jagannathpur)..... 9608096505
- **Dr. Devkant Prasad** (KVK Giridih)..... 9431411359
- **Dr. Aarti Beena Ekka** (KVK Darisai)..... 9709010792
- **Dr. Amrit Kumar Jha** (KVK Sahebganj) 9835512202
- **Sri LK Das** (KVK Palamau)..... 9431507690
- **Dr. Ansar Ahmad** (Veterinary Science) 9471182660
- **Dr. Subodh Kumar Sinha** (Veterinary Science)..... 8789167101
- **Dr. Ravinder Kumar** (Veterinary Science)..... 9031978155
- **Dr. Rakesh Ranjan** (Fishery Science) 7488500810
- **Dr. Birendra Yadav** (Agroforestry)..... 9835503195
- **Dr. Majid Ansari** (ZRS Chianki) 9430355884
- **Dr. PK Singh** (University Professor, Department of Entomology) 8986721058
- **Dr. S.N. Karamakar** (University Professor, Department of Agronomy)..... 9431929915
- **Dr. Krishna Prasad** (Junitor Scientist cum Assistant Professor, Genetics and Plant Breeding) 9934199128



GRAMIN KRISHI MAUSAM SEWA

(ग्रामीण कृषि मौसम सेवा)

INDIA METEOROLOGICAL DEPARTMENT - MINISTRY OF EARTH SCIENCE (IMD-MoES)



• बिरसा कृषि विश्वविद्यालय, कांके रँची द्वारा

झारखण्ड के किसानों को दिये जा रहे अति उपयोगी सेवा

मौसम पूर्वानुमान आधारित मौसम-कृषि परामर्श सेवा



सप्ताह में दो दिन, प्रत्येक मंगलवार एवं शुक्रवार को जिलावार परामर्श बुलेटिन जारी की जाती है।

1. अगामी 5 दिनों का जिलावार मौसम पूर्वानुमान
2. मौसम के संभावित स्थिति के अनुसार कृषि सलाह बुलेटिन
3. बुलेटिन में खेतों की तैयारी, फसल व किस्म का चयन, बोआई, रोपनी एवं हर प्रकार के प्रबन्धन की जानकारी दी जाती है जो उस समय आवश्यक हो।
4. प्रत्येक मंगलवार एवं शुक्रवार के अलावा प्रत्येक दिन यदि आवश्यक हुआ तो मोबाइल मैसेज द्वारा तत्काल सलाह दी जाती है।
5. जिला स्तरीय सेवा के अलावा फिलहाल यह सेवा प्रखण्ड स्तर पर भी रँची (काँके, रातू, अनगढ़ा एवं ओरमांडी); पूर्वी सिंहभूम (बहरागोड़ा, घाटशिला, चाकुलिया एवं धालभूमगढ़) तथा दुमका (दुमका, काठिकुंड, जामा एवं जरमुंडी) जिलों के प्रखण्डों के लिये शुरू की गई है और व्हाट्सएप ग्रुप्स से भी कृषि सलाह दी जाती है।

संचार माध्यम

- समाचार पत्र
- रेडियो
- टेलीवीजन
- क्षेत्रीय अनुसंधान केन्द्र
- कृषि विज्ञान केन्द्र
- आतंका
- जनसंपर्क तथा विभिन्न वेबसाईट्स :

www.baujharkhand.org | www.bau-eagriculture.com | www.imdagrimet.gov.in | www.cropweatheroutlook.ernet.in | www.sameti.org

इस सेवा को प्राप्त करने के लिये संपर्क करें

Dr. A. Wadood, Principal Nodal Officer : 9431371693

AMFU, Ranchi- Sanjiv Kumar : 9431354072

AMFU, Dumka- Raju Linda : 9661143150

AMFU, Darisai- Binod Kumar : 9334729740

**DEPARTMENT OF AGROMETEOROLOGY AND ENVIRONMENTAL SCIENCE
BIRSA AGRICULTURAL UNIVERSITY, KANKE, RANCHI-834006**