

Autoimmune Disease Is a Barrier Problem

What the current model gets wrong, and what fixing it actually looks like.

Dustin Lee

Here's What This Is About

The standard story about autoimmune disease goes like this: your immune system gets confused. It starts attacking your own body. So the job of medicine is to calm the immune system down. That's the whole playbook, lupus, Crohn's, MS, rheumatoid arthritis, all of it. Suppress immune activity. Use drugs to turn the response down. That story is wrong at the source. Your immune system is not confused. It is doing exactly what it is supposed to do. The problem is that something it's supposed to keep *out* is getting *in*, because the physical barrier that's supposed to stop it is broken.

That barrier is called the composite epithelial barrier. It's a layered system: a mucus gel, a charged sugar coating on your cells, the cell membrane itself, interlocking seal proteins between cells, and immune patrol inside the lining. When that system fails, bacteria and food particles cross into territory where your immune system lives and works. Your immune system responds correctly to an ongoing breach. And then, here's where it gets bad, the immune response itself damages the barrier further. The loop is now running on its own. It doesn't need a trigger anymore. It feeds itself.

The Core Point

The fix is not to suppress the immune system. The fix is to repair the barrier. There is an 8-step sequence that does that. Immunosuppression can't close the loop, it just turns down the alarm while the fire keeps burning.

What Medicine Got Wrong

Immunology has been solving the wrong problem.

The standard story goes: in autoimmune disease, your immune system targets your own tissue. T-cells misfiring. Antibodies attacking self. Genetic bad luck. Fix it by suppressing the immune system. Steroids. Methotrexate. TNF-alpha blockers, drugs that jam the signal chemicals your immune system uses. B-cell depletion, wipe out the cells making the antibodies. All of these reduce immune output. None of them repair the barrier.

The problem with this model isn't that it's imprecise. It's that it's structurally backwards. The immune system in autoimmune disease is not malfunctioning. It is finding antigens, foreign particles, somewhere those antigens should not be, and it's responding proportionally. It's doing its job. The question nobody is asking with enough force is: what's causing the breach that lets those antigens in there in the first place?

The answer is a broken barrier. Specifically, the composite epithelial barrier, the physical and biochemical structure that separates the inside of your gut from the part of your body your immune system patrols. When that structure is intact, very little gets through. When it degrades, stuff crosses constantly. Your immune system responds to an ongoing structural failure, not a programming error. Chronic inflammation is not an immune mistake. It is an accurate response to an unrepaired breach.

The Analogy

Immunosuppression is welding the pressure relief valve shut. The pipe causing the overpressure is never fixed. The barrier-engineering model says: fix the pipe.

What the Barrier Actually Is

Your gut lining isn't just a wall. It's a layered system, five layers, each doing something different. The barrier's ability to keep things out is a result of all five working together. Take away two or three of them and things start getting through.

Layer 1 - The Mucus Gel

This is the outermost layer, the one in direct contact with everything sloshing around in your gut. It's made of proteins called mucins, which are heavily coated in sugar chains that make them gel up into a thick, sticky mesh. Think of it as a pre-filter. Big particles, bacteria, and antigens get trapped in the gel before they ever reach your cells. The inner part of this gel is normally completely bacteria-free. When the gel thins out, bacteria touch your cell surface directly. That's a problem.

Layer 2 - The Glycocalyx (the charged sugar coating)

Every gut cell has a dense brush of negatively charged sugar chains sticking out from its surface. That negative charge electrostatically repels most bacterial products, including LPS, the molecule that makes gram-negative bacteria inflammatory, along with microbial proteins and enzymes called proteases. It's like a force field made of electrical charge. When this layer gets stripped, by certain bacterial enzymes or by inflammation, antigens get right up to your cell surface regardless of whether the mucus is still doing its job.

Layer 3 - The Cell Membrane

This is the actual surface of your gut cell. It's selectively permeable by design, it lets specific things through via dedicated transport proteins, but most antigens and bacterial products can't pass through it because the membrane is made of fat and those antigens are water-soluble. Oil and water don't mix. When the membrane is disrupted, by surfactants, certain fats at the wrong concentration, or oxidative damage, it loses that selectivity and starts letting things through that it shouldn't.

Layer 4 - The Tight Junctions

The gaps between adjacent cells are sealed shut by a network of interlocking proteins called claudins, occludin, and ZO proteins. This is the seal that stops things from leaking between cells instead of through them. This is what people usually mean when they say “leaky gut.” But leaky gut is not just a tight junction problem, the whole five-layer system matters. These junctions are held shut by the cell’s internal skeleton. When inflammatory signals ramp up, the skeleton contracts too hard, pulls the junction proteins apart, and creates gaps. Those gaps are the leak.

Layer 5 - The Immune Patrol (sIgA and dendritic cells)

Secretory IgA, call it sIgA, is an antibody that gets exported into the mucus layer and coats antigens before they reach your cells. It’s immune exclusion happening in the gut cavity itself, not in the tissue. Dendritic cells, immune scouts, extend arms between the gut cells to sample what’s in the gut without letting it actually cross. This is the last line before your immune system gets activated. If this layer fails, activation is all that’s left.

Five Ways the Barrier Breaks

The barrier doesn’t fail randomly. There are five specific failure modes. Know them and you know what you’re fixing.

The Seal Proteins Dissolve

The tight junction proteins degrade or get actively dissolved. A molecule called zonulin, released by gliadin (the protein in gluten) and by certain bacteria, directly forces the junctions open. Think of gliadin as a solvent sprayed on the seal. It also happens when the cell simply can’t make enough claudin protein because the amino acid or fatty acid supply is too low. No raw material, no seal.

The Mucus Gel Thins Out

The pre-filter disappears. Either the goblet cells that manufacture the gel can't keep up with demand, because they need the right amino acids and enough vitamin A, or the bacteria in a disrupted gut are eating the mucus faster than it's being made. Certain bacteria produce enzymes called mucinases that digest mucin directly. The pre-filter is being eaten alive.

The Electrostatic Force Field Goes Down

The charged sugar coat on your cell surface gets destroyed. Inflammation produces an enzyme called heparanase that clips the heparan sulfate chains. Certain bacteria and viruses produce an enzyme called neuraminidase that clips sialic acid residues, the parts of the coating that carry the charge. High-sugar diets and osmotic disruption also wreck it. Once the charge is gone, antigens that the glycocalyx would have repelled now reach your cell surface freely.

Cells Die Faster Than They're Replaced

Your gut lining replaces itself every three to five days. If inflammatory signals kill cells faster than the stem cells at the base of the gut lining can produce new ones, you get actual holes. Not metaphorical permeability. Literal gaps in the cell sheet. No amount of junction protein is going to seal a hole when there isn't a cell there to anchor it.

Bacterial Enzymes Actively Disassemble the Barrier

Dysbiotic bacteria, bacteria that shouldn't dominate but do, produce enzymes that directly snip the junction proteins from the outside. The gut normally makes inhibitors of these enzymes to fight back. But when the bacterial disruption is bad enough, the enzymes win. The barrier is being actively disassembled in real time while you're trying to hold it together.

The Immune System Is Doing Its Job

Your immune system is a sensor-and-response machine. Its job is to detect things that shouldn't be in certain places, and respond with proportional force. The space just beneath your gut lining, called the lamina propria, is exactly such a place. Immune cells there are literally waiting to respond. That's what they're there for.

When barrier failure lets antigens cross into that space, those immune cells do their job. They see LPS, flagellin, bacterial DNA, things called PAMPs, which stands for pathogen-associated molecular patterns, meaning molecular shapes that mark something as foreign, and they pull the alarm cord. TNF-alpha, IL-6, IL-17, IL-1-beta pour out. Full inflammatory cascade. Exactly as designed.

Here's where the "autoimmune" part comes in. When this keeps happening, when the barrier stays broken and antigens keep crossing for months, your adaptive immune system starts learning. It makes antibodies and T-cells specifically targeting the antigens that keep showing up. That's normal learning. The problem is that some bacterial antigens look like human tissue. Your joint proteins. Your myelin, the insulation on nerve fibers. Your thyroid. These share structural similarities with certain microbial peptides, small protein fragments from bacteria.

So your immune system, doing a perfectly rational job, ends up attacking your own tissue. Not because it's confused. Because a bacterial molecule that looks like your knee joint kept crossing into its territory for months, and it learned to treat that molecular shape as a threat. It was right about the threat. It was wrong about which thing in your body was the threat.

The Loop That Never Stops

Here's the mechanism that makes autoimmune disease self-sustaining. This is the part that explains why it doesn't just go away on its own.

The barrier breaks, for any of the five reasons above. Antigens cross. Immune sensors detect them and trigger a release of cytokines, the chemical alarm signals. Those cytokines act directly back on the barrier. TNF-alpha kills gut cells and pulls tight junctions open. IL-17 turns off claudin production, so the seal proteins stop being made. IL-1-beta stresses the cells further. The barrier degrades more. More antigens cross. The sensors fire again. The loop closes and becomes self-sustaining.

[Barrier Failure]

↓

[Antigens Cross]

↓

[Immune Sensors Activate]

↓

[Cytokines Released]

↓

[Cytokines Damage the Barrier]

↓

[Back to Barrier Failure]

Immunosuppression acts between nodes 3 and 4 only. It reduces cytokine output. It does not fix the barrier, nodes 1 and 2. So antigen keeps crossing. Sensors keep firing at reduced level. Cytokines keep damaging the barrier at a reduced rate. The loop is still running. It's just quieter.

When you stop the drug, the full loop reasserts immediately, because the barrier was never fixed. The situation driving the immune response was always still there. This is why autoimmune disease comes back when you stop immunosuppressants. It was never resolved. It was just turned down.

The 8-Step Fix

There is a specific order to fixing this. The order matters because each step depends on the previous one. Skip steps or do them out of order and you get predictable failure, not because the approach is wrong, but because you violated a dependency. Here is the sequence.

Step 1, Stop Punching Holes in the Barrier

You cannot repair something while you are still actively destroying it. Before anything else, identify and remove the inputs that are breaking the barrier right now.

- Gliadin, the protein in gluten, triggers zonulin release, which chemically forces junctions open.
- Alcohol directly dissolves the lipid structure of the cell membrane.
- NSAIDs, common pain relievers, wipe out the prostaglandin system that protects the mucus layer.
- Dysbiotic bacteria maintain a constant protease load that eats junction proteins.

This step is non-negotiable. Everything in steps 2 through 8 gets neutralized if step 1 is skipped.

Skip risk: You're filling a bucket with the tap still running. Nothing downstream works.

Step 2, Reduce the Bacterial Enzyme Attack

Dysbiotic bacteria are producing enzymes, metalloproteases, serine proteases, cysteine proteases, that clip the junction proteins from the outside. Your gut normally makes inhibitors of these enzymes; the main one is called SLPI. In chronic inflammation, SLPI levels drop. The enzymes win.

Fix this by shifting the microbial community away from the protease-producing species and by supporting SLPI levels. Certain probiotics, *Lactobacillus rhamnosus* GG, *Akkermansia muciniphila*, directly compete with protease-producing bacteria and protect the junction proteins.

Skip risk: Everything you build in steps 5 through 7 gets enzymatically shredded from the outside before it has a chance to work.

Step 3, Rebuild the Mucus Gel

The mucus is your first line of defense. If it's thin, bacteria and antigens are in direct contact with your cells regardless of how good your tight junctions are. The gel has to come first.

- The mucus backbone, a protein called MUC2, is built from threonine, which is the most quickly depleted amino acid during high mucus turnover. Supply it.
- N-acetylcysteine provides cysteine for the disulfide crosslinks that give the gel its structural integrity.
- Vitamin A drives the differentiation of goblet cells, the specific cells that manufacture mucus. Without enough vitamin A, you simply have fewer mucus-making cells.
- Zinc is needed for the enzymes that attach sugar chains to the mucin backbone, which is what makes it gel up properly.

Skip risk: Antigens contact your cell surface directly at full rate regardless of everything else you do downstream.

Step 4, Rebuild the Electrostatic Force Field

The charged sugar coating on your cell surface needs to be restored. Its charge comes from sulfate groups and sialic acid residues, specific chemical attachments that carry negative charge and repel the bad stuff.

- Glucosamine and chondroitin sulfate provide the raw material for the sulfated proteoglycan chains that make up the coating.
- N-acetylmannosamine provides the direct precursor for sialic acid synthesis. It bypasses the rate-limiting step in that synthesis pathway when sialic acid production is impaired, meaning your body can make it even when the normal route is jammed.

Skip risk: Small antigens, LPS fragments, small peptides, toxins, that slip through the mucus have direct access to your cells even if everything else is working.

Step 5, Supply the Raw Materials for Tight Junction Proteins

Claudins, occludin, and ZO proteins have to be continuously synthesized. The entire tight junction network is rebuilt every three to five days as cells turn over. You need a constant supply of building material.

- L-glutamine is the primary fuel source for gut cells and is also needed to build the proteins themselves. Under inflammatory conditions it gets used up so fast it becomes conditionally essential, meaning you can't make enough, you have to get it from outside.
- Glycine is a structural component of junction proteins.

- Claudins require palmitoylation to be properly inserted into the cell membrane, that means a fatty acid has to be attached to the protein. This requires saturated fat availability.
- Vitamin D₃ directly turns on the claudin and occludin genes at the level of DNA. No D₃, the genes don't run at full capacity.
- Butyrate, produced by gut bacteria fermenting dietary fiber, keeps the tight junction genes switched on by blocking the enzymes that switch them off. This is called HDAC inhibition.
- Zinc is a cofactor for ZO-1, the intracellular scaffold that holds all the junction proteins in their proper positions.

Skip risk: You have the signaling environment for junctions but no material to build them from. Nothing gets assembled.

Step 6, Activate the Assembly Signals

Having the raw materials isn't enough. The cell has to actively assemble those materials into a functional seal. That requires specific internal signaling pathways to be switched on.

- RhoA signaling creates the baseline tension in the cell's internal skeleton that pulls junction strands into contact across the gap between adjacent cells. No tension, no contact, no seal.
- PI3K/Akt signaling keeps the cells alive and suppresses excessive apoptosis, programmed cell death. You need cells alive long enough to do the work.
- Short-chain fatty acids, butyrate and propionate from fiber fermentation, activate these survival signals through receptors on gut cells.
- Epidermal growth factor, EGF, found in egg yolk and colostrum, activates the receptor that drives the downstream assembly signaling.

- Reducing the inflammatory cytokine burden is also part of this step. TNF-alpha continuously drives a kinase called MLCK that forces junctions open. Less TNF-alpha means junctions can actually close.

Skip risk: The materials pile up in the cell but don't get assembled into working junctions.

Like having lumber delivered but no crew to build.

Step 7, Make Sure There Are Enough Cells

The tight junction network is only as good as the cells holding it. If cells are dying faster than stem cells at the crypt, the base of the gut lining, can produce new ones, you get gaps. Actual holes in the cell sheet. No amount of junction protein seals a hole when there isn't a cell there to anchor it to.

- IGF-1, supported by adequate protein and zinc intake, drives stem cell proliferation at the crypt.
- EGF drives replication of the transit-amplifying cells that fill the villus, the finger-like projections of your gut lining that do most of the work.
- CoQ10 and riboflavin support the mitochondria that power cell division. Dividing fast is energetically expensive. You need power.
- NAC, N-acetylcysteine, reduces ER stress, which is a major driver of excessive gut cell death. The endoplasmic reticulum is the cell's protein-making factory; when it's overloaded and stressed, the cell kills itself.
- Vitamin A regulates stem cell fate decisions so new cells differentiate into the right types, goblet cells, absorptive cells, rather than the wrong ones.

Skip risk: You're trying to seal a leaking boat with a crew too small to cover the holes.

Step 8, Establish the Ecology That Keeps the Barrier Running

The barrier is not self-sustaining. It co-evolved with a specific microbial community over hundreds of millions of years, and it depends on that community for continuous

maintenance signals it cannot produce on its own. Without the ecology, the rebuilt barrier degrades again within weeks.

- *Faecalibacterium prausnitzii* and *Roseburia intestinalis* produce butyrate from dietary fiber. Without continuous butyrate production, claudin gene expression falls within hours. Within weeks, the tight junction network degrades. This isn't theoretical, animals raised without any gut bacteria have measurably compromised tight junctions.
- *Akkermansia muciniphila* lives in the mucus layer and actively stimulates goblet cells to make more mucus. Its outer membrane protein directly activates mucus secretion. Without it, the mucus thins progressively.
- *Bifidobacterium longum* primes the immune cells in the lamina propria to produce more sIgA, the antibody coat in the mucus that neutralizes antigens before they reach your cell surface.

Feed these species: prebiotic fibers, inulin, pectin, arabinoxylan, and polyphenols like quercetin and luteolin. Introduce the keystone species directly if they're absent.

Skip risk: The reconstructed barrier is a building with no maintenance crew. It falls apart again within weeks.

The Five-Layer Filter Model

Your composite barrier works like a depth-filtration stack, same principle as a water treatment plant. Each layer catches different things. The layers together achieve a separation that no single layer could achieve on its own.

The most important thing to understand about it as a system: a single layer can fail without catastrophic system failure. The remaining layers compensate. But when two or more layers fail simultaneously, the amount of antigen reaching the immune

compartment jumps sharply, often not in a straight line. A small additional failure causes a large additional breach. That's non-linear. That's the cliff.

Here's how it plays out:

- **Mucus layer fails alone.** The glycocalyx, cell membrane, tight junctions, and sIgA still block most things. Antigen flux, the rate of antigens crossing, increases marginally. The system compensates.
- **Mucus and glycocalyx fail together.** Full antigen load hits the cell membrane and tight junctions directly. Any imperfection in those layers lets things through. The system is stressed.
- **Add tight junction failure from dysbiosis-driven protease cleavage.** Now only the cell membrane and sIgA stand between the gut cavity and the immune compartment. Antigen flux is substantial and chronic. The loop activates.

This Is Autoimmune Disease Under This Model

Simultaneous failure of two or more barrier layers, producing enough antigen flux to trigger persistent immune activation. Not an immune error. A filter failure.

This also explains why partial interventions produce partial improvement but not resolution. Fix one layer, antigen flux drops but doesn't go to zero. The loop continues at reduced intensity. Symptoms improve. Disease continues. Full resolution requires all five layers to be functional simultaneously.

Why Suppressing the Immune System Doesn't Work Long-Term

Immunosuppression acts on nodes 3 and 4 of a 5-node loop. It reduces cytokine output and receptor activation. It does not touch nodes 1 and 2. Antigen keeps crossing. The

sensors keep detecting it. The cytokines keep damaging the barrier, at a reduced rate, but they keep going.

When you stop the drug, the full signal returns immediately. Because the barrier was never fixed. The situation that was driving the immune response was always still there, ticking along underneath the suppression.

There's an additional cost that doesn't get talked about enough. Long-term immune suppression also suppresses sIgA production and dendritic cell surveillance, that's Layer 5 of the barrier, the last line of luminal defense. So while you're suppressing the immune response to the ongoing breach, you're also degrading the last-layer defense of the barrier itself. The longer you suppress, the harder structural remission becomes. You're digging out of a deeper hole.

Why Diet Helps But Almost Never Finishes the Job

Dietary interventions produce real results, sometimes dramatic results. Under this model, that's entirely expected. Diet affects multiple steps of the reconstruction sequence at once. But diet almost never addresses all eight steps simultaneously, and the gaps explain why people plateau.

Take gluten elimination. It stops zonulin-mediated junction opening, part of step 1. But it does not rebuild the mucus gel (step 3), restore the glycocalyx charge (step 4), supply junction protein building blocks (step 5), activate assembly signaling (step 6), restore cell replication rate (step 7), or fix the microbial ecology (step 8). So the patient improves, genuinely, but the barrier remains partially failed. Other failure modes are still running.

"Diet works remarkably for autoimmune disease" and "diet doesn't work for autoimmune disease" are both true, about different patients with different failure mode distributions. The barrier model explains why. Patients who are deficient in the substrates a given diet

addresses will respond. Patients whose primary failure mode lies elsewhere won't. Neither group is wrong. They're just dealing with different broken layers.

The reconstruction sequence has to be followed in order. Supplying junction protein building blocks in step 5 before the protease load is controlled in step 2 means those proteins get enzymatically clipped as fast as they're made. You're building with one hand and the bacteria are tearing it down with the other. The order matters.

The Microbiome Is Infrastructure, Not a Supplement

People treat the microbiome as something you add. A probiotic capsule. A course of supplements. You take it for a few weeks and see what happens. This is the wrong frame entirely.

The microbiome is the ongoing maintenance infrastructure of the composite barrier. Your gut co-evolved with these bacterial communities over hundreds of millions of years. The barrier depends on them for specific molecular outputs it cannot produce itself. This isn't a nice-to-have. It's structural.

- **Butyrate** is produced continuously by specific bacterial species from fiber fermentation. It keeps claudin genes switched on. Stop the butyrate and claudin expression falls within hours. The tight junction network degrades within weeks. This happens in germ-free animals, animals raised without any gut bacteria. It's not hypothetical.
- **Akkermansia** physically lives in the mucus layer and actively drives mucus production. Without it, the mucus thins progressively. The pre-filter slowly disappears.
- **sIgA production** depends on bacterial interactions to reach adequate levels. Without sufficient sIgA, luminal antigen adhesion increases and the last-layer immune defense disappears.

If you do everything right in steps 1 through 7 but don't establish a stable microbial ecology, the barrier will fail again within weeks. There's no maintenance crew. The reconstructed barrier is a building without anyone keeping it up. It deteriorates. That's what buildings without maintenance do.

The goal of step 8 is not to create a permanent dependency on supplements. It's to establish an ecology that maintains itself, that produces its own butyrate, its own mucus signals, its own sIgA support. Self-sustaining. That's the end state.

The Classification System Is Built on the Wrong Thing

Current autoimmune disease names, lupus, Crohn's, MS, rheumatoid arthritis, Hashimoto's, describe the downstream immune phenotype. They tell you which tissues are being damaged and which cytokines dominate. They don't say where the problem started.

Two patients with "Crohn's disease" may have entirely different barrier failure profiles. One has primary tight junction dissolution from dysbiosis-driven proteases. The other has mucus layer depletion from goblet cell failure and vitamin A deficiency. Same diagnosis. Different failure modes. Different reconstruction requirements. The current system lumps them together and gives them the same drugs.

A barrier-engineering classification would categorize by: where the barrier failed, gut, respiratory tract, skin, blood-brain barrier; which layers failed and how; and what the microbial ecology looks like. That maps directly to a treatment protocol. The name tells you what to do.

Current names will stay useful as shorthand. They describe what the immune system is doing. But as guides to where the problem came from and how to fix it, they are misleading.

To Close

The claim is simple and specific: autoimmune disease is a barrier failure. The immune system is not broken. It's doing its job against an ongoing structural breach. The chronicity of these diseases is not immune dysfunction. It is the correct behavior of a correctly-functioning system responding to an unresolved structural failure. That distinction matters enormously for how you treat it.

The five-layer composite barrier is the actual site of pathology. When it fails across multiple layers simultaneously, chronic antigen crossing happens. The immune system responds. Its response damages the barrier more. The loop self-sustains. You get a disease that looks like it's about the immune system, because the immune system is the loudest part of the loop, but is actually about the barrier.

The 8-step Barrier Reconstruction Sequence addresses the failure in dependency order. Each step targets a specific failure mode. The order matters, dependencies are real, and skipping steps produces predictable failure. Completing the sequence in order produces a mechanistically intact barrier. That is what structural remission looks like.

Immunosuppression addresses the alarm, not the fire. When you stop it, the alarm resumes, because the fire was never put out.

The Bottom Line

A framework built on barrier physics will outperform one built on immune phenotyping when the objective is actual structural remission, not just symptom management.

Phenotyping tells you what the immune system is doing in response to a structural failure. Barrier physics tells you where the failure is and how to fix it. Those are different questions, and only one of them has an answer that ends the disease

APPENDIX

Quick Reference, What Does What

Every substance and bacterial species mentioned in this document, in one place. Plain English only.

Substance or Species	Which Barrier Layer It Targets	What It Actually Does	Which Step
L-Glutamine	Tight junctions; cell survival	Primary fuel for gut cells; also used as raw material to build junction proteins. Gets depleted fast during inflammation, you can't make enough, so you have to supply it.	Step 5
Zinc	Mucus gel; tight junctions; cell replication	Needed for the enzymes that attach sugar chains to mucin (mucus structure), for ZO-1 (the scaffold that holds junction proteins in place), and for supporting IGF-1-driven cell division.	Steps 3, 5, 7
Vitamin D ₃	Tight junctions	Directly switches on the claudin and occludin genes at the DNA level. Without enough D ₃ , the genes that make your seal proteins don't run at full output.	Step 5
Butyrate	Tight junctions; cell survival signals	Keeps tight junction genes switched on by blocking the enzymes that turn them off (HDAC inhibition). Also activates cell survival signaling through gut cell receptors. Produced continuously by specific bacteria from fiber fermentation.	Steps 5, 6, 8
N-Acetylcysteine (NAC)	Mucus gel; cell survival	Provides cysteine for the disulfide crosslinks that give the mucus gel its structural strength. Also reduces ER stress, the overload state inside cells that causes them to self-destruct.	Steps 3, 7

Substance or Species	Which Barrier Layer It Targets	What It Actually Does	Which Step
Vitamin A (Retinoic Acid)	Mucus gel; cell replication	Drives goblet cell differentiation, goblet cells are the specific cells that make mucus. Without enough vitamin A, you have fewer of them. Also regulates stem cell fate so new gut cells differentiate into the right types.	Steps 3, 7
Glucosamine	Glycocalyx (charged sugar coating)	Provides raw material for the sulfated proteoglycan chains that make up the charged surface coating on gut cells. Restoring this restores the electrostatic repulsion that blocks bacterial products.	Step 4
Chondroitin Sulfate	Glycocalyx (charged sugar coating)	Same as glucosamine, supplies the sulfated chain building blocks for the glycocalyx. Works alongside glucosamine to restore the electrical charge that repels antigens at the cell surface.	Step 4
N-Acetylmannosamine (ManNAc)	Glycocalyx (charged sugar coating)	Direct precursor for sialic acid, the specific residues on the glycocalyx that carry most of its charge. Bypasses the rate-limiting step in sialic acid synthesis when normal production is impaired.	Step 4
Threonine	Mucus gel	The amino acid backbone of MUC ₂ , the main mucus protein. It's the most quickly depleted amino acid when mucus turnover is high. You run out of it before you run out of anything else. Supply it directly.	Step 3
Glycine	Tight junctions	A structural component of junction proteins. Needed in the protein synthesis that builds the seal network during each 3-to-5-day turnover cycle.	Step 5

Substance or Species	Which Barrier Layer It Targets	What It Actually Does	Which Step
Akkermansia muciniphila	Mucus gel; protease control	Lives in the mucus layer and actively stimulates goblet cells to produce more mucus via its outer membrane protein. Competes with protease-producing bacteria. Without it, the mucus thins progressively.	Steps 2, 8
Faecalibacterium prausnitzii	Tight junctions; cell survival	Produces butyrate from fiber fermentation. This is the primary bacterial source of the butyrate that keeps claudin gene expression running. Without it, the tight junction network degrades within weeks.	Step 8
Lactobacillus rhamnosus GG	Tight junctions; protease control	Competes with protease-producing pathobionts, the bacteria actively disassembling junction proteins. Directly protects junction protein integrity. One of the most well-studied probiotics for barrier function.	Step 2
Bifidobacterium longum	Immune patrol layer (sIgA)	Primes immune cells in the lamina propria to produce more sIgA, the antibody that coats antigens in the mucus and stops them from reaching your cell surface. Restoring sIgA is restoring Layer 5 of the barrier.	Step 8
Inulin / Prebiotic Fiber	Tight junctions; mucus gel (via ecology)	Feeds the butyrate-producing bacteria, Faecalibacterium prausnitzii, Roseburia intestinalis, that maintain claudin gene expression and mucus signals. The fiber is the food supply for the maintenance crew.	Step 8
Polyphenols (Quercetin / Luteolin)	Tight junctions; ecology	Quercetin directly stabilizes tight junction proteins and inhibits the MLCK kinase that forces junctions open. Luteolin reduces inflammatory	Steps 6, 8

Substance or Species	Which Barrier Layer It Targets	What It Actually Does	Which Step
		cytokine output. Both also feed beneficial bacterial species.	
SLPI	Tight junctions (protease defense)	The gut's own natural inhibitor of the protease enzymes that dysbiotic bacteria use to clip junction proteins. In chronic inflammation, SLPI levels drop and the bacterial enzymes win. Supporting SLPI levels is part of stopping the active disassembly.	Step 2
CoQ10	Cell replication (mitochondrial support)	Supports the mitochondria, the power generators, that fuel rapid cell division. Gut lining cells divide fast. Fast division is energetically expensive. CoQ10 keeps the power on.	Step 7
Epidermal Growth Factor (EGF)	Cell replication; tight junction assembly	Activates EGFR, the receptor that drives downstream assembly signaling for tight junctions and replication of the transit-amplifying cells that fill the gut lining. Found naturally in egg yolk and colostrum.	Steps 6, 7