

Magnetism, The Shear Loop Theory of Electromagnetic Reality

A Mechanical Account of the Electromagnetic Substrate from Electron Spin to Galactic Filaments

Dustin Lee

Abstract

Magnetism is not a mysterious "second force" bolted onto electricity. It's the same substrate deformation as the electric field, just rotated 90 degrees. This paper lays out the full mechanical picture: what a magnetic field actually is (a shear loop in the electromagnetic substrate), why moving charge creates it, why field lines behave like tension loops, why iron magnetizes, how electromagnets work, how magnetic force is really shear compatibility between two deformation patterns, and why electricity and magnetism are the same thing seen from different angles. We also go out to cosmic scales and down to the biological scale. The framework presented here, the Shear Loop Theory of Electromagnetic Reality, is a theoretical interpretation built on the physical content of Maxwell's equations, special relativity, and quantum field theory, restated without the hand-waving. No mystery. No hedging. Just mechanics.

1. What Is the Substrate?

Space Is Not Empty, Get That Out of Your Head Right Now

Here's the thing that most intro physics courses get completely wrong by omission: they tell you that fields exist "in space," as if space is just an inert backdrop, an empty stage that fields happen to perform on. That's wrong. Space is the medium. The quantum vacuum, what we casually call "empty space", is a structured, physical field medium with real, measurable mechanical properties. It is not nothing. It is something, and that something has stiffness and inertia built right into its fundamental constants.

Those constants are the electric permittivity $\epsilon_0 = 8.854 \times 10^{-12}$ F/m and the magnetic permeability $\mu_0 = 4\pi \times 10^{-7}$ H/m. Think of ϵ_0 as the substrate's resistance to being electrically deformed, its stiffness against compressive or tensile strain. Think of μ_0 as its inertial resistance to magnetic (shear) deformation, how much the medium "fights back" when you try to spin it. These aren't arbitrary numbers someone cooked up. They are physical properties of the vacuum, measurable in the lab, and they tell us something deep about the structure of the medium we're operating in.

The smoking gun is this equation:

$$c = 1 / \sqrt{(\epsilon_0 \mu_0)}$$

Speed of light from vacuum mechanical properties

The speed of light, the fastest signal propagation in the universe, falls directly out of the ratio of the substrate's stiffness to its inertia. This is exactly what you get in any wave-supporting medium. The speed of sound in steel is determined by steel's elastic modulus and density. The speed of light in the vacuum is determined by ϵ_0 and μ_0 . If that doesn't tell you that space is a medium with mechanical properties, I don't know what will.

Now, here's what this is not: this is not aether theory resurrected. The 19th-century luminiferous aether was a bad model, a rigid mechanical medium with a preferred rest

frame, killed dead by Michelson-Morley and then properly buried by special relativity. The substrate described here has no preferred rest frame. It is Lorentz-invariant. Every inertial observer sees the same vacuum properties. What we're doing is taking the modern quantum field theoretic picture, in which the vacuum is a seething sea of virtual particle-antiparticle pairs, with ground-state energy density and measurable effects like the Casimir effect and the Lamb shift, and stating its mechanical implications plainly, rather than hiding them behind formalism.

Every field, electric, magnetic, gravitational, is a deformation pattern in this medium. An electric field is a particular geometry of strain. A magnetic field is a particular geometry of shear. A gravitational field, in general relativity, is spacetime curvature, literally the geometry of the medium changing. Different fields, different deformation modes, same medium. That's the picture. Keep it in your head as we go forward.

Key Concept

The quantum vacuum is not empty. It is a structured medium. ϵ_0 is its electrical stiffness. μ_0 is its magnetic inertia. $c = 1/\sqrt{(\epsilon_0\mu_0)}$ is the wave speed in that medium, the speed of light. Every electromagnetic field is a deformation of this substrate.

2. Electric Fields as Compression/Tension Deformation

Pressing a Finger Into a Gel

Put a positive point charge somewhere in the substrate. What happens? The charge deforms the medium radially outward, in every direction, symmetrically, like pressing your fingertip into the center of a ball of gel. The medium immediately surrounding the charge gets compressed. That compression propagates outward, decreasing in intensity with the square of the distance (because the same deformation energy is spreading over an ever-larger spherical surface). This is Coulomb's law, stated mechanically.

The "lines" of the electric field that you see in every textbook diagram are not mysterious invisible arrows floating in space. They are strain contours, lines of equal deformation running through the medium, pointing radially outward from the compressed region around a positive charge. The direction of the field line is the direction in which the substrate is being pushed.

A negative charge works in reverse. Instead of compressing the medium outward, it creates tension, the substrate is being pulled inward toward the charge from all directions, like a suction dimple in the gel. The field lines point inward because the deformation gradient points inward.

Why Opposite Charges Attract

Place a positive and a negative charge near each other. The positive charge is compressing the substrate outward; the negative is pulling it inward. Between them, those two deformation patterns are in direct opposition, and the medium, like any elastic material, wants to minimize its deformation energy. The lowest-energy state is when the two charges are together, because that's when the compression of the positive and the tension of the negative cancel each other out most efficiently. The medium is literally under tension between them, like a stretched rubber band, pulling them together. That tension is what we call electrostatic attraction.

Why Like Charges Repel

Put two positive charges near each other. Now you've got two outward compression patterns fighting over the same real estate between the charges. The substrate between them is being compressed from both sides simultaneously. That's a high-energy state. The medium minimizes energy by shoving the charges apart, pushing them to where each charge can deform its own territory without fighting the other. That's electrostatic repulsion. Same logic, same mechanism, opposite geometry.

This is not an analogy. The math of elastic media and the math of electrostatics are formally identical in the static case. Faraday knew this intuitively, he talked about "lines of force"

under tension and lateral pressure. Maxwell formalized it into a stress tensor. We're just calling it what it is: substrate mechanics.

3. Moving Charge Creates Perpendicular Shear

Dragging Deformation Through a Medium

This is the key section. Everything about magnetism flows from this one mechanical fact. Pay attention.

When a charge sits still, it deforms the substrate radially and symmetrically, compression/tension in all directions equally. Clean. Spherically symmetric. No preferred direction. No magnetic field. Now set the charge moving.

Here's what happens. The charge's electric deformation pattern, that radial compression field, moves with it through the substrate. It doesn't teleport; it drags. The charge pushes through the medium, and as it does, the deformation pattern has to continuously restructure itself in the medium ahead and release in the medium behind. That process of dragging a spatially extended deformation pattern through a medium creates a secondary deformation, a shear stress, perpendicular to the direction of motion.

Think about dragging your hand sideways through a bucket of thick mud. Your hand pushes mud forward and outward, sure, but there's also a circular, curling motion in the mud around your hand. The mud doesn't just pile up ahead of you; it curls around your hand in loops perpendicular to the direction you're dragging. That curling, rotational stress in the mud, that's shear. And that shear is precisely what the magnetic field is.

The Right-Hand Rule Is Just Shear Geometry

The direction of that shear, which way the loops curl, is not arbitrary. It's a direct consequence of the geometry of how a translating radial deformation creates a curl in a continuous medium. If the charge moves in the $+x$ direction, the shear loops curl around the x -axis. Point your right thumb in the direction of charge motion (conventional current

direction); your fingers curl in the direction of the magnetic field. That's the right-hand rule. It's not a memorization trick, it's the natural curl direction of the shear stress induced by translational deformation in a right-hand-consistent 3D medium. The rule is a direct physical consequence of the geometry of the shear.

The Math, Stated Plain

For a point charge q moving with velocity v , the magnetic field at a position r from the charge is given by the Biot-Savart law (in the low-velocity limit):

$$B = (\mu_0 / 4\pi) \cdot (qv \times \hat{r}) / r^2$$

Biot-Savart law for a moving point charge

Let's break that apart in plain English. $\mu_0/4\pi$ is just the substrate's magnetic inertia scaled for geometry, it sets the overall strength of the shear response. The $1/r^2$ tells you the shear intensity falls off with the square of distance, just like the electric deformation does. The key part is $v \times \hat{r}$.

v is the velocity vector, which direction the charge is moving. $\hat{r}$ is the unit vector pointing from the charge to the observation point, which direction you're looking from. The cross product $v \times \hat{r}$ gives you a vector that is perpendicular to both. It is, by construction, perpendicular to the velocity (the shear is rotated 90° from the motion) and it wraps around the direction of motion in exactly the right-hand sense.

So the equation is literally saying: the magnetic field at any point is a shear stress in the substrate, oriented perpendicular to both the charge's motion and the direction to that point, with magnitude falling off as $1/r^2$. It is the shear signature of a moving compression pattern in an elastic-like medium. That's all it is.

Current in a Wire

A wire carrying current is just a hell of a lot of moving charges, electrons drifting in one direction. Each electron contributes its own tiny shear loop to the substrate. With billions of them per cubic centimeter all moving in the same direction, those shear contributions

add up coherently. The result is a macroscopic, measurable magnetic field that wraps in cylindrical loops around the wire, exactly as you'd expect from a continuous shear stress pattern induced by translational motion through the substrate. The field circles the wire because the shear loops from every individual moving electron wrap around the direction of motion, and the net effect of all of them together is a cylindrical curl pattern.

Mechanical Analogy

A moving charge in the substrate is like dragging your hand through thick mud. The mud doesn't just displace forward, it curls in loops around your hand. Those loops are the magnetic field. The right-hand rule is just the geometry of that curl.

Why the Field Is Always Perpendicular to Velocity

One more thing worth nailing down here. The magnetic field generated by a moving charge is always perpendicular to its velocity, always. Why? Because shear stress in a continuous medium is always perpendicular to the direction of strain propagation. You can't get a shear component parallel to the motion without it being indistinguishable from the original compression/tension pattern. The medium neatly separates these two deformation modes, longitudinal (electric) and transverse (magnetic), by geometry. This is the same reason transverse and longitudinal waves in solids are distinct and don't couple except at interfaces. The geometry of the substrate enforces the perpendicularity. It's not a rule that got added to the theory; it's an inevitable consequence of the mechanics.

4. Field Lines Are Tension Loops

Why There Are No Magnetic Monopoles

Electric field lines can start and stop. A positive charge is a source, a field line begins there. A negative charge is a sink, a field line ends there. In substrate language: there are real physical locations where the medium is being compressed outward (positive charge) or sucked inward (negative charge). Those are the endpoints of strain lines. They exist because charges are real, localized objects sitting in the substrate and actively deforming it from a specific point.

Magnetic field lines are different in a fundamental way: they never start and they never stop. They always form closed loops. In substrate language, this is because magnetic fields are shear loops, rotational deformation patterns in the medium, and shear loops in a continuous medium have no source or sink. There's nowhere for a shear loop to "begin" or "end" because a shear is a circulation, not a radial strain. You can have a source of radial strain (a charge), but there's no such thing as a source of pure circulation. The curl has no divergence. This is Maxwell's second equation, $\nabla \cdot \mathbf{B} = 0$, stated mechanically: the shear deformation has no sources, so its field lines must close on themselves.

If you want a magnetic monopole, you'd need an object that is a source of pure rotational stress in the substrate without any associated translational motion, something that spins up the medium's shear from a single point, outward in all directions, like a source of curl rather than divergence. There's no known physical mechanism for that. Hence, no monopoles.

Why Field Lines Act Like Rubber Bands

Here's something Faraday noticed intuitively and that the stress tensor formulation makes rigorous: magnetic field lines behave exactly like elastic bands under tension. They want to shorten. They resist being bent. They repel each other sideways. Let's explain each of these in substrate terms.

Field lines want to shorten because they represent stored deformation energy in the substrate. A longer shear loop stores more energy. The substrate, like any elastic medium, seeks to minimize its deformation energy, so the loop naturally wants to contract to the shortest configuration consistent with the boundary conditions (i.e., consistent with the current distribution that's creating the shear). This manifests as a tension along the field line direction. Maxwell calculated this tension explicitly: it equals $B^2/(2\mu_0)$ per unit area, the magnetic pressure acting along field lines.

Why Field Lines Repel Sideways

Adjacent, parallel magnetic field lines of the same sense, meaning they wrap in the same direction, repel each other laterally. The substrate mechanics reason is elegant: if you try to push two same-sense shear loops together, their deformations add in the region between them. More deformation = more stored energy. The system minimizes energy by pushing the loops apart. Maxwell calculated a lateral pressure of $B^2/(2\mu_0)$ per unit area, equal in magnitude to the longitudinal tension, but directed outward perpendicular to the field line direction.

This is why parallel wires carrying current in the same direction attract each other (their external field lines, between the wires, are in opposite senses and cancel, releasing energy as they approach), and parallel wires carrying current in opposite directions repel (their field lines between the wires add, costing energy). We'll come back to this in Section 8. For now, understand that field line behavior is not a geometric curiosity, it's a direct readout of the substrate's attempt to minimize its deformation energy.

5. Why Magnets Have Internal Alignment Domains and Spin

Electron Spin Is a Microscopic Shear Loop

Every electron has an intrinsic angular momentum, spin, with a magnitude of $\hbar/2$, where $\hbar$ is the reduced Planck constant. Spin is one of those things quantum mechanics hands you and says "it just is what it is." But in the substrate framework, spin has a natural mechanical interpretation: it is an intrinsic, irreducible shear loop in the local substrate associated with the electron's presence. The electron isn't spinning like a ball, that picture breaks down at relativistic speeds. Instead, the electron is an excitation of the electron field that inherently carries a unit of rotational deformation in the substrate around it. A magnetic dipole moment of magnitude $\mu_B = e\hbar/(2m_e)$, the Bohr magneton, is associated with each such loop.

In most materials, electrons come in pairs with opposite spins, one spin-up, one spin-down. Their shear loops are opposite in sense and cancel. The net magnetic moment of a filled electron shell is zero. The medium around those atoms has no net shear from spin, it's a wash. These materials are diamagnetic: in an applied field, they actually develop a tiny opposing magnetization (the substrate pushes back), but the effect is weak.

Ferromagnetism: When the Loops Line Up

In ferromagnetic materials, iron, nickel, cobalt, and a handful of others, something special happens. Quantum exchange interaction, arising from the combination of the Pauli exclusion principle and Coulomb repulsion between electrons, energetically favors neighboring atoms having their unpaired electron spins aligned in the same direction. The substrate shear loops from neighboring atoms line up. They reinforce each other instead of canceling.

Why is this energetically favorable? Because aligned shear loops in adjacent regions of the substrate can share their deformation more efficiently than opposed loops. The exchange interaction is not a classical force, it's purely quantum mechanical, arising from the requirement that the total electron wavefunction be antisymmetric under particle exchange (Pauli exclusion). But its consequence in substrate language is clear: parallel spin alignment lowers the total shear deformation energy of the system.

Magnetic Domains: Competing Energy Minimizations

Here's where it gets interesting. If ferromagnetic alignment is lower energy, why doesn't the whole damn magnet just align all at once and stay that way? The answer is that there are competing energy terms. A large single-domain magnet would have a huge external magnetic field, and storing that field in the substrate outside the magnet costs energy. The system trades off exchange energy (favoring large aligned regions) against magnetostatic energy (favoring smaller regions with fields that close inside the material). The result is magnetic domains: regions within the material, typically microns to millimeters in size,

where spins are collectively aligned in one direction, but different domains point in different directions, so the total external field is partially or fully cancelled.

Domain walls are the transition regions between domains. In a domain wall, the spin direction rotates gradually from one domain's orientation to the next, typically over a few hundred atomic spacings. Moving domain walls costs energy; the wall has to drag spins through intermediate orientations that are higher in energy.

Magnetization in an External Field

When you put a piece of iron in an external magnetic field, the field tilts the energy landscape. Domains whose alignment is close to the external field direction are now in a lower-energy state; domains pointing against the field are in a higher-energy state. Domain walls move, the aligned domains grow at the expense of the misaligned ones. The substrate "votes" for the lower-energy aligned state by shifting the wall positions. If the applied field is strong enough, the material approaches single-domain alignment, that's magnetic saturation.

Permanent Magnets: Locked-In Alignment

A permanent magnet is what you get when that domain alignment gets locked in. In hard magnetic materials (alnico alloys, neodymium magnets, hard steel), domain walls are pinned, they're stuck against crystal defects, grain boundaries, and inclusions that make it energetically costly to move them. Once aligned in a strong field and then released, the walls can't relax back to a low-energy random configuration because there are too many pinning sites. The shear loops stay pointed, and you have a permanent magnet. In soft iron, the walls are free to move easily, the material magnetizes readily in a field and demagnetizes just as readily when the field is removed. Which is exactly what you want in an electromagnet core, as we'll see next.

6. Why Iron Is Magnetic And Most Stuff Isn't

Iron's Electron Configuration: The Right Cards

Iron's electron configuration is $[\text{Ar}] 3d^6 4s^2$. The argon core fills all the inner shells cleanly. The $4s^2$ gives iron its metallic bonding. The interesting action is in $3d^6$.

The $3d$ subshell has five orbitals, each of which can hold two electrons. By Hund's rule, which says electrons fill degenerate orbitals one-at-a-time with parallel spins before any pairing occurs, because parallel spins minimize electron-electron repulsion, the six $3d$ electrons in iron fill as: five orbitals get one electron each (all parallel), then the sixth electron pairs up in one of those orbitals. Net result: four unpaired electrons per iron atom, each contributing a Bohr magneton of magnetic moment. That's four uncompensated shear loops per atom. That's a significant number, more than most transition metals field at standard conditions.

The Exchange Interaction Sweet Spot

Having unpaired electrons is necessary but not sufficient for ferromagnetism. The real question is whether the quantum exchange interaction between neighboring atoms favors parallel or antiparallel spin alignment. This depends critically on the ratio of interatomic distance to the radius of the magnetic orbitals, what's sometimes called the Bethe-Slater curve.

Iron sits in a sweet spot. Its atomic spacing in the body-centered cubic (BCC) crystal structure is large enough that the exchange integral is positive, meaning parallel spin alignment between neighbors is energetically favored (ferromagnetic). This is what makes iron ferromagnetic rather than antiferromagnetic or simply paramagnetic.

Manganese is a particularly instructive case. More unpaired electrons than iron, five to iron's four, and yet it's antiferromagnetic. The shear loops are there, they're just pointing in alternating opposite directions in the crystal, canceling each other completely. Get the spacing wrong and all that magnetic potential goes to waste. Iron gets the spacing right.

The Curie Temperature

Iron's Curie temperature is 770°C (1043 K). Below this temperature, thermal energy is not enough to knock individual spin shear loops out of alignment against the exchange interaction, the loops stay ordered in domains. Above it, thermal agitation exceeds the exchange coupling energy, and the domains randomize. The long-range order collapses. Iron becomes a simple paramagnet, individual atoms still have their four unpaired shear loops, but they point every which way and average to zero. Cobalt holds its order to 1127°C and nickel to 354°C, reflecting differences in exchange coupling strength. These temperatures are set by a simple energy balance: $k_B T_C \approx J$, where J is the exchange integral energy. That's the Curie temperature in substrate terms: the temperature at which thermal agitation wins the fight against shear loop alignment.

7. How Electromagnets Work

Current in a Wire: Shear Loops on Demand

An electromagnet is the most controllable magnetic device humans have built, and the reason it works is completely transparent in the substrate picture. Run a current through a wire, and you have moving charges generating shear loops in the substrate. The loops wrap around the wire in cylindrical shells, with intensity falling off as $1/r$ from the wire's axis. One straight wire gives you a decent local field, but nothing focused.

Now bend that wire into a loop. The shear contributions from opposite sides of the loop now point in the same direction through the center of the loop, they add constructively along the axis. The field is concentrated. Bend it into another loop, stack it adjacent to the first: the contributions add again. Stack N turns of wire into a helix, a solenoid, and the shear contributions from every single turn pile up along the central axis, producing a uniform field through the interior of the coil.

Solenoid Geometry: Why It Works

Inside a solenoid, every wire loop's shear contribution threads through the same central region. Outside the solenoid, the contributions from opposite sides cancel (the field lines close on themselves outside the coil, but they're spread over a much larger volume and thus weaker). The result is a strong, nearly uniform field inside, essentially the shear loops are being focused and summed in one direction, and a weak, diffuse field outside. This is why solenoids are used in MRI machines, particle accelerators, and electromagnets: they concentrate shear in a controlled region.

The field inside an ideal solenoid is:

$$B = \mu_0 \mu_r n I$$

Field inside a solenoid with ferromagnetic core

Where n is the number of turns per unit length, I is the current, μ_0 is the vacuum permeability (the substrate's baseline magnetic inertia), and μ_r is the relative permeability of the core material. For air or vacuum, $\mu_r = 1$. For soft iron, μ_r can reach 5,000 or more. That's the iron core doing its job: responding to the applied shear field by aligning its own spin loops with the applied field, adding their contributions coherently, and multiplying the total substrate shear by a factor of up to 5,000. More turns n means more shear contributions summed along the axis. More current I means more moving charge per unit time, more shear per loop. Both linear relationships, as you'd expect from a system where contributions simply add.

Hysteresis: Soft vs. Hard Iron

Soft iron has freely movable domain walls. Apply a field, the walls shift, the domains align, you get a strong electromagnet. Kill the current, the walls relax back toward their equilibrium random configuration, the external shear that was holding them in place is gone, and the substrate inside the iron returns toward its undeformed state. The iron loses most of its magnetization. Perfect for an electromagnet you need to switch on and off.

Hard iron or steel has domain walls pinned by crystal defects and impurities. Apply a field, force the walls past their pinning sites with enough energy, the domains align, and now they stay aligned because moving those walls back requires energy that thermal agitation at room temperature can't supply. The shear loops are locked in. You've made a permanent magnet. The hysteresis loop, the plot of magnetization vs. applied field, has a wide, fat area for hard materials (high coercivity, lots of energy needed to demagnetize) and a narrow loop for soft materials (low coercivity, easy to switch). That area enclosed by the hysteresis loop is literally the energy per unit volume dissipated as heat every time you cycle the magnetization, and in transformer cores, minimizing it matters enormously.

8. Magnetic Force as Shear Compatibility

Force Is the Substrate Minimizing Energy

Here's the unifying principle for all magnetic forces: when two magnetic fields interact, the force between their sources is the substrate's attempt to reach its lowest total deformation energy state. There's no mysterious action at a distance. There's a medium between the objects, and that medium has energy stored in it, and it pushes the objects toward the configuration that minimizes that stored energy. Screw the "action at a distance" story, that was a placeholder for people who hadn't figured out field theory yet.

Parallel Currents Attract

Take two parallel wires carrying current in the same direction. Each wire generates shear loops in the substrate that wrap around it. Between the wires, the shear loop from the left wire (going, say, downward in the gap) and the shear loop from the right wire (going upward in the gap, from its opposite side) are in opposite senses, they partially cancel. Less total shear deformation in the gap means less stored energy in the gap. The substrate can lower its energy further by narrowing the gap, by pulling the wires together. The wires attract. This isn't a metaphor; this is the direction of the Maxwell stress tensor in the gap region.

Anti-Parallel Currents Repel

Reverse one wire's current. Now the shear loops between the wires are in the same sense, they add. More total deformation in the gap means more stored energy. The substrate minimizes energy by widening the gap, pushing the wires apart. The wires repel. Same medium, same mechanism, opposite result because the geometry of the shear addition flipped.

The Lorentz Force: Shear Pushing Charge

Now consider a single charged particle moving through an existing magnetic field. The force on it is the Lorentz force:

$$\mathbf{F} = q(\mathbf{v} \times \mathbf{B})$$

Lorentz force on a moving charge in a magnetic field

In substrate language: the moving charge has its own shear deformation pattern (Section 3). The external magnetic field is an existing shear pattern in the substrate. The two shear patterns interact, they either add or subtract in different regions around the charge, creating an asymmetric total deformation around the charge. The substrate pushes the charge in the direction that reduces total deformation energy, which works out to be perpendicular to both the charge's velocity and the external field direction. That perpendicularity, the cross product, is the substrate's geometry working out to where the reduction of energy lies.

The force is always perpendicular to velocity for a profound reason: shear energy is symmetric with respect to motion along the field. The charge can't extract energy from the magnetic field by moving along it, there's no energy gradient in that direction. Only perpendicular motion can lower the combined shear energy. This is why magnetic forces can change the direction of a charged particle's motion but can never change its speed: they do no work. Work requires a force component parallel to velocity. Magnetic force has none. The kinetic energy stays constant; only the velocity direction rotates. This is the principle behind cyclotrons, mass spectrometers, and the auroral displays over Earth's poles, all

situations where charged particles spiral along magnetic field lines, their speeds unchanged, their directions constantly rotated by the substrate's shear geometry.

Key Insight

Magnetic force is always perpendicular to velocity because shear energy in the substrate has no gradient along the direction of motion. This is why magnetic fields cannot do work on charges, they redirect, never accelerate. The cyclotron, the aurora, and the MRI gradient coil all rely on this single mechanical fact.

9. Electricity and Magnetism Are the Same Deformation Rotated

Maxwell's Equations in Substrate Language

Maxwell's four equations are the complete description of electromagnetic field dynamics. Let's look at the two that show the deep connection between electric and magnetic fields:

$$\partial B / \partial t = -\nabla \times E$$

Faraday's law: changing electric field drives magnetic field

$$\partial E / \partial t = c^2 \nabla \times B$$

Ampere-Maxwell law: changing magnetic field drives electric field

Plain English: a changing electric deformation in the substrate, compression/tension pattern that's shifting over time, spins up a perpendicular shear. And a changing shear pattern spins up a perpendicular compression/tension. They generate each other. Continuously. Alternately. This is how an electromagnetic wave propagates through the substrate: the electric deformation creates a magnetic shear, that shear creates a new electric deformation shifted in space and time, which creates a new shear, and so on, two deformation modes leapfrogging through the medium at speed c , exactly as you'd calculate from the substrate's mechanical properties ϵ_0 and μ_0 .

Special Relativity Seals the Deal

Here's where it gets genuinely beautiful, and where the substrate picture gets its hardest confirmation. Consider a positive charge sitting still in a lab. An observer in the lab measures a pure electric field, compression radiating outward in all directions. No magnetic field. The charge isn't moving; no shear is being dragged through the medium.

Now boost to a frame moving relative to the lab. In this new frame, the charge is moving. And in this frame, there IS a magnetic field, the shear loops we described in Section 3. Same physical charge. Same physical substrate deformation. Two different observers. Two different field readings. Where did the magnetic field "come from"?

It was there all along. The electric and magnetic fields are not two separate physical things, they are components of a single object, the electromagnetic field tensor $F_{\mu\nu}$, a 4×4 antisymmetric matrix in spacetime. When you change your velocity relative to the field source, you're rotating your perspective through spacetime, and the rotation mixes the electric and magnetic components of $F_{\mu\nu}$ exactly the way a rotation in space mixes the x and y components of a vector. The magnetic field is the electric field seen from a moving frame. They are the same substrate deformation, decomposed differently based on your velocity through spacetime.

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

Electromagnetic field tensor from the 4-potential

Written out, the tensor $F_{\mu\nu}$ looks like this: the electric field components live in the time-space cross terms ($F_{0i} = E_i/c$), and the magnetic field components live in the space-space terms ($F_{ij} = -\epsilon_{ijk}B_k$). One object. Six independent components. Three of them look like electric field to a stationary observer; three look like magnetic field. Change your velocity, and the split between those six components changes, more E, less B, or vice versa, but the total object $F_{\mu\nu}$ Lorentz-transforms as a tensor and remains invariant.

This Isn't a Metaphor, It's the Actual Math

Let me be direct: the unification of electricity and magnetism through special relativity is one of the most rigorously confirmed facts in all of physics. It's not a theoretical nicety. It has been tested to extraordinary precision. The magnetic force between two parallel current-carrying wires can be derived entirely from the electric Coulomb force plus special relativity, length contraction of the moving charge distribution changes the local charge density as seen from the frame of the other wire's electrons, and that changed charge density produces an electric force that is precisely equal to what we call the magnetic force. No new physics. Just the same physics looked at from a moving frame.

In the substrate picture, the unification is complete. The quantum vacuum has one set of mechanical properties, ϵ_0 for longitudinal deformation resistance, μ_0 for transverse shear resistance. Electric fields are longitudinal deformations of this medium. Magnetic fields are transverse shear deformations. Special relativity is the statement that the medium's behavior is the same in all inertial frames, which forces the mixing of E and B components under boosts. Maxwell's equations are the dynamical equations of the medium. The Lorentz force law tells you how sources (charges) respond to the medium's deformation state. That's the whole theory. One medium. One set of deformation modes. One unified framework.

10. Cosmic Magnetism

The Universe Is Threaded With Shear Loops at Every Scale

The substrate doesn't stop being interesting just because we've left the laboratory. Magnetic fields operate at every scale in the cosmos, from planetary dynamos to galactic filaments, and the same shear-loop mechanics applies throughout. Let's go through them.

Planetary Magnetic Fields: Convective Dynamos

Earth has a global magnetic field with a surface intensity of about 25–65 microtesla, roughly dipolar, with its axis tilted about 11° from the geographic rotation axis. Where does it come from? The outer core of the Earth is a liquid iron-nickel alloy, convecting vigorously as heat

flows outward from the inner core. Moving conductive fluid is moving charge, the free electrons in the metal drift with the bulk fluid flow. Moving charge in a conductor creates shear loops in the substrate. The rotation of the Earth, via the Coriolis effect, organizes the convective flows into helical patterns. Helical flows in a conducting fluid are a particularly efficient geometry for self-sustaining dynamo action: the shear loops from the fluid flow induce magnetic field; that field modifies the flow via the Lorentz force; the modified flow sustains the field. A self-sustaining magnetohydrodynamic dynamo.

Mars is the instructive counterexample. Mars shows remnant crustal magnetization, ancient rock formations that preserve a record of a once-active global field, but has no global field today. The reason: Mars is smaller than Earth and cooled faster. Its core solidified. No liquid iron outer core means no convective fluid flow means no moving charge means no dynamo. The shear loops stopped being generated, and the field decayed. The Martian magnetosphere collapsed, the solar wind stripped away much of the atmosphere, and Mars became the cold desert it is today. Earth's liquid iron outer core is, in a very real sense, what makes complex life on the surface possible.

Stellar Magnetic Fields: The Solar Dynamo

The Sun generates its magnetic field through a similar dynamo mechanism, but in a plasma rather than liquid metal. The Sun is not uniformly rotating: the equator rotates in about 25 days, the poles in about 35 days. This differential rotation shears magnetic field lines (stretching the substrate's shear loops), converting poloidal field (north-south loops) into toroidal field (east-west loops), a process called the Ω effect.

The critical layer is the tachocline, a thin shear zone between the rigidly rotating radiative core and the differentially rotating convective envelope, at about 70% of the solar radius. This is where the most intense dynamo action occurs. The combination of differential rotation, turbulent convection, and the Coriolis effect drives the solar 11-year activity cycle, in which magnetic field complexity grows, peaks, and then simplifies and reverses polarity.

Sunspots are where intense magnetic flux tubes, tight bundles of shear loops, break through the photosphere. The concentrated field inhibits convection (convection is the dominant heat transport mechanism near the surface; the field's shear stiffens the plasma against the convective overturn), so the region cools by about 1,500 K relative to surrounding photosphere and appears dark by contrast. Sunspot pairs are connected by arching loop structures, the flux tube breaking through the surface, arching overhead in the corona, and re-entering the surface at the conjugate spot. Solar flares and coronal mass ejections occur when these loop structures become tangled and reconnect, releasing stored substrate shear energy in violent, explosive bursts.

Pulsars: The Most Intense Magnetic Objects Known

At the extreme end of the scale are pulsars, rapidly rotating neutron stars. A neutron star is the remnant of a core-collapse supernova: roughly 1.4 solar masses compressed into a sphere about 20 km in diameter. Ordinary stellar magnetic flux is conserved during the collapse (the substrate shear loops are squeezed into a vastly smaller volume), amplifying the surface field from a typical stellar value of $\sim 10^2$ gauss to magnetar-class fields of 10^{14} – 10^{15} gauss, the most intense magnetic environments observed in the universe.

A pulsar rotates up to hundreds of times per second. The entire object, a quantum degenerate, rigid body, is rotating at this rate, and its enormous magnetic field rotates with it. The rapidly rotating magnetic dipole drives an electromagnetic wave (via the dipole radiation mechanism), losing rotational energy and gradually slowing down. The rotating field also strips charged particles from the surface and accelerates them to relativistic velocities along the field lines, producing the coherent radio beams we detect as regular pulses when the beam sweeps past Earth.

Galactic Magnetic Fields: Large-Scale Substrate Structure

Galaxies are not just stars and gas, they are permeated by large-scale coherent magnetic fields, typically of order 1–10 microgauss, threaded through the interstellar medium. These fields are coherent over kiloparsec scales, meaning the substrate shear has the same

orientation over distances of thousands of light-years. The Milky Way's field roughly follows the spiral arm structure.

These galactic fields matter enormously for astrophysics. They channel the flow of ionized gas, suppressing compression perpendicular to field lines and facilitating it along them. They regulate cosmic ray propagation, charged particles from supernova remnants spiral along galactic field lines and diffuse through the galaxy, and the field structure determines where they go and how long they stay. They influence star formation rates: magnetized gas is harder to collapse gravitationally than unmagnetized gas, because the field provides a pressure (that Maxwell stress again) opposing compression perpendicular to field lines.

The origin of galactic magnetic fields is still debated. The leading view involves a small-scale seed field, possibly primordial, from phase transitions in the early universe, or from the first generation of stars and supernovae, that was subsequently amplified by galactic dynamo action over cosmic time. Primordial magnetic fields, if they exist, would represent shear patterns locked into the substrate from the earliest moments of the universe's existence, cosmological-scale deformations present since the plasma epoch. The substrate has been structured with shear loops since very shortly after the Big Bang.

11. Biological Magnetism

Life Learned to Read the Substrate

Here's something that should give you pause: living things have been reading Earth's magnetic field, the substrate's ambient shear state, for hundreds of millions of years, long before humans figured out what magnetic fields even were. Biology got there first. And the mechanisms are, frankly, astonishing.

Magnetoreception: A Quantum Compass in a Bird's Eye

The European robin (*Erithacus rubecula*) migrates thousands of kilometers between Europe and sub-Saharan Africa, navigating with extraordinary precision. Laboratory experiments have demonstrated that robins use Earth's magnetic field as a compass,

disrupt the field in the lab and the bird gets confused; restore it and the bird orients correctly. They're sensitive to field inclination (the angle the field makes with horizontal) rather than polarity, they can't tell north from south by the substrate shear alone, but they can tell equatorward from poleward. This is a genuine biological magnetometer.

The leading mechanism for avian magnetic sensing is the cryptochrome hypothesis. Cryptochrome proteins in the retinal cells of migratory birds may exploit radical pair chemistry, a quantum mechanical process. When a photon is absorbed by a flavin cofactor in the cryptochrome, it triggers an electron transfer reaction that produces a radical pair: two molecules each containing an unpaired electron. The spin states of these electron pairs are quantum mechanically entangled. Here's the key: Earth's weak magnetic field (~50 microtesla) is enough to influence the rate at which the radical pair's spins convert between singlet and triplet states, because the energy splitting between those states (due to the magnetic field) is comparable to other relevant energy scales in the radical pair chemistry. The yield of chemical products from the reaction depends on this spin dynamics, and that yield difference produces a spatially varying signal in the retina that the bird's brain interprets as field direction. A quantum compass, embedded in a protein, reading substrate shear at the cellular level.

Magnetite: Mechanical Transducers in Biology

A second, completely different mechanism exists: biological magnetite. Magnetite, Fe_3O_4 , is a ferrimagnetic iron oxide with a strong permanent magnetic moment. Chains of magnetite crystals have been found in the magnetotactic bacteria that swim along magnetic field lines; in the beaks of homing pigeons, where they may act as a mechanical transducer (field-aligned crystals pulling on mechanoreceptor hair cells); in the abdomens of honeybees; and even in the ethmoid bone of the human skull.

In 2019, a study from Caltech (Wang et al.) used EEG to show measurable alpha-wave suppression in human subjects exposed to a rotating magnetic field, suggesting humans may retain some residual neural response to field changes, a vestigial magnetoreceptive circuit, possibly mediated by magnetite crystals somewhere in the nervous system. We

don't consciously "feel" the field, but the brain may still be registering it at some level we haven't learned to attend to.

Sea turtles navigate open ocean to return to the exact beach where they were hatched. Sharks detect the electric fields of prey using electroreceptors (the ampullae of Lorenzini) but also use magnetic field-induced electric potentials from swimming through the Earth's field. Salmon use magnetic imprinting during their outward migration and field-based navigation for their return. The substrate's ambient shear state is, for many animals, as real and readable a feature of the environment as light or sound.

MRI: The Most Direct Human Application

Magnetic resonance imaging is the most direct and dramatic application of magnetic physics to biology, and it works in three stages that are all pure substrate mechanics.

First, a powerful external field (1.5–7 tesla in clinical machines) aligns hydrogen nuclear spins, protons in the body's water molecules, with the field. The substrate shear physically tips the proton spin angular momentum into alignment with the applied field direction, like compass needles aligning in a field.

Second, a radiofrequency pulse at the Larmor frequency, the resonant frequency at which protons precess around the field direction, knocks them out of alignment. They absorb the RF energy and tip perpendicular to the field.

Third, as the protons precess back toward alignment (relaxation), they emit RF signals at the Larmor frequency. The key physics: the Larmor frequency $f = \gamma B_0 / (2\pi)$ depends on the local field strength B_0 . By superimposing gradient fields, small, spatially varying additions to the main field, the machine makes the precession frequency a known function of position. Decoding the frequencies of the emitted RF signals gives you the spatial distribution of protons, i.e., the water distribution in tissue, which maps anatomy with extraordinary precision.

Iron in Blood: Paramagnetism and fMRI

Hemoglobin carries iron as Fe^{2+} in the heme group. In deoxyhemoglobin (hemoglobin without oxygen), iron has four unpaired electrons and is paramagnetic, it weakly augments a local magnetic field. In oxyhemoglobin (with oxygen bound), the unpaired electrons are consumed in the binding, and the molecule becomes diamagnetic, it weakly opposes the field. This difference in magnetic susceptibility between oxygenated and deoxygenated blood is exploited by functional MRI's BOLD (Blood Oxygen Level Dependent) signal. Active brain regions consume oxygen, changing the local ratio of oxy- to deoxyhemoglobin, which changes the local magnetic susceptibility, which slightly distorts the local precession frequencies of water protons, which shows up as a signal change in the MRI image. Your brain activity is visible in an MRI scanner because of the magnetic properties of iron. The substrate shear state of blood is a readout of neural activity.

Conclusion

It Was Always Mechanics

Let's take stock. Magnetism is not mysterious. It never was. The mystery was a symptom of not asking "why" hard enough, and being satisfied with mathematical descriptions that stopped short of mechanical explanation.

The quantum vacuum is not empty space. It is a structured medium with real mechanical properties, stiffness ϵ_0 and inertia μ_0 , and the speed of light is simply the wave propagation speed in that medium. Electric fields are longitudinal deformation of that medium: compression outward from positive charges, tension inward toward negative charges. Magnetic fields are transverse shear deformation: the perpendicular curl induced when you drag a radial deformation pattern through the medium by setting a charge in motion.

Field lines are not helpful diagrams. They are physical contours of stress in the substrate, under tension along their length and in lateral pressure against adjacent lines of the same sense. They close on themselves because shear loops in a continuous medium have no

source or sink, there are no magnetic monopoles, not because of an arbitrary rule, but because there's no such thing as a source of pure curl in a continuous elastic medium.

Iron is magnetic because its 3d electrons are configured by Hund's rule and quantum exchange interaction to lock their spin-shear loops in parallel alignment, at atomic spacings that favor ferromagnetic over antiferromagnetic ordering. Electromagnets work because current-driven shear loops add coherently in a solenoid geometry, and a soft-iron core multiplies the field by aligning its own spin loops with the applied shear. Magnetic force is the substrate seeking its minimum-energy deformation state, always, at every scale.

And electricity and magnetism are not two forces. They are one deformation, one medium, sliced differently depending on your velocity through spacetime. Maxwell's equations describe the dynamics of the medium. Special relativity describes how different observers decompose the same deformation tensor into E and B components. The Lorentz force law describes how sources respond to the medium's deformation state.

The cosmos is woven through with shear loops at every scale: electron spin, atomic magnetic moments, ferromagnetic domains, Earth's dynamo, the Sun's activity cycle, pulsar magnetospheres, galactic filaments, and possibly primordial fields from the earliest moments of the universe. Living organisms evolved to read these shear patterns, in bird retinas, in magnetite crystals, in the iron of blood, hundreds of millions of years before we built our first compass.

The substrate is real. The fields are real deformations of it. The mechanics is what it is. There's no magic here, just a medium we're still learning to fully describe, behaving exactly as you'd expect a structured, wave-supporting, Lorentz-invariant elastic continuum to behave. That's magnetism. That's all of it.

References

1. Maxwell, J. C. (1865). A Dynamical Theory of the Electromagnetic Field. *Philosophical Transactions of the Royal Society of London*, 155, 459–512. The foundational paper deriving the displacement current and predicting electromagnetic waves from the mechanical properties of the "luminiferous medium."
2. Feynman, R. P., Leighton, R. B., and Sands, M. (1964). *The Feynman Lectures on Physics, Vol. II: Mainly Electromagnetism and Matter*. Addison-Wesley. Chapters 1–6 cover the field concept, Maxwell's equations, and the mechanical stress tensor with characteristic Feynman clarity.
3. Jackson, J. D. (1999). *Classical Electrodynamics* (3rd ed.). Wiley. The standard graduate-level reference for electromagnetic theory. Chapter 11 covers the covariant formulation and the electromagnetic field tensor $F_{\mu\nu}$.
4. Griffiths, D. J. (2017). *Introduction to Electrodynamics* (4th ed.). Cambridge University Press. The definitive undergraduate text. Sections 5.1–5.4 cover the Biot-Savart law; Section 10.3 covers the relativistic transformation of fields.
5. Purcell, E. M., and Morin, D. J. (2013). *Electricity and Magnetism* (3rd ed.). Cambridge University Press. Uniquely derives magnetic force from Coulomb's law plus special relativity in Chapter 5, the best direct demonstration of the electric-magnetic unification.
6. Kittel, C. (2005). *Introduction to Solid State Physics* (8th ed.). Wiley. Chapters 12–13 cover ferromagnetism, magnetic domains, exchange interaction, and the Curie temperature in detail.
7. Milonni, P. W. (1994). *The Quantum Vacuum: An Introduction to Quantum Electrodynamics*. Academic Press. Covers the physical reality of the quantum vacuum, virtual particle pairs, the Casimir effect, and the Lamb shift.
8. Ritz, T., Adem, S., and Schulten, K. (2000). A Model for Photoreceptor-Based Magnetoreception in Birds. *Biophysical Journal*, 78(2), 707–718. The original paper proposing the cryptochrome radical-pair mechanism for avian magnetic sensing.

9. Wiltschko, W., and Wiltschko, R. (2005). Magnetic Orientation and Magnetoreception in Birds and Other Animals. *Journal of Comparative Physiology A*, 191(8), 675–693. Comprehensive review of behavioral and physiological evidence for biological magnetoreception.
10. Wang, C. X., Hilburn, I. A., Wu, D. A., Mizuhara, Y., Cousté, C. P., Abrahams, J. N., Bernstein, S. A., Matani, A., Shimojo, S., and Kirschvink, J. L. (2019). Transduction of the Geomagnetic Field as Evidenced from alpha-Band Activity in the Human Brain. *eNeuro*, 6(2), ENEURO.0483-18.2019. The Caltech EEG study documenting measurable neural responses to magnetic field rotation in humans.
11. Kirschvink, J. L., Kobayashi-Kirschvink, A., and Woodford, B. J. (1992). Magnetite Biomineralization in the Human Brain. *Proceedings of the National Academy of Sciences*, 89(16), 7683–7687. Reports the discovery of magnetite crystals in the human ethmoid bone and brain tissue.
12. Vallée, J. P. (2011). The Milky Way's Magnetic Field. *New Astronomy Reviews*, 55(3–4), 23–90. Review of observational evidence for galactic magnetic field structure, strength, and coherence scale.
13. Zweibel, E. G., and Heiles, C. (1997). Magnetic Fields in Galaxies and Beyond. *Nature*, 385, 131–136. Overview of cosmic magnetic field origins, dynamo amplification, and the case for primordial seed fields.
14. Heger, A., Woosley, S. E., and Spruit, H. C. (2005). Presupernova Evolution of Differentially Rotating Massive Stars Including Magnetic Fields. *Astrophysical Journal*, 626(1), 350–363. Discusses magnetic field evolution in stellar interiors and the conservation of flux in core collapse, the origin of pulsar field strengths.