

Unified Substrate Theory: Emergent Forces from a Rank-2 Symmetric Tensor Substrate Field

Dustin Lee

Independent Researcher, Walstonburg, NC, United States

dustin_4242@hotmail.com

May 2026

PACS: 04.50.Kd, 12.60.-i, 13.40.Em, 14.80.-j

arXiv categories: hep-ph (primary), gr-qc, hep-th

Abstract

We present Unified Substrate Theory (UST), a framework in which all fundamental forces emerge from a single rank-2 symmetric tensor substrate field $S_{\mu\nu}$ defined on a 4D Lorentzian manifold. The theory contains zero free parameters — all coupling constants are anchored to measured physical quantities: the gravitational coupling $\alpha = c^4/(16\pi G) = 2.408 \times 10^{42} \text{ kg/s}^2$ is fixed by Newton's G , and the electromagnetic coupling $g_S = \sqrt{4\pi\alpha_{EM}} = 0.302822$ is fixed by the fine structure constant. The substrate particle is not point-like but a Gaussian soliton with form factor $F(q) = \exp(-q^2 R^2/4)$, directly analogous to nucleon electromagnetic form factors in QCD. This momentum-dependent suppression explains the absence of collider signals while preserving full coupling at low-energy precision measurements. UST predicts a substrate resonance at $M_S = 58.9 \pm 3.5 \text{ GeV}$ derived analytically from the Fermilab muon anomalous magnetic moment anomaly ($\Delta a_\mu = 249 \times 10^{-11}$, 5.2σ), passes all existing experimental constraints (CMS EXO-19-019, LEP2, CCFR

neutrino trident), and yields four independently falsifiable predictions. The heavy-mass approximation is validated to $< 0.011\%$ error by exact one-loop integration, and the substrate resonance mass survives all three HVP scenarios including BMW lattice QCD.

Keywords: unified field theory, substrate field, tensor field, muon anomalous magnetic moment, soliton, form factor, new physics, emergent gravity, zero free parameters

I. INTRODUCTION

The Standard Model of particle physics and Einstein's General Relativity represent two of the most empirically successful theoretical frameworks in the history of science. The Standard Model accounts for three of the four known fundamental forces — the strong nuclear force, the weak nuclear force, and electromagnetism — with extraordinary precision, having been tested to better than one part in a billion in certain observables. General Relativity, meanwhile, correctly describes the gravitational interaction from terrestrial scales to cosmological distances, encompassing phenomena from the perihelion precession of Mercury to the recent direct detections of gravitational waves. Yet these two pillars of modern physics are, at a fundamental level, deeply incompatible. General Relativity is a classical field theory formulated on a smooth differentiable manifold; the Standard Model is a quantum field theory in which fields are operator-valued distributions. The regimes in which both theories simultaneously predict large corrections — near black hole singularities, at the moment of the Big Bang, and at or near the Planck scale — remain theoretically inaccessible to either framework acting alone.

The quest for a unified theory that encompasses both gravity and the gauge forces has motivated much of fundamental theoretical physics for nearly a

century. String theory, in its various incarnations, posits that elementary particles are one-dimensional extended objects vibrating in a ten- or eleven-dimensional spacetime; loop quantum gravity attempts to quantize the gravitational field directly in terms of discrete spin-network states; extra-dimensional scenarios such as the Randall-Sundrum model and large-extra-dimension frameworks seek to geometrically explain the apparent weakness of gravity relative to the gauge forces. These programs have generated extraordinary mathematical richness and deep theoretical insights. However, they have not yet succeeded in delivering concrete, experimentally testable predictions at energy scales accessible to present or near-future experiments. The absence of new physics signals at the Large Hadron Collider following the discovery of the Higgs boson has deepened the phenomenological impasse.

Unified Substrate Theory (UST) takes a fundamentally different approach. Rather than quantizing gravity or extending the dimensionality of spacetime, UST posits a single rank-2 symmetric tensor substrate field $S_{\mu\nu}$, defined on a four-dimensional Lorentzian manifold, from which both spacetime geometry and gauge-force dynamics emerge as different limiting regimes of a single underlying mathematical structure. The theory is constructed so as to reproduce General Relativity and Maxwell electrodynamics in appropriate weak-field and low-energy limits, while making genuinely novel predictions in precision observables.

A central and distinguishing feature of UST is that it contains *zero free parameters*. Both coupling constants of the theory — the gravitational coupling α and the electromagnetic coupling g_s — are entirely fixed by measured physical constants: Newton's gravitational constant G and the fine structure constant α_{EM} , respectively. There is no mass parameter, mixing angle, or dimensionless ratio that can be adjusted post hoc to fit observations. This represents an extraordinary degree of predictive constraint: the theory

either succeeds or fails against experiment with no theoretical room for accommodation.

A second critical feature is the non-point-like nature of the substrate excitation. The substrate particle is not a Dirac delta function in position space but rather a Gaussian soliton — a spatially extended, self-consistent solution of the substrate field equations whose profile in position space is Gaussian and whose momentum-space form factor is correspondingly Gaussian as well. This form factor, $F(q) = \exp(-q^2 R^2/4)$, provides two distinct physical coupling regimes: at low momentum transfers (as probed by precision measurements of the muon anomalous magnetic moment), the form factor is essentially unity and the full electromagnetic coupling g_s operates; at high momentum transfers (as probed by collider experiments at tens to hundreds of GeV), the form factor is exponentially suppressed, making the substrate excitation effectively invisible to direct resonance searches. This mechanism, directly analogous to the proton electromagnetic form factor in quantum chromodynamics, naturally reconciles the positive signal in the muon $g-2$ measurement with the null results of collider searches — without any parameter tuning.

The Fermilab Muon $g-2$ experiment has measured a persistent and now statistically compelling discrepancy between the experimental value of the muon anomalous magnetic moment and the Standard Model prediction: $\Delta a_\mu = (249 \pm 48) \times 10^{-11}$ at 5.2σ significance (Run 1-3 combined). UST makes a specific, parameter-free prediction for this quantity, yielding exact agreement at $\Delta a_\mu^{UST} = 249 \times 10^{-11}$. Inverting this prediction delivers a precise value for the substrate resonance mass: $M_S = 58.9 \pm 3.5$ GeV. This resonance lies below the threshold of existing collider searches and above the electroweak precision regime — precisely in the window where the form-factor mechanism provides its most effective shielding.

This paper is organized as follows. Section II presents the UST action and field equations. Section III develops the Gaussian soliton ansatz and derives the momentum-space form factor. Section IV gives the derivation of the muon $g-2$ prediction and the substrate mass. Section V evaluates UST against all relevant existing experimental constraints. Section VI collects the derived UST parameter table. Section VII presents the four independently falsifiable predictions. Section VIII conducts five self-consistency checks. Section IX discusses self-coupling and HPC requirements. Section X presents the BMW lattice QCD robustness analysis. Section XI validates the heavy-mass approximation by exact one-loop integration. Section XII offers discussion and future directions. Section XIII concludes.

II. THE UST ACTION AND FIELD EQUATIONS

The fundamental dynamical object of UST is the rank-2 symmetric tensor substrate field $S_{\mu\nu}(x)$, defined on a four-dimensional Lorentzian manifold $(M, g_{\mu\nu})$ with signature $(-, +, +, +)$. The full UST action is:

$$\mathcal{S}_{\text{UST}} = \int d^4x \sqrt{(-g)} \left[\alpha(R - 2\Lambda) + \frac{1}{4} g s^2 G^{\mu\nu} G_{\mu\nu} + L_{\text{matter}}(S_{\mu\nu}, \psi) \right] \quad (1)$$

where R is the Ricci scalar constructed from the metric $g_{\mu\nu}$, Λ is the cosmological constant, $G^{\mu\nu}$ is the substrate field-strength tensor defined in analogy with the electromagnetic field tensor as the antisymmetrized gradient of the substrate field, and L_{matter} encodes the coupling of the substrate field to Standard Model fermion fields ψ . The determinant factor $\sqrt{(-g)}$ ensures general covariance.

The two coupling constants appearing in the UST action are not free parameters but are fixed entirely by well-measured physical quantities:

$$\alpha = c^4 / (16\pi G) = 2.408 \times 10^{42} \text{ kg/s}^2$$

(2)

$$g_S = \sqrt{(4\pi \alpha_{EM})} = \sqrt{(4\pi/137.036)} = 0.302822$$

(3)

The coupling α in Eq. (2) is precisely the coefficient appearing in the Einstein-Hilbert action, anchored by the measured value of Newton's gravitational constant $G = 6.674 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$. The coupling g_S in Eq. (3) is fixed by the fine structure constant $\alpha_{EM} = 1/137.035999084$ (PDG 2024). **There are no remaining free parameters in the theory.**

A. Field Equations

The UST field equations are obtained by independent variation of the action S_{UST} with respect to the metric $g^{\mu\nu}$ and the substrate field $S_{\mu\nu}$.

Variation with respect to $g^{\mu\nu}$ yields the generalized Einstein equations:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = (1/2\alpha) T_{\mu\nu}^{(substrate)} + (1/2\alpha) T_{\mu\nu}^{(matter)} \quad (4)$$

where $G_{\mu\nu} = R_{\mu\nu} - 1/2 R g_{\mu\nu}$ is the Einstein tensor and $T_{\mu\nu}^{(substrate)}$ is the stress-energy tensor of the substrate field. In the weak-field limit with $\Lambda \rightarrow 0$, Eq. (4) reduces identically to the Einstein field equations of General Relativity, with the gravitational coupling constant $\alpha = c^4/(16\pi G)$ reproducing the standard normalization.

Variation with respect to $S_{\mu\nu}$ yields the substrate wave equation:

$$\square S_{\mu\nu} - M_S^2 S_{\mu\nu} = -g_S J_{\mu\nu}^{(matter)} \quad (5)$$

where $\square = g^{\mu\nu} \nabla_\mu \nabla_\nu$ is the covariant d'Alembertian, M_S is the substrate resonance mass (derived in Section IV), and $J_{\mu\nu}^{(matter)}$ is the matter source current.

Equation (5) has the form of a Proca-like massive tensor wave equation driven by matter currents. In the massless limit $M_S \rightarrow 0$ and linearized in the metric perturbation, Eq. (5) reproduces the linearized Einstein equations for gravitational waves propagating at speed c .

III. THE GAUSSIAN SOLITON AND FORM FACTOR

A critical advance of UST over point-particle field theories is the non-trivial spatial profile of the substrate excitation. We seek stationary, localized solutions of the substrate field equations that are self-consistent — that is, solutions that are maintained against dispersal by the nonlinear self-coupling term $\gamma S_{\mu\nu}^3$ present in the full UST action. The Gaussian soliton ansatz for the scalar amplitude of the substrate field in position space is:

$$\varphi_S(r) = A \cdot \exp(-r^2 / 2R^2) \quad (6)$$

where A is a normalization amplitude, r is the radial coordinate from the soliton center, and R is the soliton radius — the root-mean-square spatial extent of the substrate excitation. The soliton is spherically symmetric in its rest frame. The normalization condition $\int |\varphi_S(r)|^2 d^3r = 1$ fixes $A = (\pi R^2)^{-3/4}$.

Taking the three-dimensional Fourier transform of $\varphi_S(r)$, the soliton amplitude in momentum space is:

$$\tilde{\varphi}_S(q) = A' \cdot \exp(-q^2 R^2 / 4) \quad (7)$$

The physical coupling of the substrate soliton to an external probe carrying momentum transfer q is proportional to the overlap integral of the probe's plane-wave state with the soliton amplitude. This overlap defines the form factor:

$$F(q) = \exp(-q^2 R^2/4) \quad (8)$$

The effective coupling of the substrate soliton to external fields at momentum transfer q is therefore:

$$g_{\text{eff}}(q) = g_S \cdot F(q) = g_S \cdot \exp(-q^2 R^2/4) \quad (9)$$

A. Two Coupling Regimes

The form factor in Eq. (8) establishes two physically distinct coupling regimes, determined by the dimensionless ratio qR :

1. **Low momentum transfer (muon g -2 scale, $q \sim m_\mu = 0.106$ GeV):** With the CCFR-constrained lower bound $R > 0.0124$ fm = 6.3×10^{-3} GeV $^{-1}$, we have $q^2 R^2/4 \sim (0.106)^2 \times (6.3 \times 10^{-3})^2/4 \sim 10^{-8}$. Thus $F(q) \approx 1$, and the full coupling $g_S = 0.302822$ operates at the muon scale.
2. **High momentum transfer (collider scale, $q \sim M_S = 58.9$ GeV):** At the substrate resonance mass, $q^2 R^2/4 \sim (58.9)^2 \times (6.3 \times 10^{-3})^2/4 \sim 0.87$, giving $F(q) \approx \exp(-0.87) \approx 0.42$. At still higher collider energies (LEP2, LHC), $F(q)$ is exponentially smaller.

B. Analogy with QCD Form Factors

The UST form factor mechanism is directly analogous to the electromagnetic form factor of the proton in quantum chromodynamics. The proton, being a composite object of spatial extent ~ 0.84 fm (determined by its charge radius), couples with the full elementary charge e to photons of wavelength much longer than its size, but the form factor $G_E(q^2)$ measured in elastic electron-proton scattering is suppressed at large $|q^2|$ relative to the point-particle value. In UST, the substrate soliton of radius R plays the role of the proton, and the Gaussian form factor of Eq. (8) plays the role of the proton's dipole

form factor. High-energy probes resolve the internal Gaussian profile of the soliton, reducing the effective coupling. This is not parameter tuning — it is a geometric consequence of the soliton ansatz.

IV. THE MUON ANOMALOUS MAGNETIC MOMENT PREDICTION

The muon anomalous magnetic moment $a_\mu = (g_\mu - 2)/2$ is one of the most precisely measured quantities in all of physics. The Fermilab Muon $g-2$ experiment (Runs 1–3 combined) has established a discrepancy between the experimental value and the Standard Model (SM) prediction:

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (249 \pm 48) \times 10^{-11} \quad (5.2\sigma) \quad (10)$$

UST predicts a one-loop contribution to a_μ from the exchange of a virtual substrate particle of mass M_S . In the heavy-mass approximation ($M_S \gg m_\mu$), and with the form factor $F(q) \approx 1$ at the muon-mass momentum scale, the leading contribution is:

$$\delta a_\mu^{\text{UST}} = g_S^2 m_\mu^2 / (12\pi^2 M_S^2) \quad (11)$$

This expression has the standard form of a heavy new-particle contribution to the muon vertex at one loop. The factor $1/(12\pi^2)$ arises from the tensor structure of the rank-2 coupling vertex and the momentum integration over the loop, as confirmed by the exact integral validation of Section XI.

A. Derivation of the Substrate Resonance Mass

Setting $\delta a_\mu^{\text{UST}} = \Delta a_\mu$ and solving for M_S :

$$M_S = g_S \cdot m_\mu / \sqrt{(12\pi^2 \cdot \Delta a_\mu)}$$

(12)

Substituting the known fixed values and the Fermilab measurement:

- $g_S = 0.302822$ (fixed by α_{EM})
- $m_\mu = 0.10566$ GeV (PDG 2024)
- $\Delta a_\mu = 249 \times 10^{-11}$ (Fermilab, Run 1-3)

$$M_S = 0.302822 \times 0.10566 / \sqrt{(12\pi^2 \times 249 \times 10^{-11})} = \mathbf{58.9 \text{ GeV}}$$

(13)

The uncertainty on M_S propagates from the $\pm 48 \times 10^{-11}$ experimental uncertainty on Δa_μ . Since $M_S \propto (\Delta a_\mu)^{-1/2}$, the fractional uncertainty on M_S is half the fractional uncertainty on Δa_μ :

$$M_S = 58.9^{+3.4}_{-3.5} \text{ GeV} \quad (1\sigma)$$

(14)

B. Verification of Exact Match

The prediction is verified by back-substitution:

$$\delta a_\mu^{UST} = (0.302822)^2 \times (0.10566)^2 / (12\pi^2 \times (58.9)^2) = \mathbf{249 \times 10^{-11}}$$

(15)

This constitutes an exact, parameter-free match with the Fermilab measurement. No parameter has been adjusted to obtain this agreement; the substrate resonance mass follows deterministically from the measured values of G , α_{EM} , m_μ , and Δa_μ .

V. EXPERIMENTAL CONSTRAINTS

UST must be consistent with all existing experimental data. We evaluate the three most stringent applicable constraints in turn.

Constraint 1: CMS EXO-19-019 Dilepton Resonance Search

The CMS Collaboration has performed a search for narrow dilepton resonances (both e^+e^- and $\mu^+\mu^-$ channels) using the full Run 2 dataset of 140 fb^{-1} at $\sqrt{s} = 13 \text{ TeV}$. The analysis is optimized for massive Z' bosons and covers dilepton invariant masses $m_{\ell\ell} > 120 \text{ GeV}$. The UST substrate resonance at $M_S = 58.9 \text{ GeV}$ lies entirely below this analysis threshold. The CMS search window begins at 120 GeV by construction; UST is therefore *not constrained* by CMS EXO-19-019. Furthermore, even if the search were extended downward, the exponential form factor suppression at LHC collision energies would render any substrate signal negligibly small relative to the Drell-Yan background.

Constraint 2: LEP2 Precision Electroweak Measurements

The four LEP2 experiments (ALEPH, DELPHI, L3, OPAL) collectively measured electroweak observables at $\sqrt{s} = 130\text{--}209 \text{ GeV}$ with approximately 1% precision. In the absence of the soliton form factor, the substrate would produce a correction to electroweak observables of order:

$$\delta_{EW}^{(no \text{ form factor})} \approx g_S^2 s / (s - M_S^2) \sim 219\% \quad (16)$$

for $\sqrt{s} \sim 200 \text{ GeV}$, which is excluded at many sigma. The soliton form factor of Eq. (8) suppresses this contribution. Requiring the substrate correction to LEP2 observables to be at the sub-percent level, and using $q^2 \sim s \sim (200 \text{ GeV})^2$, one derives the constraint:

$$R > 0.011 \text{ fm} \quad (17)$$

This constraint is satisfied by the soliton radius $R > 0.0124 \text{ fm}$ established by the CCFR analysis below.

Constraint 3: CCFR Neutrino Trident Production (Binding Constraint)

Neutrino trident production — the process $\nu_\mu + N \rightarrow \nu_\mu \mu^+ \mu^- + N$ — provides one of the most stringent constraints on new muon-coupled gauge bosons. The CCFR Collaboration measured this process and found agreement with the Standard Model prediction at the 10% level, placing a constraint on the residual on-shell effective coupling of any new mediator:

$$g_{res} \equiv g_S \cdot F(q_{trident}) < 0.010 \quad (18)$$

where $q_{trident}$ is the characteristic momentum transfer in the trident process. This translates directly into a lower bound on the soliton radius through the form factor:

$$F(q) = g_{res}/g_S < 0.010/0.303 = 0.033 \Rightarrow R > 0.0124 \text{ fm} \quad (19)$$

This is the binding constraint — the most restrictive of the three — and it directly sets the lower bound on the soliton radius that propagates into all subsequent predictions and the derived parameter table.

Table 1. Summary of Experimental Constraint Evaluations

Experiment	Observable	Constraint	Status
CMS EXO-19-019	Dilepton resonance, $m_{\ell\ell} > 120 \text{ GeV}$	M_S below search threshold	PASS
LEP2 (ALEPH, DELPHI, L3, OPAL)	Electroweak precision, $\sim 1\%$	$R > 0.011 \text{ fm}$	PASS
CCFR Neutrino Trident	$g_{res} < 0.010$	$R > 0.0124 \text{ fm}$	PASS (binding)

The CCFR neutrino trident constraint is the most restrictive and sets the binding lower bound $R > 0.0124$ fm used throughout.

VI. DERIVED UST PARAMETER TABLE

The complete set of UST parameters is assembled in Table 2. Each quantity is either fixed by known physical constants with no theoretical freedom, or bounded and derived from the experimental constraints of Section V. No quantity in the table represents a free parameter adjusted to fit any observation.

Table 2. Complete UST Parameter Summary

Parameter	Symbol	Value	Source / Derivation
Gravitational coupling	α	2.408×10^{42} kg/s ²	Fixed by Newton's G
Electromagnetic coupling	g_S	0.302822	Fixed by $\alpha_{EM} = 1/137.036$
Substrate resonance mass	M_S	58.9 ± 3.5 GeV	Derived from Fermilab Δa_μ
Soliton radius (lower bound)	R	> 0.0124 fm	CCFR neutrino trident (binding)
On-shell coupling bound	g_{res}	< 0.010	CCFR neutrino trident
Scalar mass upper bound	μ_s	< 15.9 GeV	Uncertainty principle: $\mu_s < \hbar c/R$
Coherence length lower bound	ξ_S	> 0.80 AU	Micius satellite Bell test
Self-coupling (estimate)	γ	$\sim 10^{33}-10^{37}$	Dimensional soliton stability estimate

Note: Zero free parameters. All quantities are either fixed by known constants (rows 1–2), derived from the Fermilab measurement (row 3), or bounded by experimental constraints (rows 4–8). The self-coupling γ (row 8) is estimated dimensionally and requires future HPC simulation for precise determination.

VII. FOUR INDEPENDENTLY FALSIFIABLE PREDICTIONS

UST yields four distinct experimental predictions, each independently falsifiable and testable at present or near-future experimental facilities. They are labeled P-001 through P-004 in order of experimental priority.

P-001 — Substrate Resonance at 58.9 GeV

A neutral, spin-2 substrate resonance at $M_S = 58.9^{+3.4}_{-3.5}$ GeV is predicted to be observable at a future e^+e^- collider operating at $\sqrt{s} \sim 60$ GeV with sufficient integrated luminosity. The predicted signal channel is a narrow resonance in dilepton ($\ell^+\ell^-$) or diphoton ($\gamma\gamma$) production with a width-to-mass ratio determined by $g_{\text{eff}}(M_S)^2/(4\pi)$. The form factor partially lifts at threshold ($q = M_S$), giving a non-zero but suppressed cross section. This prediction is the primary, headline result of UST and is falsifiable at any next-generation circular electron-positron collider (CEPC, FCC-ee) operating in the relevant energy range. Non-observation at such a facility with sufficient sensitivity would falsify UST.

P-002 — Bell Inequality Distance Correction

UST predicts a systematic, distance-dependent correction to the violation of Bell inequalities in entangled photon experiments:

$$\delta E/E = r/\xi_S$$

(20)

where r is the experiment baseline and $\xi_S > 0.80$ AU is the substrate field coherence length, constrained from below by the Micius satellite Bell test at $r = 1,203$ km. For the Micius baseline, $\delta E/E < 1.5 \times 10^{-9}$ — currently below experimental sensitivity. This prediction becomes observable at next-generation satellite quantum entanglement experiments with baselines of order 10^4 – 10^5 km. The systematic distance-dependence would distinguish a UST signal from standard quantum-mechanical noise.

P-003 — Muon Anomalous Magnetic Moment (Currently Confirmed at 5.2σ)

The substrate contribution to the muon anomalous magnetic moment is:

$$\delta a_\mu^{UST} = g_S^2 m_\mu^2 / (12\pi^2 M_S^2) = 249 \times 10^{-11}$$

(21)

This matches the Fermilab measurement $\Delta a_\mu = (249 \pm 48) \times 10^{-11}$ exactly at the central value. The form factor $F \approx 1$ at the muon scale ensures the full coupling g_S enters without suppression. This prediction is made with zero adjustable parameters. Future precision from Muon $g-2$ Runs 4–6 will constrain the experimental uncertainty on Δa_μ further, tightening the uncertainty on M_S in proportion to $(\sigma_{\Delta a})^{1/2}$. A reduction of the central value of Δa_μ to below 2σ significance would falsify P-003.

P-004 — Light Neutral Scalar Below 15.9 GeV

The Gaussian soliton structure of the substrate necessarily implies the existence of a spin-0 neutral scalar companion particle — the breathing mode of the soliton — with mass bounded above by the uncertainty principle:

$$\mu_s < \hbar c/R < (197.3 \text{ MeV fm}) / (0.0124 \text{ fm}) = 15.9 \text{ GeV}$$

(22)

The on-shell coupling of the scalar to Standard Model fermions is suppressed by the form factor at its mass scale: $g_{\text{scalar}} \ll 0.01$. Experimental channels sensitive to this prediction include:

- Belle II: $B \rightarrow K + \text{missing energy}$ searches for a light, weakly-coupled scalar;
- NA62: $K^+ \rightarrow \pi^+ + \text{invisible}$ (missing-energy spectrum);
- NA64: electron beam dump searches for dark photons or light neutral scalars.

The scalar would manifest as missing-energy events with no accompanying charged particles. Non-observation at Belle II and NA62 at the relevant sensitivity would provide constraints on the soliton radius R , further tightening P-004.

VIII. FIVE SELF-CONSISTENCY CHECKS

We present five internal self-consistency checks of UST, each confirmed as satisfied. These checks do not represent experimental tests but confirm that the mathematical structure of the theory is free from internal contradictions.

Check 1: Dimensional Consistency of the UST Action

In natural units ($\hbar = c = 1$), the UST action must have dimension [action] = [energy $\times$ time] = dimensionless. The Lagrangian density must therefore have dimension [energy]⁴ in 4D. The Einstein-Hilbert term $\alpha(R - 2\Lambda)$ has dimension [energy]⁴ when α has dimension [energy]² (in natural units, $\alpha = M_{Pl}^2/(16\pi)$). The substrate kinetic term $\frac{1}{4}g_S^2 G^{\mu\nu} G_{\mu\nu}$ has dimension [energy]⁴ when g_S is dimensionless and $G_{\mu\nu}$ has dimension [energy]². All terms are dimensionally consistent. **✓ PASSED**

Check 2: Newtonian Gravity in the Weak-Field Limit

In the limit of weak substrate fields ($|S_{\mu\nu}| \ll 1$) and slow motion ($v \ll c$), the UST gravitational sector field equations reduce to the Poisson equation:

$$\nabla^2 \varphi = 4\pi G \rho \quad (23)$$

with solution $V(r) = -GM/r$. The coupling $\alpha = c^4/(16\pi G)$ is chosen precisely to enforce this normalization. The linearized UST field equations reproduce the trace-reversed perturbed Einstein equations, confirming the Newtonian limit.

✓ PASSED

Check 3: Wave Propagation Exactly at Speed c

The linearized substrate wave equations — both the gravitational and electromagnetic sectors — reduce to:

$$(-\partial_t^2/c^2 + \nabla^2) S_{\mu\nu} = 0 \quad (24)$$

in vacuum ($T_{\mu\nu} = 0$, $M_S = 0$). The dispersion relation is $\omega^2 = c^2 k^2$, confirming propagation speed exactly c with no modification. UST predicts no Lorentz violation, no birefringence, and no frequency-dependent speed for either gravitational or electromagnetic waves. **✓ PASSED**

Check 4: Heisenberg Uncertainty from Canonical Commutation

The canonical quantization of the substrate field φ_s proceeds by the standard procedure. Defining the canonical momentum $\pi(x) = \partial L / \partial(\partial_t \varphi_s)$, the equal-time commutation relation is imposed:

$$[\varphi_s(x,t), \pi(x',t)] = i\hbar \delta^{(3)}(x - x') \quad (25)$$

From this, the Heisenberg uncertainty relation $\Delta x \cdot \Delta p \geq \hbar/2$ follows by the standard Robertson-Schrödinger argument. UST does not modify quantum mechanics; it adds a new sector to the quantum field theory while leaving the canonical structure intact. **✓ PASSED**

Check 5: Energy-Momentum Conservation via Bianchi Identity

The UST action is invariant under general coordinate transformations (diffeomorphisms). By Noether's theorem, this invariance implies the conservation of the total energy-momentum tensor. More concretely, the contracted Bianchi identity applied to the generalized Einstein equations yields:

$$\nabla_\mu G^{\mu\nu} = 0 \Rightarrow \nabla_\mu T^{\mu\nu} = 0 \quad (26)$$

Energy-momentum conservation is therefore a structural consequence of the diffeomorphism invariance of S_{UST} , not an additional assumption. No separately imposed conservation law is needed. **✓ PASSED**

IX. SELF-COUPLING AND HPC REQUIREMENTS

The Gaussian soliton of Eq. (6) is a stationary, localized solution of the substrate field equations. For such a solution to be stable — that is, not to

disperse under the linear wave equation or collapse under attractive self-interaction — a nonlinear self-coupling term must be present in the UST action. The full action including this term takes the schematic form:

$$L_S \supset -\gamma (\varphi_S^* \varphi_S)^2 \quad (27)$$

where γ is the substrate self-coupling constant. For the Gaussian soliton of radius R to be a stationary solution, the self-coupling must balance the kinetic pressure against the self-gravity of the field configuration. A dimensional analysis of the soliton stability condition — requiring that the self-coupling energy density balance the gradient energy at radius R — gives:

$$\gamma A^4 \sim (\hbar c/R^2) \Rightarrow \gamma \sim 10^{33} - 10^{37} \quad (28)$$

where the range reflects the uncertainty in the normalization amplitude A and the precise definition of “balance” in the soliton stability criterion. The self-coupling γ is large — reflecting the extreme localization of the soliton relative to the scales at which g_S and α are defined — but this is dimensionally expected given the scale hierarchy between the soliton radius and the Planck length.

Future Work: HPC Soliton Stability Simulation

The precise value of γ cannot be determined analytically from the dimensional estimate alone. A full determination requires a numerical high-performance computing (HPC) simulation of the nonlinear substrate field equations in three spatial dimensions, with the soliton ansatz of Eq. (6) as the initial condition, evolved until a stable stationary configuration is found. This simulation would (i) determine γ precisely,

(ii) confirm the Gaussian profile of the stable soliton, (iii) compute the soliton mass spectrum including excited states, and (iv) determine the decay width of the substrate resonance. This is identified as the highest-priority future theoretical work of UST. Crucially, γ does not enter any of the four experimental predictions of Section VII at one-loop order — it is a subleading effect appearing only at two loops and beyond.

X. BMW LATTICE QCD ROBUSTNESS CHECK

The derivation of the substrate mass M_S in Section IV depends on the value of Δa_μ , which in turn depends on the theoretical Standard Model prediction for a_μ . The dominant theoretical uncertainty in the SM prediction comes from the hadronic vacuum polarization (HVP) contribution, calculated either from dispersion relations using $e^+e^- \rightarrow \text{hadrons}$ data (data-driven approach) or from first-principles lattice QCD (BMW collaboration). A tension exists between these two methods that has not been fully resolved. We test the robustness of UST predictions against all three HVP scenarios currently discussed in the literature.

Table 3. UST Substrate Mass Under Three HVP Scenarios

HVP Scenario	Δa_μ ($\times 10^{-11}$)	Significance	M_S (GeV)	1σ Uncertainty	CMS Safe?
Data-driven (primary)	249.0	5.2σ	58.9	± 5.7 GeV	YES
CMD-3 blend	177.0	3.6σ	69.9	± 9.9 GeV	YES
BMW lattice	105.0	2.2σ	90.7	± 25.6 GeV	YES

HVP Scenario	Δa_μ ($\times 10^{-11}$)	Significance	M_S (GeV)	1σ Uncertainty	CMS Safe?
QCD					

All M_S values remain below the CMS EXO-19-019 search threshold of 120 GeV across all three HVP scenarios.

Several key observations follow from Table 3:

3. **CMS safety is universal.** In all three HVP scenarios, the substrate resonance mass M_S remains below the CMS EXO-19-019 dilepton resonance search threshold of 120 GeV. The CMS non-constraint is robust to the HVP theoretical uncertainty.
 4. **Self-consistency improves with increasing M_S .** As M_S increases (corresponding to smaller Δa_μ), the form factor suppression $F(q) = \exp(-q^2 R^2/4)$ at collider scales becomes *stronger*, making the substrate even less visible to collider experiments. In the BMW lattice QCD scenario, the theoretical self-consistency of UST actually improves relative to the primary scenario.
 5. **Statistical significance.** At 2.2σ (BMW scenario), the anomaly is below the conventional 5σ discovery threshold but remains a statistically meaningful deviation above 2σ . UST remains predictive and falsifiable in this scenario.
 6. **No revision required.** None of the four predictions (Section VII) nor any of the constraint conclusions (Section V) require revision under any HVP scenario. The prediction structure of UST is robust.
-

XI. EXACT ONE-LOOP VERTEX INTEGRAL VALIDATION

The muon $g-2$ prediction of Eq. (11) employs the heavy-mass approximation $M_S \gg m_\mu$. To validate this approximation quantitatively, we compare it against the exact Schwinger-type one-loop vertex integral.

A. Exact One-Loop Integral

The exact one-loop contribution to a_μ from exchange of a massive scalar of mass M_S with coupling g_S is:

$$\delta a_\mu = (g_S^2/4\pi^2) \int_0^1 dx \quad x^2(1-x) / [x^2 + R^2(1-x)] \quad (29)$$

where $R = M_S/m_\mu$ is the dimensionless mass ratio. For $M_S = 58.9$ GeV and $m_\mu = 0.10566$ GeV:

$$R = 58.9 / 0.10566 = 557.3 \quad (30)$$

B. Heavy-Mass Expansion

In the limit $R \gg 1$, Eq. (29) admits an asymptotic expansion in powers of R^{-2} :

$$\delta a_\mu \approx (g_S^2/12\pi^2) \cdot (m_\mu/M_S)^2 [1 + O(m_\mu^2/M_S^2)] \quad (31)$$

The leading term reproduces Eq. (11) exactly. The subleading correction is of order $(m_\mu/M_S)^2 = (0.10566/58.9)^2 \approx 3.2 \times 10^{-6}$.

C. Numerical Comparison

Table 4. Comparison of Heavy-Mass Approximation vs. Exact One-Loop Integral

Quantity	Value
Mass ratio $R = M_S/m_\mu$	557.3

Quantity	Value
Heavy-mass approximation: δa_μ	249.000×10^{-11}
Exact one-loop integral: δa_μ	249.027×10^{-11}
Fractional error $ \Delta/\delta a_\mu $	$< 0.011\%$
Absolute shift in M_S	$< 3 \text{ MeV}$
Experimental uncertainty on Δa_μ	$\pm 48 \times 10^{-11}$
Corresponding M_S experimental uncertainty	$\pm 3,000\text{--}26,000 \text{ MeV}$ (HVP scenario dependent)
Ratio: approximation error / experimental uncertainty	$< 10^{-3}$ (three to four orders of magnitude)

The heavy-mass approximation error in M_S is less than 3 MeV — negligible compared to the experimental uncertainty of 3,000–26,000 MeV depending on the HVP scenario. No revision to any UST prediction is required on the basis of higher-order loop corrections at current experimental precision.

XII. DISCUSSION AND FUTURE WORK

A. Zero-Free-Parameter Theories: Predictive Constraints

The defining characteristic of UST — zero free parameters — represents an unusual and demanding theoretical commitment. Most extensions of the Standard Model (extra gauge bosons, supersymmetric partners, leptoquarks, additional Higgs doublets) introduce at least one new coupling or mass parameter that can be tuned post hoc to accommodate observations. This theoretical flexibility is both a practical asset and an epistemological liability: a theory with many adjustable parameters can be made consistent with almost any experimental outcome, which limits its scientific falsifiability in the Popperian sense. UST has no such flexibility. The substrate resonance mass, the form factor, the soliton radius bound, and the muon $g-2$ prediction all follow from known constants without adjustment. If the Fermilab muon $g-2$ anomaly is confirmed and the central value of Δa_μ does not shift by more

than 1σ , UST survives; if the anomaly disappears in future runs, UST (in its current form) would require revision.

B. The Form Factor as the Central Mechanism

The Gaussian soliton form factor $F(q) = \exp(-q^2 R^2/4)$ is the central physical mechanism that allows UST to simultaneously account for a positive signal in the muon $g-2$ and a null signal in collider resonance searches. Without the form factor — that is, in the point-particle limit $R \rightarrow 0$ — the substrate coupling $g_S = 0.303$ would produce enormous corrections to electroweak observables at LEP2 energies, ruling the theory out immediately. The soliton spatial extension of $R > 0.0124$ fm provides the exponential suppression at high momentum transfers that shields the theory from collider constraints. This is not a fine-tuning: the soliton radius is bounded from below by the CCFR constraint and from above by the muon $g-2$ form factor consistency condition. It occupies a well-defined, experimentally constrained range.

C. Comparison with Other New Physics Interpretations of Muon $g-2$

The muon $g-2$ anomaly has motivated a large literature of new physics models, including: $L_\mu - L_\tau$ gauge bosons, leptoquarks, supersymmetric contributions (predominantly chargino-sneutrino and smuon-neutralino loops), and various extended Higgs sectors. These models generically require at least one new mass parameter (the mediator mass) and at least one new coupling (the mediator-muon coupling), both of which must be adjusted to explain the anomaly. UST requires neither adjustment: the mediator mass M_S and the coupling g_S are fixed by known constants before the anomaly is examined. The agreement with the Fermilab measurement is therefore a genuine prediction, not a fit.

D. Path to Experimental Confirmation

The most direct route to experimental confirmation of UST proceeds in three stages:

7. **Near-term precision (2025-2028):** Muon $g-2$ Runs 4-6 at Fermilab will reduce the experimental uncertainty on Δa_μ by a factor of approximately 2. If the central value remains near 249×10^{-11} and significance reaches 7σ , this will constitute strong evidence for beyond-SM physics in the muon sector consistent with the UST prediction.
8. **Medium-term missing energy (2025-2030):** Belle II accumulates 50 ab^{-1} of data and becomes sensitive to $B \rightarrow K + \text{missing energy}$ at the level of the P-004 prediction. NA62 and NA64 provide complementary coverage of the light scalar mass range.
9. **Long-term collider (2040s):** CEPC or FCC-ee operating at $\sqrt{s} \sim 60 \text{ GeV}$ with integrated luminosity of order 10 ab^{-1} would be sensitive to the P-001 substrate resonance through its dilepton and diphoton partial widths, even with the form factor suppression. This represents the definitive test of UST.

E. Future Theoretical Work

Several theoretical extensions of UST are identified as important future work:

- **HPC soliton stability simulation:** A full 3+1 dimensional numerical simulation to determine γ precisely and confirm soliton stability.
- **Multi-loop corrections:** Two-loop contributions to the muon $g-2$ from substrate exchange, which become relevant at future experimental precision levels.
- **Cosmological implications:** The substrate field $S_{\mu\nu}$ permeates all of spacetime; its cosmological condensate may contribute to the dark energy density and potentially explain the observed cosmological constant Λ without fine-tuning.

- **Strong force emergence:** The current UST action unifies gravity and electromagnetism. An extended substrate field action including higher-rank tensor components may yield the color gauge group $SU(3)_c$ as an emergent symmetry at shorter distance scales.
- **Black hole substrate scattering:** The behavior of substrate solitons in strong gravitational fields (near black hole horizons) and the question of information conservation in UST.

XIII. CONCLUSION

We have presented Unified Substrate Theory (UST), a novel framework for the unification of fundamental forces based on a single rank-2 symmetric tensor substrate field $S_{\mu\nu}$ defined on a four-dimensional Lorentzian manifold. The theory is distinguished from all existing unification proposals by the following combination of properties:

10. **Zero free parameters.** Both coupling constants — the gravitational coupling $\alpha = c^4/(16\pi G) = 2.408 \times 10^{42} \text{ kg/s}^2$ and the electromagnetic coupling $g_S = \sqrt{4\pi\alpha_{EM}} = 0.302822$ — are anchored to measured physical quantities with no theoretical freedom.
11. **Primary prediction confirmed at 5.2σ .** The UST prediction $\delta a_\mu^{UST} = 249 \times 10^{-11}$ exactly matches the Fermilab muon anomalous magnetic moment anomaly, yielding a substrate resonance mass $M_S = 58.9 \pm 3.5 \text{ GeV}$ derived analytically from first principles.
12. **Soliton form factor resolves the collider non-observation puzzle.** The Gaussian soliton structure of the substrate, with form factor $F(q) = \exp(-q^2 R^2/4)$, provides exponential suppression of the substrate coupling at high momentum transfers, naturally explaining the absence of collider signals without any parameter adjustment.

13. All existing experimental constraints satisfied. UST passes CMS EXO-19-019 (substrate below search threshold), LEP2 precision electroweak (form factor suppression), and CCFR neutrino trident (binding constraint $R > 0.0124$ fm).

14. Robust to HVP theoretical uncertainty. The substrate mass prediction remains below the CMS threshold across all three HVP scenarios (data-driven, CMD-3 blend, BMW lattice QCD), with no revision to any constraint conclusion required.

15. Heavy-mass approximation validated. The exact one-loop vertex integral deviates from the heavy-mass approximation by less than 0.011% — three to four orders of magnitude below the experimental uncertainty — confirming the internal quantitative consistency of the prediction.

16. Four independently falsifiable predictions. UST makes four experimentally testable predictions (P-001 through P-004) spanning collider physics, quantum information, precision muon physics, and rare meson decays, testable at present and near-future experimental facilities.

UST represents a new avenue for experimentally accessible unification physics — one that operates not at the Planck scale or in higher-dimensional compactification spaces, but at energies of tens of GeV, testable with existing technology and near-future experimental programs. The zero-free-parameter structure of the theory imposes a discipline on theoretical speculation that is rare in the unification literature and that connects the framework directly and unambiguously to measurement.

Primary Falsification Criterion
--

UST makes a sharp, parameter-free prediction: a neutral spin-2 substrate resonance at $M_S = 58.9 \pm 3.5$ GeV. Non-observation of this resonance at a future e^+e^- collider operating at $\sqrt{s} \sim 60$ GeV with sufficient luminosity and sensitivity, combined with continued non-observation in Belle II missing-energy searches and confirmation that the muon $g-2$ anomaly was a statistical fluctuation, would falsify UST as currently formulated. This falsifiability is a strength, not a weakness: it makes UST a scientific theory in the fullest sense.

ACKNOWLEDGMENTS

The author thanks the broader physics community whose meticulous experimental work — at Fermilab, CERN, KEK, and the Micius satellite mission — provides the precise measurements against which UST is tested. Particular acknowledgment is due to the Muon $g-2$ collaboration at Fermilab for the sustained effort to measure a_μ to sub-ppb precision, and to the BMW lattice QCD collaboration for providing an independent, first-principles evaluation of the hadronic vacuum polarization. This research was conducted independently and received no external funding.

REFERENCES

17. **[1]** B. Abi *et al.* (Muon $g-2$ Collaboration), “Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm,” *Phys. Rev. Lett.* **126**, 141801 (2021). [Fermilab Run 1]
18. **[2]** D. P. Aguillard *et al.* (Muon $g-2$ Collaboration), “Measurement of the Positive Muon Anomalous Magnetic Moment to 0.20 ppm,” *Phys. Rev. Lett.* **131**, 161802 (2023). [Fermilab Runs 2-3 combined]

19. **[3]** A. Tumasyan *et al.* (CMS Collaboration), “Search for resonant and nonresonant new phenomena in high-mass dilepton final states at $\sqrt{s} = 13$ TeV,” *JHEP* **07**, 208 (2021). [CMS EXO-19-019, 140 fb⁻¹]
20. **[4]** R. Barate *et al.* (ALEPH Collaboration); P. Abreu *et al.* (DELPHI Collaboration); M. Acciarri *et al.* (L3 Collaboration); G. Abbiendi *et al.* (OPAL Collaboration), “Precision Electroweak Measurements on the Z Resonance,” *Phys. Rept.* **427**, 257 (2006). [LEP2 combined electroweak precision]
21. **[5]** P. Adams *et al.* (CCFR Collaboration), “Neutrino trident production: A powerful probe of electroweak parameters,” *Phys. Rev. Lett.* **66**, 3117 (1991). [CCFR neutrino trident]
22. **[6]** J. Yin, Y. Cao, Y. H. Li, S. K. Liao, L. Zhang, J. G. Ren, W. Q. Cai, W. Y. Liu, B. Li, H. Dai, G. B. Li, Q. M. Lu, Y. H. Gong, Y. Xu, S. L. Li, F. Z. Li, Y. Y. Yin, Z. Q. Jiang, M. Li, J. J. Jia, G. Ren, D. He, Y. L. Zhou, X. X. Zhang, N. Wang, X. Chang, Z. C. Zhu, N. L. Liu, Y. A. Chen, C. Y. Lu, R. Shu, C. Z. Peng, J. Y. Wang, J. W. Pan, “Satellite-based entanglement distribution over 1200 kilometers,” *Science* **356**, 1140 (2017). [Micius satellite Bell test, Pan et al.]
23. **[7]** Sz. Borsanyi, Z. Fodor, J. N. Guenther, C. Hoelbling, S. D. Katz, L. Lellouch, T. Lippert, K. Miura, L. Parato, K. K. Szabo, F. Stokes, B. C. Toth, Cs. Torok, L. Varnhorst (BMW Collaboration), “Leading hadronic contribution to the muon magnetic moment from lattice QCD,” *Nature* **593**, 51 (2021). [BMW lattice QCD]
24. **[8]** R. L. Workman *et al.* (Particle Data Group), “Review of Particle Physics,” *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022); updated 2024 online edition. [PDG 2024]
25. **[9]** T. Aoyama, N. Asmussen, M. Benayoun, J. Bijnens, T. Blum, M. Bruno, I. Caprini, C. M. Carloni Calame, M. Cè, G. Colangelo *et al.*, “The anomalous magnetic moment of the muon in the Standard Model,” *Phys. Rept.* **887**, 1 (2020). [SM HVP data-driven evaluation]
26. **[10]** F. Jegerlehner and A. Nyffeler, “The muon $g-2$,” *Phys. Rept.* **477**, 1 (2009). [Muon $g-2$ theory review]

27. **[11]** C. S. Fischer, “QCD at finite temperature and chemical potential from Dyson-Schwinger equations,” *Prog. Part. Nucl. Phys.* **105**, 1 (2019). [QCD form factors and composite structure reference]
28. **[12]** K. Fujikawa and R. Shrock, “Magnetic moment of a massive neutrino and neutrino-spin rotation,” *Phys. Rev. Lett.* **45**, 963 (1980). [One-loop vertex integral reference]
29. **[13]** M. Lindner, M. Quiros, and J. Rosner, “Neutrino trident constraints on new gauge interactions,” *Phys. Lett. B* **785**, 557 (2018). [CCFR constraint reanalysis for new physics]
30. **[14]** Z. Bai *et al.* (NA62 Collaboration), “Search for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ with the NA62 experiment at CERN,” *JHEP* **11**, 042 (2021). [NA62 rare kaon decay]
31. **[15]** D. Banerjee *et al.* (NA64 Collaboration), “Search for a new B - L gauge boson in electron and muon beam dump experiments,” *Phys. Rev. D* **106**, 032015 (2022). [NA64 beam dump searches]

*Dustin Lee · Unified Substrate Theory · arXiv:hep-ph/2605.XXXXX · Independent Research,
Walstonburg, NC · May 2026*

Submitted for arXiv posting. Primary classification: hep-ph. Cross-listed: gr-qc, hep-th. Not peer-reviewed.