An Attempt on Problem 4 from Abouzaid et al. [2026]

Unique Subedi and Gemini

subedi@umich.edu

Comments in red are my addition. Rest of the content is copied more or less directly from Gemini's response. See the chat log at https://gemini.google.com/share/3543f5fde84b.

See problems on https://1stproof.org/.

1 Problem 4 from Abouzaid et al. [2026]

Let $p(x)$ and $q(x)$ be two monic polynomials of degree n:

$$p(x) = \sum_{k=0}^{n} a_k x^{n-k}, \qquad q(x) = \sum_{k=0}^{n} b_k x^{n-k},$$

where $a_0 = b_0 = 1$. Define $p \boxplus_n q(x)$ to be the polynomial

$$(p \boxplus_n q)(x) = \sum_{k=0}^{n} c_k x^{n-k},$$

where the coefficients c_k are given by the formula

$$c_k = \sum_{i+j=k} \frac{(n-i)!(n-j)!}{n!(n-k)!} a_i b_j, \qquad k = 0, 1, \ldots, n.$$

For a monic polynomial $p(x) = \prod_{i \le n}(x - \lambda_i)$, define

$$\Phi_n(p) := \sum_{i \le n} \left(\sum_{j \ne i} \frac{1}{\lambda_i - \lambda_j} \right)^2,$$

and define $\Phi_n(p) := \infty$ if p has a multiple root.

Is it true that if $p(x)$ and $q(x)$ are monic real-rooted polynomials of degree n, then

$$\frac{1}{\Phi_n(p \boxplus_n q)} \ge \frac{1}{\Phi_n(p)} + \frac{1}{\Phi_n(q)}\,?$$

2 Warmup: Gemini Solution for $n = 2$.

Gemini's gave this solution pretty much immediately. I verified the details and it looks correct.

2.1 The Functional $\Phi_2(p)$ and the Discriminant

Let $p(x)$ be a monic polynomial of degree 2 with roots λ_1, λ_2:

$$p(x) = x^2 + a_1 x + a_2 = (x - \lambda_1)(x - \lambda_2).$$

The functional $\Phi_2(p)$ is defined as:

$$\Phi_2(p) = \sum_{i \neq j} \frac{1}{(\lambda_i - \lambda_j)^2} = \frac{1}{(\lambda_1 - \lambda_2)^2} + \frac{1}{(\lambda_2 - \lambda_1)^2} = \frac{2}{(\lambda_1 - \lambda_2)^2}.$$

Recall that the discriminant of a quadratic $p(x)$ is $\Delta_p = (\lambda_1 - \lambda_2)^2 = a_1^2 - 4a_2$.Thus, we can rewrite the functional as:

$$\Phi_2(p) = \frac{2}{\Delta_p} \implies \frac{1}{\Phi_2(p)} = \frac{\Delta_p}{2}.$$

2.2 Calculating the Convolution Coefficients

Let $q(x) = x^2 + b_1 x + b_2$ be another monic polynomial.We compute the coefficients c_k of the convolution $(p \boxplus_2 q)(x) = x^2 + c_1 x + c_2$ using the formula:

$$c_k = \sum_{i+j=k} \frac{(2-i)!(2-j)!}{2!(2-k)!} a_i b_j.$$

(Note: $a_0 = b_0 = 1$).For $k = 1$:

$$c_1 = \frac{1!2!}{2!1!} a_1 b_0 + \frac{2!1!}{2!1!} a_0 b_1 = a_1 + b_1.$$

For $k = 2$:

$$c_2 = \frac{0!2!}{2!0!} a_2 b_0 + \frac{1!1!}{2!0!} a_1 b_1 + \frac{2!0!}{2!0!} a_0 b_2 = a_2 + \frac{1}{2} a_1 b_1 + b_2.$$

Thus, the convolved polynomial is:

$$(p \boxplus_2 q)(x) = x^2 + (a_1 + b_1)x + \left(a_2 + b_2 + \frac{1}{2} a_1 b_1\right).$$

2.3 Discriminant of the Convolution

Now we calculate the discriminant of the new polynomial $p \boxplus_2 q$:

$$\Delta_{p \boxplus q} = c_1^2 - 4c_2 = (a_1 + b_1)^2 - 4\left(a_2 + b_2 + \frac{1}{2} a_1 b_1\right).$$

Expanding the terms:

$$\Delta_{p \boxplus q} = (a_1^2 + 2a_1 b_1 + b_1^2) - 4a_2 - 4b_2 - 2a_1 b_1.$$

Notice that the cross-term $2a_1 b_1$ cancels out:

$$\Delta_{p \boxplus q} = (a_1^2 - 4a_2) + (b_1^2 - 4b_2).$$

This reveals a beautiful additivity property for $n = 2$:

$$\Delta_{p \boxplus q} = \Delta_p + \Delta_q.$$

2.4 Conclusion

Substituting the discriminants back into the functional relation derived in Step 1:

$$\frac{1}{\Phi_2(p \boxplus_2 q)} = \frac{\Delta_{p\boxplus q}}{2} = \frac{\Delta_p + \Delta_q}{2} = \frac{\Delta_p}{2} + \frac{\Delta_q}{2}.$$

$$\frac{1}{\Phi_2(p \boxplus_2 q)} = \frac{1}{\Phi_2(p)} + \frac{1}{\Phi_2(q)}.$$

Thus, for the $n = 2$ case, the inequality holds as an equality.

3 Gemini's Solution for general n.

This approach initially looked promising, but there is a huge bug. The proof uses a lemma about score function for distributions with smooth density functions and applies it to distribution of unitary matrices on $n \times n$. However, the set of unitary matrices has Lebesgue measure zero in the space of $n \times n$ matrices and therefore does not have a density with respect to Lebesgue measure.

I am sharing this solution because I initially approached this problem with the idea that it might be possible to compute the roots of all the polynomials involved using a brute-force method. This quickly becomes too complicated for humans, but I was curious whether an LLM might be able to handle it. Instead, Gemini suggested a different perspective: it connected the polynomial $p \boxplus_n q$ to the (expected) characteristic polynomial of a certain random matrix and proposed using tools from probability theory and random matrix theory. I found this approach quite beautiful, because it suddenly brings a wide range of probabilistic tools into play, rather than relying on a purely brute-force algebraic computation. Even though Gemini did not produce a fully rigorous proof, the connection reframed the problem within a richer mathematical framework. Often, discovering such connections is the most challenging part of research.

3.1 General Theory of Score Functions and Convolution

We first establish a general inequality for the Fisher information of the sum of independent random variables. Let X, Y be independent random variables (or random vectors/matrices) with smooth probability densities p_X and p_Y.

Definition 1 (Score Function). Let X be a random vector with a smooth probability density function p_X. The score function Ξ_X is defined as the gradient of the log-density:

$$\Xi_X(u) := \nabla \log p_X(u) = \frac{\nabla p_X(u)}{p_X(u)}.$$

The Fisher Information $I(X)$ is the expected squared Frobenius norm of the score:

$$I(X) := \mathbb{E}[\|\Xi_X(X)\|_F^2].$$

Lemma 1 (Projection Lemma). Let X and Y be independent random vectors (or matrices). Let $Z = X + Y$. The score function of the sum is the conditional expectation of the score of the summand:

$$\Xi_Z(Z) = \mathbb{E}[\Xi_X(X) \mid Z].$$

*Proof.*Since X and Y are independent, the probability density of the sum Z, denoted p_Z, is given by the convolution of the marginal densities p_X and p_Y:

$$p_Z(z) = (p_X * p_Y)(z) = \int p_X(z - y) p_Y(y)\, dy.$$

We compute the score function $\Xi_Z(z) = \frac{\nabla p_Z(z)}{p_Z(z)}$. First, we differentiate the convolution integral with respect to z:

$$\nabla p_Z(z) = \nabla_z \int p_X(z-y)p_Y(y)\,dy = \int (\nabla p_X)(z-y)p_Y(y)\,dy.$$

Using the definition $\nabla p_X(u) = \Xi_X(u)p_X(u)$, we substitute this into the integrand:

$$\nabla p_Z(z) = \int \Xi_X(z-y)p_X(z-y)p_Y(y)\,dy.$$

Dividing both sides by $p_Z(z)$ yields the expression for the score:

$$\Xi_Z(z) = \int \Xi_X(z-y)\frac{p_X(z-y)p_Y(y)}{p_Z(z)}\,dy.$$

We now identify the fraction inside the integral. The joint density of (X, Y) is $p_{X,Y}(x, y) = p_X(x)p_Y(y)$ due to independence. The conditional density of Y given $Z = z$ (where $X = z - y$) is:

$$p_{Y|Z}(y|z) = \frac{p_{X,Y}(z-y, y)}{p_Z(z)} = \frac{p_X(z-y)p_Y(y)}{p_Z(z)}.$$

Thus, the integral can be rewritten as an expectation with respect to this conditional density:

$$\Xi_Z(z) = \int \Xi_X(z-y)p_{Y|Z}(y|z)\,dy = \mathbb{E}[\Xi_X(X) \mid Z = z]. \quad \square$$

Lemma 2 (Fisher Information Inequality). For any $t \in [0, 1]$,

$$I(X + Y) \leq t^2 I(X) + (1-t)^2 I(Y).$$

Proof. From Lemma 1, $\Xi_Z = \mathbb{E}[\Xi_X \mid Z]$. By symmetry, $\Xi_Z = \mathbb{E}[\Xi_Y \mid Z]$.We write Ξ_Z as a convex combination:

$$\Xi_Z = \mathbb{E}[t\Xi_X + (1-t)\Xi_Y \mid Z].$$

Applying Jensen's Inequality for conditional expectations ($\|\mathbb{E}[U|V]\|^2 \leq \mathbb{E}[\|U\|^2]$) to the Fisher Information $I(Z) = \mathbb{E}[\|\Xi_Z\|^2]$:

$$I(Z) \leq \mathbb{E}[\|t\Xi_X + (1-t)\Xi_Y\|^2].$$

Expanding the square:

$$= t^2\mathbb{E}\|\Xi_X\|^2 + (1-t)^2\mathbb{E}\|\Xi_Y\|^2 + 2t(1-t)\mathbb{E}\langle\Xi_X, \Xi_Y\rangle.$$

Since X and Y are independent, Ξ_X and Ξ_Y are independent. The expectation of a score function is zero ($\mathbb{E}[\Xi_X] = \int \nabla p_X = 0$). Thus, the cross-term vanishes.

$$I(Z) \leq t^2 I(X) + (1-t)^2 I(Y). \quad \square$$

3.2 Application to Polynomials and Random Matrices

We now apply the general theory to the specific random matrices defining the operation $\boxplus_n$.Let Λ_p and Λ_q be diagonal matrices containing the roots of p and q. We define the independent random matrices:

$$A := U\Lambda_p U^*, \qquad B := V\Lambda_q V^*.$$

Let $C = A + B$. The polynomial $p \boxplus_n q$ is the expected characteristic polynomial of C. (see Theorem 1.2 of Marcus et al. [2022])

Derivation of the Explicit Score Function. The random matrix A is defined as $A = U\Lambda_p U^*$, where $\Lambda_p = \mathrm{diag}(\lambda_1, \ldots, \lambda_n)$ is fixed and U is Haar-distributed. The probability distribution of A is singular, supported entirely on the unitary orbit of Λ_p. To derive the score function rigorously, we consider the volume measure on the space of Hermitian matrices $\mathcal{H}_n$.By the Weyl Integration Formula, the Lebesgue measure dM on $\mathcal{H}_n$ decomposes into radial (eigenvalue) and angular (eigenvector) components:

$$dM = C_n \cdot \underbrace{\prod_{1 \le i < j \le n} (\lambda_i - \lambda_j)^2}_{\mathcal{J}(\Lambda)} \, d\Lambda \, dU,$$

where $\mathcal{J}(\Lambda)$ is the Jacobian of the spectral map (the squared Vandermonde determinant).The score function Ξ_A is defined as the gradient of the log-probability density, $\Xi_A(M) = \nabla_M \log p(M)$. For the orbital measure, the density $p(M)$ is induced by the change of variables. The "effective potential" acting on the eigenvalues is determined by the logarithm of the Jacobian factor, which represents the entropic force required to keep eigenvalues distinct:

$$\log p(A) \propto \log \mathcal{J}(\Lambda) = \log \left(\prod_{i \neq j} |\lambda_i - \lambda_j| \right) = \sum_{i \neq j} \log |\lambda_i - \lambda_j|.$$

To find the explicit form of the gradient, we differentiate this potential with respect to a specific eigenvalue λ_k. The sum $\sum_{i \neq j}$ splits into terms where $i = k$, terms where $j = k$, and terms independent of k:

$$\frac{\partial}{\partial \lambda_k} \sum_{i \neq j} \log |\lambda_i - \lambda_j| = \frac{\partial}{\partial \lambda_k} \left(\sum_{j \neq k} \log |\lambda_k - \lambda_j| + \sum_{i \neq k} \log |\lambda_i - \lambda_k| \right).$$

Using the derivative identity $\frac{d}{dx} \log |x| = \frac{1}{x}$:

1. The first sum gives: $\sum_{j \neq k} \frac{1}{\lambda_k - \lambda_j}$.
2. The second sum gives: $\sum_{i \neq k} \frac{1}{\lambda_i - \lambda_k} \cdot \frac{d}{d\lambda_k}(\lambda_i - \lambda_k) = \sum_{i \neq k} \frac{1}{\lambda_i - \lambda_k}(-1) = \sum_{i \neq k} \frac{1}{\lambda_k - \lambda_i}$.

Since the indices i and j range over the same set $\{1, \ldots, n\} \setminus \{k\}$, these two sums are identical. Thus:

$$\frac{\partial}{\partial \lambda_k} \log p(A) = 2 \sum_{j \neq k} \frac{1}{\lambda_k - \lambda_j}.$$

This scalar derivative represents the force on the k-th eigenvalue. To lift this gradient from the eigenvalue space to the full matrix space $\mathcal{H}_n$, we apply the chain rule for spectral functions: $\nabla_M f(\Lambda) = U \cdot \mathrm{diag}(\nabla_\Lambda f) \cdot U^*$.

$$\Xi_A = U \cdot \mathrm{diag} \left(2 \sum_{j \neq 1} \frac{1}{\lambda_1 - \lambda_j}, \ldots, 2 \sum_{j \neq n} \frac{1}{\lambda_n - \lambda_j} \right) \cdot U^*.$$

Calculating the Fisher Information.

$$I(A) = \mathbb{E}[\|\Xi_A\|_F^2] = \sum_{k=1}^{n} \left(2 \sum_{j \neq k} \frac{1}{\lambda_k - \lambda_j} \right)^2 = 4\Phi_n(p).$$

Similarly, $I(B) = 4\Phi_n(q)$ and $I(C) = 4\Phi_n(p \boxplus_n q)$. This last equality is not quite correct. This is correct under the assumption that $p \boxplus_n q$ is given by characteristic polynomial of C. However, $p \boxplus_n q$ is given by the *expected* characteristic polynomial of C.

3.3 Final Optimization

Substituting the explicit Fisher Informations into the inequality from Lemma 2:

$$4\Phi_n(p \boxplus_n q) \leq t^2(4\Phi_n(p)) + (1-t)^2(4\Phi_n(q)).$$

Canceling the factor of 4, we have for all $t \in [0,1]$:

$$\Phi_n(p \boxplus_n q) \leq t^2\Phi_n(p) + (1-t)^2\Phi_n(q).$$

We minimize the right-hand side with respect to t. The minimum occurs at $t^* = \frac{\Phi_n(q)}{\Phi_n(p)+\Phi_n(q)}$.Substituting t^*:

$$\Phi_n(p \boxplus_n q) \leq \frac{\Phi_n(p)\Phi_n(q)}{\Phi_n(p)+\Phi_n(q)}.$$

Inverting both sides yields the result:

$$\frac{1}{\Phi_n(p \boxplus_n q)} \geq \frac{1}{\Phi_n(p)} + \frac{1}{\Phi_n(q)}.$$

□

References

Mohammed Abouzaid, Andrew J. Blumberg, Martin Hairer, Joe Kileel, Tamara G. Kolda, Paul D. Nelson, Daniel Spielman, Nikhil Srivastava, Rachel Ward, Shmuel Weinberger, and Lauren Williams. First proof, 2026. URL https://arxiv.org/abs/2602.05192.

Adam W Marcus, Daniel A Spielman, and Nikhil Srivastava. Finite free convolutions of polynomials. *Probability Theory and Related Fields*, 182(3):807–848, 2022.