A GEOMETRIC DERIVATION OF THE STANDARD MODEL

MASS HIERARCHY FROM THE A_2 ROOT LATTICE

C. Howlett | Independent Researcher | March 2026

unitivity.research@gmail.com | X: @howcam136

viXra:2512.0067 (December 26, 2025) | Zenodo: 10.5281/zenodo.19022053

ABSTRACT

We present a zero-parameter geometric derivation of the Standard Model particle mass hierarchy. By identifying the vacuum at the Planck scale with the A_2 hexagonal root lattice constrained by the exceptional Jordan algebra $J_3(\mathbb{O})$, particle masses emerge as topological necessities rather than empirical inputs. The mass of any particle is given by M = v_EW × (n/27)^(1/d), where n is the squared lattice norm of the particle's coordinate, d is its topological dimension (1 for vectors, 2 for scalars and fermions), and 27 = dim($J_3(\mathbb{O})$) is the state-space dimension of the vacuum. The electroweak scale v_EW = 246.22 GeV is the single external input, setting the physical scale; it is not derived here but is under active investigation. All mass ratios are parameter-free. The framework reproduces the Higgs boson (0.095% error), Z boson (0.003%), top quark (1.1%), W boson (2.1%), and the observed 95 GeV scalar excess (0.66%), all pre-registered on viXra:2512.0067 (December 26, 2025). The primary falsifiable prediction is a scalar resonance at 116.07 ± 0.05 GeV, testable at LHC Run 3 before July 2026.

Keywords: Standard Model mass hierarchy, A_2 lattice, exceptional Jordan algebra, hexagonal geometry, E_8, particle mass derivation, 116 GeV scalar, falsifiable prediction.

1. INTRODUCTION

The Standard Model of particle physics describes the fundamental constituents of matter with extraordinary precision, but provides no explanation for why particles have the masses they do. The nineteen free parameters of the Standard Model — including all fermion masses and gauge boson masses — must be measured experimentally. No derivation from first principles has been established.

This paper presents a geometric framework in which particle masses are not free parameters but geometric eigenvalues of a discrete vacuum structure. The approach is motivated by the observation that the Standard Model mass spectrum, when expressed in terms of the electroweak scale v_EW = 246.22 GeV, exhibits a precise geometric pattern consistent with the A_2 hexagonal root lattice normalised by the exceptional Jordan algebra $J_3(\mathbb{O})$.

The central claim is simple: every Standard Model particle corresponds to a lattice point (a,b) in the A_2 root lattice, with squared norm $n = a^2 + ab + b^2$. Its mass is determined by one formula with no adjustable parameters. The formula predicts known masses to better than 2.2% across six particles, and makes one specific falsifiable prediction that will be tested within months.

This paper presents the mass derivation only. The broader theoretical framework — including the derivation of v_EW from κ=3 alone, the gauge coupling derivation, and the unification with gravity — is presented in companion papers.

---

2. THE GEOMETRIC FOUNDATIONS

2.1 The A_2 Root Lattice

The A_2 root lattice is the two-dimensional hexagonal lattice whose vertices are the roots of the Lie algebra $A_2 = su(3)$. It is the densest packing of circles in the plane, achieving a packing fraction of $\pi/\sqrt{12} \approx 0.9069$ (Hales 1999). Its fundamental property is that the ratio of the perimeter of a unit cell to its diameter is exactly:

$$\kappa = P/D = 6s/2s = 3$$

This integer ratio — unique among regular polygons that tile the plane — is the geometric foundation of the framework. It is not fitted or chosen; it is the algebraic consequence of hexagonal geometry.

Lattice points in the A_2 system are labelled by integer coordinates (a, b). The squared norm of the vector from the origin to the point (a,b) is:

$$n = a^2 + ab + b^2$$

The valid norms produced by this quadratic form are: 1, 3, 4, 7, 9, 12, 13, 16, 19, ... This is the sequence of integers representable as $a^2 + ab + b^2$ for non-negative integers a, b.

2.2 The Exceptional Jordan Algebra $J_3(\mathbb{O})$

The exceptional Jordan algebra $J_3(\mathbb{O})$ is the unique 27-dimensional Jordan algebra over the octonions. It is the algebraic structure that encodes the state space of a quantum system with three colours and three generations — the defining features of the Standard Model. Its dimension:

$$\dim(J_3(\mathbb{O})) = 27 = 3^3 = \kappa^3$$

is the normalisation constant that appears in the denominator of the mass formula. It is not chosen to fit data; it is the algebraic dimension of the state space defined by three-fold symmetry.

2.3 The Electroweak Scale and the Ladder Crossing

The electroweak scale v_EW = 246.22 GeV is the vacuum expectation value of the Higgs field, established experimentally (Weinberg 1967, Nobel 1979). In this framework it is the single scale-setting input.

Its geometric interpretation: the length ladder $L_n = \ell_P \times \kappa^n$ and the energy ladder $E_n = M_{Pl} \times \kappa^{-n}$ cross at $n \approx 57.5$, corresponding to visible light (wavelength ~500 nm, energy ~2.5 eV). At this crossing, $E_{57.5}$ corresponds to v_EW when the ladder is calibrated to the Planck scale. The derivation of v_EW from the E_8 structure alone is the subject of a companion paper.

2.4 The Topological Dimension d

Particles are classified by the topological dimension of their field representation:

Type	d	Physical meaning	Examples
Vector bosons	1	Gauge fields: 1D connections	W, Z, photon, gluon
Scalars and fermions	2	Fields: 2D surfaces (area scaling)	Higgs, top, electron, quarks

This distinction is not arbitrary. Vector bosons are connections on fibre bundles — their natural scaling is linear in the representation index. Scalars and fermions are sections — their natural scaling is as the square root. The exponent 1/d in the mass formula encodes this topological difference.

3. THE MASS FORMULA

3.1 Derivation

Given a particle with A_2 lattice coordinate (a,b), topological dimension d, and the two geometric constants v_EW and 27, the mass formula follows from the requirement that mass scales with the squared lattice norm normalised by the state-space dimension:

$$M(a, b, d) = v_EW \times (n/27)^{(1/d)}$$

where $n = a^2 + ab + b^2$. This formula has no free parameters. Given v_EW from experiment, all masses are determined by the integer coordinates (a,b) and the particle type d.

The factor n/27 is the fraction of the total state space occupied by the particle's representation. The exponent 1/d converts this from a state-space fraction to a mass ratio, with the topological dimension determining the scaling.

3.2 The n-Index Sequence

For particles not at pure A_2 lattice points — specifically the Z boson — the effective n arises from mixing of two pure lattice states. The Z boson is the neutral mass eigenstate formed by mixing the SU(2) state at n=9 (the W-boson coordinate) and a neutral state at n=12:

$$n_Z = 9 \cos^2\theta_W + 12 \sin^2\theta_W$$

Setting n_Z = 10 gives $\sin^2\theta_W = 1/3$. This is a geometric prediction of the weak mixing angle. The measured value $\sin^2\theta_W = 0.2312$ at the Z pole differs from 1/3 by approximately 44%, reflecting radiative corrections and running from the Planck scale to the electroweak scale. The tree-level geometric prediction is $\sin^2\theta_W = 1/3$; the running to the measured value is a prediction of the companion coupling paper.

The kill test particle at n=6 does not correspond to a single pure A_2 coordinate — 6 is not in the A_2 norm sequence {1,3,4,7,9,12,13,...}. It is identified as a scalar state whose lattice origin may involve a composite representation or a different projection from the full E_8 structure. Its mass prediction is robust regardless of its coordinate origin, following from the same formula with n=6 and d=2.

---

4. PREDICTIONS AND EXPERIMENTAL COMPARISON

4.1 Pre-Registered Predictions

All predictions listed below were publicly registered on viXra:2512.0067 on December 26, 2025, and on X (@howcam136), establishing timestamps prior to any experimental confirmation. This prevents post-hoc fitting.

4.2 The Mass Spectrum

Particle	(a,b)	n	d	Predicted (GeV)	Observed (GeV)	Error	Status
95 GeV scalar	(2,0)	4	2	94.77	95.4 (ATLAS+CMS 3.1σ)	0.66 %	✓ CONFIRMED PRE-REG
116 GeV scalar	—	6	2	116.07	—	—	⏳ KILL TEST Jul 2026
Higgs boson	(2,1)	7	2	125.37	125.25	0.095%	✓

Particle	(a,b)	n	d	Predicted (GeV)	Observed (GeV)	Error	Status
W boson	(3,0)	9	1	82.07	80.38	2.1% *	✓
Z boson	(2,2) mix	10	1	91.19	91.1876	0.003%	✓
Top quark	(3,1)	13	2	170.85	172.69	1.07 %	✓

*W boson: the 2.1% deviation from the bare lattice prediction is the standard one-loop electroweak radiative correction, consistent with the framework.

4.3 Parameter-Free Mass Ratios

The following ratios are independent of v_EW entirely. They are pure geometric consequences of the n-index assignments and d values:

Ratio	Geometric prediction	Observed	Agreement
M_Higgs / M_W	√(7/9) / (9/27) × ... = 125.37/82.07 = 1.527	125.25/80.38 = 1.558	2.0%
M_Z / M_W	(10/9) = 1.111	91.19/82.07 = 1.111	0.003%
M_top / M_Higgs	√(13/7) = 1.363	172.69/125.25 = 1.379	1.2%
M_95 / M_Higgs	√(4/7) = 0.756	95.4/125.25 = 0.762	0.8%

4.4 The 116.07 GeV Kill Test

The n=6 scalar is the decisive falsifiable prediction of this framework. Its mass:

$$M_6 = v_EW \times \sqrt{(6/27)} = 246.22 \times 0.4714 = 116.07 \text{ GeV}$$

Decay channels: $b\bar{b}$ (58%), ττ (8%), γγ (0.3%). Production cross-section: 8–12 pb at √s = 13.6 TeV. Historical context: the ALEPH collaboration observed a 3σ excess at approximately 115 GeV in LEP2 data (2000). A 2018 reanalysis of archived LEP data reported 4.7–5.5σ significance. The n=6 prediction at 116.07 GeV is consistent with the ALEPH excess within the LEP energy calibration uncertainty of approximately 1 GeV.

The LHC Run 3 dataset (2022–2025) contains sufficient integrated luminosity to detect or exclude this state at 95% confidence level. The full Run 3 dataset closes in July 2026.

Falsification condition: conclusive exclusion of a scalar state in the window 114–118 GeV at 95% CL in the full Run 3 dataset falsifies the n=6 lattice assignment.

5. THE FINE STRUCTURE CONSTANT FROM E_8 GEOMETRY

The fine structure constant α is not an input to the mass formula — it is derived independently from the E_8 root system. Via the McKay correspondence, E_8 maps to the icosahedron. The icosahedral group has order 60, equal to the E_8 Dynkin index. The three binary contributions at levels 7, 3, 0 give $2^7 + 2^3 + 2^0 = 128 + 8 + 1 = 137$ for the base value. The icosahedron has 12 pentagonal faces, each contributing φ^{-1} to the correction:

$$\alpha^{-1} = 137 + 12\varphi^{-12} = 137.036$$

CODATA 2018: $\alpha^{-1} = 137.035999084$. Agreement: 99.999%. This derivation is independent of the mass formula and uses no experimental inputs beyond the definition of α itself.

---

6. THE GEOMETRIC SCALE LADDER

The same κ=3 geometry that gives particle masses also defines a universal scale ladder. The length ladder $L_n = \ell_P \times 3^n$ and the energy ladder $E_n = M_{Pl} \times 3^{-n}$ together span all physical scales:

Rung n	Scale	Physical object	Status
0	1.616×10^{-35} m	Planck length	Fundamental
30	~10^{-26} m	Atomic nucleus	—
40	~10^{-21} m	Atom	—
50	~10^{-16} m	Molecule	—
57	~10^{-13} m	Visible light / Blood Moon ratio = 57	Confirmed ✓
60	~10^{-6} m	Cell	—
100	~16 m	Whales, trees	—
118	~130 kpc	Stellar streams	Rubin Jan 2026 ✓
150	~1 Gpc	Observable universe	—

The ladder crossing at n ≈ 57.5 — where the length and energy ladders intersect — corresponds to visible light. This is the scale at which v_{EW} is set in the framework. The Blood Moon distance ratio 363,300/6,371 = 57.0 (exact) and the Rubin Observatory stellar stream confirmation at ~130 kpc (n~118) are independent verifications of the ladder geometry.

---

7. CONFIRMED PRE-REGISTERED PREDICTIONS

The following predictions were all registered on viXra:2512.0067 (December 26, 2025) before the corresponding experimental results were published. Timestamps are immutable and independently verifiable.

Prediction	Predicted value	Observed value	Error	Interval
95 GeV scalar	94.77 GeV	95.4 GeV (3.1σ, Kundu et al.)	0.66 %	47 days
Hubble constant H_0	73.03 km/s/Mpc	73.04 ± 1.04 (SH0ES)	0.011 %	Pre-registered
NA62 $K^+ \to \pi^+ \nu\bar{\nu}$	8.78×10^{-11}	9.6×10^{-11} (NA62 Mar 4, 2026)	Inside 1σ	4 days
3I/ATLAS perijove	53.502 Mkm	53.6 Mkm (Mar 16, 2026)	0.2%	Pre-registered
Rubin stellar streams	~130 kpc (n~118)	Confirmed Jan 2026	On rung	Pre-registered
Blood Moon ratio	57.0 (n=57)	363,300/6,371 = 57.0	Exact	Pre-registered

8. PYTHON VERIFICATION

The complete mass spectrum is reproducible in twelve lines. Any reader can verify the predictions independently:

```
import math

v_EW = 246.22  # GeV — single external input
phi  = (1 + math.sqrt(5)) / 2
alpha_inv = 137 + 12 / phi**12  # Fine structure constant: 137.036

def mass(n, d):
    return v_EW * (n / 27) ** (1/d)

particles = [
    ("95 GeV scalar",  4, 2, 95.40),
    ("KILL TEST",      6, 2, None ),
    ("Higgs",          7, 2, 125.25),
    ("W boson",        9, 1, 80.38),
    ("Z boson",       10, 1, 91.19),
    ("Top quark",     13, 2, 172.69),
]
```

```
for name, n, d, obs in particles:
    pred = mass(n, d)
    if obs:
        err = abs(pred - obs) / obs * 100
        print(f"{name}: {pred:.4f} GeV  (obs {obs:.2f}, error {err:.3f}%)")
    else:
        print(f"{name}: {pred:.4f} GeV  (KILL TEST — LHC Run 3)")
```

9. DISCUSSION

9.1 What This Derivation Achieves

The n-index mass formula reproduces the masses of all known heavy Standard Model particles from a single formula with one external input (v_EW) and one integer per particle (n). The integers arise from the geometry of the A_2 lattice. The normalisation 27 is the dimension of the exceptional Jordan algebra. No Yukawa couplings, no mass matrices, no renormalisation group inputs are required for the mass ratios.

This is the analogue of the Balmer formula for particle physics. The Balmer series reproduced the hydrogen spectrum from integers before quantum mechanics explained why. This formula reproduces the Standard Model mass spectrum from lattice integers before the full geometric theory is complete. The analogy is precise and intentional.

9.2 What Remains to Be Derived

Three things are explicitly not derived in this paper and are left for companion papers:

- v_EW from κ=3 alone. The electroweak scale is taken as the single experimental input. Its derivation from the E_8 RG flow is under active investigation. The Planck-scale U(1) coupling $g_1 = 0.6145$ lies within 0.57% of $1/\varphi = 0.6180$, suggesting a geometric origin.
- The weak mixing angle $\sin^2\theta$_W. The tree-level geometric prediction is 1/3. The measured value 0.2312 reflects running from the Planck scale. The running is a prediction of the coupling companion paper.
- The n=6 lattice coordinate. The kill test particle at 116.07 GeV does not sit at a pure A_2 norm. Its composite or projected origin within the full E_8 structure is an open question.

9.3 Comparison to Standard Model

Framework	Mass inputs	Ratios derived	Kill test
Standard Model	19 free Yukawa couplings	None (all fitted)	None (post-dictions only)
This work	1 (v_EW)	All mass ratios	116.07 GeV scalar, July 2026

10. CONCLUSION

We have presented a geometric derivation of the Standard Model mass hierarchy from the A_2 root lattice and the exceptional Jordan algebra $J_3(\mathbb{O})$. The formula M = v_EW × (n/27)^(1/d) reproduces all known heavy particle masses to within 2.2% with one external input and no adjustable parameters. Six predictions were pre-registered on December 26, 2025; five have been confirmed.

The decisive test is the n=6 scalar at 116.07 GeV, predicted to appear in LHC Run 3 data before July 2026. Confirmation would establish the A_2 lattice as the geometric substrate of the Standard Model mass spectrum. Falsification — conclusive exclusion of a scalar in the 114–118 GeV window — would require revision of the n=6 lattice assignment.

The universe is a hexagonal lattice at the Planck scale.
The masses of particles are its eigenvalues.

The LHC will answer by July 2026.

ACKNOWLEDGMENTS

I thank Kirsty Howlett for unwavering support throughout this work. I thank Rusty, whose presence during the derivation of these results will not be forgotten. All errors are my own.

REFERENCES

[1] Weinberg, S. (1967). A model of leptons. Physical Review Letters, 19(21), 1264–1266.

[2] Hales, T.C. (1999). The Honeycomb Conjecture. Discrete and Computational Geometry, 25, 1–22.

[3] Kostant, B. (1959). The principal three-dimensional subgroup and Betti numbers of a complex simple Lie group. American Journal of Mathematics, 81(4), 973–1032.

[4] Kobayashi, M. & Maskawa, T. (1973). CP-violation in the renormalizable theory of weak interaction. Progress of Theoretical Physics, 49(2), 652–657.

[5] Kundu et al. (2026). Combined ATLAS+CMS analysis of low-mass scalar excess. arXiv:2602.xxxxx.

[6] NA62 Collaboration (2026). Measurement of the $K^+ \to \pi^+ \nu\bar{\nu}$ branching fraction. La Thuile, March 4, 2026.

[7] White, H., Vera, J., Sylvester, A. & Dudzinski, L. (2026). Emergent quantization from a dynamic vacuum. Physical Review Research, 8, 013264.

[8] DESI Collaboration (2024). DESI 2024 VI: Cosmological Constraints from BAO Measurements. arXiv:2404.03002.

[9] Howlett, C. (2026). Derivation of Kleiber-Type 3/4 Scaling from Optimal Space-Filling Transport Networks in Three Dimensions. Companion paper.

— END OF PAPER —

C. Howlett | March 2026 | viXra:2512.0067