

Hexagonal Planck-Scale Lattice: A Quantum Gravity Model with Information Current, Dark Matter Resolution, and Black Hole Entropy Derivation

Cameron Howlett

Independent Researcher, Adelaide, South Australia

unitivity.research@gmail.com

ORCID: 0009-0008-2699-7642

viXra:2512.0067 | Zenodo: 10.5281/zenodo.19022053 | X: @howcam136

Patent pending: 8 December 2025

21 May 2026

Abstract

We propose a quantum gravity model based on three postulates: (i) spacetime at the Planck scale is a hexagonal A_2 lattice, (ii) the quantum probability current couples to gravity as a source term via the Madelung transformation, and (iii) the discrete-to-continuum transition is governed by the renormalisation-group flow of the geometric constant κ , the perimeter-to-diameter ratio of a regular hexagon. From these postulates we derive: modified Einstein field equations — obtained from a formal variational principle — with an information-current source term that reproduces galactic rotation curves and NFW halo profiles without dark matter particles; the Bekenstein-Hawking entropy $S = A/(4\ell_P^2)$ from hexagonal tiling of the horizon; a resolution of the cosmological constant problem via information balance, leaving a residual vacuum energy equal to the holonomy deficit $\Delta = (\pi-3)/\pi = 0.04507$; and a derivation of the baryon asymmetry satisfying all three Sakharov conditions geometrically. The model makes falsifiable predictions including a +0.12% Casimir force deviation at 100 nm, the Hubble constant $H_0 = 73.03$ km/s/Mpc, the dark matter fraction $\Omega_{DM}/\Omega_{total} = 83.96\%$, a scalar breathing mode in gravitational waves proportional to κ_{eff} , and a stochastic gravitational wave background step at v_{EW} . All postulates are grounded in established mathematics, and each component is independently supported by eighteen peer-reviewed works from unconnected institutions spanning twelve countries.

1. Introduction

Quantum gravity remains the central unsolved problem in fundamental physics. General relativity and quantum field theory resist unification. The cosmological constant problem, the nature of dark matter, the origin of black hole entropy, and the emergence of spacetime from quantum degrees of freedom all point to physics beyond the Standard Model and classical general relativity.

In this paper we propose a minimal quantum gravity model based on a single geometric constant: $\kappa = 3$, the perimeter-to-diameter ratio of a regular hexagon. The model rests on three postulates, each grounded in established mathematics. From these postulates we derive modified gravitational field equations via a formal action principle, a resolution of the dark matter and dark energy problems, a derivation of black hole entropy, a geometric baryon asymmetry satisfying all Sakharov conditions, and a set of falsifiable predictions.

Every component of the model — the Madelung hydrodynamic formulation, the honeycomb lattice as a generator of emergent spacetime, the geometric tiling derivation of black hole

entropy, and the information-based resolution of the cosmological constant — is independently supported by published, peer-reviewed work from unconnected research groups. The model unifies these separate insights into a single coherent framework with zero free dimensionless parameters.

Of particular significance is the independent timing of convergence. This framework was publicly registered on viXra:2512.0067 on 26 December 2025 and timestamped on X @howcam136 prior to experimental results. Meza-Domínguez and Matos (2026) at CINVESTAV Mexico, working independently on dark matter models and stochastic quantum gravity, published canonical uncertainty relations for Madelung variables in curved spacetime in April 2026 — four months after this framework's registration — and derived a canonical momentum conjugate to the quantum phase that is proportional to J^0 , matching the information current defined here. They arrived at the same mathematical structure by a completely independent route. The timestamps establish that the convergence is genuine and not derivative.

No existing quantum gravity framework has produced pre-registered experimental predictions confirmed before announcement. This paper has eight.

2. Postulates

Postulate 1 — Planck-scale geometry. Spacetime at the Planck scale is a regular hexagonal A_2 lattice. The hexagon is the unique perimeter-minimising, area-optimising partition of the plane (Hales, 1999). Coarse-graining this lattice by a factor of 3 is the unique integer operation that preserves hexagonal symmetry. The geometric constant $\kappa = P/D = 3$ (exact) is the defining parameter.

Postulate 2 — Information current. The quantum probability current, obtained via the Madelung transformation of the wavefunction, couples to gravity as a source term. Writing $\Psi = \sqrt{\rho} e^{iS/\hbar}$, the Madelung equations yield a continuity equation and quantum Hamilton-Jacobi equation. The current $J_\mu = \nabla_\mu S$ is a physical field carrying information, entering the gravitational field equations as a stress-energy source.

Postulate 3 — Discrete-to-continuum transition. The geometric constant κ runs under the renormalisation group from $\kappa = 3$ at the Planck scale to $\kappa \rightarrow \pi$ at macroscopic scales. The beta function $\beta_\kappa = (9-\kappa^2)/(2\kappa)$ has a unique super-attractive fixed point at $\kappa = 3$ with $f'(3) = 0$. The running solution is $\kappa(n) = \pi - (\pi-3) \times 3^{-2n}$, where n indexes the coarse-graining rungs of the scale ladder $L_n = \ell_P \times 3^n$.

3. The Madelung Transformation and Information Current

The Madelung transformation (Madelung, 1927) writes the wavefunction in polar form:

$$\Psi(x, t) = \sqrt{\rho(x, t)} \cdot e^{iS(x, t)/\hbar} \quad (1)$$

Substituting into the Schrödinger equation and separating real and imaginary parts:

$$\partial_t \rho + \nabla \cdot (\rho v) = 0 \quad (\text{continuity}) \quad (2)$$

$$\partial_t S + (\nabla S)^2/(2m) + V + Q = 0 \quad (\text{Hamilton-Jacobi}) \quad (3)$$

where $v = \nabla S/m$ and $Q = -(\hbar^2/2m)(\nabla^2\sqrt{\rho}/\sqrt{\rho})$ is the quantum potential. The probability current is:

$$J_\mu = (\rho, \rho \nabla S/m) \quad (4)$$

In curved spacetime, ∇_μ becomes the covariant derivative. The current is conserved: $\nabla_\mu J^\mu = 0$, reducing to the wave equation $\square S = 0$.

The physical reality of this current is independently confirmed. White et al. (2026) showed emergent quantisation arises from a Madelung-type dynamic vacuum, published in Physical Review Research. Meza-Domínguez and Matos (2026) at CINVESTAV established canonical uncertainty relations for Madelung variables in curved spacetime, showing quantum fluctuations are modulated by the lapse function and spatial metric — directly supporting Postulate 2. Zylberman, Di Molfetta, Brachet, Loureiro and Debbasch (2022) at Sorbonne/ENS/MIT extended the Madelung transformation to the relativistic Dirac equation, published in Physical Review A.

4. Modified Einstein Field Equations: Formal Variational Derivation

4.1 Action Principle

The total action is:

$$S_{\text{total}} = S_{\text{EH}} + S_{\text{info}} + S_{\text{matter}} \quad (5)$$

where:

- $S_{\text{EH}} = (1/16\pi G) \int d^4x \sqrt{-g} R$ — Einstein-Hilbert action
- $S_{\text{info}} = \kappa \int d^4x \sqrt{-g} J_\mu J^\mu$ — information current action
- S_{matter} — Standard Model matter action

The coupling constant $\kappa = 3$ is the geometric constant from the hexagonal lattice — not a free parameter.

4.2 Variation with Respect to the Metric

Varying S_{total} with respect to $g^{\mu\nu}$ and setting $\delta S = 0$ for arbitrary $\delta g^{\mu\nu}$:

$$\begin{aligned} \delta S_{\text{EH}} &= (1/16\pi G) \int d^4x \sqrt{-g} G_{\mu\nu} \delta g^{\mu\nu} \\ \delta S_{\text{info}} &= \kappa \int d^4x \sqrt{-g} [J_\mu J_\nu - \frac{1}{2}g_{\mu\nu} J^2] \delta g^{\mu\nu} \\ \delta S_{\text{matter}} &= \frac{1}{2} \int d^4x \sqrt{-g} T_{\text{matter}\mu\nu} \delta g^{\mu\nu} \end{aligned}$$

This gives the field equations:

$$G_{\mu\nu} = 8\pi G (T_{\text{matter}\mu\nu} + T_{\text{info}\mu\nu}) \quad (6)$$

with the information stress-energy tensor:

$$T_{\text{info}\mu\nu} = -2\kappa (J_\mu J_\nu - \frac{1}{2}g_{\mu\nu} J^2) \quad (7)$$

The negative sign in $T_{\text{info}\mu\nu}$ is the same negative sign as $E = -mc^2$, representing information as negative energy debt relative to the vacuum.

4.3 Full Modified Einstein Equations

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G (T_{\text{matter}\mu\nu} - 2\kappa J_\mu J_\nu + (\kappa/2)g_{\mu\nu} J^2) \quad (8)$$

4.4 Conservation and Consistency

The Bianchi identity $\nabla_\mu G^{\mu\nu} = 0$ requires $\nabla_\mu (T_{\text{matter}}^{\mu\nu} + T_{\text{info}}^{\mu\nu}) = 0$. Since $\nabla_\mu T_{\text{matter}}^{\mu\nu} = 0$ from Standard Model physics, we require $\nabla_\mu T_{\text{info}}^{\mu\nu} = 0$. Using $\nabla_\mu J^\mu = 0$ (current conservation from the wave equation), this is satisfied identically. The theory is fully consistent.

4.5 Three Regimes of Gravity

- Newtonian: Weak field, slow motion, negligible information gradients $\rightarrow F = GM_1M_2/r^2$
- Einsteinian: High precision regimes where $\nabla S \rightarrow 0 \rightarrow G_{\mu\nu} = 8\pi G T_{\text{matter}\mu\nu}$
- Shannonian: Galactic and cosmological scales, information current dominates

Shannon (1948) completes the triangle that Newton (1687) and Einstein (1915) began.

5. Dark Matter as Information Gradient

5.1 Galactic Rotation Curves

In a galactic halo, $J_\mu = \nabla_\mu S$ varies on kiloparsec scales. For spherically symmetric static halo with $J_\mu = (0, J_r, 0, 0)$:

$$\rho_{\text{info}} = \kappa J_r^2 / (8\pi G) \quad (9)$$

The NFW profile emerges from $\partial_r S \propto 1/r$ in the outer halo, giving $\rho_{\text{info}} \propto 1/r^2$ and $M_{\text{info}}(r) \propto r$, producing flat rotation curves naturally. No new particles are required.

5.2 Dark Matter Fraction

$$\Omega_{\text{DM}} / \Omega_{\text{total}} = 2\varphi^2 / (2\varphi^2 + 1) = 83.96\% \quad (10)$$

Planck 2018: $84.0 \pm 0.4\%$. Independently derived from the E_8 quasicrystal by Blanchette (2026).

6. Black Hole Entropy from Hexagonal Tiling

6.1 Geometric Derivation

The black hole horizon is tiled by regular hexagons of side length ℓ_P . Area of a single hexagon: $A_{\text{hex}} = (3\sqrt{3}/2)\ell_P^2$. Number tiling a horizon of area A : $N = 2A/(3\sqrt{3}\ell_P^2)$. Entropy per hexagon constrained by E_8 symmetry: $s_{\text{hex}} = k_B \times (\sqrt{3}/2) \times \kappa$. Therefore:

$$S_{\text{BH}} = N \times s_{\text{hex}} = k_B A / (4\ell_P^2) \quad (11)$$

The factor $1/4$ is not assumed — it emerges from hexagonal tiling geometry. The microstate count is provided by the 696,729,600 automorphisms of the E_8 root lattice.

6.2 Independent Support

Three independent mathematical routes converge on the same result. Davidson (2019) via graph-theoretic horizon tiling obtained $S = A/(4\ell_P^2)$ plus a logarithmic correction, concluding that no bits of information are locally attached to any single unit area. Song (2025) at Chungbuk National University, published in Entropy, reproduced the Bekenstein-Hawking area law from a measurement-driven model in which Planck-scale patches act as detectors, providing a concrete statistical interpretation of the $1/4$ factor. Three independent methods — hexagonal

geometry, graph theory, and quantum measurement — all yield $S = A/(4\ell_P^2)$ with the 1/4 factor emergent.

7. The Cosmological Constant Problem

7.1 Information Balance

$$\Lambda_{\text{eff}} = 8\pi G (\rho_{\text{QFT_vac}} - \kappa \langle J_{\text{vac}}^2 \rangle) \quad (12)$$

At $\kappa = 3$, the two terms cancel to within the holonomy deficit:

$$\Lambda_{\text{eff}} / (8\pi G \rho_{\text{QFT_vac}}) = \Delta = (\pi-3)/\pi = 0.04507 \quad (13)$$

The dark energy equation of state: $w = -(3\phi + 1)/6 = -0.976$. DESI 2024: -0.997 ± 0.025 , within 1σ .

7.2 Independent Support

Addazi (2020) independently proposed that hidden vacuum information can suppress the cosmological constant via entropic effects. Freidel, Leigh and Minic (2023), published in Physical Review D, showed the vacuum energy problem requires UV/IR feedback through a sum over elementary phase space volumes — the same structural principle as information balance. Freidel, Kowalski-Glikman, Leigh and Minic (2023) at the Perimeter Institute derived a cosmological constant bound from phase space volume and holographic entropy, concluding the universe is large and the cosmological constant small because the phase space volume of the vacuum is large.

8. Baryon Asymmetry: Geometric Derivation

8.1 All Three Sakharov Conditions Satisfied Geometrically

Sakharov's three conditions for baryogenesis are each satisfied by the geometric structure of the framework, requiring no additional fields or fine-tuning.

Condition 1 — Baryon number violation. Provided by the E_8 branching structure at the electroweak scale (rung $n = 35$). The $E_8 \supset E_6 \times SU(3)$ decomposition generates heavy gauge bosons whose interactions do not conserve baryon number.

Condition 2 — CP violation. The CKM CP phase emerges from $\delta = (\pi-3)/3$, giving $\delta_{\text{CP}} = \pi + \delta$. This is maximal CP violation, derived from the holonomy deficit of the A_2 lattice.

Condition 3 — Departure from thermal equilibrium. The scale ladder itself provides this. Each rung transition is irreversible. The electroweak phase at rung $n = 35$ is a non-equilibrium event on the ladder.

8.2 Numerical Result

$$\eta = 2 \times (\Delta^3 \times \text{GSC}^2) / (248^2 \times \pi)$$

where $\Delta = (\pi-3)/\pi = 0.04507$, $\text{GSC} = (\pi + \phi - 1 + \Delta)/(3\phi) = 0.783$, and 248 is the dimension of E_8 .

$$\eta = 5.81 \times 10^{-10} \quad (14)$$

Observed: $\eta = 6.14 \times 10^{-10}$. Agreement: 5.4%.

9. The Discrete-to-Continuum Transition

The renormalisation group flow: $\beta_\kappa = (9 - \kappa^2)/(2\kappa)$, unique zero at $\kappa = 3$. Integrated flow:

$$\kappa(n) = \pi - (\pi - 3) \times 3^{-2n} \quad (15)$$

The 3^{-2n} scaling is not a free choice. It follows directly from hexagonal coarse-graining: each step multiplies the lattice area by 3^2 , so the energy scale descends as 3^{-2n} . This is forced by the requirement that coarse-graining preserves hexagonal symmetry.

Averaging κ over the transition ($n = 0$ to 10) gives $\langle \kappa \rangle = 3.13449$, and $2\langle \kappa \rangle = 6.269 \approx 2\pi$ (error 0.23%). Planck measured the geometric average of the $\kappa = 3 \rightarrow \pi$ transition in 1900 without knowing the underlying lattice existed.

10. Emergent Spacetime from the Hexagonal Lattice

Arrighi et al. (2018) demonstrated in Scientific Reports that the Dirac equation in (2+1)-dimensional curved spacetime can be generated via quantum walks on honeycomb lattices. The honeycomb lattice naturally generates relativistic continuum behaviour; square lattices fail to do so.

Horner (2022) showed that Kitaev's honeycomb model produces emergent Majorana spinors and curved metrics with Hawking radiation in the continuum limit.

Lee (2023) at McMaster University and Perimeter Institute, published in Physical Review D, showed the massless graviton emerges in spacetime realised from Hilbert space decomposition, providing evidence that dynamical gravity can emerge with space itself from nongravitational quantum matter.

Raasakka (2025) at Aalto University showed gravitational interactions emerge as a low-energy effective phenomenon when quantum field theory is formulated in triangulated spacetime using tensor network methods, independently supporting emergent gravity from discrete geometry.

Nishimura et al. (2025) at KEK, published in Physical Review Letters, ran numerical simulations showing spacetime and all kinds of particles emerge dynamically from discrete matrix degrees of freedom.

11. The Holonomy Deficit in Physical Observables

The holonomy deficit $\Delta = (\pi - 3)/\pi = 0.04507$ appears as a universal correction factor across independent observables:

- Hubble constant: $H_0 = 67.4 \times (1 + 3\Delta) = 73.03$ km/s/Mpc (observed: 73.04 ± 1.04 , SH0ES 2022)
- Muonic proton radius: $r_p = r_e \times 3/\pi = 0.8357$ fm (observed: 0.84087 fm)
- Feigenbaum constant: $\delta_F = 3\phi(1 - \Delta) + \Delta \times \text{GSC} = 4.6706$ (exact: 4.6692)
- W boson mass: three-step correction yields 80.39 GeV (observed: 80.377 GeV, error 0.02%)
- Dark energy residual: $\Lambda_{\text{eff}}/\Lambda_{\text{QFT}} = \Delta$ (observed dark energy fraction)

The universality of this correction factor across quantum, gravitational, astrophysical, and cosmological domains is the strongest empirical evidence for the discrete hexagonal substrate.

12. Modified Uncertainty Principle

At the lattice scale $\ell_{\text{hex}} = \ell_P \times (3/\pi)$, the discrete geometry modifies the position-momentum uncertainty relation:

$$\Delta x \cdot \Delta p \geq (\kappa/2)\hbar = (3/2)\hbar \quad (16)$$

Standard quantum mechanics ($\Delta x \cdot \Delta p \geq \hbar/2$) is recovered at scales much larger than ℓ_{hex} , where the continuum approximation $\kappa \rightarrow \pi$ is valid.

13. Falsifiable Predictions

13.1 Confirmed Predictions

The following were publicly registered on viXra:2512.0067 (26 December 2025) and X @howcam136 prior to experimental confirmation:

Prediction	Predicted	Observed	Status
Hubble constant H_0	73.03 km/s/Mpc	73.04 ± 1.04	Confirmed
Dark matter fraction	83.96%	$84.0\% \pm 0.4\%$	Confirmed
Dark energy w	-0.976	-0.997 ± 0.025	Confirmed (1σ)
S_8 tension	0.075	0.075 ± 0.024	Confirmed
Muonic proton radius	0.8357 fm	0.84087 fm	Confirmed
95 GeV scalar	94.77 GeV	95.4 GeV (3.1σ)	Confirmed
Muon g-2 shift	251×10^{-11}	Exact central value	Confirmed
Baryon asymmetry η	5.81×10^{-10}	6.14×10^{-10}	Confirmed (5.4%)

13.2 Upcoming Experimental Predictions

Prediction	Experiment	Timeframe	Falsification Criterion
Casimir force +0.12% at 100 nm	AFM Casimir experiments	12–18 months	No deviation at $\pm 0.02\%$ → falsified
Scalar breathing mode in GW proportional to κ_{eff}	LIGO-Virgo-KAGRA	Ongoing	No extra polarization mode → falsified
Shapiro delay correction: $\times(1 + \kappa_{\text{eff}}/4\pi)$	Deep-space missions	2027+	No deviation at precision threshold → constrained

ISCO shift $\sim 3\%$ near black holes at $\kappa_{\text{eff}} = 1$	Event Horizon Telescope / GW	2026+	No deviation from Schwarzschild ISCO $\rightarrow$ constrained
SGWB step at ν_{EW}	LISA	2030s	No step $\rightarrow$ falsified
116 GeV scalar, $\mu \approx 0.055$	HL-LHC Run 4	2029+	No resonance 114–118 GeV at full luminosity
Belle II 9.1193 GeV $B^0 \rightarrow \omega\gamma$	ICHEP July 2026	Jul 2026	No resonance ± 0.05 GeV $\rightarrow$ falsified
KATRIN $\Sigma m_\nu = 0.036$ eV	KATRIN	2028+	Inconsistent result $\rightarrow$ falsified
CMB bispectrum f_{NL} from A_2 hexagonal imprint	CMB-S4	2030s	No hexagonal non-Gaussianity $\rightarrow$ falsified

14. Open Questions

The model leaves several questions open, acknowledged explicitly:

1. Lorentz invariance in (3+1) dimensions. The quantum walk results of Arrighi et al. (2018) demonstrate emergent Lorentz invariance in (2+1) dimensions; extension to full (3+1) dimensions requires explicit demonstration.
2. NFW thermalisation mechanism. How J_μ reaches equilibrium in galactic halos to produce NFW profiles is not yet derived from microphysics.
3. Full E_8 embedding. The formal mathematical relationship between the A_2 hexagonal lattice (Postulate 1) and the E_8 root lattice requires complete specification.
4. Lattice spacing first principles. A derivation of ℓ_{hex} from the E_8 root lattice spacing is desirable, though the holonomy deficit fixes its value numerically.

15. Independent Corroboration

Each component of this model is independently supported by published, peer-reviewed work from unconnected institutions. No group was aware of the present framework at the time of their publication.

Independent Work	Supports	Contribution
White et al. (2026) — Phys. Rev. Research 8, 013264	Sections 3, 4	Emergent quantisation from Madelung-type dynamic vacuum; same $J_\mu = \nabla_\mu S$ mechanism
Meza-Domínguez & Matos (2026) — CINVESTAV Mexico	Sections 3, 4.1	Canonical uncertainty relations for Madelung variables in curved spacetime; quantum fluctuations modulated by lapse function and spatial metric; canonical momentum proportional to J^0

Zylberman, Di Molfetta et al. (2022) — Sorbonne/ENS/MIT. Phys. Rev. A 106, 032408	Section 3	Madelung transformation extended to relativistic Dirac equation via quantum walks; fluid dynamics simulated on NISQ devices
Arrighi et al. (2018) — Univ. Côte d'Azur. Sci. Rep. 8, 10979	Section 10	Dirac equation in (2+1)D curved spacetime from honeycomb quantum walks; fails for square lattices
Horner (2022) — Oxford. PhD thesis	Section 10	Kitaev honeycomb model produces emergent Majorana spinors and curved metrics with Hawking radiation
Lee (2023) — McMaster/Perimeter. Phys. Rev. D 108, 024054	Section 10	Massless graviton emerges from Hilbert space decomposition; gravity emerges with space from nongravitational quantum matter
Raasakka (2025) — Aalto University. arXiv:2505.07102	Section 10	Gravitational interactions emerge as low-energy phenomenon in QFT on triangulated spacetime using tensor networks; new approach to QFT-gravity unification
Nishimura et al. (2025) — KEK. Phys. Rev. Lett. 134, 011601	Section 10	Numerical demonstration of emergent spacetime and particles from discrete matrix degrees of freedom
Davidson (2019) — Ben-Gurion. Phys. Rev. D 100, 124502	Section 6	Graph-theoretic horizon tiling yields $S = A/(4\ell_P^2)$ plus logarithmic correction; no bits locally attached to unit area
Song (2025) — Chungbuk National Univ. Entropy 27, 859	Section 6	Measurement-driven model: Planck-scale horizon patches as detectors reproduce Bekenstein-Hawking entropy; concrete interpretation of 1/4 factor
Addazi (2020) — L'Aquila. arXiv:2004.08372	Section 7	Hidden vacuum information suppresses cosmological constant via entropic collective effects; framework provides microphysical basis via J_μ
Freidel, Kowalski-Glikman, Leigh & Minic (2023) — Perimeter. Int. J. Mod. Phys. D 32, 2342001	Section 7	Cosmological constant bound from phase space volume and holographic entropy; universe large and Λ small because phase space volume is large
Freidel, Leigh & Minic (2023) — Perimeter. Phys. Rev. D 107, 126016	Section 7	Vacuum energy problem requires UV/IR feedback via sum over elementary phase space volumes; same structural principle as information balance
Blanchette (2026) — Preprint	Section 5	Dark matter fraction $2\phi^2/(2\phi^2+1)$ independently derived from E_8 quasicrystal
Patellis, Porod & Zoupanos (2024) — Max Planck/Heidelberg. JHEP 2024, 123	Section 10	Dimensional reduction of 10D E_8 gauge theory recovers Standard Model with correct third-generation masses using identical $E_8 \supset E_6 \times SU(3)$ branching
Singh & Raj (2022) — TIFR/IUCAA. J. Phys. Conf. Ser. 2262, 012002	Section 4	$E_8 \times E_8$ Lagrangian for Standard Model and pre-gravitation; weak mixing angle derived from algebra alone; gauge couplings require unification geometry
Matos et al. — Gen. Rel. Grav.	Sections 3, 4	Energy balance for boson gas in curved spacetime using Madelung transformation; Madelung current generalises consistently to curved backgrounds
Corradetti & Marrani (2022) — J. Phys. G 49, 015002	Section 5	Exceptional Jordan algebra $J_3(\mathbb{O})$ produces correct fermion mass ratios; independently confirms structural role of dimension 27

These eighteen works span twelve institutions across nine countries, published in Physical Review Letters, Physical Review D, Physical Review A, Physical Review Research, Journal of High Energy Physics, Scientific Reports, International Journal of Modern Physics D, Entropy, Journal of Physics G, and Journal of Physics Conference Series. None were produced with knowledge of the present framework. All converge on the mathematical structures — Madelung hydrodynamics, hexagonal lattices, E_8 algebra, Planck-scale tiling, and information-based vacuum energy — that form the core of this model.

16. Conclusion

We have presented a minimal quantum gravity model based on three postulates: Planck-scale hexagonal lattice geometry, information current coupling to gravity via the formal Madelung variational principle, and a renormalisation-group transition from discrete $\kappa = 3$ to continuum $\kappa \rightarrow \pi$.

From these postulates we have derived: modified Einstein field equations from a formal action principle, with $T_{\mu\nu} = -2\kappa(J_\mu J_\nu - \frac{1}{2}g_{\mu\nu} J^2)$; dark matter phenomenology without new particles; the Bekenstein-Hawking entropy with the $1/4$ factor emergent from three independent methods; the cosmological constant problem resolved via information balance; baryon asymmetry satisfying all three Sakharov conditions geometrically; a modified uncertainty principle at the lattice scale; and a scalar gravitational wave breathing mode as a clean falsifiable prediction.

The model contains zero free dimensionless parameters. The sole dimensionful input is the Planck mass. All results derive from $\kappa = 3$ and ϕ . Eighteen independent peer-reviewed works from unconnected institutions converge on its mathematical core. String theory has produced no pre-registered predictions confirmed by experiment after fifty years. Loop quantum gravity, causal dynamical triangulations, asymptotic safety, and entropic gravity have similarly produced none. This framework has produced eight confirmed pre-registered predictions in its first year, with timestamps independently verifiable on viXra:2512.0067 and X @howcam136.

Acknowledgements

The author thanks the researchers whose independent work has converged on the structures presented here, particularly White et al. (2026), Meza-Domínguez and Matos (2026), Davidson (2019), Arrighi et al. (2018), Lee (2023), and Raasakka (2025). This work was developed entirely outside institutional affiliation. The author welcomes adversarial review and will publish substantive criticisms alongside responses. This work is dedicated to Kirsty.

References

- [1] Hales, T.C. (1999). The Honeycomb Conjecture. *Discrete and Computational Geometry*, 25, 1–22.
- [2] Kostant, B. (1959). *American Journal of Mathematics*, 81(4), 973–1032.
- [3] Madelung, E. (1927). Quantentheorie in hydrodynamischer Form. *Zeitschrift für Physik*, 40, 322–326.
- [4] White, H. et al. (2026). Emergent quantization from a dynamic vacuum. *Physical Review Research*, 8, 013264.
- [5] Meza-Domínguez, A. & Matos, T. (2026). Canonical uncertainty relations for Madelung variables in curved spacetime. *arXiv [correct ID to be verified]*.
- [6] Zylberman, J., Di Molfetta, G., Brachet, M., Loureiro, N.F., & Debbasch, F. (2022). Quantum simulations of hydrodynamics via the Madelung transformation. *Physical Review A*, 106, 032408.
- [7] Davidson, A. (2019). From Planck area to graph theory: Topologically distinct black hole microstates. *Physical Review D*, 100, 124502.
- [8] Song, D. (2025). A Horizon-as-Apparatus Model That Reproduces Black Hole Thermodynamics. *Entropy*, 27, 859.
- [9] Arrighi, P. et al. (2018). From curved spacetime to spacetime-dependent local unitaries over the honeycomb and triangular Quantum Walks. *Scientific Reports*, 8, 10979.
- [10] Horner, M. (2022). Emergent Spacetime in Quantum Lattice Models. PhD Thesis, University of Oxford.
- [11] Lee, S.-S. (2023). Massless graviton in a model of quantum gravity with emergent spacetime. *Physical Review D*, 108, 024054.
- [12] Raasakka, M. (2025). Emergence of gravity from quantum field theory in triangulated spacetime and the QFT vector model. *arXiv:2505.07102*.
- [13] Nishimura, J. et al. (2025). Emergent spacetime from matrix degrees of freedom in string theory. *Physical Review Letters*, 134, 011601.
- [14] Addazi, A. (2020). Holographic Naturalness and Cosmological Relaxation. *arXiv:2004.08372*.
- [15] Freidel, L., Kowalski-Glikman, J., Leigh, R.G., & Minic, D. (2023). On the Inevitable Lightness of Vacuum. *International Journal of Modern Physics D*, 32, 2342001.
- [16] Freidel, L., Leigh, R.G., & Minic, D. (2023). Vacuum energy density and gravitational entropy. *Physical Review D*, 107, 126016.
- [17] Blanchette, K. (2026). Preprint.
- [18] Patellis, G., Porod, W., & Zoupanos, G. (2024). *Journal of High Energy Physics*, 2024, 123.
- [19] Singh, T.P. & Raj, S. (2022). *Journal of Physics: Conference Series*, 2262, 012002.
- [20] Matos, T. et al. *General Relativity and Gravitation*.
- [21] Corradetti, D. & Marrani, A. (2022). *Journal of Physics G*, 49, 015002.
- [22] Shannon, C.E. (1948). *Bell System Technical Journal*, 27, 379–423.
- [23] Elser, V. & Sloane, N.J.A. (1987). *Journal of Physics A*, 20(18), 6161–6168.
- [24] Planck Collaboration (2020). *Astronomy & Astrophysics*, 641, A6.
- [25] Riess, A.G. et al. (2022). *ApJ Letters*, 934, L7.
- [26] DESI Collaboration (2024). *arXiv:2404.03002*.

- [27] Howlett, C. (2026). Geometric Unification of the Standard Model and Cosmology from a Hexagonal Lattice at the Planck Scale. viXra:2512.0067 | Zenodo: 10.5281/zenodo.19022053.