

MPC WITH NO MODEL UNCERTAINTY

BASIL, GANNON, "MPC", SPRINGER 2016

SCATOLINI, "MPC COURSE", SIGRA SUMMER SCHOOL 2011

FROM LINEAR QUADRATIC OPTIMAL CONTROL TO MPC

LTI, DT systems without uncertainties and disturbances

$$\begin{cases} x(k+1) = A x(k) + B u(k) \\ y(k+1) = C x(k+1) \end{cases} \quad \begin{matrix} x(k) \in \mathbb{R}^{n_x} \\ y(k) \in \mathbb{R}^{n_y} \end{matrix}, u(k) \in \mathbb{R}^{n_u}$$

ASSUMPTION (A, B) STABILIZABLE

(A, C) OBSERVABLE

$x(k)$ MEASURED AT TIME k

LINEAR CONSTRAINTS

$$F x(k) + G u(k) \leq b$$

m_c : number of constraints

$$F \in \mathbb{R}^{m_c \times m_x}$$

$$G \in \mathbb{R}^{m_c \times m_u}$$

$$b \in \mathbb{R}^{m_c}$$

$F = 0 \Rightarrow$ input constraints $G u(k) \leq b$

$G = 0 \Rightarrow$ state constraints $F x(k) \leq b$

Non stationary constraints F_k, G_k

FEASIBLE PAIRS

$$(x(k), u(k)) \Leftrightarrow F x(k) + G u(k) \leq b$$

FEASIBLE SEQUENCE

$$\{(x(0), u(0)), (x(1), u(1)), \dots\}$$

$\Leftrightarrow (x(k), u(k))$ is feasible, $k = 0, 1, \dots$

REGULATION PROBLEM

- Design of a controller that steers the system state to the origin ($x(k) \rightarrow 0$)
- QUADRATIC COST, INFINITE HORIZON, UNCONSTRAINED CASE

$$J(x(0), \{u(0), u(1), \dots\}) = \sum_{k=0}^{\infty} (\|x(k)\|_Q^2 + \|u(k)\|_R^2)$$

$$- \|v\|_S^2 = v^T S v$$

$$- Q \in \mathbb{R}^{n_x \times n_x}, \quad R \in \mathbb{R}^{m_u \times m_u}$$

ASSUMPTION

$$R \succ 0$$

SYMMETRIC, POSITIVE DEFINITE (eigs are real and strictly positive)

$$Q \succeq 0$$

"

,

"

SEMI-DEFINITE (eigs are real and non-negative)

INFINITE HORIZON UNCONSTRAINED QUADRATIC PROBLEM

$$J^*(x(k)) = \min_{\{u(k), u(k+1), \dots\}} J(x(k), \{u(k), u(k+1), \dots\})$$

$$\left(\nabla! \text{ the constrained infinite-h problem has no closed-form solution} \right)$$

$$\begin{aligned} & \min_{\{ \quad \}} J(x(k), \{ \quad \}) \\ & \text{s.t.} \quad F x(k+i) + G u(k+i) \leq b, \quad i=0,1,\dots \end{aligned}$$

• LINEAR STATE FEEDBACK CONTROL $u(k) = K x(k), \quad K \in \mathbb{R}^{m_u \times m_x}$

\Rightarrow CLOSED-LOOP DYNAMICS

$$\begin{cases} x(k+1) = A x(k) + B u(k) \\ u(k) = K x(k) \end{cases}$$

$$x(k+1) = (A + BK) x(k)$$

$$\Rightarrow \begin{cases} x(k) = (A + BK)^k x(0) \\ u(k) = K (A + BK)^k x(0), \quad k=0,1,\dots \end{cases}$$

$$\Rightarrow J(x(0), \{u(0), u(1), \dots\}) = J(x(0))$$

$\leftarrow u(k)$ can be expressed
as function of $x(0)$

$$\Rightarrow J(x(0)) = x(0)^T W x(0) = \|x(0)\|_W^2$$

$$x(k) = (A+BK)^k x(0)$$

$$\leftarrow u(k) = K(A+BK)^k x(0)$$

with
$$W = \sum_{k=0}^{\infty} \left[\left((A+BK)^k \right)^T Q (A+BK)^k + \right. \\ \left. + \left((A+BK)^k \right)^T K^T R K (A+BK)^k \right] =$$

$$= \sum_{k=0}^{\infty} \left[\left((A+BK)^k \right)^T (Q + K^T R K) (A+BK)^k \right] \in \mathbb{R}^{n_x \times n_x}$$

• If $(A+BK)$ is strictly stable ($|\text{eigs}| < 1$) $\Rightarrow W$ has a finite number of elements

$$\cdot \begin{cases} R > 0 \\ Q \geq 0 \end{cases} \Rightarrow J(x) > 0, W > 0$$

LEMMA 1 (Lyapunov matrix equation)

W is the unique positive definite solution of

$$W = (A+BK)^T W (A+BK) + Q + K^T R K \iff (A+BK) \text{ is strictly stable}$$

THEOREM 1 (DISCRETE TIME ALGEBRAIC RICCATI EQUATION)

the optimal gain matrix K^* minimizing $J(x(0)) = \|x(0)\|_W^2$, for any $x(0) \in \mathbb{R}^{n_x}$ is

$$K^* = (B^T W B + R)^{-1} B^T W A$$

LQR

$$u(k) = K^* x(k)$$

OPTIMAL CONTROL LAW FOR THE DISCRETE TIME
LTI SYSTEM UNDER QUADRATIC COST FUNCTION
WITHOUT CONSTRAINTS

$$J^*(x(0)) = \|x(0)\|_W^2$$

OPTIMAL COST WHICH ONLY DEPENDS ON $x(0)$

MPC

DUAL MODE PREDICTION PARADIGM

- Presence of constraints

- Finite horizon \Rightarrow RECEDING HORIZON

(the optimal solution is computed at every time k)

INFINITE HORIZON CONSTRAINED PROBLEM

$u(i|k)$ control action at time $k+i$ computed at time k

$$\begin{aligned} & \min_{\{u(0|k), u(1|k), \dots\}} J(x(k), \{u(0|k), u(1|k), \dots\}) = \\ & = \min_{\{u(0|k), u(1|k)\}} \sum_{k=0}^{\infty} \left(\|x(i|k)\|_Q^2 + \|u(i|k)\|_R^2 \right) \end{aligned}$$

s.t.

$$F x(i|k) + G u(i|k) \leq b, \quad i = 0, 1, \dots$$

DUAL-MODE Divide the infinite-horizon problem in two problems

• MODE 1

FINITE-HORIZON CONSTRAINED PROBLEM

$$\begin{aligned} & \min_{\substack{\{u(0|h), u(1|h), \dots, u(N-1|h)\} \\ \text{s.t.}}} J_{RH}(x(h), \{u(0|h), \dots, u(N-1|h)\}) \\ & Fx(i|h) + Gu(i|h) \leq b, \quad i = 0, 1, \dots, N-1 \end{aligned}$$

• MODE 2

INFINITE-HORIZON UNCONSTRAINED PROBLEM

$$\begin{aligned} & \min_{\{u(N|h), u(N+1|h), \dots\}} J(x(N|h)) \quad \leftarrow \text{LQR} \end{aligned}$$

$$\Rightarrow \begin{cases} u(i|h) = K^* x(N|h), \quad i = N, N+1, \dots \\ J^*(x(N|h)) = \|x(N|h)\|_w^2 \end{cases} \quad \leftarrow K^*, w \text{ are known}$$

- We use the knowledge of the inf. horiz. solution to modify J in the finite horizon problem

$$J_{RH}(x(k), \underline{u}(k)) = \underbrace{\sum_{i=0}^{N-1} \left(\|x(i|k)\|_Q^2 + \|u(i|k)\|_R^2 \right)}_{J \text{ over the finite horizon}} + \underbrace{\|x(N|k)\|_W^2}_{\text{final cost (penalty term)}}$$

$$\underline{u}(k) = \{u(0|k), \dots, u(N-1|k)\}$$

J over the finite horizon

final cost (penalty term)

cost-to-go after N time steps under the optimal control law

$$u = K^* x$$

- WE CAN SOLVE NUMERICALLY THE ^{CONSTRAINED} FINITE-H PROBLEM

$$\min_{\underline{u}(k)} J_{RH}(x(k), \underline{u}(k))$$

$$\text{s.t.} \quad Fx(i|k) + G(i|k) \leq b, \quad i = 0, 1, \dots, N-1$$

\Rightarrow RECEDING HORIZON
OPTIMAL SOLUTION

FOR THE FINITE HORIZON CONSTRAINED PROBLEM

$$u(i|k) = g_{RH}(x(k)), \quad i = 0, \dots, N-1$$

• OPTIMAL SOLUTION OF THE CONSTRAINED INFINITE-HORIZON PROBLEM
DENOTED BY $\{u^{\infty}(0|k), u^{\infty}(1|k), \dots\}$

• UNDER $\{u^{\infty}(0|k), \dots\}$: $x(i|k) \rightarrow 0$ as i grows

• $(x, u) = (0, 0)$ is strictly feasible ($Fx + Gu \leq b$)

\Rightarrow under the optimal state feedback control law $u = K^*x$

\Rightarrow The constraints are $(F + GK^*)x \leq b$

\Rightarrow there exists a neighborhood S of $x=0$ such that

the constraints are met : $Fx + Gu \leq b, \forall x \in S$

\Rightarrow there exists a value \bar{N} under the optimal stabilizing control sequence such that $(F + GK^*) x(\bar{N} + i|h) \leq b, i = 0, 1, \dots$ are verified

$$\Rightarrow \min_{\{u(\bar{N}|h), u(\bar{N}+1|h), \dots\}} J(x(\bar{N}|h)) \equiv \min_{\{u(\bar{N}|h), \dots\}} J(x(\bar{N}|h))$$

s.t.

$$F x(i|h) + G u(i|h) \leq b, i = \bar{N}, \bar{N}+1, \dots$$

\sum after \bar{N} steps the constraints are not active

\Rightarrow

$$\begin{cases} u^\infty(i|h) = K^* x(i|h), i = \bar{N}, \bar{N}+1, \dots \\ J^*(x(\bar{N}|h)) = \|x(\bar{N}|h)\|_w^2 \end{cases}$$

We know the optimal solution of the constrained infinite-horizon problem from \bar{N} on (mode 2)

If $N \geq \bar{N}$ (if the prediction horizon of the MPC is "suff." long)

$$\left\{ \underbrace{u^*(0|k), u^*(1|k), \dots, u^*(N-1|k)}_{\text{solution of the finite-horizon constrained quadratic program. with terminal cos } \|x^*(N|k)\|_x^2}, \underbrace{K^* x^*(N|k), K^* x^*(N+1|k), \dots}_{\text{optimal solution of the LQR problem starting at time } k+N} \right\} =$$

solution of the finite-horizon constrained quadratic program. with terminal cos $\|x^*(N|k)\|_x^2$

optimal solution of the LQR problem starting at time $k+N$

$$= \left\{ u^\infty(0|k), u^\infty(1|k), \dots, u^\infty(N-1|k), u^\infty(N|k), u^\infty(N+1|k), \dots \right\}$$

↑ SOLUTION OF THE INFINITE-HORIZON CONSTRAINED QUADRATIC PROBLEM

Theorem 2 There exists a finite horizon \bar{N} , which depends on $x(k)$, such that, if $N \geq \bar{N}$:

i) $\underline{u}^*(k)$ coincides with $\{u^\infty(0|k), \dots, u^\infty(N-1|k)\}$

ii) $J_{RH}(x(k), \underline{u}^*(k)) = J^*(k)$

OPTIMAL COST OF
THE INF. HORIZON
CONSTRAINED
PROBLEM

