

计算线代

面向计算的线性代数

第一版

计线

面向计算的线性代数

第一版

张津睿
新域, 中国



Self Publishers Worldwide
Seattle San Francisco New York
London Paris Rome Beijing Barcelona

This book was typeset using L^AT_EX software.
本书给那些爱我与恨我的人。

-1 前言

现今市面上的线性代数教材非常充足，像我这样一个人来写一本教材必然会贻笑大方。但是我并不能放下那些我有时候自己脑海中的古怪想法和天马行空的路线。我是看面向算子的线性代数入门的，我也感觉非常庆幸能够在学生时代读到那样的好书。

这本书将基于一种面向计算的方式来讲解一般意义上的线性代数。线性代数，线性是一个非常广阔的话题，我非常惶恐地尝试将我在这片非常神奇的领域所学到的东西分享在这本书中。

这本书会回答所有我曾经感到非常困惑和未能够得到任意一位老师满意答案的问题。包括行列式为什么这么定义行向量和列向量到底有什么区别，转置究竟在干什么，为什么转置有和逆一样的性质（乘积的转置等于分别转置但是交换相乘），迹又是什么，为什么单单把对角线元素加起来就得到迹，这种搞笑的计算方法为什么有那么多神奇的性质，都是那里来的？

In this book, the main goal is to answer all the questions about those are not well pondered in the traditional linear algebra course of any kind. In the computational matrix theory there are plenty ways of decompose the matrix, such as LU, QR, and the most fundamental SVD decomposition. While in the more math oriented linear course the main topic are always in the most abstract way as they move forward. And the main concepts such as the Transpose, dule space, determinant, trace and all other fundamental but always treated just in a forced memorize level.

The main goal is to explain all the concepts in a more mathematical natrual and fun way, even the sudden expose of a certain defination would be a very hard problem to those first expose to the abstract algebra world

student just like me. So I'll try my best to get every defination and notion in a smooth and more reasonalbe way.

Table of Contents

-1 前言	v
1 0 杂谈	1
1.1 张量计算方法	1
1.1.1 (a,b)-张量 $\forall a, b \in \mathbb{N}$ 的例子	1
1.1.2 (0,1)-张量列向量	2
1.1.3 (1,0)-张量行向量	3
1.1.4 (1,1)-张量线性变换	3
1.1.5 (2,0)-张量双线性函数	3
1.1.6 练习	3
1.2 外代数计算方法	3
1.2.1 几何代数举例	3
1.2.2 二向量和三向量	3
1.2.3 点积楔积和三维向量场的通量	3
1.2.4 练习	3
2 1 张量(向量) 空间	5
2.1 张量基础	5
2.1.1 基底, 坐标分量, 协变与逆变	5
2.1.2 练习	5
3 2 使用数组表示	7
4 3 度量与度规张量	9

5 Re:1 向量空间 **11**

6 Re:2 线性映射与作为表示的矩阵 **13**

7 Re:3 内(外)积空间 **15**

7.1 内积, 作为一种双线性函数 15

7.1.1 内积, 带度量的空间 15

7.1.2 内积作为度量 15

7.1.3 练习 17

7.2 外积作为度量的可能性 17

7.2.1 几何代数举例 17

7.2.2 二向量和三向量 17

7.2.3 点积楔积和三维向量场的通量 17

7.2.4 练习 17

Index **19**

第 1 章 0 杂谈

数学并不能防止你受锤，但是在你的黄金时代认真地学习数学能防止你睡着，让你醒着受锤，清醒地感受疼痛，和绝望。但同时也给予你正确认识和描述世界的语言与改变世界的工具，希望你能用你手里的利剑斩开你的前路，希望创造，毁灭的力量就在你的手里了。

本章并不会证明地讲解若干中学代数之后可能继续发展代数的方向。

这里面主要还是以坐标计算为主，希望开发若干和高维空间解析几何和物理有密切关系的代数运算方法。主要以计算方法为主，并不会牵涉公理化的进世代代数方法。

1.1 张量计算方法

1.1.1 (a,b)-张量 $\forall a, b \in \mathbb{N}$ 的例子

$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ 是一个(1,1)-张量，有一个协变部分和一个逆变部分，这种张量其实就是矩阵。矩阵是线性变换的表示。

$\begin{bmatrix} 1 \\ 3 \end{bmatrix}$ 是一个(0,1)-张量，这种张量有一个逆变部分，也就是之后线性代数中的列向量。

$\begin{bmatrix} 1 & 3 \end{bmatrix}$ 是一个(1,0)-张量，这种张量有一个协变部分，也是之后线代中的协向量，对偶向量，行向量，线性泛函。

1.1.2 (0,1)-张量列向量

列向量没什么好说的，你写一个大括号，中间放一系列东西就行，也许不是数也行。

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

这样的向量就是列向量。

这样的一系列数是没有意义的。聚个例子，我有三个苹果一个桃子可以记作 $\begin{bmatrix} 3 \\ 1 \end{bmatrix}$ 同时我要是有三个房子一个汽车，我也可以记成 $\begin{bmatrix} 3 \\ 1 \end{bmatrix}$ 要是没有单位或者说“基底”来确定每一个分量代表的意义这个一系列数就没啥意义（或者至少在纯粹的数学以外，很难找到意义）

所以这两个 $\begin{bmatrix} 3 \\ 1 \end{bmatrix}$ 应该写成一些基向量的多少倍的和的形式。比如把苹果作为基底 \vec{e}_1 ，桃子作为基底 \vec{e}_2 ，房子记作基底 \vec{e}_1 ，汽车记作基底 \vec{e}_2 。对第一种情况就是 $3\vec{e}_1 + 1\vec{e}_2$ 对第二种情况就是 $3\vec{e}_1 + 1\vec{e}_2$ 我们先把这个东西记成一个更简洁的形式。第一种情况 $\begin{bmatrix} \vec{e}_1 & \vec{e}_2 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = 3\vec{e}_1 + 1\vec{e}_2$ 第二种情况 $\begin{bmatrix} \vec{e}_1 & \vec{e}_2 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = 3\vec{e}_1 + 1\vec{e}_2$ 这两个列向量的分量一样，但是因为基底不同可以有不同的意义。

textbook in linear algebra is very abundant, include the linear operator oriented approach, which is the currently best one I have ever seen.

This book will base on a computational oriented approach to the subject of the so called linear algebra. Linear algebra is such a vast topic that I'm having the most frightened heart to output what I have learnt so far in this amazing land.

In this book, the main goal is to answer all the questions about those are not well pondered in the traditional linear algebra course of any kind. In the computational matrix theory there are plenty ways of decompose the matrix, such as LU, QR, and the most fundamental SVD decomposition. While in the more math oriented linear course the main topic are always in the most abstract way as they move forward. And the main concepts such as the Transpose, dule space, determinant, trace and all other fundamental

but always treated just in a forced memorize level.

The main goal is to explain all the concepts in a more mathematical natural and fun way, even the sudden expose of a certain definition would be a very hard problem to those first expose to the abstract algebra world student just like me. So I'll try my best to get every definition and notion in a smooth and more reasonable way.

1.1.3 $(1,0)$ -张量行向量

1.1.4 $(1,1)$ -张量线性变换

1.1.5 $(2,0)$ -张量双线性函数

1.1.6 练习

1.2 外代数计算方法

1.2.1 几何代数举例

1.2.2 二向量和三向量

1.2.3 点积楔积和三维向量场的通量

1.2.4 练习

第 2 章 1 张量(向量) 空间

就像「0 杂谈」中说过的那样，单独写一个堆有一定结构的分量组成各种类型的张量并没有什么意义，除非我们在某种基底的意义下讨论个张量。

2.1 张量基础

2.1.1 基底, 坐标分量, 协变与逆变

2.1.2 练习

第 3 章 2 使用数组表示

第 4 章 3 度量与度规张量

第 5 章 Re:1 向量空间

第 6 章 Re:2 线性映射与作为表示的矩阵

第7章 Re:3 内(外)积空间

这一章我们将会回答那些我曾经非常希望知道答案的问题，关于行列式，迹，转置，对偶与各种算子（矩阵）分解的相关结论和性质。我希望当看完这本书全书，和之后这一章后读者能够有一种近乎直观的感受感受到这些非常繁琐繁多的性质其实可以非常显然与直观。

7.1 内积，作为一种双线性函数

7.1.1 内积，带度量的空间

$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ 是一个(1,1)-张量，有一个协变部分和一个逆变部分，这种张量其实就是矩阵。矩阵是线性变换的表示。

$\begin{bmatrix} 1 \\ 3 \end{bmatrix}$ 是一个(0,1)-张量，这种张量有一个逆变部分，也就是之后线性代数中的列向量。

$\begin{bmatrix} 1 & 3 \end{bmatrix}$ 是一个(1,0)-张量，这种张量有一个协变部分，也是之后线代中的协向量，对偶向量，行向量，线性泛函。

7.1.2 内积作为度量

列向量没什么好说的，你写一个大括号，中间放一列东西就行，也许

不是数也行。 $\begin{bmatrix} a \\ b \\ c \end{bmatrix}$

这样的向量就是列向量。

这样的一列数是没有意义的。聚个例子，我有三个苹果一个桃子可以记作 $\begin{bmatrix} 3 \\ 1 \end{bmatrix}$ 同时我要是有三个房子一个汽车，我也可以记成 $\begin{bmatrix} 3 \\ 1 \end{bmatrix}$ 要是没有单位或者说“基底”来确定每一个分量代表的意义这个一列数就没啥意义（或者至少在纯粹的数学以外，很难找到意义）

所以这两个 $\begin{bmatrix} 3 \\ 1 \end{bmatrix}$ 应该写成一些基向量的多少倍的和的形式。比如把苹果作为基底 \vec{e}_1 ，桃子作为基底 \vec{e}_2 ，房子记作基底 \vec{e}_1 ，汽车记作基底 \vec{e}_2 。对第一种情况就是 $3\vec{e}_1 + 1\vec{e}_2$ 对第二种情况就是 $3\vec{e}_1 + 1\vec{e}_2$ 我们先把这个东西记成一个更简洁的形式。第一种情况 $\begin{bmatrix} \vec{e}_1 & \vec{e}_2 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = 3\vec{e}_1 + 1\vec{e}_2$ 第二种情况 $\begin{bmatrix} \vec{e}_1 & \vec{e}_2 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = 3\vec{e}_1 + 1\vec{e}_2$ 这两个列向量的分量一样，但是因为基底不同可以有不同的意义。

textbook in linear algebra is very abundant, include the linear operator oriented approach, which is the currently best one I have ever seen.

This book will base on a computational oriented approach to the subject of the so called linear algebra. Linear algebra is such a vast topic that I'm having the most firghtened heart to output what I have learnt so far in this amazing land.

In this book, the main goal is to answer all the questions about those are not well pondered in the traditional linear algebra course of any kind. In the computational matrix theory there are plenty ways of decompose the matrix, such as LU, QR, and the most fundamental SVD decomposition. While in the more math oriented linear course the main topic are always in the most abstract way as they move forward. And the main concepts such as the Transpose, dule space, determinant, trace and all other fundamental but always treated just in a forced memorize level.

The main goal is to explain all the concepts in a more mathematical natrual and fun way, even the sudden expose of a certain defination would be a very hard problem to those first expose to the abstract algebra world student just like me. So I'll try my best to get every defination and notion in a smooth and more reasonalbe way.

7.1.3 练习

7.2 外积作为度量的可能性

7.2.1 几何代数举例

7.2.2 二向量和三向量

7.2.3 点积楔积和三维向量场的通量

7.2.4 练习

索引

A1, 3, 5, 17

A2, 3, 5, 17

B1, 3, 5, 17

B2, 3, 5, 17

C1, 3, 5, 17

C2, 3, 5, 17

D1, 3, 5, 17

D2, 3, 5, 17

E1, 3, 5, 17

E2, 3, 5, 17

F1, 3, 5, 17

F2, 3, 5, 17

G1, 3, 5, 17

G2, 3, 5, 17

H1, 3, 5, 17

H2, 3, 5, 17

I1, 3, 5, 17

I2, 3, 5, 17

J1, 3, 5, 17

J2, 3, 5, 17

K1, 3, 5, 17

K2, 3, 5, 17

L1, 3, 5, 17

L2, 3, 5, 17

M1, 3, 5, 17

M2, 3, 5, 17

N1, 3, 5, 17

N2, 3, 5, 17

O1, 3, 5, 17

O2, 3, 5, 17

P1, 3, 5, 17

P2, 3, 5, 17

Q1, 3, 5, 17

Q2, 3, 5, 17

R1, 3, 5, 17

R2, 3, 5, 17

S1, 3, 5, 17

S2, 3, 5, 17

T1, 3, 5, 17

T2, 3, 5, 17

U1, 3, 5, 17

U2, 3, 5, 17

V1, 3, 5, 17

V2, 3, 5, 17

W1, 3, 5, 17

W2, 3, 5, 17

X1, 3, 5, 17

X2, 3, 5, 17

Y1, 3, 5, 17

Y2, 3, 5, 17

Z1, 3, 5, 17

Z2, 3, 5, 17