

# Roots: From Rick to Recent Research

Brian Harbourne

Cather Professor of Mathematics  
Fellow of the AMS  
Department of Mathematics  
University of Nebraska-Lincoln (UNL)

Joy in Algebraic Geometry:  
A conference in honor of Rick Miranda for his 70th birthday  
September 28, 2023  
Levico Terme

<https://unlblh.github.io/BrianHarbourne/index.html>

# Introduction


Root systems have a history of appearing in algebraic geometry in interesting ways.

They have come up (at least tangentially) in Rick's work and they have come up in some recent research, in a way related to a problem Rick has worked on.

I want to tell you about this recent research, but first let me highlight some of Rick's related work.

So let's go way back...

Rick, Jeanne and I shared a grad school office in the 70s



Me

Rick

Jeanne



MIT, 2-229

The office is no longer there. James Simons gave MIT a ton of money to get rid of it.

We aren't the same either! Can you spot Rick?

From 1976:



Thanks to Barbara Peskin for the photo!


# In those days Rick worked on GIT for cubic pencils

**Thesis:** “On the stability of rational elliptic surfaces with section”. '79

**1st paper:** “On the stability of pencils of cubic curves”. Amer. J. '80

**Theorem:** A pencil is stable (GIT sense) if and only if it contains a smooth member and every fiber of the elliptic surface obtained by blowing up base points is reduced.


The non-stable fibers are reducible, classified by Dynkin diagrams of affine root systems:



## Rick and Ulf Persson

“On extremal rational elliptic surfaces,” Math. Z. '86

Results: Rick and Ulf studied cubic pencils with finitely many sections.



*extremal* = finitely many  $(-1)$ -curves

$\sigma$  = number of  $(-1)$ -curves

*Jacobian* = has sections so no multiple fibers

This paper gives a nice formula for  $\sigma$  in terms of the fibers  $F$ .

$r_F^2$  = number of reduced components of  $F$

**Theorem:**  $\sigma = \prod_F r_F$

## The non-Jacobian case...Rick and Me

“Exceptional curves on rational numerically elliptic surfaces,” J. Alg. '90

A non-Jac. extremal rational elliptic surf. has unique multiple fiber.

$m$  = multiplicity of multiple fiber

$r_F(t)$  = power series defined in terms of the Dynkin diagram of  $F$

$\pi(t) = \prod_F r_F(t)$  = the Hadamard product:  $\sum_i a_i t^i \sum_i b_i t^i = \sum_i a_i b_i t^i$

$(\pi(t))_m$  = coefficient of  $t^m$  in  $\pi(t)$

**Theorem:** Except in special cases

$$\sigma = \left( \prod_F r_F(t) \right)_m .$$

(Results are given in all cases, but the full result is more complicated.)

# Interpolation problems

Ciro, Quim and Olivia have all done work on this with Rick.





## Interpolation problems (continued)

$R_d$ : all forms on  $\mathbb{P}^2$  of degree  $d$ , so  $R = \mathbb{C}[\mathbb{P}^2] = \bigoplus_d R_d$ .

$\overline{mP} = m_1 P_1 + \cdots + m_s P_s \subset \mathbb{P}^2$ : scheme defined by all forms vanishing to order  $\geq m_i$  at general points  $P_i$ .

$[I(\overline{mP})]_d = I(\overline{mP}) \cap R_d$ , so  $I(\overline{mP}) \subset R$  is the ideal of  $\overline{mP}$ .

**Problem:** Given  $\overline{m} = (m_1, \dots, m_s)$  and  $d$ , find  $\dim[I(\overline{mP})]_d$ .

There's a Cremona group  $G$  which acts to reduce the data  $(d, m_1, \dots, m_s)$  to the case (\*)  $d \geq m_1 + m_2 + m_3$  with  $m_1 \geq m_2 \geq \cdots \geq m_s \geq 0$ .

**SHGH Conjecture:** Given (\*), then

$$\dim[I(\overline{mP})]_d = \max \left\{ 0, \dim R_d - \sum_{m_i > 0} \binom{m_i + 1}{2} \right\}.$$

This group  $G$  is the Weyl group of a Dynkin diagram. See "On the Kantor group of a set of points in a plane", PLMS '37,

**Patrick Du Val**, right, for the case  $s \leq 8$ .



## SHGH: Current status



B. Segre  
1961



BH  
1986



A. Gimigliano  
1987



A. Hirschowitz  
1989

**Theorem** All four versions are equivalent. (Ciro and Rick, 2001)

Situation understood for  $s \leq 9$ . (Castelnuovo, 1891)

The conjecture is true when  $12 \geq m_1 = \dots = m_s$  (Ciro and Rick, 2000)  
and for  $m_1 = \dots = m_{10}$  when  $\frac{d}{m_1} < \frac{117}{37}$  (Ciro, Rick, Quim, Olivia, 2011).

But it's still open!

Idea: Could it help to study versions of a more general problem?

## Unexpectedness: Cook, H., Migliore, Nagel, Compositio '18

Original SHGH problem:  $n = 2$ ,  $R_d = [I(Z)]_d$  for  $Z = \emptyset$ ,  $\overline{mP} \subset \mathbb{P}^2$ ,  $P_i$  general. We say  $Z$  has *unexpected curves* in degree  $d$  and multiplicity  $\overline{m}$  if

$$\dim[I(Z \cup \overline{mP})]_d > \max\{0, \dim[I(Z)]_d - \sum_i \binom{m_i + 1}{2}\}.$$

The SHGH Conjecture accounts for all known unexpectedness.


New problem: any  $n$ ,  $Z = q_1 + \cdots + q_r \subset \mathbb{P}^n$  any given points  $q_i$ ,  $mP \subset \mathbb{P}^n$ ,  $P$  general. We say  $Z$  has *unexpected hypersurfaces* in degree  $d$  and multiplicity  $m$  if

$$\dim[I(Z \cup mP)]_d > \max\{0, \dim[I(Z)]_d - \binom{m + n - 1}{n}\}.$$

Problem (CHMN): To understand when  $Z, m, d$  is unexpected.



## First example (Di Gennaro, Ilardi, Vallès, '14)



$Z$  is a set of 9 points in  $\mathbb{P}^2$  consisting of:

4 general points (red) which give a pencil of conics;

3 points (white), singular points of the singular conics;

2 points (black) where a singular conic meets the line through the singular points of the other 2 singular conics.


**Fact:**  $Z$  has an unexpected quartic with a general triple point; we expect no such quartic. What does this have to do with roots?



R. Di Gennaro



G. Ilardi



J. Vallès

# Root systems and unexpectedness

See H., Migliore, Nagel, Teitler, Mich. J. '21

Projectivizing the  $B_3$  root system gives the 9 points;  $Z = Z(B_3)$ !

Projectivizing  $D_4$  gives 12 points  $Z(D_4) \subset \mathbb{P}^3$  with two unexpected cones: a cubic and a quartic.


Projectivizing  $F_4$  gives 24 points  $Z(F_4) \subset \mathbb{P}^3$  with two unexpected cones: a quartic and a sextic.

Projectivizing  $H_4$  gives 60 points  $Z(H_4) \subset \mathbb{P}^3$  with two unexpected cones: a sextic and decic (see Wiśniewska-Zięba, [arXiv:2107.08107](https://arxiv.org/abs/2107.08107)).



Teitler and a new recruit

Trying to understand unexpectedness is an expanding area of research. For example ...



P. Wiśniewska



M. Zięba

## General projections to a complete intersection: Geproci

**Definition:** We say a finite set  $Z \subset \mathbb{P}^3$  is  $(a, b)$ -geproci if its image  $\overline{Z}$  under projection to a plane from a general point  $P \in \mathbb{P}^3$  is an  $(a, b)$ -complete intersection.

**Theorem** (Levico Terme Working Group, 2018): A finite noncoplanar  $Z \subset \mathbb{P}^3$  is  $(a, b)$ -geproci if  $|Z| = ab$  and  $Z$  has unexpected cones of degrees  $a \leq b$  with no common components.

**Corollary:**  $Z(D_4)$ ,  $Z(F_4)$  and  $Z(H_4)$  are geproci!

**Open Problems:** (1) What other kinds of sets are geproci?

(2) If  $Z \subset \mathbb{P}^3$  is noncoplanar and  $(a, b)$ -geproci with  $3 \leq a \leq b$ , must  $Z$  have unexpected cones of degrees  $a$  and  $b$ ?

(3) Does every noncoplanar geproci set  $Z$  have a nontrivial matroid (e.g., does it have subsets of 3 collinear points)?

# Levico Terme Working Group, 2018



Alessandra  
Bernardi



Luca  
Chiantini




Graham  
Denham



Giuseppe  
Favacchio



Brian  
Harbourne



Juan  
Migliore



Tomasz  
Szemberg



Justyna  
Szpond




# Classifying geproci sets is an inverse scattering problem

Studying inverse scattering problems has led to remarkable advances in scientific knowledge. Here we propose carrying this idea over to classification problems in algebraic geometry.


Inverse scattering Problems (ISP):  
try to discern structure from  
projected or reflected data.

Idea: classify structures  
algebro-geometrically based on  
properties of projected images.




# Some examples of ISP


Echolocation (biology):



Rutherford scattering (physics; led to Bohr model of atom):




X-ray crystallography (chem/bio; led to DNA double helix model):



## More examples

Tomography (medicine):



GePro- $\mathcal{P}$  (math): Pick a property  $\mathcal{P}$  and classify finite point sets  $Z \subset \mathbb{P}^n$  whose General Projections  $\overline{Z}$  to a hyperplane satisfy  $\mathcal{P}$ .

**Example 1:** Say  $\mathcal{P}$  means “ $\overline{Z}$  is Gorenstein”. Then a set  $Z$  of  $n + 1$  general points in  $\mathbb{P}^n$  is gepro- $\mathcal{P}$  since the image  $\overline{Z}$  is a set of  $n + 1$  general points in  $H$ , which is Gorenstein.

**Open Problem 1:** Classify gepro-Gorenstein sets  $Z$ .

Geproci is when  $\mathcal{P}$  means “ $\overline{Z}$  is a complete intersection”. Every geproci set is also gepro-Gorenstein but not conversely.

**Open Problem 2:** Classify geproci sets in  $\mathbb{P}^n$  for  $n \geq 3$ .

## Some history: Polizzi and Panov


We know no interesting examples of geproci sets in  $\mathbb{P}^n$  for  $n > 3$ .

A geproci set  $Z$  in a plane  $H \subset \mathbb{P}^3$  is called *degenerate*; it is just the complete intersection of two curves in  $H$ .


**Question 1** (F. Polizzi 2011): Is every geproci set in  $\mathbb{P}^3$  degenerate?

**Answer** (D. Panov, 2011): No!


Nondegenerate geproci sets are given by  $(a, b)$ -grids (i.e., sets  $Z$  of  $ab$  points where  $Z = A \cap B$  and  $A$  consists of  $a$  skew lines and  $B$  consists of  $b$  skew lines).



Francesco  
Polizzi



and Dimitri  
Panov



## What we now know: POLITUS and Kettinger

Call a geproci  $Z \subset \mathbb{P}^3$  *trivial* if it is a grid or degenerate.

Call a nontrivial geproci  $Z$  a *half-grid* when  $\overline{Z}$  is the complete intersection of two curves if one curve is a union of lines.

Most known examples over  $\mathbb{C}$  are half-grids, including  $Z(D_4)$  and  $Z(F_4)$ . The POLITUS research group has found many more half-grids closely related to the combinatorics of skew lines in  $\mathbb{P}^3$ .

We know only 3 examples over  $\mathbb{C}$  of nontrivial geproci non-half-grids:  $Z(H_4)$  and two examples used in quantum mechanics, a  $(5, 8)$ -geproci and a  $(10, 12)$ -geproci. Things are different in positive characteristics.

**Theorem**(Kettinger [arXiv:2307.04857](https://arxiv.org/abs/2307.04857)): Let  $\mathbb{F}$  be a finite field,  $q = |\mathbb{F}|$ . Then the points of  $\mathbb{P}_{\mathbb{F}}^3$  are  $(q + 1, q^2 + 1)$ -geproci.

Moreover, for odd  $q \geq 7$  there are nontrivial non-half-grid  $(q + 1, d)$ -geproci sets  $Z \subset \mathbb{P}_{\mathbb{F}}^3$  whenever  $q + 1 < d < \frac{q^2 + 1}{2} - 6$ .

## Jake Kettinger

Jake's results are closely related to the notion of a *spread* (a topic in combinatorics), namely the  $(q+1)b$  points on a set of  $b \leq q^2 + 1$  skew lines in  $\mathbb{P}_{\mathbb{F}}^3$  over a finite field  $\mathbb{F}$  of order  $q$ . Thanks to the Hopf fibration,  $\mathbb{P}_{\mathbb{F}}^3$  is itself a spread. Jake's nontrivial non-half-grids come from maximal partial spreads.



# Main POLITUS references

POLITUS 1: [arXiv:2209.04820](https://arxiv.org/abs/2209.04820)


POLITUS 2: [arXiv:2308.00761](https://arxiv.org/abs/2308.00761)



Łucja and Karolina Farnik, Tomasz Szemberg, Justyna Szpond, Luca Chiantini, Giuseppe Favacchio, Juan Migliore, Me.

# Combinatorics of skew lines? POLITUS 2

Any set  $\mathcal{L} = \{L_1, L_2, L_3, \dots, L_b\}$  of skew lines  $L_i \subset \mathbb{P}^3$  has an associated groupoid  $G(\mathcal{L})$  generated by the maps  $\gamma_{ijk} : L_i \rightarrow L_j$  where, given  $p \in L_i$ ,  $\gamma_{ijk}(p)$  is the point  $q \in L_j$  such that  $\overline{pq}$  is contained in the unique quadric  $Q$  containing  $L_i, L_j, L_k$ .



If  $p \in \cup_i L_i$ , the images  $\phi(p)$  of  $p$  under all the maps  $\phi \in G(\mathcal{L})$  for which  $\phi(p)$  is defined is the *orbit* of  $p$  under the groupoid.

**Theorem** (POLITUS 2) A geproci half-grid of  $a$  points on each of the  $b$  lines of  $\mathcal{L}$  is a finite union of orbits. Conversely, if  $Z$  is a finite union of finite orbits consisting of  $a \geq b - 1 \geq 3$  points on each of the  $b$  lines, then  $Z$  is a half-grid  $(a, b)$ -geproci set.



## Some open problems...

1. Every grid in  $\mathbb{P}^3$  is contained in a quadric. If  $Z$  is geproci and contained in a quadric, is  $Z$  a grid?
2. Are there nontrivial non-half-grids over  $\mathbb{C}$  than the 3 we know?
3. The nontrivial non-half-grid  $(5, 8)$ -geproci is Gorenstein. Are there other Gorenstein geproci sets?
4. Is there a finite set of points in  $\mathbb{P}^n$  that is a nontrivial geproci set when  $n > 3$ ?

Thanks to Rick for giving us a reason to gather here.  
Happy 70th!

MFO Concert: 2010

