

# 90% Confidence Level Upper Bound

Brief discussion of Feldman & Cousins

10-31-2024

# Outline

- Parameter Estimataion ..... 3
- Confidence Interval (CI) ..... 5
- Standard CIs ..... 12
- Feldman & Cousins CI ..... 15
- F&C CI construction ..... 22
- Bayesian ..... 29

# Outline

Outline .....	1	Example acceptance region for $\mu = 0.5$ .....	27
Parameter Estimataion .....	3	Bayesian .....	29
Poisson Distribution .....	4	Bayes Theorem .....	30
Confidence Interval (CI) .....	5	Summary: Bayesian Credible Interval .....	32
CI Definition .....	6		
CI Construction: Confidence Belt .....	8		
Standard CIs .....	12		
Acceptance region? .....	13		
Example CIs .....	14		
Feldman & Cousins CI .....	15		
Question: which one to use? .....	16		
Problem: “flip-flop” .....	17		
Standard Poisson (90% Upper) CI .....	18		
Comparison: look at small n .....	19		
Standard Poisson (90% central) CI .....	20		
Comparison: look at large n .....	21		
F&C CI construction .....	22		
Maximum Likelihood .....	23		
Likelihood Ratio .....	25		

# Parameter Estimataion

---

- Expect an average of (real)  
 $\mu \geq 0$  neutrinos per time

- Expect an average of (real)  
 $\mu \geq 0$  neutrinos per time
- Probability of seeing (integer)  
 $n \geq 0$  neutrinos per time?

- Expect an average of (real)  
 $\mu \geq 0$  neutrinos per time
- Probability of seeing (integer)  
 $n \geq 0$  neutrinos per time?
- Poisson Distribution:

$$\Pr(n|\mu) = \frac{e^{-\mu} \mu^n}{n!}$$

- Prob. of data  $n$ , given parameter  $\mu$ ; aka the *likelihood*.

# Poisson Distribution


## Parameter Estimataion

- Expect an average of (real)  
 $\mu \geq 0$  neutrinos per time
- Probability of seeing (integer)  
 $n \geq 0$  neutrinos per time?
- Poisson Distribution:

$$\Pr(n|\mu) = \frac{e^{-\mu} \mu^n}{n!}$$

- Prob. of data  $n$ , given parameter  $\mu$ ; aka the *likelihood*.

- Example:





# Confidence Interval (CI)

---

- Goal: estimate parameter  $\mu$  whose true value is  $\mu_t$

- Goal: estimate parameter  $\mu$  whose true value is  $\mu_t$
- Make a measurement  $x$ ; first measurement yields  $x_0$ .

- Goal: estimate parameter  $\mu$  whose true value is  $\mu_t$
- Make a measurement  $x$ ; first measurement yields  $x_0$ .
- Construct an interval (discussed later)  $[\mu_l, \mu_u]$

- Goal: estimate parameter  $\mu$  whose true value is  $\mu_t$
- Make a measurement  $x$ ; first measurement yields  $x_0$ .
- Construct an interval (discussed later)  $[\mu_l, \mu_u]$ 
  - $\mu_l = \mu_l(x_0)$ : lower bound associated w/ this 1<sup>st</sup> measurement

- Goal: estimate parameter  $\mu$  whose true value is  $\mu_t$
- Make a measurement  $x$ ; first measurement yields  $x_0$ .
- Construct an interval (discussed later)  $[\mu_l, \mu_u]$ 
  - $\mu_l = \mu_l(x_0)$ : lower bound associated w/ this 1<sup>st</sup> measurement
  - $\mu_u = \mu_u(x_0)$ : upper bound associated w/ this 1<sup>st</sup> measurement

- Goal: estimate parameter  $\mu$  whose true value is  $\mu_t$
- Make a measurement  $x$ ; first measurement yields  $x_0$ .
- Construct an interval (discussed later)  $[\mu_l, \mu_u]$ 
  - $\mu_l = \mu_l(x_0)$ : lower bound associated w/ this 1<sup>st</sup> measurement
  - $\mu_u = \mu_u(x_0)$ : upper bound associated w/ this 1<sup>st</sup> measurement
- Repeat experiment; get outcome  $x_1 \rightarrow$  construct  $[\mu_l(x_1), \mu_u(x_1)]$

- Goal: estimate parameter  $\mu$  whose true value is  $\mu_t$
- Make a measurement  $x$ ; first measurement yields  $x_0$ .
- Construct an interval (discussed later)  $[\mu_l, \mu_u]$ 
  - $\mu_l = \mu_l(x_0)$ : lower bound associated w/ this 1<sup>st</sup> measurement
  - $\mu_u = \mu_u(x_0)$ : upper bound associated w/ this 1<sup>st</sup> measurement
- Repeat experiment; get outcome  $x_1 \rightarrow$  construct  $[\mu_l(x_1), \mu_u(x_1)]$
- More experiments; get a bunch of intervals. *i.e.* we get a set

$$C \equiv \{[\mu_l(x_0), \mu_u(x_0)], [\mu_l(x_1), \mu_u(x_1)], [\mu_l(x_2), \mu_u(x_2)] \dots\}$$



- $C \equiv \{[\mu_l, \mu_u], [\mu_l, \mu_u], [\mu_l, \mu_u] \dots\}$
- The set  $C$  has the property that

$$P([\mu_l, \mu_u] \ni \mu_t) = \alpha\%$$

- $C \equiv \{[\mu_l, \mu_u], [\mu_l, \mu_u], [\mu_l, \mu_u] \dots\}$
- The set  $C$  has the property that

$$P([\mu_l, \mu_u] \ni \mu_t) = \alpha\%$$

- In words:
  - the true value  $\mu_t$  is *fixed* (although unknown)

- $C \equiv \{[\mu_l, \mu_u], [\mu_l, \mu_u], [\mu_l, \mu_u] \dots\}$
- The set  $C$  has the property that

$$P([\mu_l, \mu_u] \ni \mu_t) = \alpha\%$$

- In words:
  - the true value  $\mu_t$  is *fixed* (although unknown)
  - Some members of  $C$  would *cover* this true value

- $C \equiv \{[\mu_l, \mu_u], [\mu_l, \mu_u], [\mu_l, \mu_u] \dots\}$
- The set  $C$  has the property that

$$P([\mu_l, \mu_u] \ni \mu_t) = \alpha\%$$

- In words:
  - the true value  $\mu_t$  is *fixed* (although unknown)
  - Some members of  $C$  would *cover* this true value
  - e.g. say  $\mu_t = 2$ , and maybe in  $C$  there is  $[1, 3]$ .

- $C \equiv \{[\mu_l, \mu_u], [\mu_l, \mu_u], [\mu_l, \mu_u] \dots\}$
- The set  $C$  has the property that

$$P([\mu_l, \mu_u] \ni \mu_t) = \alpha\%$$

- In words:
  - the true value  $\mu_t$  is *fixed* (although unknown)
  - Some members of  $C$  would *cover* this true value
  - e.g. say  $\mu_t = 2$ , and maybe in  $C$  there is  $[1, 3]$ .
- The members of  $C$  are called *confidence intervals*.

# CI Construction: Confidence Belt

Confidence Interval (CI)

- Recall *likelihood*  $\Pr(x|\mu)$   
(ie. probability of data,  
assume known  
parameter)

# CI Construction: Confidence Belt


Confidence Interval (CI)

- Recall *likelihood*  $\Pr(x|\mu)$   
(ie. probability of data,  
assume known  
parameter)
- Take for example  $\mu = 4$ .  
Expect 4 neutrinos per  
time.

# CI Construction: Confidence Belt

Confidence Interval (CI)

- Recall *likelihood*  $\Pr(x|\mu)$   
(ie. probability of data,  
assume known  
parameter)
- Take for example  $\mu = 4$ .  
Expect 4 neutrinos per  
time.






# CI Construction: Confidence Belt

Confidence Interval (CI)


- Recall *likelihood*  $\Pr(x|\mu)$   
(ie. probability of data,  
assume known  
parameter)
- Take for example  $\mu = 4$ .  
Expect 4 neutrinos per  
time.
- Select a region  $[x_l, x_u]$   
such that the probability  
of measuring  $x \in [x_l, x_u]$   
is, say, 80%.



# CI Construction: Confidence Belt

Confidence Interval (CI)


- Recall *likelihood*  $\Pr(x|\mu)$   
(ie. probability of data,  
assume known  
parameter)
- Take for example  $\mu = 4$ .  
Expect 4 neutrinos per  
time.
- Select a region  $[x_l, x_u]$   
such that the probability  
of measuring  $x \in [x_l, x_u]$   
is, say, 80%.



# CI Construction: Confidence Belt

Confidence Interval (CI)


- Recall *likelihood*  $\Pr(x|\mu)$   
(ie. probability of data,  
assume known  
parameter)
- Take for example  $\mu = 4$ .  
Expect 4 neutrinos per  
time.
- Select a region  $[x_l, x_u]$   
such that the probability  
of measuring  $x \in [x_l, x_u]$   
is, say, 80%.



# CI Construction: Confidence Belt

Confidence Interval (CI)


- But of course  $\mu_t$  is unknown.



# CI Construction: Confidence Belt

Confidence Interval (CI)


- But of course  $\mu_t$  is unknown.
- The true flux  $\mu_t$  could be, say,  $\mu = 5$



# CI Construction: Confidence Belt

Confidence Interval (CI)


- But of course  $\mu_t$  is unknown.
- The true flux  $\mu_t$  could be, say,  $\mu = 5$
- So, construct another acceptance region  $[x_l, x_u]$  for  $\mu = 5$



# CI Construction: Confidence Belt

Confidence Interval (CI)


- But of course  $\mu_t$  is unknown.
- The true flux  $\mu_t$  could be, say,  $\mu = 5$
- So, construct another acceptance region  $[x_l, x_u]$  for  $\mu = 5$
- Rinse and repeat



# CI Construction: Confidence Belt

Confidence Interval (CI)

- But of course  $\mu_t$  is unknown.
- The true flux  $\mu_t$  could be, say,  $\mu = 5$
- So, construct another acceptance region  $[x_l, x_u]$  for  $\mu = 5$
- Rinse and repeat






# CI Construction: Confidence Belt

Confidence Interval (CI)


- Make a measurement, get result  $x_0 = 6$



# CI Construction: Confidence Belt

Confidence Interval (CI)


- Make a measurement, get result  $x_0 = 6$
- The probability of  $x_0$  falling in the acceptance region (red) is 80%, by construction



# CI Construction: Confidence Belt

Confidence Interval (CI)


- Make a measurement, get result  $x_0 = 6$
- The probability of  $x_0$  falling in the acceptance region (red) is 80%, by construction
- The **confidence interval**  $[\mu_l, \mu_u]$  from this experiment is the vertical intercept.



# CI Construction: Confidence Belt

Confidence Interval (CI)


- Make some more measurements



# CI Construction: Confidence Belt

Confidence Interval (CI)

- Make some more measurements
- Get some more confidence intervals.



# CI Construction: Confidence Belt


Confidence Interval (CI)

- Make some more measurements
- Get some more confidence intervals.

- Have a set

$$C = \{CI_1, CI_2, CI_3, CI_4, CI_5\}$$

- 80% of this set would cover the true value,  $\mu_t$ .



# Standard CIs

---

- Recall acceptance region:

$$\Pr(n \in [n_1, n_2] \mid \mu_{\text{fixed}}) = 90\%$$

---

<sup>1</sup>See Feldman & Cousins Section II.B




# Acceptance region?

- Recall acceptance region:

$$\Pr(n \in [n_1, n_2] \mid \mu_{\text{fixed}}) = 90\%$$

- Complete freedom in choosing a 90% range.



---


<sup>1</sup>See Feldman & Cousins Section II.B

# Acceptance region?

- Recall acceptance region:

$$\Pr(n \in [n_1, n_2] \mid \mu_{\text{fixed}}) = 90\%$$

- Complete freedom in choosing a 90% range.
- A choice<sup>1</sup> leads to the **upper limit**.



---


<sup>1</sup>See Feldman & Cousins Section II.B

# Acceptance region?

- Recall acceptance region:

$$\Pr(n \in [n_1, n_2] \mid \mu_{\text{fixed}}) = 90\%$$

- Complete freedom in choosing a 90% range.
- A choice<sup>1</sup> leads to the **upper limit**.
- Another common choice leads to the two-sided **central interval**.




---

<sup>1</sup>See Feldman & Cousins Section II.B

# Example CIs

## Standard CIs




Gaussian likelihood with known  $\sigma = 1$

$$\Pr(x \mid \mu) = \frac{1}{\sqrt{2\pi}} \cdot e^{-(x-\mu)^2/2}$$


standard 90% upper limit

# Example CIs

## Standard CIs



standard 90% upper limit





standard 90% central CI

# Feldman & Cousins CI

---

# Question: which one to use?


Feldman & Cousins CI




standard 90% upper limit

standard 90% central CI

# Problem: “flip-flop”




- One **cannot** pick the type of interval after-the-fact. The Choice is made before we perform the experiment. Otherwise we would be “flip-flopping”.








# Problem: “flip-flop”




- One **cannot** pick the type of interval after-the-fact. The Choice is made before we perform the experiment. Otherwise we would be “flip-flopping”.
- It turns out that flip-flopping leads to invalid intervals



# Problem: “flip-flop”



- One **cannot** pick the type of interval after-the-fact. The Choice is made before we perform the experiment. Otherwise we would be “flip-flopping”.
- It turns out that flip-flopping leads to invalid intervals
- Feldman & Cousins’s approach removes the possibility of (or motivation to commit) flip-flopping.




Poisson likelihood with background


$$\mathcal{L} \equiv \text{Pr}(n \mid \mu) = \frac{(\mu + b)^n e^{-(\mu+b)}}{n!}$$

# Comparison: look at small n


Feldman & Cousins CI



standard 90% upper CI




Feldman & Cousins




# Comparison: look at large n

Feldman & Cousins CI




standard 90% central CI



Feldman & Cousins

# F&C CI construction

---



- Recall acceptance region:

$$\Pr(n \in [n_1, n_2] \mid \mu_{\text{fixed}}) = 90\%$$

- Complete freedom in choosing how to construct the acceptance regions.
- Consider likelihood: Poisson with background  $b$ :

$$\mathcal{L} \equiv \Pr(n \mid \mu) = \frac{(\mu + b)^n e^{-(\mu+b)}}{n!}$$

- F&C propose to compute a likelihood ratio  $R$ 
  - This needs a “best fit”  $\mu_{\text{best}} \equiv \max(0, n - b)$



## Derivation

- Likelihood is a Poisson in this case.

$$\mathcal{L} \equiv \text{Pr}(n \mid \mu) = \frac{(\mu + b)^n e^{-(\mu+b)}}{n!}$$

- Find maximum (fixing  $n$ , vary  $\mu$ ):

$$\left. \frac{d\mathcal{L}}{d\mu} \right|_{\mu=\mu_{\text{best}}} = 0$$

- Result: “best fit”  $\mu = \mu_{\text{best}} = n - b$
- Require physical  $\mu \geq 0 \Rightarrow \mu_{\text{best}} = \max(0, n - b)$

- Do this for representative values of  $\mu$ ; say we start with  $\mu = 0.5$ 
  - As an example background,  $b = 3$
  - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
  - 
  - 
  - 
  -

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0					

- Do this for representative values of  $\mu$ ; say we start with  $\mu = 0.5$ 
  - As an example background,  $b = 3$
  - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
  - For  $n = 0$ , compute  $\Pr(n \mid \mu = 0.5)$
  - 
  - 
  -

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.03				

- Do this for representative values of  $\mu$ ; say we start with  $\mu = 0.5$ 
  - As an example background,  $b = 3$
  - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
  - For  $n = 0$ , compute  $\Pr(n \mid \mu = 0.5)$
  - For  $n = 0$ , compute  $\mu_{\text{best}} = 0$
  - 
  -

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.03	0			

- Do this for representative values of  $\mu$ ; say we start with  $\mu = 0.5$ 
  - As an example background,  $b = 3$
  - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
  - For  $n = 0$ , compute  $\Pr(n \mid \mu = 0.5)$
  - For  $n = 0$ , compute  $\mu_{\text{best}} = 0$
  - For  $n = 0$ , compute  $\Pr(n \mid \mu = \mu_{\text{best}})$
  -

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.03	0	0.05		


- Do this for representative values of  $\mu$ ; say we start with  $\mu = 0.5$ 
  - As an example background,  $b = 3$
  - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
  - For  $n = 0$ , compute  $\Pr(n \mid \mu = 0.5)$
  - For  $n = 0$ , compute  $\mu_{\text{best}} = 0$
  - For  $n = 0$ , compute  $\Pr(n \mid \mu = \mu_{\text{best}})$
  - Divide likelihoods to get  $R$ .

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.03	0	0.05	0.607	

- As a reminder, this is still just for  $\mu = 0.5$  (and example  $b = 3$ )

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.03	0	0.05	0.607	


- We will see that the acceptance region for  $\mu = 0.5$  is this:



- As a reminder, this is still just for  $\mu = 0.5$  (and example  $b = 3$ )

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.03	0	0.05	0.607	

- We will see that the acceptance region for  $\mu = 0.5$  is this:





- As a reminder, this is still just for  $\mu = 0.5$  (and example  $b = 3$ )

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.03	0	0.05	0.607	

- To construct the region, make a new row for  $n = 1$

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.030	0	0.050	0.607	
1	0.106	0	0.149	0.708	

# Example acceptance region for $\mu = 0.5$

F&C CI construction

And then for a bunch of other  $n$ .

# Example acceptance region for $\mu = 0.5$

F&C CI construction

$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.030	0	0.050	0.607	
1	0.106	0	0.149	0.708	
2	0.185	0	0.224	0.826	
3	0.216	0	0.224	0.963	
4	0.189	1	0.195	0.966	
5	0.132	2	0.175	0.753	
6	0.077	3	0.161	0.480	
7	0.039	4	0.149	0.259	

# Example acceptance region for $\mu = 0.5$

F&C CI construction


$n$	$\Pr(n \mu)$	$\mu_{\text{best}}$	$\Pr(n \mu_{\text{best}})$	$R$	rank
0	0.030	0	0.050	0.607	6
1	0.106	0	0.149	0.708	5
2	0.185	0	0.224	0.826	3
3	0.216	0	0.224	0.963	2
4	0.189	1	0.195	0.966	1
5	0.132	2	0.175	0.753	4
6	0.077	3	0.161	0.480	7
7	0.039	4	0.149	0.259	

## Example acceptance region for $\mu = 0.5$

- Start adding  $\Pr(n|\mu)$  in the second column based on the rank.
- Stop when total probability exceeds 90%.
- The  $n$ 's that contribute to the sum are the ones included in the acceptance region.


## Example acceptance region for $\mu = 0.5$

- Start adding  $\Pr(n|\mu)$  in the second column based on the rank.
- Stop when total probability exceeds 90%.
- The  $n$ 's that contribute to the sum are the ones included in the acceptance region.
- Acceptance region for  $\mu = 0.5$  is therefore  $n \in [0, 6]$



## Example acceptance region for $\mu = 0.5$

- Start adding  $\Pr(n|\mu)$  in the second column based on the rank.
- Stop when total probability exceeds 90%.
- The  $n$ 's that contribute to the sum are the ones included in the acceptance region.
- Acceptance region for  $\mu = 0.5$  is therefore  $n \in [0, 6]$
- Next, construct the acceptance region for other  $\mu$  as well.



# Bayesian

---



- Goal: estimate some parameter  $\mu$

- Goal: estimate some parameter  $\mu$
- *e.g.* the flux of neutrinos at some energy  $E = 10^{21}$  eV.

- Goal: estimate some parameter  $\mu$
- *e.g.* the flux of neutrinos at some energy  $E = 10^{21}$  eV.
  - see  $n$  neutrinos in some time window; want to estimate the flux.

# Bayes Theorem

- Goal: estimate some parameter  $\mu$
- e.g. the flux of neutrinos at some energy  $E = 10^{21}$  eV.
  - see  $n$  neutrinos in some time window; want to estimate the flux.

- Bayes Theorem

$$\Pr(\mu \mid n) = \frac{\Pr(n \mid \mu) \cdot \Pr(\mu)}{\Pr(n)}$$

Recall *Likelihood*  $\frac{e^{-\mu} \mu^n}{n!}$

# Bayes Theorem

- Goal: estimate some parameter  $\mu$
- e.g. the flux of neutrinos at some energy  $E = 10^{21}$  eV.
  - see  $n$  neutrinos in some time window; want to estimate the flux.

- Bayes Theorem

$$\Pr(\mu \mid n) = \frac{\Pr(n \mid \mu) \cdot \Pr(\mu)}{\Pr(n)}$$

Recall *Likelihood*  $\frac{e^{-\mu} \mu^n}{n!}$

*evidence*: typically a normalization and ignored

# Bayes Theorem

- Goal: estimate some parameter  $\mu$
- e.g. the flux of neutrinos at some energy  $E = 10^{21}$  eV.
  - see  $n$  neutrinos in some time window; want to estimate the flux.

- Bayes Theorem

$$\Pr(\mu \mid n) = \frac{\Pr(n \mid \mu) \cdot \Pr(\mu)}{\Pr(n)}$$

Recall *Likelihood*  $\frac{e^{-\mu} \mu^n}{n!}$

evidence: typically a normalization and ignored

*Posterior*, encodes possible values of  $\mu_t$  as a probability dist.

The “truth”  $\mu_t$  is **not** fixed from the Bayesian perspective

$$\Pr(\mu \mid n) = \frac{\Pr(n \mid \mu) \cdot \Pr(\mu)}{\Pr(n)}$$

$$\Pr(\mu \mid n) = \frac{\Pr(n \mid \mu) \cdot \Pr(\mu)}{\Pr(n)} \longleftarrow \text{prior}$$

- If we specify a prior, we get the posterior.



$$\Pr(\mu \mid n) = \frac{\Pr(n \mid \mu) \cdot \Pr(\mu)}{\Pr(n)} \longleftarrow \text{prior}$$

- If we specify a prior, we get the posterior.
- “uniform prior”  $\Pr(\mu) = 1$

$$\Pr(\mu \mid n) = \frac{\Pr(n \mid \mu) \cdot \Pr(\mu)}{\Pr(n)} \longleftarrow \text{prior}$$

- If we specify a prior, we get the posterior.
- “uniform prior”  $\Pr(\mu) = 1 \implies \Pr(\mu \mid n) = \Pr(n \mid \mu)$

$$\Pr(\mu \mid n) = \frac{\Pr(n \mid \mu) \cdot \Pr(\mu)}{\Pr(n)} \longleftarrow \text{prior}$$

- If we specify a prior, we get the posterior.
- “uniform prior”  $\Pr(\mu) = 1 \implies \Pr(\mu \mid n) = \Pr(n \mid \mu)$
- Suppose we measure  $n = 0$  event, then the posterior is

$$\Pr(\mu \mid n = 0) = \Pr(n = 0 \mid \mu) = \frac{e^{-\mu} \cdot \mu^n}{n!} = e^{-\mu}$$

# Summary: Bayesian Credible Interval

Bayesian

- Want an upper bound on  $\mu$

# Summary: Bayesian Credible Interval

- Want an upper bound on  $\mu$
- A choice: find a  $\mu_{\max}$  such that

$$\int_0^{\mu_{\max}} \text{Posterior} = 90\%$$

# Summary: Bayesian Credible Interval

- Want an upper bound on  $\mu$
- A choice: find a  $\mu_{\max}$  such that

$$\int_0^{\mu_{\max}} \text{Posterior} = 90\%$$

$$\Rightarrow \int_0^{\mu_{\max}} e^{-\mu} = 0.9$$

$$\Rightarrow \mu_{\max} = \ln(10) \approx 2.3$$

# Summary: Bayesian Credible Interval

- Want an upper bound on  $\mu$
- A choice: find a  $\mu_{\max}$  such that

$$\int_0^{\mu_{\max}} \text{Posterior} = 90\%$$

$$\Rightarrow \int_0^{\mu_{\max}} e^{-\mu} = 0.9$$

$$\Rightarrow \mu_{\max} = \ln(10) \approx 2.3$$

# Summary: Bayesian Credible Interval

Bayesian

- Want an upper bound on  $\mu$
- A choice: find a  $\mu_{\max}$  such that

$$\int_0^{\mu_{\max}} \text{Posterior} = 90\%$$

$$\Rightarrow \int_0^{\mu_{\max}} e^{-\mu} = 0.9$$

$$\Rightarrow \mu_{\max} = \ln(10) \approx 2.3$$

- so we “estimate with 90% confidence that  $\mu \leq 2.3$ ” base on a non-detection.

