

90% Confidence Level Upper Bound

Brief discussion of Feldman & Cousins

10-31-2024

Outline

Outline	1	Bayes Theorem	16
Parameter Estimataion	2	Summary: Bayesian Credible	
Poisson Distribution	3	Interval	18
Confidence Interval (CI)	4		
CI Definition	5		
CI Construction: Confidence			
Belt	7		
Acceptance Region	11		
Maximum Likelihood	12		
Derivation (skip me!)	13		
Likelihood Ratio	14		
Bayesian	15		

Parameter Estimataion

- Expect an average of μ counts per time, with $\mu \in \mathbb{R} \geq 0$

- Expect an average of μ counts per time, with $\mu \in \mathbb{R} \geq 0$
- Probability of getting $n \in \mathbb{N}$ counts per time?

- Expect an average of μ counts per time, with $\mu \in \mathbb{R} \geq 0$
- Probability of getting $n \in \mathbb{N}$ counts per time?
- Poisson Distribution:

$$\Pr(n|\mu) = \frac{e^{-\mu} \mu^n}{n!}$$

- Prob. of data n , given parameter μ ; aka the *likelihood*.

Poisson Distribution

Parameter Estimataion

- Expect an average of μ counts per time, with $\mu \in \mathbb{R} \geq 0$


- Probability of getting $n \in \mathbb{N}$ counts per time?

- Poisson Distribution:

$$\Pr(n|\mu) = \frac{e^{-\mu} \mu^n}{n!}$$

- Prob. of data n , given parameter μ ; aka the *likelihood*.

- Example:



Confidence Interval (CI)

- Goal: estimate parameter μ whose true value is μ_t

- Goal: estimate parameter μ whose true value is μ_t
- Make a measurement x ; suppose the first measurement yields x_0 .

- Goal: estimate parameter μ whose true value is μ_t
- Make a measurement x ; suppose the first measurement yields x_0 .
- Construct an interval (discussed later) $[\mu_l, \mu_u]$

- Goal: estimate parameter μ whose true value is μ_t
- Make a measurement x ; suppose the first measurement yields x_0 .
- Construct an interval (discussed later) $[\mu_l, \mu_u]$
 - $\mu_l = \mu_l(x_0)$: lower bound associated w/ this 1st measurement

- Goal: estimate parameter μ whose true value is μ_t
- Make a measurement x ; suppose the first measurement yields x_0 .
- Construct an interval (discussed later) $[\mu_l, \mu_u]$
 - $\mu_l = \mu_l(x_0)$: lower bound associated w/ this 1st measurement
 - $\mu_u = \mu_u(x_0)$: upper bound associated w/ this 1st measurement

- Goal: estimate parameter μ whose true value is μ_t
- Make a measurement x ; suppose the first measurement yields x_0 .
- Construct an interval (discussed later) $[\mu_l, \mu_u]$
 - $\mu_l = \mu_l(x_0)$: lower bound associated w/ this 1st measurement
 - $\mu_u = \mu_u(x_0)$: upper bound associated w/ this 1st measurement
- Repeat experiment; get outcome $x_1 \rightarrow$ construct $[\mu_l(x_1), \mu_u(x_1)]$

- Goal: estimate parameter μ whose true value is μ_t
- Make a measurement x ; suppose the first measurement yields x_0 .
- Construct an interval (discussed later) $[\mu_l, \mu_u]$
 - $\mu_l = \mu_l(x_0)$: lower bound associated w/ this 1st measurement
 - $\mu_u = \mu_u(x_0)$: upper bound associated w/ this 1st measurement
- Repeat experiment; get outcome $x_1 \rightarrow$ construct $[\mu_l(x_1), \mu_u(x_1)]$
- More experiments; get a bunch of intervals. *i.e.* we get a set

$$C \equiv \{[\mu_l(x_0), \mu_u(x_0)], [\mu_l(x_1), \mu_u(x_1)], [\mu_l(x_2), \mu_u(x_2)] \dots\}$$

- The set C has the property that

$$P([\mu_l, \mu_u] \ni \mu) = \alpha \quad \forall \text{ allowed } \mu$$

- The set C has the property that

$$P([\mu_l, \mu_u] \ni \mu) = \alpha \quad \forall \text{ allowed } \mu$$

- In words:
 - pick a *fixed* value of μ ; say we pick the truth μ_t .

- The set C has the property that

$$P([\mu_l, \mu_u] \ni \mu) = \alpha \quad \forall \text{ allowed } \mu$$

- In words:
 - pick a *fixed* value of μ ; say we pick the truth μ_t .
 - of the members in C , a fraction α would cover μ_t .

- The set C has the property that

$$P([\mu_l, \mu_u] \ni \mu) = \alpha \quad \forall \text{ allowed } \mu$$

- In words:
 - pick a *fixed* value of μ ; say we pick the truth μ_t .
 - of the members in C , a fraction α would cover μ_t .
 - others won't.

- The set C has the property that

$$P([\mu_l, \mu_u] \ni \mu) = \alpha \quad \forall \text{ allowed } \mu$$

- In words:
 - pick a *fixed* value of μ ; say we pick the truth μ_t .
 - of the members in C , a fraction α would cover μ_t .
 - others won't.
- The members of C are called *confidence intervals*.

CI Construction: Confidence Belt

Confidence Interval (CI)

- Consider probability

$$\Pr(x|\mu)$$

CI Construction: Confidence Belt

Confidence Interval (CI)

- Consider probability
 $\Pr(x|\mu)$
- Take for example $\mu = 4$,
consider $\Pr(x|\mu = 4)$

- Consider probability $\Pr(x|\mu)$
- Take for example $\mu = 4$, consider $\Pr(x|\mu = 4)$
- Select a region $[x_l, x_u]$ such that the probability of measuring $x \in [x_l, x_u]$ is, say, 80%.

- Consider probability $\Pr(x|\mu)$
- Take for example $\mu = 4$, consider $\Pr(x|\mu = 4)$
- Select a region $[x_l, x_u]$ such that the probability of measuring $x \in [x_l, x_u]$ is, say, 80%.
- that is,


$$\Pr(x \in [x_l, x_u] | \mu = 4) = 80\%$$

CI Construction: Confidence Belt

Confidence Interval (CI)

- Consider probability $\Pr(x|\mu)$
- Take for example $\mu = 4$, consider $\Pr(x|\mu = 4)$
- Select a region $[x_l, x_u]$ such that the probability of measuring $x \in [x_l, x_u]$ is, say, 80%.
- that is,

$$\Pr(x \in [x_l, x_u] | \mu = 4) = 80\%$$




CI Construction: Confidence Belt

Confidence Interval (CI)

- Consider probability $\Pr(x|\mu)$
- Take for example $\mu = 4$, consider $\Pr(x|\mu = 4)$
- Select a region $[x_l, x_u]$ such that the probability of measuring $x \in [x_l, x_u]$ is, say, 80%.
- that is,

$$\Pr(x \in [x_l, x_u] | \mu = 4) = 80\%$$




CI Construction: Confidence Belt

Confidence Interval (CI)

- Take another value μ , say $\mu = 5$
- Get another acceptance region $[x_l, x_u]$ associated with this value

$$\Pr(x \in [x_l, x_u] | \mu = 5) = 80\%$$




CI Construction: Confidence Belt

Confidence Interval (CI)

- Take another value μ , say $\mu = 5$
- Get another acceptance region $[x_l, x_u]$ associated with this value

$$\Pr(x \in [x_l, x_u] | \mu = 5) = 80\%$$

- Rinse and repeat




CI Construction: Confidence Belt

Confidence Interval (CI)

- Take another value μ , say $\mu = 5$
- Get another acceptance region $[x_l, x_u]$ associated with this value

$$\Pr(x \in [x_l, x_u] | \mu = 5) = 80\%$$

- Rinse and repeat




CI Construction: Confidence Belt

Confidence Interval (CI)

- Take another value μ , say $\mu = 5$
- Get another acceptance region $[x_l, x_u]$ associated with this value

$$\Pr(x \in [x_l, x_u] | \mu = 5) = 80\%$$

- Rinse and repeat




suppose this is μ_t

CI Construction: Confidence Belt

Confidence Interval (CI)


- Make a measurement, get result x_0
- The probability of x_0 falling in the region is 80%, by construction



CI Construction: Confidence Belt

Confidence Interval (CI)

- Make a measurement, get result x_0
- The probability of x_0 falling in the region is 80%, by construction
- The **confidence interval** $[\mu_l, \mu_u]$ from this experiment is the vertical intercept.



CI Construction: Confidence Belt


Confidence Interval (CI)

- Make some more measurements
- Get some more confidence intervals.


- Have a set

$$C = \{CI_1, CI_2, CI_3, CI_4, CI_5\}$$

- 80% of this set would cover the true value, μ_t .



Acceptance Region



- Recall acceptance region:

$$\Pr(n \in [n_1, n_2] \mid \mu_{\text{fixed}}) = 80\%$$

- Complete freedom in choosing how to construct the acceptance regions.
- Consider Poisson with background b :

$$\mathcal{L} \equiv \Pr(n \mid \mu) = \frac{(\mu + b)^n e^{-\mu+b}}{n!}$$

- F&C propose to compute a likelihood ratio R
 - This needs a “best fit” $\mu_{\text{best}} \equiv \max(0, n - b)$

Maximum Likelihood

Derivation (skip me!)

- Likelihood is a Poisson in this case.

$$\mathcal{L} \equiv \text{Pr}(n \mid \mu) = \frac{(\mu + b)^n e^{-\mu+b}}{n!}$$

- Find maximum (fixing n , vary μ):

$$\left. \frac{d\mathcal{L}}{d\mu} \right|_{\mu=\mu_{\text{best}}} = 0$$

- Result: “best fit” $\mu = \mu_{\text{best}} = n - b$
- Require physical $\mu \geq 0 \Rightarrow \mu_{\text{best}} = \max(0, n - b)$

- Do this for representative values of μ ; say we start with $\mu = 0.5$
 - As an example background, $b = 3$
 - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
 -
 -
 -
 -

n	$\text{Pr}(n \mu)$	μ_{best}	$\text{Pr}(n \mu_{\text{best}})$	R	rank
0					

- Do this for representative values of μ ; say we start with $\mu = 0.5$
 - As an example background, $b = 3$
 - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
 - For $n = 0$, compute $\Pr(n \mid \mu = 0.5)$
 -
 -
 -

n	$\Pr(n \mu)$	μ_{best}	$\Pr(n \mu_{\text{best}})$	R	rank
0	0.03				

- Do this for representative values of μ ; say we start with $\mu = 0.5$
 - As an example background, $b = 3$
 - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
 - For $n = 0$, compute $\Pr(n \mid \mu = 0.5)$
 - For $n = 0$, compute $\mu_{\text{best}} = 0$
 -
 -

n	$\Pr(n \mu)$	μ_{best}	$\Pr(n \mu_{\text{best}})$	R	rank
0	0.03	0			

- Do this for representative values of μ ; say we start with $\mu = 0.5$
 - As an example background, $b = 3$
 - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
 - For $n = 0$, compute $\Pr(n \mid \mu = 0.5)$
 - For $n = 0$, compute $\mu_{\text{best}} = 0$
 - For $n = 0$, compute $\Pr(n \mid \mu = \mu_{\text{best}})$
 -

n	$\Pr(n \mu)$	μ_{best}	$\Pr(n \mu_{\text{best}})$	R	rank
0	0.03	0	0.05		

- Do this for representative values of μ ; say we start with $\mu = 0.5$
 - As an example background, $b = 3$
 - $\Rightarrow \mu_{\text{best}} \equiv \max(0, n - b) = \max(0, n - 3)$
- Procedure:
 - For $n = 0$, compute $\Pr(n \mid \mu = 0.5)$
 - For $n = 0$, compute $\mu_{\text{best}} = 0$
 - For $n = 0$, compute $\Pr(n \mid \mu = \mu_{\text{best}})$
 - Divide likelihoods to get R .

n	$\Pr(n \mu)$	μ_{best}	$\Pr(n \mu_{\text{best}})$	R	rank
0	0.03	0	0.05	0.607	

Bayesian


- The problem is that the parameter λ is what we want to figure out.

- The problem is that the parameter λ is what we want to figure out.
 - e.g. the flux of neutrinos at some energy $E = 10^{21}$ eV.

- The problem is that the parameter λ is what we want to figure out.
 - e.g. the flux of neutrinos at some energy $E = 10^{21}$ eV.
- What we can do is measure n and *estimate* λ

- The problem is that the parameter λ is what we want to figure out.
 - e.g. the flux of neutrinos at some energy $E = 10^{21}$ eV.
- What we can do is measure n and *estimate* λ
- Bayes Theorem

$$\Pr(\lambda \mid n) = \frac{\Pr(n \mid \lambda) \cdot \Pr(\lambda)}{\Pr(n)}$$


Posterior

- The problem is that the parameter λ is what we want to figure out.
 - e.g. the flux of neutrinos at some energy $E = 10^{21}$ eV.
- What we can do is measure n and *estimate* λ
- Bayes Theorem

$$\underset{\substack{\uparrow \\ \textit{Posterior}}}{\text{Pr}(\lambda \mid n)} = \frac{\text{Pr}(n \mid \lambda) \cdot \text{Pr}(\lambda)}{\text{Pr}(n)}$$

\swarrow Likelihood $\frac{e^{-\lambda} \lambda^n}{n!}$

- The problem is that the parameter λ is what we want to figure out.
 - e.g. the flux of neutrinos at some energy $E = 10^{21}$ eV.
 - What we can do is measure n and *estimate* λ
 - Bayes Theorem
- The diagram shows the Bayes Theorem equation with three labels and arrows:

 - Posterior**: An arrow points from the label to the term $\Pr(\lambda | n)$, which is enclosed in a light blue box.
 - Likelihood**: An arrow points from the label to the term $\Pr(n | \lambda)$, which is enclosed in a light red box. To the right of this term is the expression $\frac{e^{-\lambda} \lambda^n}{n!}$.
 - evidence**: An arrow points from the label to the denominator term $\Pr(n)$.

The full equation is:
$$\Pr(\lambda | n) = \frac{\Pr(n | \lambda) \cdot \Pr(\lambda)}{\Pr(n)}$$
- Evidence is typically just a normalization and ignored. Let's call it 1 :)

$$\Pr(\lambda \mid n) = \frac{\Pr(n \mid \lambda) \cdot \Pr(\lambda)}{\Pr(n)}$$

$$\Pr(\lambda \mid n) = \frac{\Pr(n \mid \lambda) \cdot \Pr(\lambda)}{\Pr(n)} \longleftarrow \text{prior}$$

- If we specify a prior, we get the posterior.

$$\Pr(\lambda \mid n) = \frac{\Pr(n \mid \lambda) \cdot \Pr(\lambda)}{\Pr(n)} \longleftarrow \text{prior}$$

- If we specify a prior, we get the posterior.
- “uniform prior” $\Pr(\lambda) = 1$

$$\Pr(\lambda \mid n) = \frac{\Pr(n \mid \lambda) \cdot \Pr(\lambda)}{\Pr(n)} \longleftarrow \text{prior}$$

- If we specify a prior, we get the posterior.
- “uniform prior” $\Pr(\lambda) = 1 \implies \Pr(\lambda \mid n) = \Pr(n \mid \lambda)$

$$\Pr(\lambda \mid n) = \frac{\Pr(n \mid \lambda) \cdot \Pr(\lambda)}{\Pr(n)} \longleftarrow \text{prior}$$

- If we specify a prior, we get the posterior.
- “uniform prior” $\Pr(\lambda) = 1 \implies \Pr(\lambda \mid n) = \Pr(n \mid \lambda)$
- Suppose we measure $n = 0$ event, then the posterior is

$$\Pr(\lambda \mid n = 0) = \Pr(n = 0 \mid \lambda) = \frac{e^{-\lambda} \cdot \lambda^n}{n!} = e^{-\lambda}$$

Summary: Bayesian Credible Interval

- Want an upper bound on λ

Summary: Bayesian Credible Interval

- Want an upper bound on λ
- A choice: find a λ_{\max} such that

$$\int_0^{\lambda_{\max}} \text{Posterior} = 90\%$$

Summary: Bayesian Credible Interval

- Want an upper bound on λ
- A choice: find a λ_{\max} such that

$$\int_0^{\lambda_{\max}} \text{Posterior} = 90\%$$

$$\Rightarrow \int_0^{\lambda_{\max}} e^{-\lambda} = 0.9$$

$$\Rightarrow \lambda_{\max} = \ln(10) \approx 2.3$$

Summary: Bayesian Credible Interval

- Want an upper bound on λ
- A choice: find a λ_{\max} such that

$$\int_0^{\lambda_{\max}} \text{Posterior} = 90\%$$

$$\Rightarrow \int_0^{\lambda_{\max}} e^{-\lambda} = 0.9$$

$$\Rightarrow \lambda_{\max} = \ln(10) \approx 2.3$$

Summary: Bayesian Credible Interval


- Want an upper bound on λ
- A choice: find a λ_{\max} such that

$$\int_0^{\lambda_{\max}} \text{Posterior} = 90\%$$

$$\Rightarrow \int_0^{\lambda_{\max}} e^{-\lambda} = 0.9$$

$$\Rightarrow \lambda_{\max} = \ln(10) \approx 2.3$$

- so we “estimate with 90% confidence that $\lambda \leq 2.3$ ” base on a non-detection.



Summary: Bayesian Credible Interval

- Let μ denote the unknown parameter we wish to estimate.
- Let x_0 denote the outcome of a single measurement.
- Assume that we know how the measurement outcome depends on the parameter, $x = x(\mu)$.
 - e.g. if the neutrino flux is very small, then oftentimes a measurement reports a non-detection.
 - In other words, we know the *likelihood*, $P(x_0|\mu)$.
- From the Bayesian perspective, we can flip things around and say that the parameter is a function of the measurement, $P(\mu|x_0)$, provided that we state our prior beliefs about the parameter, $P(\mu)$.