

ASSIGNMENT-2

UNNATI GUPTA

Download all python codes from

<https://github.com/unnatigupta2320/Assignment-2/tree/master/codes>

and latex-tikz codes from

<https://github.com/unnatigupta2320/Assignment-2/tree/master>

1 QUESTION No. 2.36

Construct a quadrilateral $MIST$ where $MI = 3.5$, $IS = 6.5$, $\angle M = 75^\circ$, $\angle I = 105^\circ$ and $\angle S = 120^\circ$.

2 SOLUTION

1) Let us assume vertices of given quadrilateral $MIST$ as $\mathbf{M}, \mathbf{I}, \mathbf{S}$ and \mathbf{T} .

2) According to given data:

$$\angle M = 75^\circ, \angle I = 105^\circ, \angle S = 120^\circ \quad (2.0.1)$$

$$\|\mathbf{M} - \mathbf{I}\| = 3.5, \|\mathbf{I} - \mathbf{S}\| = 6.5 \quad (2.0.2)$$

$$\mathbf{M} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{I} = \begin{pmatrix} 3.5 \\ 0 \end{pmatrix} \quad (2.0.3)$$

3) For this quadrilateral $MIST$ we have,

$$\angle M + \angle I = 75^\circ + 105^\circ = 180^\circ, \quad (2.0.4)$$

$$\Rightarrow MT \parallel IS (\because MI \text{ being the transversal})$$

4) As, sum of adjacent angle on same side is 180° only when lines are **parallel**. Now, ST is another transversal on these parallel lines then, we get:

$$\Rightarrow \angle S + \angle T = 180^\circ \quad (2.0.5)$$

$$\Rightarrow \angle T = 60^\circ \quad (2.0.6)$$

5) Now taking sum of all the angles given and (2.0.6) we get the sum as 360° . So construction of given quadrilateral is possible.

6) For finding coordinates of \mathbf{S} :-

$$\mathbf{S} = \mathbf{I} + \lambda \mathbf{m} \quad (2.0.7)$$

$$\|\mathbf{S} - \mathbf{I}\| = |\lambda| \times \left\| \begin{pmatrix} \cos 75^\circ \\ \sin 75^\circ \end{pmatrix} \right\| \quad (2.0.8)$$

$$\Rightarrow \|\mathbf{S} - \mathbf{I}\| = |\lambda| \quad (2.0.9)$$

$$\Rightarrow |\lambda| = 6.5 \quad (2.0.10)$$

$$\Rightarrow \lambda = 6.5 \quad (2.0.11)$$

Now putting value of λ in (2.0.7) and solving we get,

$$\Rightarrow \mathbf{S} = \begin{pmatrix} 3.5 \\ 0 \end{pmatrix} + 6.5 \begin{pmatrix} \cos 75^\circ \\ \sin 75^\circ \end{pmatrix} \quad (2.0.12)$$

$$\Rightarrow \mathbf{S} = \begin{pmatrix} 5.18 \\ 6.27 \end{pmatrix} \quad (2.0.13)$$

7) For finding value of MT :-

In $\triangle MTS$, Let $\angle TSM = \phi$, $\angle TMS = \omega$, $\angle MTS = \beta$, Now using sine formula in $\triangle MTS$ we have

$$\frac{\sin \phi}{MT} = \frac{\sin \omega}{ST} = \frac{\sin \beta}{SM} \quad (2.0.14)$$

$$\frac{\sin 95.47^\circ}{MT} = \frac{\sin 60^\circ}{8.14} \quad (2.0.15)$$

$$\Rightarrow MT = 9.35 \quad (2.0.16)$$

8) For finding coordinates of \mathbf{T} :-

$$\mathbf{T} = \mathbf{M} + \mu \mathbf{m} = \mu \mathbf{m} (\because \mathbf{M} = \mathbf{0}) \quad (2.0.17)$$

Using inner products of vectors in quadrilateral $MIST$,

$$\frac{(S - T)^\top (S - I)}{\|S - T\| \times \|S - I\|} = \cos 120^\circ \quad (2.0.18)$$

$$\frac{(S - T)^\top (M - T)}{\|S - T\| \times \|M - T\|} = \cos 60^\circ \quad (2.0.19)$$

9) Now, dividing (2.0.18) and (2.0.19), we get:

$$\frac{(S - T)^\top (S - I)}{(S - T)^\top (M - T)} \times \frac{\|M - T\|}{\|S - I\|} = \frac{\cos 120^\circ}{\cos 60^\circ} \quad (2.0.20)$$

10) On solving (2.0.17) and (2.0.20) using (2.0.16), we have

$$\Rightarrow \mu = 9.35 \quad (2.0.21)$$

$$\Rightarrow \mathbf{T} = 9.35 \begin{pmatrix} \cos 75^\circ \\ \sin 75^\circ \end{pmatrix} \quad (2.0.22)$$

$$\Rightarrow \mathbf{T} = \begin{pmatrix} 2.42 \\ 9.63 \end{pmatrix} \quad (2.0.23)$$

11) Now, the vertices of given Quadrilateral MIST can be written as,

$$\mathbf{M} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{I} = \begin{pmatrix} 3.5 \\ 0 \end{pmatrix}, \mathbf{S} = \begin{pmatrix} 5.18 \\ 6.27 \end{pmatrix}, \mathbf{T} = \begin{pmatrix} 2.42 \\ 9.63 \end{pmatrix} \quad (2.0.24)$$

12) On constructing the given quadrilateral we, get:




Fig. 2.1: Quadrilateral MIST