

# ASSIGNMENT-13

Unnati Gupta

Download all python codes from

[https://github.com/unnatigupta2320/  
Assignment\\_13](https://github.com/unnatigupta2320/Assignment_13)

and latex-tikz codes from

[https://github.com/unnatigupta2320/  
Assignment\\_13](https://github.com/unnatigupta2320/Assignment_13)

## 1 QUESTION No-6.20

An unbiased dice is thrown twice. Let the event A be 'odd number on the first throw' and B be 'Odd number on second throw'. Check the independence of event A and B.

## 2 SOLUTION

**Lemma 2.1.** *Two events are independent if knowing one event occurred doesn't change the probability of the other event.*

$\therefore$  A and B are said to be independent if and only if:-

$$\Pr(AB) = \Pr(A) \Pr(B) \quad (2.0.1)$$

- 1) Let  $X_0$  and  $X_1$  be the random variables representing the numbers we get when a dice is thrown for first and second time respectively.

$$X_0 \in \{1, 2, 3, 4, 5, 6\} \quad (2.0.2)$$

$$X_1 \in \{1, 2, 3, 4, 5, 6\} \quad (2.0.3)$$

- 2) Also, the probability

$$\Pr(X = i) = \begin{cases} \frac{1}{6} & 1 \leq i \leq 6 \\ 0 & \text{otherwise} \end{cases} \quad (2.0.4)$$

- 3) According to question, the events are:-

Events	Description
A	Odd number on first throw
B	Odd number on second throw
AB	Odd Numbers appears on both throws

- 4) For the event **A**:-

- The probability of odd number on first throw is-

$$\Pr(A) = \sum_{i=1,3,5} \Pr(X_0 = i) \quad (2.0.5)$$

- So,

$$\Pr(A) =$$

$$\Pr(X_0 = 1) + \Pr(X_0 = 3) + \Pr(X_0 = 5) \quad (2.0.6)$$

$$\Rightarrow \Pr(A) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} \quad (2.0.7)$$

$$\Rightarrow \Pr(A) = \frac{3}{6} \quad (2.0.8)$$

$$\Rightarrow \Pr(A) = \frac{1}{2} \quad (2.0.9)$$

- 5) For the event **B**:-

- The probability of odd number on second throw is-

$$\Pr(B) = \sum_{i=1,3,5} \Pr(X_1 = i) \quad (2.0.10)$$

- So,

$$\Pr(B) =$$

$$\Pr(X_1 = 1) + \Pr(X_1 = 3) + \Pr(X_1 = 5) \quad (2.0.11)$$

$$\Rightarrow \Pr(B) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} \quad (2.0.12)$$

$$\Rightarrow \Pr(B) = \frac{3}{6} \quad (2.0.13)$$

$$\Rightarrow \Pr(B) = \frac{1}{2} \quad (2.0.14)$$

- 6) For the event **AB**:-

- The probability that odd numbers appears on

both throw is-

$$\Pr(AB) = \Pr(A|B) \Pr(B) \quad (2.0.15)$$

$$\implies \Pr(AB) = \frac{1}{2} \times \frac{1}{2} \quad (2.0.16)$$

$$\implies \Pr(AB) = \frac{1}{4} \quad (2.0.17)$$

7) Now to check whether the events are **independent**, we use Lemma (2.1).

$$\implies \Pr(AB) = \Pr(A) \Pr(B) \quad (2.0.18)$$

8) Putting values from (2.0.9) and (2.0.14) we get,

$$\implies \Pr(AB) = \frac{1}{2} \times \frac{1}{2} \quad (2.0.19)$$

$$\implies \Pr(AB) = \frac{1}{4} \quad (2.0.20)$$

This is equal to value in equation (2.0.17).

Hence, the events are **independent**.