1

$$\|G(x) - G(x^*)\| \le L \|x - x^*\|$$

$$\tilde{s}_k = G_*^{1/2} s_k, \quad \tilde{y}_k = G_*^{-1/2} y_k, \quad \tilde{B}_k = G_*^{-1/2} B_k G_*^{1/2},$$

$$\cos \tilde{\theta}_k = \frac{\tilde{s}_k^T \tilde{B}_k \tilde{s}_k}{\|\tilde{s}_k\| \|\tilde{B}_k \tilde{s}_k}, \quad \tilde{q}_k = \frac{\tilde{s}_k^T \tilde{B}_k \tilde{s}_k}{\|\tilde{s}_k\|^2}$$

$$\tilde{M}_k = \frac{\|\tilde{y}_k\|^2}{\tilde{y}_k^T \tilde{s}_k}, \quad \tilde{m}_k = \frac{\tilde{y}_k^T \tilde{s}_k}{\tilde{s}_k^T \tilde{b}_k}.$$

$$B_{k+1}^- = \tilde{B}_k - \frac{\tilde{B}_k \tilde{s}_k \tilde{s}_k^T \tilde{B}_k}{\tilde{s}_k^T \tilde{B}_k \tilde{s}_k} + \frac{\tilde{y}_k \tilde{y}_k^T}{\tilde{y}_k^T \tilde{s}_k}.$$

$$\psi(\tilde{B}_{k+1}^-) = \psi(\tilde{B}_k + (\tilde{M}_{k+1} - \ln(\tilde{m}_k) - 1) + \left[1 - \frac{\tilde{q}_k}{\cos^2 \tilde{\theta}_k} + \ln\left(\frac{\tilde{q}_k}{\cos^2 \tilde{\theta}_k}\right)\right] + \ln\left(\cos^2 \tilde{\theta}_k\right)$$

$$y_k - \tilde{G}_* s_k = (\tilde{G}_k - \tilde{G}_*) s_k,$$

$$\tilde{y}_k - \tilde{s}_k = G_*^{-1/2}(\tilde{G}_k - \tilde{G}_*) G_*^{-1/2} \tilde{s}_k.$$

$$\|\tilde{y}_k - \tilde{s}_k\| \le \|G_*^{-1/2}\|^2 \|\tilde{s}_k\| \|\tilde{G}_k - \tilde{G}_*\| \le \|G_*^{-1/2}\|^2 \|\tilde{s}_k\| L \epsilon_k,$$

$$\epsilon_k = \max\{\|x_{k+1} - x^*\|, \|x_k - x^*\|\}.$$

$$\|\tilde{y}_k - \tilde{s}_k\| \le \tilde{c}_k,$$

$$(2)$$

 $\mathbf{2}$

$$\|\tilde{y_k} - \tilde{s_k}\| \le \bar{c}\epsilon_k \|\tilde{s_k}\|, \quad \|\tilde{s_k} - \tilde{y_k}\| \le \bar{c}\epsilon_k \|\tilde{s_k}\|$$

$$(1 - \bar{c}\epsilon_k)\|\tilde{s}_k\| \le \|\tilde{y}_k\| \le (1 + \bar{c}\epsilon_k)\|\tilde{s}_k\|$$
(3)

(2)

$$(1 - \bar{c}\epsilon_k)^2 \|\tilde{s}_k\|^2 - 2\tilde{y}_k^T \tilde{s}_k + \|\tilde{s}_k\|^2 \le \|\tilde{y}_k\|^2 - 2\tilde{y}_k^T \tilde{s}_k + \|\tilde{s}_k\|^2 \le \bar{c}^2 \epsilon_k^2 \|\tilde{s}_k\|^2,$$
$$2\tilde{y}_k^T \tilde{s}_k \ge (1 - 2\bar{c}\epsilon_k + \bar{c}^2 \epsilon_k^2 + 1 - \bar{c}^2 \epsilon_k^2) \|\tilde{s}_k\|^2 = 2(1 - \bar{c}\epsilon_k) \|\tilde{s}_k\|^2$$

$$\tilde{m_k} = \frac{\tilde{y_k}^T \tilde{s_k}}{\|\tilde{s_k}\|^2} \ge 1 - \bar{c}\epsilon_k \tag{4}$$

$$\tilde{M}_k = \frac{\|\tilde{y}_k\|^2}{\tilde{y}_k^T \tilde{s}_k} \le \frac{1 + \bar{c}\epsilon_k}{1 - \bar{c}\epsilon_k} \tag{5}$$

$$\tilde{M}_k \le 1 + \frac{2\bar{c}}{1 - \bar{c}\epsilon_k} \epsilon_k \le 1 + \bar{c}\epsilon_k \tag{6}$$

$$\frac{-x}{1-x} - \ln(1-x) = h\left(\frac{1}{1-x}\right) \le 0$$
$$\ln(1-\bar{c}\epsilon_k) \ge \frac{-\bar{c}\epsilon_k}{1-\bar{c}\epsilon_k} \ge -2\bar{c}\epsilon_k$$

$$\ln(\tilde{m_k}) \ge \ln(1 - \bar{c}\epsilon_k) \ge -2\bar{c}\epsilon_k \ge -2c\epsilon_k \tag{7}$$

$$0 < \psi(\tilde{B}_{k+1}) \le \psi(\tilde{B}_k) + 3c\epsilon_k + \ln(\cos^2(\tilde{\theta}_k)) + \left[1 - \frac{\tilde{q}_k}{\cos^2(\tilde{\theta}_k)} + \ln\left(\frac{\tilde{q}_k}{\cos^2(\tilde{\theta}_k)}\right)\right]$$
(8)

$$\sum_{j=0}^{\infty} \left(\ln \left(\frac{1}{\cos^2 \tilde{\theta_j}} \right) - \left[1 - \frac{\tilde{q_j}}{\cos^2 \tilde{\theta_j}} + \ln \left(\frac{\tilde{q_j}}{\cos^2 \tilde{\theta_j}} \right) \right] \right) \le \psi(\tilde{B_0}) + 3c \sum_{j=0}^{\infty} \epsilon_j < +\infty$$

$$\lim_{j \to \infty} \ln \left(\frac{1}{\cos^2 \tilde{\theta_j}} \right) = 0, \qquad \qquad \lim_{j \to \infty} \left(1 - \frac{\tilde{q_j}}{\cos^2 \tilde{\theta_j}} + \ln \left(\frac{\tilde{q_j}}{\cos^2 \tilde{\theta_j}} \right) \right) = 0$$

$$\lim_{j \to \infty} \cos \tilde{\theta}_j = 1, \qquad \lim_{j \to \infty} \tilde{q}_j = 1 \tag{9}$$

$$\frac{\|G_*^{-1/2}(B_k - G_*)s_k\|^2}{\|G_*^{1/2}s_k\|} = \frac{\frac{\|(\tilde{B}_k - I)\tilde{s}_k\|^2}{\|\tilde{s}_k\|^2}}{\|\tilde{s}_k\|^2}$$

$$= \frac{\frac{\|\tilde{B}_k\tilde{s}_k\|^2 - 2\tilde{s}_k^T\tilde{B}_k\tilde{s}_k + \tilde{s}_k^T\tilde{s}_k}{\tilde{s}_k^T\tilde{s}_k}}{\tilde{s}_k^T\tilde{s}_k}$$

$$= \frac{\tilde{q}_k^2}{\cos\tilde{\theta}_k^2} - 2\tilde{q}_k + 1.$$

$$\therefore \lim_{k \to \infty} \frac{\|(B_k - G_*)s_k\|}{\|s_k\|} = 0$$