


Linear Algebra Reformed for 21st C Application

A. J. Roberts *

University of Adelaide
South Australia, 5005

November 29, 2016



* <http://orcid.org/0000-0001-8930-1552>

Contents

1	Vectors	11
1.1	Vectors have magnitude and direction	13
1.2	Adding and stretching vectors	23
1.3	The dot product determines angles and lengths	38
1.4	The cross product	62
1.5	Use MATLAB/Octave for vector computation .	78
1.6	Summary of vectors	88
2	Systems of linear equations	93
2.1	Introduction to systems of linear equations . .	96
2.2	Directly solve linear systems	105
2.3	Linear combinations span sets	137
2.4	Summary of linear equations	147
3	Matrices encode system interactions	152
3.1	Matrix operations and algebra	154
3.2	The inverse of a matrix	186
3.3	Factorise to the singular value decomposition .	230
3.4	Subspaces, basis and dimension	268
3.5	Project to solve inconsistent equations	308
3.6	Introducing linear transformations	374
3.7	Summary of matrices	406

4 Eigenvalues and eigenvectors of symmetric matrices 426

4.1 Introduction to eigenvalues and eigenvectors . . 429

4.2 Beautiful properties for symmetric matrices . . 456

4.3 Summary of symmetric eigen-problems 487

5 Approximate matrices 493

5.1 Measure changes to matrices 494

5.2 Regularise linear equations 540

5.3 Summary of matrix approximation 565

6 Determinants distinguish matrices 570

6.1 Geometry underlies determinants 571

6.2 Laplace expansion theorem for determinants . . 590

6.3 Summary of determinants 618

7 Eigenvalues and eigenvectors in general 622

7.1 Find eigenvalues and eigenvectors of matrices . 631

7.2 Linear independent vectors may form a basis . 681

7.3 Diagonalisation identifies the transformation . . 716

7.4 Summary of general eigen-problems 744

Preface

Traditional courses in linear algebra make considerable use of the reduced row echelon form (RREF), but the RREF is an unreliable tool for computation in the face of inexact data and arithmetic. The [Singular Value Decomposition] SVD can be regarded as a modern, computationally powerful replacement for the RREF.¹ *Cleve Moler, MathWorks (2006)*

The Singular Value Decomposition (SVD) is sometimes called the *jewel in the crown* of linear algebra. Traditionally the SVD is introduced and explored at the end of several linear algebra courses. Question: Why were students required to wait until the end of the course, if at all, to be introduced to beauty and power of this jewel? Answer: limitations of hand calculation.

This book establishes a new route through linear algebra, one that reaches the SVD jewel in linear algebra's crown very early, in Section 3.3. Thereafter its beautiful power both explores many modern applications and also develops traditional linear algebra concepts, theory, and methods. No rigour is lost in this new route: indeed, this book demonstrates that most theory is better proved with an SVD rather than with the traditional RREF. This new route through linear algebra becomes available by the ready availability of ubiquitous computing in the 21st century.

As so many other disciplines use the SVD, it is not only important that mathematicians understand what it is, but also teach it thoroughly in linear algebra and matrix analysis courses.

(Turner et al. 2015, p.30)

¹ <http://au.mathworks.com/company/newsletters/articles/professor-svd.html> [9 Jan 2015]

Comment: In this draft, the blue text comments such as this are notes to developers about related references, about reasons for decisions, about more exercises, about possible future extensions, and so on. Such comments are not intended for the published version, they are only notes for myself and developers.

the consensus curriculum is still an unwitting prisoner of history
George Cobb, 2007

Comment: Outstanding tasks for a first version include:

- finalising the scope of the book and of applications;
- exercises throughout on higher level interpretative and integrative aspects;
- exercises on computing that do not require a computer, e.g., interpretation;
- checking the staged use of abstract symbolism;
- potentially more applications that involve ‘real’ data, especially in Chapter 5 on approximating matrices;
- adapt information, especially some uses of the SVD, from the book by Mark Holmes (2016) “Introduction to scientific computing and data analysis” Springer;
- refining the index;
- section/chapter summaries, and possibly concept maps, including summary of some application/modelling;
- short videos of procedures, examples, proofs;
- and continual checking for typos and clarity.

Aims for students

How should mathematical sciences departments reshape their curricula to suit the needs of a well-educated workforce in the twenty-first century?

... The mathematical sciences themselves are changing as the needs of big data and the challenges of modeling complex systems reveal the limits of traditional curricula. *Bressoud et al. (2014)*

Linear algebra is packed with compelling results for application in science, engineering and computing, and with answers for the twenty-first century needs of big data and complex systems. This book provides the conceptual understanding of the essential linear algebra of vectors and matrices for modern engineering and science. The traditional linear algebra course has been reshaped herein to meet modern demands.

Crucial is to inculcate the terms and corresponding relationships that you will most often encounter later in professional life, often when using professional software. For example, the manual for the engineering software package Fluent most often invokes the linear algebra terms of diagonal, dot product, eigenvalue, least square, orthogonal, projection, principal axes, symmetric, unit vector. Engineers need to know these terms. What such useful terms mean, their relationships, and use in applications are central to the mathematical development in this book: you will see them introduced early and used often.

For those who proceed on to do higher mathematics, the development also provides a great solid foundation of key concepts, relationships and transformations necessary for higher mathematics.

Important for all is to develop facility in manipulating, interpreting and transforming between visualisations, algebraic forms, and vector-matrix representations—of problems, working, solutions and interpretation. In particular, one overarching aim of the book is to encourage your formulation, thinking and operation at the crucial system-wide level of matrix/vector operations.

In view of ubiquitous computing, throughout this book explicitly integrates computer support for developing concepts

and their relations. The central computational tools to understand are the operation $\mathbf{A} \backslash$ for solving straightforward linear equations; the function `svd()` for difficult linear equation; the function `svd()` for approximation; and function `eig()` for probing structures. This provides a framework to understand key computational tools to effectively utilise the so-called third arm of science: namely, computation.

Throughout the book examples (many graphical) introduce and illustrate the concepts and relationships between the concepts. Working through these will help form the mathematical relationships essential for application. Interspersed throughout the text are questions labelled “Activity”: these aim to help form and test your understanding of the concepts being developed.

Also included are many varied applications, described to varying levels of details. These applications indicate how the mathematics will empower you to answer many practical challenges in engineering and science.

The main contribution of mathematics to the natural sciences is not in formal computations . . . , but in the investigation of those non-formal questions where the exact setting of the question (what are we searching for and what specific models must be used) usually constitute half the matter.

. . . Examples teach no less than rules, and errors, more than correct but abstruse proofs. Looking at the pictures in this book, the reader will understand more than learning by rote dozens of axioms
(*Arnold 2014, p.xiii*)

Background for teachers

Depending upon the background of your students, your class should pick up the story somewhere in the first two or three chapters. Some students will have previously learnt some vector material in just 2D and 3D, in which case refresh the concepts in n D as presented in the first chapter.

As a teacher you can use this book in several ways.

- One way is as a reasonably rigorous mathematical development of concepts and interconnections by invoking its definitions, theorems, and proofs, all interleaved with examples.
- Another way is as the development of practical techniques and insight for application orientated science and engineering students via the motivating examples to appropriate definitions, theorems and applications.
- Or any mix of these two.

One of the aims of this book is to organise the development of linear algebra so that if a student only studies part of the material, then s/he still obtains a powerful and useful body of knowledge for science or engineering.

The book typically introduces concepts in low dimensional cases, and subsequently develops the general theory of the concept. This is to help focus the learning, empowered by visualisation, while also making a preformal connection to be strengthened subsequently. People are not one dimensional; knowledge is not linear. Cross-references make many connections, explicitly recalling earlier learning (although sometimes forward to material not yet ‘covered’).

information that is recalled grows stronger with each retrieval . . . spaced practice is preferable to massed practice. *Halpern & Hakel (2003) [p.38]*

One characteristic of the development is that the concept of linear independence does not appear until relatively late, namely in Chapter 7. This is good for several reasons. First, orthogonality is much more commonly invoked in science and engineering than is linear independence. Second, it is well documented that students struggle with linear independence:

there is ample talk in the math ed literature of classes hitting a ‘brick wall’, when linear (in)dependence is studied in the middle of such a course
Uhlig (2002)

Consequently, here we learn the more specific orthogonality before the more abstract linear independence. Many modern applications are made available by the relatively early

introduction of orthogonality.

In addition to many exercises at the end of every section, throughout the book are questions labelled “Activity”. These are for the students to do to help form the concepts being introduced with a small amount of work (currently, each version of this book invokes a different (random) permutation of the answers). These activities may be used in class to foster active participation by students (perhaps utilising clickers or web tools such as that provided by <http://www.quizsocket.com>). Such active learning has positive effects (Pashler et al. 2007).

On visualisation

All Linear Algebra courses should stress visualization and geometric interpretation of theoretical ideas in 2- and 3-dimensional spaces. Doing so highlights “algebraic and geometric” as “contrasting but complementary points of view,”

(Schumacher et al. 2015, p.38)

Throughout, this book also integrates visualisation. This visualisation reflects the fundamentally geometric nature of linear algebra. It also empowers learners to utilise different parts of their brain and integrate the knowledge together from the different perspectives. Visualisation also facilitates greater skills at interpretation and modelling so essential in applications. But as commented by Fara (2009) [p.249] “just like reading, deciphering graphs and maps only becomes automatic with practice.” Lastly, visual exercise questions develops understanding without a learner being able to defer the challenge to online tools, as yet.

Visual representations are effective because they tap into the capabilities of the powerful and highly parallel human visual system. We like receiving information in visual form and can process it very efficiently: around a quarter of our brains are devoted to vision, more than all our other senses combined ... researchers (especially those from mathematic backgrounds) see visual notations as being informal, and that serious analysis can only take place at the level of their semantics. However, this is a misconception: visual languages are no less formal

than textual ones

Moody (2009)

On integrated computation

Cowen argued that because “no serious application of linear algebra happens without a computer,” computation should be part of every beginning Linear Algebra course. ... While the increasing applicability of linear algebra does not require that we stop teaching theory, Cowen argues that “it should encourage us to see the role of the theory in the subject as it is applied.”

(Schumacher et al. 2015, p.38)

We need to empower students to use computers to improve their understanding, learning and application of mathematics; not only integrated in their study but also in their later professional career.

One often expects it should be easy to sprinkle a few computational tips and tools throughout a mathematics course. This is not so—extra computing is difficult. There are two reasons for the difficulty: first, the number of computer language details that have to be learned is surprisingly large; second, for students it is a genuine intellectual overhead to learn and relate both the mathematics and the computations.

Consequently, this book chooses a computing language where it is as simple as reasonably possible to perform linear algebra operations: MATLAB/Octave appears to answer this criteria.² Further, we are as ruthless as possible in invoking herein the smallest feasible set of commands and functions from MATLAB/Octave so that students have the minimum to learn. Most teachers will find many of their favourite commands are missing—this omission is all to the good in focussing upon useful mathematical development aided by only essential integrated computation.

This book does not aim to teach computer programming: there

² To compare popular packages, just look at the length of expressions students have to type in order to achieve core computations: MATLAB/Octave is almost always the shortest (Nakos & Joyner 1998, e.g.). (Of course be wary of this metric: e.g., APL would surely be too concise!)

is no flow control, no looping, no recursion, nor function definitions. The aim herein is to use short sequences of declarative assignment statements, coupled with the power of vector and matrix data structures, to learn core mathematical concepts, applications and their relationships in linear algebra.

The internet is now ubiquitous and pervasive. So too is computing power: students can execute MATLAB/Octave not only on laptops, but also on tablets and smart phones, perhaps using university or public servers, octave-online.net, MATLAB-Online or MATLAB-Mobile. We no longer need separate computer laboratories. Instead, expect students to access computational support simply by reaching into their pockets or bags.

long after Riemann had passed away, historians discovered that he had developed advanced techniques for calculating the Riemann zeta function and that his formulation of the Riemann hypothesis—often depicted as a triumph of pure thought—was actually based on painstaking numerical work.

Donoho & Stodden (2015)

Linear algebra for statisticians

This book forms an ideal companion to modern statistics courses. The recently published Curriculum Guidelines for Undergraduate Programs in Statistical Science by [Horton et al. \(2014\)](#) emphasises that linear algebra courses must provide “matrix manipulations, linear transformations, projections in Euclidean space, eigenvalues/eigenvectors, and matrix decompositions”. These are all core topics in this book, especially the statistically important SVD factorisation (Chaps. 3 and 5). Furthermore, this book explicitly makes the recommended “connections between concepts in these mathematical foundations courses and their applications in statistics” ([Horton et al. 2014](#), p.12).

Moreover, with the aid of some indicative statistical applications along with the sustained invocation of “visualization” and “basic programming concepts”, this book helps to underpin the requirement to “Encourage synthesis of theory, methods, computation, and applications” ([Horton et al. 2014](#), p.13).

Acknowledgements

I acknowledge with thanks the work of many others who inspired much design and details here, including the stimulating innovations of calculus reform ([Hughes-Hallett et al. 2013](#), e.g.), the comprehensive efforts behind recent reviews of undergraduate mathematics and statistics teaching ([Alpers et al. 2013](#), [Bressoud et al. 2014](#), [Turner et al. 2015](#), [Horton et al. 2014](#), [Schumacher et al. 2015](#), [Bliss et al. 2016](#), e.g.), and the books of [Anton & Rorres \(1991\)](#), [Davis & Uhl \(1999\)](#), [Holt \(2013\)](#), [Larson \(2013\)](#), [Lay \(2012\)](#), [Nakos & Joyner \(1998\)](#), [Poole \(2015\)](#), [Will \(2004\)](#). I also thank the entire L^AT_EX team, especially Knuth, Lamport, Feuersänger, and the AMS.

Bibliography

Alpers, B., Demlova, M., Fant, C.-H., Gustafsson, T., Lawson, D., Mustoe, L., Olsson-Lehtonen, B., Robinson, C. & Velichova, D. (2013), A framework for mathematics curricula in engineering education, Technical report, European Society for Engineering Education (SEFI).

<http://sefi.htw-aalen.de/curriculum.htm>

Anton, H. & Rorres, C. (1991), *Elementary linear algebra. Applications version*, 6th edn, Wiley.

Arnold, V. I. (2014), *Mathematical understanding of nature*, Amer. Math. Soc.

Berry, M. W., Dumais, S. T. & O'Brien, G. W. (1995), 'Using linear algebra for intelligent information retrieval', *SIAM Review* **37**(4), 573–595.

<http://epubs.siam.org/doi/abs/10.1137/1037127>

Bliss, K., Fowler, K., Galluzzo, B., Garfunkel, S., Giordano, F., Godbold, L., Gould, H., Levy, R., Libertini, J., Long, M., Malkevitch, J., Montgomery, M., Pollak, H., Teague, D., van der Kooij, H. & Zbiek, R. (2016), GAIMME—Guidelines for Assessment and Instruction in Mathematics Modeling Education, Technical report, SIAM and COMAP.

http://www.siam.org/reports/gaimme.php?_ga=1.192190826.137003948.1386128871

Bressoud, D. M., Friedlander, E. M. & Levermore, C. D. (2014), 'Meeting the challenges of improved post-secondary education in the mathematical sciences', *Notices of the AMS* **61**(5), 502–3.

Chartier, T. (2015), *When life is linear: from computer graphics to bracketology*, Math Assoc Amer.

<http://www.maa.org/press/books/>

[when-life-is-linear-from-computer-graphics-to-bracketology](http://www.maa.org/press/books/when-life-is-linear-from-computer-graphics-to-bracketology)

Cowen, C. C. (1997), On the centrality of linear algebra in the curriculum, Technical report, Mathematical Association of

America.

<http://www.maa.org/centrality-of-linear-algebra>

Cuyt, A. (2015), Approximation theory, in N. J. Higham, M. R. Dennis, P. Glendinning, P. A. Martin, F. Santosa & J. Tanner, eds, 'Princeton Companion to Applied Mathematics', Princeton, chapter IV.9, pp. 248–262.

Davis, B. & Uhl, J. (1999), *Matrices, Geometry and Mathematics*, Wolfram Research.

Donoho, D. L. & Stodden, V. (2015), Reproducible research in the mathematical sciences, in N. J. Higham, M. R. Dennis, P. Glendinning, P. A. Martin, F. Santosa & J. Tanner, eds, 'Princeton Companion to Applied Mathematics', Princeton, chapter VIII.5, pp. 916–925.

Driscoll, T. A. & Maki, K. L. (2007), 'Searching for rare growth factors using multicanonical monte carlo methods', *SIAM Review* **49**(4), 673–692.

<http://link.aip.org/link/?SIR/49/673/1>

Fara, P. (2009), *Science: a four thousand year history*, OUP.

Gorodetski, V. I., Popyack, L. J. & Samoilov, V. (2001), SVD-based approach to transparent embedding data into digital images, in V. I. Gorodetski, V. A. Skormin & L. J. Popyack, eds, 'Information assurance in computer networks: methods, models, and architectures for network security', Vol. 2052 of *Lecture Notes in Computer Science*, Springer, pp. 263–274.

Halpern, D. F. & Hakel, M. D. (2003), 'Applying the science of learning to the university and beyond: Teaching for long-term retention and transfer', *Change: The Magazine of Higher Learning* **35**(4), 36–41.

Hannah, J. (1996), 'A geometric approach to determinants', *The American Mathematical Monthly* **103**(5), 401–409.

<http://www.jstor.org/stable/2974931>

Higham, N. J. (1986), 'Computing the polar decomposition—with applications', *SIAM Journal on Scientific and Statistical Computing* **7**(4), 1160–1174.

<http://dx.doi.org/10.1137/0907079>

Higham, N. J. (1996), *Accuracy and stability of numerical algorithms*, SIAM.

- Higham, N. J. (2015a), Numerical linear algebra and matrix analysis, in N. J. Higham, M. R. Dennis, P. Glendinning, P. A. Martin, F. Santosa & J. Tanner, eds, 'Princeton Companion to Applied Mathematics', Princeton, chapter IV.10, pp. 263–281.
- Higham, N. J. (2015b), What is applied mathematics?, in N. J. Higham, M. R. Dennis, P. Glendinning, P. A. Martin, F. Santosa & J. Tanner, eds, 'Princeton Companion to Applied Mathematics', Princeton, chapter I.1, pp. 1–8.
- Holt, J. (2013), *Linear algebra with applications*, Freeman.
- Hopcroft, J. & Kannan, R. (2014), Foundations of data science, Technical report, [<https://www.researchgate.net/publication/265013479>].
- Horton, N., Chance, B., Cohen, S., Grimshaw, S., Hardin, J., Hesterberg, T., Hoerl, R., Malone, C., Nichols, R. & Nolan, D. (2014), Curriculum guidelines for undergraduate programs in statistical science, Technical report, American Statistical Association.
<http://www.amstat.org/education/curriculumguidelines.cfm>
- Hughes-Hallett, D., Gleason, A. M. & McCallum, et al., W. G. (2013), *Calculus: single and multivariable*, 6th edn, Wiley.
- Kleiber, M. (1947), 'Body size and metabolic rate', *Physiological Reviews* **27**, 511–541.
- Kress, R. (2015), Integral equations, in N. J. Higham, M. R. Dennis, P. Glendinning, P. A. Martin, F. Santosa & J. Tanner, eds, 'Princeton Companion to Applied Mathematics', Princeton, chapter IV.4, pp. 200–208.
- Larson, R. (2013), *Elementary linear algebra*, 7th edn, Brooks/Cole Cengage learning.
- Lay, D. C. (2012), *Linear Algebra and its Applications*, 4th edn, Addison–Wesley.
- Lichman, M. (2013), 'UCI machine learning repository', [online].
<http://archive.ics.uci.edu/ml>
- Mandelbrot, B. B. (1982), *The fractal geometry of nature*, W. H. Freeman.

- Moody, D. L. (2009), ‘The “physics” of notations: Towards a scientific basis for constructing visual notations in software engineering’, *IEEE Trans. Soft. Engrg.* **35**(5), 756–778.
- Nakos, G. & Joyner, D. (1998), *Linear algebra with applications*, Brooks/Cole.
- Pashler, H., Bain, P. M., Bottge, B. A., Graesser, A., Koedinger, K., McDaniel, M. & Metcalfe, J. (2007), Organizing instruction and study to improve student learning: IES practice guide, Technical report, National Center for Education Research, Institute of Education Sciences, U.S. Dept of Education.
<http://files.eric.ed.gov/fulltext/ED498555.pdf>
- Poole, D. (2015), *Linear algebra: A modern introduction*, 4th edn, Cengage Learning.
- Quarteroni, A. & Saleri, F. (2006), *Scientific Computing with MATLAB and Octave*, Vol. 2 of *Texts in Computational Science and Engineering*, 2nd edition edn, Springer. (Available in pdf via the library of the University of Adelaide.).
- Roulstone, I. & Norbury, J. (2013), *Invisible in the storm: the role of mathematics in understanding weather*, Princeton.
- Schonefeld, S. (1995), ‘Eigenpictures: Picturing the eigenvector problem’, *The College Mathematics Journal* **26**(4), 316–319.
<http://www.jstor.org/stable/2687037>
- Schumacher, C. S., Siegel, M. J. & Zorn, P. (2015), 2015 CUPM curriculum guide to majors in the mathematical sciences, Technical report, The Mathematical Association of America.
<http://www.maa.org/programs/faculty-and-departments/curriculum-department-guidelines-cupm>
- Trefethen, L. N. & Bau, III, D. (1997), *Numerical linear algebra*, SIAM.
- Turner, P. R., Crowley, J. M., Humpherys, J., Levy, R., Socha, K. & Wasserstein, R. (2015), Modeling across the curriculum II. report on the second SIAM-NSF workshop, Alexandria, VA, Technical report, [http://www.siam.org/reports/modeling_14.pdf].
- Uhlrig, F. (2002), A new unified, balanced, and conceptual

approach to teaching linear algebra, Technical report, Department of Mathematics, Auburn University, <http://www.auburn.edu/~uhligfd/TLA/download/tlateach.pdf>.

Will, T. (2004), Introduction to the singular value decomposition, Technical report, [<http://www.uwlax.edu/faculty/will/svd>].

Index

- \in , 270
 λ , 429
 σ , 235
 θ , 39
 $|\cdot|$, **18**
 $()$, 79
 $+$, $-$, $*$, 79, 81, 169
 $.$, $*$, 199
 $.$, $/$, 199
 $.$, \wedge , 108
 $/$, 79
 $=$, 79, 108
 $[\dots]$, 79, 108
 \wedge , 79
 2-norm, 502
 2 d.p., 110

 $A \setminus$, 108
 $\text{acos}()$, 41, 79
 addition, **25**, **157**, 170, 175
 adjacency matrix, 455
 adolescent, 652–654
 adult, 652–654, 666
 age structure, 652–667
 age structured population, 161, 165
 angle, **39**, 40–42, 56, 66, 206, 225, 662
 ans , 79
 approximate solution, **311**, 316
 arc-cos, 79
 area, 66
 Arrow, Kenneth, 315
 artificial intelligence, 362
 associative law, 31, 32, 175, 176
 augmented matrix, **117**, 117, 118, 123
 average, 501, 522
 axis , 501

 basin area, 362
 basis, **640**, **641**, **694**, 698, 701, 704, 707, 714, 715
 best straight line, 316
 body-centered cubic, 42
 bond angles, 40
 bounded, 577
 brackets, 108, **154**
 bulls eye, 495, 508

 \mathbb{C} , **12**
 canonical form, **477**, 486
 Cardano, Gerolamo, 12
 Cauchy–Schwarz inequality, 47, **48**, 48, 50, 60, 505
 chaos, 252
 characteristic equation, **445**, 453, 623, **631**, **640**
 characteristic polynomial, **631**, 632–637, 640, 671, 672, 741
 chemistry, 40
 closed, 270
 coastlines, 362
 coefficient, **97**, **137**, 633
 $\text{colormap}()$, 326
 column space, **275**, 276, 284, 292, 302, 304, 306, 310, 330, 338, 349
 column vector, 107, 111, **155**, 168, 210, 275, 299, 710
 commutative law, 23, 31, 32, 46, 70, 175
 complex conjugate, 462
 complex eigenvalue, 461, 624, 629, 640, 660, 662
 complex eigenvector, 624
 complex numbers, **12**, 449, 462
 components, **15**, 20, **155**
 composition, **394**, 404, 405
 computed tomography, 326, 365, 563, 564

- computer algebra, 79
- cond, 243, 251, 265, 266, 322, 393
- condition number, 108, **243**, 243–253, 264, 298, 393, 552, 561, 710
- conic section, **472**, 484, 485, 617
- consistent, 102, 123, 140, 316
- constant term, **97**, 126, 633
- contradiction, 247, 358, 462, 468
- coordinate axes, 484, 485
- coordinate system, 13, 681, 703
- coordinate vector, **704**
- coordinates, **704**
- cosine, 41
- cosine rule, 38, 43, 59
- cross product, **65**, 62–77
- cross product direction, 66
- csvread(), 501, 521
- CT scan, 326, 553
- cumprod(), 537
- data mining, 362
- data reduction, 514
- data scaling, 522, 524, 526, 542, 549, 554, 563
- De Moivre's theorem, 662
- decimal places, 110
- Descartes, 31
- det(), 590
- determinant, 73, **189**, 445, 570, **573**, 574, 580, 584, 590, 601, 608, 671
- diag, 198–205, 218, 222, 236, 244, 366, 396, 403, 418, 437, 443, 469, 491, 509, 718
- diag(), 199
- diagonal entries, **198**
- diagonal matrix, **198**, 197–205, 218, 433, 469, 574, 601, 717, 718
- diagonalisable, **717**, 718, 721, 722, 730
- diagonalisation, 716–743
- difference, **25**, **158**, 170
- differential equation, 727, 730, 731
- differential equations, 727–738
- dim, 291, 353, 698, 724, 751
- dimension, **291**, 291–299, 305, 436, 514, **698**, 724, 740
- direction vector, **29**
- discrepancy principle, 561
- discriminant, 48, 49
- displacement vector, **13**, 13, 20
- distance, **27**, 316, 493
- distributive law, 32, 33, 46, 70, 175, 176
- dolphin, 679
- dot product, **39**, 40, 79, 167, **206**, 206, 527, 532
- dot(), 79
- double subscript, **155**
- Duhem, Pierre, 308
- e_j , **26**
- eig(), 437, 638
- eigen-problem, 623, 657
- eigenspace, **434**, 436, 440, 445, 454, 623, 629, 640, 641, 673, 724, 740, 741
- eigenvalue, **429**, 433, 434, 436, 437, 440, 445, 452, 454, 456, 458, 462, 463, 468, 469, 477, 478, 623, 625, 629, 631, 633, 636–638, 640, 641, 650, 659, 669, 670, 672, 680, 681, 688, 718, 722, 724, 730
- eigenvector, **429**, 434, 437, 445, 446, 452, 458, 463, 468, 477, 623, 625, 640, 641, 659, 669, 680, 681, 688, 718, 730
- eigshow(), 253, 432, 502, 624
- El Nino, 285, 292
- elementary row operation, **118**, 123
- elements, **155**
- elephant, 677
- ellipsis, 12
- empirical orthogonal functions, 514
- ensemble of simulations, 290
- entries, **155**
- equal, **15**, **156**
- equation of the plane, **54**, 54
- Error using, 171, 173
- Error using, 83

- `error:`, 83
- error: operator, 171, 173
- Euler, 455, 499
- Euler's formula, 450, 736
- `exp()`, 326
- experimental error, 542
- exponential, 326
- `eye()`, 169
- factorial, 610, 611
- factorisation, 230
- female, 652, 654
- Feynman, Richard, 542
- floating point, **79**
- `format long`, 647
- fractal, 535, 536
- free variable, **121**, 121, 123, 242, 324, 701
- Galileo Galilei, 268
- Gauss–Jordan elimination, 123, 124
- general solution, **659**, 660, 676, 681, 701, 729–731, 742, 743
- giant mouse lemur, 678
- global positioning system, 98, 104, 114, 115, 135
- GPS, 98, 104, 114, 115, 135
- Hack's exponent, 362
- Hankel matrix, 287
- `hankel()`, 286, 307
- Hawking, Stephen, 202
- `hilb()`, 546
- Hilbert matrix, 546, 563
- homogeneous, **126**, 128, 247, 278, 298, 434, 445, 640, 691, 701, 710
- Hotelling transform, 514
- hyper-cube, **573**, 576
- hyper-volume, **573**
- identity matrix, **155**, 160, 161, 168, 169, 186
- identity transformation, 396
- iff, 434
- image compression, 494–514
- `imagesc()`, 326
- Impossibility Theorem, 315
- `imread()`, 501
- in, \in , 270
- inconsistent, 102, **242**, 310, 311, 542
- inconsistent equations, 308–373
- inconsistent system, 330–333, 339
- induction, 255, 256, 468, 469
- Inf**, 108
- infant, 652
- inference, 361
- inferring, 93
- infinitely many solutions, 102, 125, 128
- infinity, 108, 243
- initial condition, 733, 735, 742
- initial population, 663
- initial value, 657, 659, 660, 667
- inner product, **39**
- integer, **11**
- integral equations, 159
- interchanging two columns, 592
- interchanging two rows, 118, 592
- intersection, 348
- `inv()`, 217
- inverse, **187**, 189, 191, 590
- inverse cosine, 79
- inverse matrix, 217
- inverse transformation, **396**, 405
- invertible, **187**, 189, 191, 193, 194, 200, 210, 227, 247, 265, 298, **396**, 399, 445, 456, 458, 570, 583, 659, 710, 717
- Iris, 516
- i*, **26**
- idempotent, 588
- identical columns, 592
- identical rows, 592
- j*, **26**
- jewel in the crown, 4, 230
- jpeg, 494
- juvenile, 652–654, 666

- k***, **26**
- Karhunen–Loève transform, **514**
- Kepler’s law, **319**
- kitten, **666**
- Kleiber’s power law, **362**
- knowledge discovery, **362**
- Lagrange multiplier, **365**
- latent semantic indexing, **524–533**
- leading one, **121**, **121**
- least square, **311**, **338**, **359**, **385**, **559**, **561**
- left triangular, **601**
- Leibnitz, Gottfried, **105**
- length, **18**, **19**, **21**, **27**, **47**, **48**, **66**, **79**, **206**, **225**, **493**, **501**
- Leslie matrix, **161**, **165**, **184**, **195**, **460**
- life expectancy, **316**
- linear combination, **137**, **137–141**, **143–145**, **659**, **682**, **683**, **687**, **704**, **730**
- linear dependence, **681–715**
- linear equation, **97**, **102**, **106**, **116–118**, **121**, **123**, **126**, **128**, **140**, **191**, **226**, **258**, **324**, **542**
- linear independence, **641**, **681–715**
- linear transformation, **375**, **374–402**, **584**, **585**
- linearly dependent, **684**, **681–715**
- linearly independent, **624**, **684**, **681–715**, **718**, **722**
- linguistic vector, **16**
- `log()`, **199**, **326**
- `log10()`, **199**, **321**
- logarithm, **326**
- Lovelace, Ada, **536**, **537**
- lower triangular, **601**
- lower-left triangular, **601**
- magnitude, **18**, **21**, **79**, **662**
- Markov chain, **161**
- Matlab, **78–86**
- MATLAB, **2**, **8**, **9**, **11**, **78**, **79**, **81–84**, **86**, **87**, **105**, **107**, **108**, **108–110**, **119**, **123**, **129**, **130**, **136**, **154**, **159**, **168**, **169**, **169–173**, **181**, **184**, **185**, **191**, **197**, **199**, **199**, **201**, **215–218**, **237**, **237**, **240**, **242–245**, **248**, **250**, **261**, **263–265**, **284**, **286**, **287**, **302**, **304**, **307**, **316**, **319**, **322**, **323**, **325**, **326**, **326**, **328**, **329**, **340**, **364**, **368**, **370**, **403**, **432**, **437**, **437**, **440**, **444**, **453**, **455**, **466**, **468**, **470**, **482**, **483**, **486**, **501**, **501–504**, **520**, **524**, **527**, **530**, **531**, **535–537**, **563**, **590**, **624**, **631**, **638**, **641**, **644**, **646–648**, **650**, **672–675**, **690**, **712**, **715**, **723**, **726**, **727**, **733**, **740**, **741**, **743**
- matrix, **106**, **154**
- matrix multiplication, **176**
- matrix norm, **501**, **502**, **502–505**, **507–509**, **513**, **514**, **520**
- matrix power, **165**, **165**, **176**, **179**
- matrix product, **164**
- matrix-vector form, **106**, **139**
- matrix-vector product, **159**
- maximum, **256**
- mean, **522**
- `mean()`, **501**, **522**
- minor, **597**, **608**
- Moler, Cleve, **4**, **323**, **590**
- Moore–Penrose inverse, **386**
- multiplicity, **436**, **437**, **440**, **454**, **636**, **637**, **638**, **640**, **671**, **672**, **674**, **722**, **724**, **741**
- NaN, **108**
- natural numbers, **11**
- n*D-cube, **573**, **573**, **576**
- n*D-parallelepiped, **576**
- n*D-volume, **573**, **573**, **576**, **577**
- negative, **25**
- nilpotent, **588**
- no solution, **102**, **125**, **242**
- non-diagonalisable matrix, **722**
- non-homogeneous, **126**
- non-orthogonal coordinates, **716**

- nonconformant arguments, 83
- nonlinear equation, 96, **97**, 99
- `norm()`, 78, 79, 501, 503, 504
- normal equation, 322, 557
- normal vector, **54**, 53–55, 61, 63, 64, 69, 72, 75
- not a number, 108
- null, 278, 349
- nullity, **294**, 294, 295, 299, 305, 710
- nullspace, **278**, 293, 294, 304–306, 349, 701

- Occam's razor, 553
- Octave, 2, 8, 9, 11, **79**, 78–87, 105, 107, **108**, 108–110, 119, 123, 129, 130, 136, 154, 159, 168, **169**, 169–173, 181, 184, 185, 191, 197, **199**, 199, 201, 215–218, **237**, 237, 240, 242–245, 248, 250, 261, 263–265, 284, 286, 287, 302, 304, 307, 316, 319, 322, 323, 325, **326**, 326, 328–330, 340, 364, 368, 370, 403, **437**, 437, 440, 444, 453, 455, 466, 468, 470, 482, 483, 486, **501**, 501, 503, 504, 520, 524, 527, 530, 531, 535–537, 563, 590, 631, 638, 641, 644, 646–648, 650, 672–675, 690, 712, 715, 723, 726, 727, 733, 740, 741, 743
- ones matrix, 169
- `ones()`, 108, 169
- orangutan, 653
- orbital period, 319
- orthogonal, **51**, 52, 53, 66, 206, 207, 463, 477, 625, 681
- orthogonal basis, 454
- orthogonal complement, **346**, 346, 348, 349, 369, 370, 372, 373
- orthogonal decomposition, **358**, 372, 373
- orthogonal matrix, **208**, 208–215, 225, 227, 228, 235, 251, 437, 469, 505, 574, 717
- orthogonal projection, **331**, **333**, 330–359, 366–368
- orthogonal set, **207**, 206–207, 222–224
- orthogonal vectors, 51–53
- orthogonally diagonalisable, **469**, 470, 483, 484, 717
- orthonormal, 235, 477
- orthonormal basis, **280**, 279–292, 304, 306, 333, 341, 368, 526, 530, 681, 694, 696, 698, 701, 712
- orthonormal set, **207**, 206–208, 210, 222–224, 227, 280, 686

- parallelepiped, 378
- parallelepiped volume, 71–74
- parallelogram, 378
- parallelogram area, 62, 66, 74
- parameter, **29**, **56**
- parametric equation, **29**, 29, 30, 48, **56**, 55–58, 138, 144
- parentheses, **155**
- partial fraction, 112
- particular solution, 729, 733, 735, 742
- PCA, 514
- perp, 354, 356, 357, 372
- perpendicular, **51**, 51, 53
- perpendicular component, **354**, 356, 371, 372
- pigeonhole principle, 468
- pixel pattern, 511
- planets, 319
- player rating, 310, 312, 339
- Poincaré, Henri, 252
- polar form, 662
- `poly()`, 631
- population, 652–667
- population model, 665
- population modelling, 622, 716
- position vector, **13**, 13, 15, 20, 29, 56
- precision, 79, 184
- principal axes, **477**, 486
- principal component analysis, 514–533, 538

- principal components, **519**, 514–526
- principal vector, **519**, 514–524, 526
- proj, 331–335, 338–343, 354, 357, 359, 366, 367, 379, 422, 423
- proper orthogonal decomposition, 514
- pseudo-inverse, **386**, 384–393, 403, 404
- Punch, John, 553
- Pythagoras, 11

- QR factorisation, 237, 641
- qr-code, 80, 109
- quadratic equation, 472
- quadratic form, **476**, 477, 478
- quit, 79
- quote, 169

- \mathbb{R} , **12**
- `randn()`, 169, 648
- random matrix, 169, 426
- random perturbation, 675
- range, 275
- rank, **244**, 244, 245, 247, 251, 264, 266, 281, 292, 295, 299, 307, 311, 391, 404, 496, 509, 542, 698, 701, 710
- rank theorem, **295**, 305
- rational function, 136
- rational numbers, 11
- rcond, 243
- `rcond()`, 107, 108, 188
- real numbers, **12**
- Recordé, Robert, 15
- reduced row echelon form, 4, **121**, 121, 123, 150
- reflection, 206, **208**
- regularisation, 542–561
- regularisation parameter, 557, 561
- relative error, 561
- repeated eigenvalue, **636**, 647–651, 674, 722
- `reshape()`, 326
- Richardson, L. F., 362
- right triangular, 601
- right-angles, 15, 26, 40, 44, 51, 52
- river length, 362
- \mathbb{R}^n , 16
- rotation, 206, **208**
- rotation and/or reflection, **208**, 210
- row space, **275**, 284, 292, 302, 304, 306, 349, 350
- row vector, **155**, 210, 275, 299, 519, 710

- scalar, **12**, 25, 29, 32, 39, 56
- scalar multiplication, **25**, 175
- scalar product, **159**
- scalar triple product, **72**, 73, 76
- `scatter()`, 501, 517
- `scatter3()`, 501
- searching, 528
- `semilogy()`, 501
- sensitivity, 647–651, 674, 675
- serval, 665
- Seven Bridges of Königsberg, 455
- shear transformation, 398
- Sierpinski carpet, 536
- Sierpinski network, 440
- Sierpinski triangle, 535
- significant digits, 79
- Singular Spectrum Analysis, 285, 306
- singular value, **235**, 242–244, 247, 281, 298, 468, 496, 501, 502, 509, 540, 542, 543, 559, 561, 581, 669, 698, 701, 710
- singular value decomposition, 4, 9, 153, 230, 231, 234, **235**, 235–237, 240, 242–246, 253–256, 258–262, 264, 266, 267, 281, 284, 285, 287, 292, 293, 295, 304, 306–311, 316, 319, 322, 323, 325, 329–331, 336, 339, 359, 360, 364–366, 368, 387, 388, 403, 426, 427, 468–470, 483, 486, 493, 494, 496, 498, 502, 504, 507, 509, 513, 514, 519, 524, 526–528, 530, 533–538, 542, 543, 553, 559, 561, 563, 570, 580–582, 631, 641, 668, 669, 680, 698, 701, 711

- singular vector, **235**, 281, 669
- size, **15**, 15, **154**, 168
- `size()`, 79, 169
- smallest change, 309–311, 314, **316**
- smallest solution, 323, **324**, 559
- Southern Oscillation Index, 285
- span, **141**, 141–143, 145, 146, 151, 273–276, 278, 281, 282, 291, 294, 301, 302, 331, 334, 338, 339, 343, 346, 347, 349, 350, 354, 355, 358, 367, 370–372, 423, 424, 436, 446, 463, 490, 640, 694, 697, 698, 721, 748, 749
- species, 516
- spectral norm, 502
- square matrix, **106**, 107, 108, 154, 169, 187, 210, 247, 298, 429, 434, 437, 443, 445, 456, 458, 469, 470, 573, 578, 580, 581, 583, 601, 631, 640, 659, 710, 716–718, 722, 730
- square-root, 484
- standard basis, 703, 704, 706, 707, 714, 715
- standard coordinate system, **14**, 15
- standard deviation, 501, 522, 524
- standard matrix, **381**, 386, 394, 399, 401, 402
- standard unit vector, **26**, 26, 51, 52, 65, 71, 74, 76, 142, 206, 207, 291, 381, 703, 706, 714
- `std()`, 501, 522
- steganography, 537
- stereo pair, **24**, 203, 209, 214, 223, 228, 298
- subspace, **270**, 268–280, 288, 291, 299, 301, 302, 333, 341, 346, 348, 353, 354, 356, 358, 434, 694, 696, 698, 701, 704
- subtraction, **158**, 170
- sum, **25**, **157**, 170
- SVD, 4, 9, 153, 230, 231, 234–237, 240, 242–246, 253–256, 258–262, 264, 266, 267, 281, 284, 285, 287, 292, 293, 295, 304, 306–311, 316, 319, 322, 323, 325, 329–331, 336, 339, 359, 360, 364–366, 368, 387, 388, 403, 426, 427, 468–470, 483, 486, 493, 494, 496, 498, 502, 504, 507, 509, 513, 514, 519, 524, 526–528, 530, 533–538, 542, 543, 553, 559, 561, 563, 570, 580–582, 631, 641, 668, 669, 680, 698, 701, 711
- `svd()`, 237
- `svds()`, 501, 522, 524, 526, 530
- symmetric matrix, **167**, 168, 179, 184, 426–487, 650, 651, 669, 717, 724
- system, **97**, 106, 117, 118, 123
- table tennis, 310
- Tasmanian Devil, 184, 676
- Tikhonov regularisation, **557**, 556–561, 563, 564
- tipping point, 290
- tomography, 326
- trace, **633**, 670, 671
- transformation, 374, 375
- transpose, **166**, 166–169, 173, 179, 184, 185, 699
- triangle inequality, 47, **48**, 50, 60, 505
- triangular matrix, 600, **601**, 601, 625, 629, 719
- triple product, scalar, 72
- unique solution, 102, 107, 125, 191, 243, 247, 298, 710
- unit cube, 378
- unit square, 377
- unit vector, **18**, 53, 206, 207, 427, 477, 478
- upper triangular, 601
- upper-right triangular, 601
- van de Snepscheut, Jan L. A., 107
- variance, 524
- vector, **15**, 18

vector product, [65](#)
vectors, [13](#)
velocity vector, [14](#)
volume, parallelepiped, [71](#), [76](#)

walking gait, [306](#)
weather forecasts, [290](#)
Wikipedia, [12](#), [135](#), [315](#), [319](#), [326](#), [556](#),
[653](#), [665](#)
Wiles, Andrew, [253](#)
wine recognition, [521](#)
word vector, [22](#), [42](#), [60](#), [525](#), [529](#)
work, [44](#)

X-ray, [553](#)

zero, [126](#), [324](#), [456](#), [468](#), [543](#)
zero column, [592](#)
zero matrix, [155](#), [168](#), [169](#)
zero row, [592](#)
zero vector, [16](#), [17](#), [19](#), [348](#), [434](#)
`zeros()`, [169](#)

Answers to selected activities

No further Activity answers collected.

Answers to selected exercises

: No further answers collected.

in Draft