Invariant manifold of your dynamical system

A. J. Roberts, University of Adelaide http://orcid.org/0000-0001-8930-1552

6:36 A.M., June 5, 2023

Throughout and generally: the lowest order, most important, terms are near the end of each expression.

off echo;

The specified dynamical system

$$\dot{u}_1 = -e_0 \varepsilon^2 u_1 - e_0 \varepsilon \exp(0) s_0 + \varepsilon u_1 u_2 + s_0 u_2 + u_2$$
$$\dot{u}_2 = e_0 \varepsilon^2 u_1 + e_0 \varepsilon \exp(0) s_0 - \varepsilon u_1 u_2 - s_0 u_2 - 2u_2$$

Invariant subspace basis vectors

$$\vec{e}_1 = \{\{1, 0\}, \exp(0)\}\$$

 $\vec{z}_1 = \{\{1, 0\}, \exp(0)\}\$
off echo;

The invariant manifold These give the location of the invariant manifold in terms of parameters s_i .

$$u_1 = O(\varepsilon^4) + s_1$$

$$u_2 = (2e_0^2\varepsilon^3s_0)/(s_0^4 + 8s_0^3 + 24s_0^2 + 32s_0 + 16) + (-2e_0\varepsilon^3s_1^2)/(s_0^3 + 6s_0^2 + 12s_0 + 8) + (2e_0\varepsilon^2s_1)/(s_0^2 + 4s_0 + 4) + (e_0\varepsilon s_0)/(s_0 + 2) + O(\varepsilon^4)$$

Invariant manifold ODEs The system evolves on the invariant manifold such that the parameters evolve according to these ODEs.

$$\begin{split} \dot{s}_1 &= e_0^2 \varepsilon^4 \big(-4s_1 s_0^2 + 2s_1 s_0 + 4s_1 \big) / \big(s_0^5 + 10 s_0^4 + 40 s_0^3 + 80 s_0^2 + 80 s_0 + \\ 32 \big) + e_0^2 \varepsilon^3 \big(2s_0^2 + 2s_0 \big) / \big(s_0^4 + 8s_0^3 + 24s_0^2 + 32 s_0 + 16 \big) + \big(-2e_0 \varepsilon^4 s_1^3 \big) / \big(s_0^4 + 8s_0^3 + 24s_0^2 + 32 s_0 + 16 \big) + \big(2e_0 \varepsilon^3 s_1^2 \big) / \big(s_0^3 + 6s_0^2 + 12 s_0 + 8 \big) + \big(-2e_0 \varepsilon^2 s_1 \big) / \big(s_0^2 + 4s_0 + 4 \big) + \big(-e_0 \varepsilon s_0 \big) / \big(s_0 + 2 \big) + O(\varepsilon^5) \end{split}$$

Normals to isochrons at the slow manifold Use these vectors: to project initial conditions onto the slow manifold; to project non-autonomous forcing onto the slow evolution; to predict the consequences of modifying the original system; in uncertainty quantification to quantify effects on the model of uncertainties in the original system. The normal vector $\vec{z_i} := (z_{i1}, \ldots, z_{in})$

$$\begin{split} z_{11} &= e_0^2 \varepsilon^4 (10s_0^2 + 8s_0 - 4)/(s_0^6 + 12s_0^5 + 60s_0^4 + 160s_0^3 + 240s_0^2 + 192s_0 + 64) + \\ &(-6e_0 \varepsilon^4 s_1^2 s_0)/(s_0^5 + 10s_0^4 + 40s_0^3 + 80s_0^2 + 80s_0 + 32) + e_0 \varepsilon^3 (4s_1 s_0 + 2s_1)/(s_0^4 + 8s_0^3 + 24s_0^2 + 32s_0 + 16) + e_0 \varepsilon^2 (-2s_0 - 2)/(s_0^3 + 6s_0^2 + 12s_0 + 8) + O(\varepsilon^5) + 1 \\ z_{12} &= e_0^2 \varepsilon^4 (10s_0^3 + 17s_0^2 - 8)/(s_0^7 + 14s_0^6 + 84s_0^5 + 280s_0^4 + 560s_0^3 + 672s_0^2 + 448s_0 + 128) + e_0 \varepsilon^4 (-6s_1^2 s_0^2 + 6s_1^2 s_0 + 16s_1^2)/(s_0^6 + 12s_0^5 + 60s_0^4 + 160s_0^3 + 240s_0^2 + 192s_0 + 64) + e_0 \varepsilon^3 (4s_1 s_0^2 + s_1 s_0 - 6s_1)/(s_0^5 + 10s_0^4 + 40s_0^3 + 80s_0^2 + 80s_0 + 32) + e_0 \varepsilon^2 (-2s_0^2 - 3s_0)/(s_0^4 + 8s_0^3 + 24s_0^2 + 32s_0 + 16) + (-\varepsilon^4 s_1^4)/(s_0^5 + 10s_0^4 + 40s_0^3 + 80s_0^2 + 80s_0 + 32) + (\varepsilon^3 s_1^3)/(s_0^4 + 8s_0^3 + 24s_0^2 + 32s_0 + 16) + (-\varepsilon^2 s_1^2)/(s_0^3 + 6s_0^2 + 12s_0 + 8) + (\varepsilon s_1)/(s_0^2 + 4s_0 + 4) + (O(\varepsilon^5) s_0 + 2O(\varepsilon^5) + s_0 + 1)/(s_0 + 2) + (\varepsilon^5 s_0^2 + 12s_0 + 8) + (\varepsilon s_1)/(s_0^2 + 4s_0 + 4) + (O(\varepsilon^5) s_0 + 2O(\varepsilon^5) + s_0 + 1)/(s_0 + 2) + (\varepsilon^5 s_0^2 + 12s_0 + 8) + (\varepsilon^5 s_0^2 + 12s_0 + 8) + (\varepsilon^5 s_0^2 + 4s_0 + 4) + (O(\varepsilon^5) s_0 + 2O(\varepsilon^5) + s_0 + 1)/(s_0 + 2) + (\varepsilon^5 s_0^2 + 2s_0^2 + 8s_0^2 + 8s_0$$