



Lecture 10

Power Series (冪級数)

&

Taylor Series (テイラー級数)

What you will learn in Lecture 10

10.1 Power Series (冪級数)

10.2 Taylor Series (テイラー級数)

10.1 Power Series (冪級数)

10.1 Power Series (冪級数)

Power Series

The notion of a **power series** is important in the study of **analytic functions**.

An infinite series

$$\sum_{k=0}^{\infty} a_k (z - z_0)^k = a_0 + a_1 (z - z_0) + a_2 (z - z_0)^2 + \cdots \quad (6.1.11)$$

where the coefficients (係数) a_k are complex constants, is called a **power series** in $z - z_0$.

The power series (6.1.11) is said to be **centered at z_0** ;
the complex point z_0 is referred to as the **center of the series**.

10.1 Power Series (冪級数)

Radius of Convergence (収束半径)

Every **complex power series** (6.1.11) has a **radius of convergence**, which is the circle centered at z_0 of largest radius $R > 0$ for which (6.1.11) converges at every point within the circle $|z - z_0| = R$.

Power series: (i) converges absolutely at all points z within its circle of convergence, i.e. for all z satisfying $|z - z_0| < R$;
and (ii) diverges at all points z exterior (外側の) to the circle, i.e. for all z satisfying $|z - z_0| > R$.

10.1 Power Series (冪級数)

Radius of Convergence

The radius of convergence can be:

- (i) $R = 0$ (in which case (6.1.11) converges only at its center $z = z_0$),
- (ii) R is a finite positive number (in which case (6.1.11) converges at all interior points of the circle $|z - z_0| = R$), or
- (iii) $R = \infty$ (in which case (6.1.11) converges for all z).

10.1 Power Series (冪級数)

EXAMPLE (例題) 6.1.5 Circle of Convergence

Evaluate the convergence condition of the power series $\sum_{k=1}^{\infty} \frac{z^{k+1}}{k}$.

Solution (解答):

By the ratio test (6.1.9)
$$\lim_{n \rightarrow \infty} \left| \frac{\frac{z^{n+2}}{n+1}}{\frac{z^{n+1}}{n}} \right| = \lim_{n \rightarrow \infty} \frac{n}{n+1} |z| = \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{1}{n}} |z| = |z|$$

Thus the series converges absolutely for $|z| < 1$. The circle of convergence is $|z| = 1$ and the radius of convergence is $R = 1$. Note that on the circle of convergence $|z| = 1$, the series does not converge absolutely since $\sum_{k=1}^{\infty} \frac{1}{k}$ is the well-known divergent harmonic series.

Bear in mind this *does not say* that the series diverges on the circle of convergence. In fact, at

$z = -1$, $\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k}$ is the convergent alternating harmonic series. Indeed, it can be shown that

the series converges at all points on the circle $|z| = 1$ except at $z = 1$.

10.1 Power Series (冪級数)

Notice

It should be clear from **Theorem 6.4** in Lecture 9 and **Example 6.1.5** that for a power series $\sum_{k=0}^{\infty} a_k (z - z_0)^k$, the limit (6.1.9) depends only on the coefficients a_k . Thus, if

(i) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L \neq 0$, the radius of convergence is $R = \frac{1}{L}$. (6.1.12)

(ii) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0$, then the series diverges $R = \infty$. (6.1.13)

(iii) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty$, the test is inconclusive $R = 0$. (6.1.14)

10.1 Power Series (冪級数)

Notice

Similar conclusions can be made for **the root test (6.1.10)** by using

$$\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} \quad (6.1.15)$$

For example, if $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L \neq 0$, then $R = \frac{1}{L}$

10.1 Power Series (冪級数)

EXAMPLE (例題) 6.1.6 Radius of Convergence by Ratio Test

For the power series $\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k!} (z - 1 - i)^k$, find its radius of convergence and the condition that it converges absolutely.

Solution (解答):

We note that $a_n = \frac{(-1)^{n+1}}{n!}$ then

$$\lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+2}}{(n+1)!}}{\frac{(-1)^{n+1}}{n!}} \right| = \lim_{n \rightarrow \infty} \frac{1}{n+1} = 0$$

Hence by (6.1.13) the radius of convergence is ∞ ;
the power series with center $z_0 = 1 + i$ converges absolutely
for all z , that is, for $|z - 1 - i| < \infty$.

10.1 Power Series (冪級数)

EXAMPLE (例題) 6.1.7 Radius of Convergence by Root Test

For the power series $\sum_{k=1}^{\infty} \left(\frac{6k+1}{2k+5}\right)^k (z - 2i)^k$, find its radius of convergence and the condition that it converges absolutely.

Solution (解答):

We note that $a_n = \left(\frac{6n+1}{2n+5}\right)^n$ then

$$\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \lim_{n \rightarrow \infty} \frac{6n+1}{2n+5} = 3$$

By reasoning similar to that leading to (6.1.12), we conclude that the radius of convergence of the series is $R = \frac{1}{3}$. The circle of convergence is $|z - 2i| = \frac{1}{3}$; the power series converges absolutely for $|z - 2i| < \frac{1}{3}$.

10.1 Power Series (冪級数)

Sums, products and ratios of power series

Let $\sum_{k=0}^{\infty} a_k(z - z_0)^k$ and $\sum_{k=0}^{\infty} b_k(z - z_0)^k$ be two convergent power series whose limits are S and T respectively at a given point z . Then the following properties hold:

- (i) $cS = \sum_{k=0}^{\infty} ca_k(z - z_0)^k, \forall c \in \mathbb{C};$
- (ii) $S + T = \sum_{k=0}^{\infty} (a_k + b_k)(z - z_0)^k, \forall c \in \mathbb{C};$
- (iii) If S or T converges absolutely, then

$$ST = \sum_{k=0}^{\infty} c_k(z - z_0)^k$$

where $c_n = \sum_{k=0}^n a_k b_{n-k}$; furthermore, if the two series S and T converge absolutely, the series converges absolutely;

- (iv) If (a) $b_0 \neq 0$,
 - (b) the series $H = \sum_{k=0}^{\infty} d_k(z - z_0)^k$, where the coefficients d_k are obtained by solving the equations $\sum_{k=0}^n a_k b_{n-k} = a_n, n = 0, 1, \dots$, converges,
 - (c) T or H converges absolutely,
 - (d) $T \neq 0$,

then $\frac{S}{T} = \sum_{k=0}^{\infty} d_k(z - z_0)^k$

10.2 Taylor Series (テイラー級数)

10.2 Taylor Series (テイラー級数)

Notice: A power series *defines* or *represents* a function f .

Differentiation and Integration of Power Series

Theorem 6.6 Continuity

A power series $\sum_{k=0}^{\infty} a_k (z - z_0)^k$ represents a continuous function f within its circle of convergence $|z - z_0| = R$.

10.2 Taylor Series (テイラー級数)

Differentiation and Integration of Power Series

Theorem 6.7 Term-by-Term (項別に) Differentiation

A power series $\sum_{k=0}^{\infty} a_k (z - z_0)^k$ can be differentiated term-by-term within its circle of convergence $|z - z_0| = R$.

Differentiating a power series term-by-term gives,

$$\frac{d}{dz} \sum_{k=0}^{\infty} a_k (z - z_0)^k = \sum_{k=0}^{\infty} a_k \frac{d}{dz} (z - z_0)^k = \sum_{k=1}^{\infty} a_k k (z - z_0)^{k-1}$$

Note that the summation index in the last series starts with $k = 1$ because the term corresponding to $k = 0$ is zero.

10.2 Taylor Series (テイラー級数)

Differentiation and Integration of Power Series

Theorem 6.8 Term-by-Term Integration

A power series $\sum_{k=0}^{\infty} a_k (z - z_0)^k$ **can be integrated term-by-term within its circle of convergence $|z - z_0| = R$** , for every contour C lying entirely within the circle of convergence.

This theorem gives that

$$\int_C \sum_{k=0}^{\infty} a_k (z - z_0)^k dz = \sum_{k=0}^{\infty} a_k \int_C (z - z_0)^k dz$$

whenever C lies in the interior of $|z - z_0| = R$.

10.2 Taylor Series (テイラー級数)

Taylor Series

Suppose a power series represents a function f within $|z - z_0| = R$, that is,

$$f(z) = \sum_{k=0}^{\infty} a_k (z - z_0)^k = a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \cdots \quad (6.2.1)$$

It follows from Theorem 6.7 that the derivatives of f are the series

$$f'(z) = \sum_{k=1}^{\infty} a_k k (z - z_0)^{k-1} = a_1 + 2a_2(z - z_0) + 3a_3(z - z_0)^2 + \cdots \quad (6.2.2)$$

$$f''(z) = \sum_{k=2}^{\infty} a_k k(k-1)(z - z_0)^{k-2} = 2 \cdot 1 \cdot a_2 + 3 \cdot 2 \cdot a_3(z - z_0) + \cdots \quad (6.2.3)$$

$$f'''(z) = \sum_{k=3}^{\infty} a_k k(k-1)(k-2)(z - z_0)^{k-3} = 3 \cdot 2 \cdot 1 \cdot a_3 + 4 \cdot 3 \cdot 2 \cdot a_4(z - z_0) + \cdots \quad (6.2.4)$$

and so on.

10.2 Taylor Series (テイラー級数)

Taylor Series

Since the power series (6.2.1) represents a differentiable function f within its circle of convergence $|z - z_0| = R$, where R is either a positive number or infinity (無限大), we conclude that *a power series represents an analytic function* within its circle of convergence.

There is a relationship between the coefficients a_k in (6.2.1) and the derivatives of f . Evaluating (6.2.1), (6.2.2), (6.2.3), and (6.2.4) at $z = z_0$ we have

$$f(z_0) = a_0, \quad f'(z_0) = 1! a_1, \quad f''(z_0) = 2! a_2, \quad f'''(z_0) = 3! a_3,$$

In general, $f^{(n)}(z_0) = n! a_n$, or

$$a_n = \frac{f^{(n)}(z_0)}{n!}, \quad n \geq 0 \tag{6.2.5}$$

10.2 Taylor Series (テイラー級数)

Taylor Series

When $n = 0$ in (6.2.5), we interpret the **zero-order derivative** as $f(z_0)$ and $0! = 1$ so that the formula gives $a_0 = f(z_0)$.

Substituting (代入する) (6.2.5) into (6.2.1), we have

$$f(z) = \sum_{k=0}^{\infty} \frac{f^{(k)}(z_0)}{k!} (z - z_0)^k \quad (6.2.6)$$

This series is called **the Taylor series for f centered at z_0** .

10.2 Taylor Series (テイラー級数)

Maclaurin Series (マクローリン展開)

A Taylor series with center $z_0 = 0$,

$$f(z) = \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} z^k \quad (6.2.7)$$

is referred to as **a Maclaurin series**.

10.2 Taylor Series (テイラー級数)

Question

If we are given a function f that is analytic in some domain D ,
can we represent it by a power series of the form (6.2.6) or (6.2.7)?

Check the answer in Theorem 6.9.

Theorem 6.9 Taylor's Theorem

Let f be analytic within a domain D and let z_0 be a point in D .

Then f has the series representation

$$f(z) = \sum_{k=0}^{\infty} \frac{f^{(k)}(z_0)}{k!} (z - z_0)^k \quad (6.2.8)$$

which is valid for the largest circle C with center at z_0 and radius R that lies entirely within D .

10.2 Taylor Series (テイラー級数)

Some Important Maclaurin Series

$$e^z = 1 + \frac{z}{1!} + \frac{z^2}{2!} + \cdots = \sum_{k=0}^{\infty} \frac{z^k}{k!} \quad (6.2.12)$$

$$\sin z = z - \frac{z^3}{3!} + \frac{z^5}{5!} - \cdots = \sum_{k=0}^{\infty} (-1)^k \frac{z^{2k+1}}{(2k+1)!} \quad (6.2.13)$$

$$\cos z = 1 - \frac{z^2}{2!} + \frac{z^4}{4!} - \cdots = \sum_{k=0}^{\infty} (-1)^k \frac{z^{2k}}{(2k)!} \quad (6.2.14)$$

10.2 Taylor Series (テイラー級数)

EXAMPLE (例題) 6.2.1 Radius of Convergence

Suppose the function $f(z) = \frac{3-i}{1-i+z}$ is expanded in a Taylor series with center $z_0 = 4 - 2i$. What is its radius of convergence R ?

Solution (解答):

Observe that the function is analytic at every point except at $z = -1 + i$, which is an isolated singularity of f . The distance from $z = -1 + i$ to $z_0 = 4 - 2i$ is

$$|z - z_0| = \sqrt{(-1 - 4)^2 + (1 - (-2))^2} = \sqrt{34}$$

This last number is the radius of convergence R for the Taylor series centered at $4 - 2i$.

10.2 Taylor Series (テイラー級数)

EXAMPLE (例題) 6.2.2 Maclaurin Series

Find the Maclaurin expansion of $f(z) = \frac{1}{(1-z)^2}$.

Solution (解答):

We could, of course, begin by computing the coefficients using (6.2.8). However, recall from (6.1.6) in Lecture 9 that for $|z| < 1$,

$$\frac{1}{1-z} = 1 + z + z^2 + z^3 + \dots \quad (6.1.6)$$

If we differentiate both sides of the last result with respect to z , then

$$\frac{d}{dz} \frac{1}{1-z} = \frac{d}{dz} 1 + \frac{d}{dz} z + \frac{d}{dz} z^2 + \frac{d}{dz} z^3 + \dots$$

or

$$\frac{1}{(1-z)^2} = 0 + 1 + 2z + 3z^2 + \dots = \sum_{k=1}^{\infty} k z^{k-1}$$

Since we are using Theorem 6.7, the radius of convergence of the last power series is the same as the original series, $R = 1$.

10.2 Taylor Series (テイラー級数)

EXAMPLE (例題) 6.2.3 Taylor Series

Expand $f(z) = \frac{1}{1-z}$ in a Taylor series with center $z_0 = 2i$.

Solution (解答):

We again use the geometric series (6.1.6) in Lecture 9. By adding and subtracting $2i$ in the denominator of $1/(1-z)$, we can write

$$\frac{1}{1-z} = \frac{1}{1-z+2i-2i} = \frac{1}{1-2i-(z-2i)} = \frac{1}{1-2i} \frac{1}{1-\frac{z-2i}{1-2i}}$$

We now write $\frac{1}{1-\frac{z-2i}{1-2i}}$ as a power series by using (6.1.6) with that z replaced by the expression $\frac{z-2i}{1-2i}$

$$\frac{1}{1-z} = \frac{1}{1-2i} \left[1 + \frac{z-2i}{1-2i} + \left(\frac{z-2i}{1-2i} \right)^2 + \left(\frac{z-2i}{1-2i} \right)^3 + \cdots \right] \quad (6.2.17)$$

Because the distance from the center $z_0 = 2i$ to the nearest singularity $z = 1$ is $\sqrt{5}$, we conclude that the circle of convergence for (6.2.17) is $|z_0 - 2i| = \sqrt{5}$. This can be verified by the ratio test of the Lecture 9.

Review for Lecture 10

- (Complex) Power Series
- (Complex) Taylor Series
- (Complex) Maclaurin Series

Assignment

Please Check <https://github.com/uoaworks/ComplexAnalysisAY2018>

Reading Materials: Section 6.1, Section 6.2, Textbook

References

- [1] A first course in Complex Analysis with application, Dennis G. Zill and Patrick D. Shanahan, Jones and Bartlett Publishers, Inc. 2003
- [2] Wikipedia