



Lecture **6**

Complex Integrals (複素積分)

What you will learn in Lecture 6

***6.1** Review of **Real Line Integral (実・線積分)**

6.2 **Complex Integral (複素積分)**

Real Line Integral

(実・線積分)

in the Cartesian Plane



Complex Integral

(複素積分)

in the Complex Plane

***6.1 Review of Real Line Integral**

(実・線積分)


Notice: In all lecture notes, the contents marked with * are not in the scope of the final examination.

*6.1 Review of Real Line Integral (実・線積分)


Domain (定義域) $x \in \mathbf{R}$




$(x, f(x))$



Domain (定義域) $(x, y) \in \mathbf{D}$



$(x, y, f(x, y))$



*6.1 Review of Real Line Integral (実・線積分)

One-Variable Calculus -- Definite integral (定積分) of f

$$\int_a^b f(x)dx = \lim_{\|\Delta x\| \rightarrow 0} \sum_{k=1}^n f(x_k^*)\Delta x = \lim_{\|\Delta x\| \rightarrow 0} [f(x_1^*)\Delta x + f(x_2^*)\Delta x + \cdots + f(x_n^*)\Delta x]$$


One-Variable Function

relationship, called function


$$y = f(x)$$

Independent variable

dependent variable





$$Area(\text{面積}) = \int_a^b f(x)dx$$



$\Delta x_1 = \Delta x_2 = \cdots = \Delta x_n = \Delta x$,
i.e. equal interval (等間隔)

*6.1 Review of Real Line Integral (実・線積分)

Recall: what is Line integral (線積分)?



*6.1 Review of Real Line Integral (実・線積分)

Line Integral (線積分)

If f is defined on a smooth (滑らか) or piecewise-smooth (区分的滑らか) curve C , then the line integral of f along C is

$$\int_C f(x, y) ds = \lim_{\|\Delta s_{max}\| \rightarrow 0} \sum_{k=1}^n f(x_k^*, y_k^*) \Delta s_k$$

if this limits exists. (Here the norm $\|\Delta s_{max}\|$ defines the length of the longest subinterval (部分区間))

How to compute Line Integral?

By introducing Arc length (弧長) $L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$, we have



$$\int_C f(x, y) ds = \int_C f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

*6.1 Review of Real Line Integral (実・線積分)

Arc Length (弧長)


$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$|\overrightarrow{P_1P_2}| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$



*6.1 Review of Real Line Integral (実・線積分)

Insight (洞察) of Line Integrals



In fact, if $f(x, y) > 0$,

$\int_C f(x, y) ds$ represents the area (面積) of one side of the "curtain".




6.2 Complex Integral (複素積分)

6.2 Complex Integral (複素積分)

Q: How to represent curves in the complex plane?

6.2 Complex Integral (複素積分)

Parametrization (パラメータ表示)
of **Real curve** (実・曲線)



Parametrization (パラメータ表示)
of **Complex curve** (複素・曲線)





Figure 5.15 $z(t) = x(t) + iy(t)$ as a position vector

$$z(t) = x(t) + iy(t), a \leq t \leq b \quad (5.2.1)$$

The points z on the curve C is expressed by a complex-valued function of a real variable t . This is called a parametrization of C .

6.2 Complex Integral (複素積分)



Parametrization (パラメータ表示)
of **Complex curve** (複素・曲線)

$$z(t) = x(t) + iy(t), a \leq t \leq b$$

For example,

$$\text{if } x(t) = \cos t, y(t) = \sin t, 0 \leq t \leq 2\pi$$


we will have

$$z(t) = \cos t + i \sin t, 0 \leq t \leq 2\pi$$


which is a parametrization of the circle C .

6.2 Complex Integral (複素積分)


Piecewise-smooth (区分的滑らか) curve



A smooth curve C



A piecewise-smooth
curve $C = C_1 \cup C_2$



A piecewise-smooth curve
 $C = C_1 \cup C_2 \cup \cdots \cup C_5$

6.2 Complex Integral (複素積分)

Suppose the derivative of

$$z(t) = x(t) + iy(t), a \leq t \leq b \quad (5.2.1)$$

is

$$z'(t) = x'(t) + iy'(t)$$




Figure 5.16 $z'(t) = x'(t) + iy'(t)$ as a tangent vector (接ベクトル)

6.2 Complex Integral (複素積分)

Smooth (滑らか) Curve

A curve C in the complex plane is called **smooth** if $z'(t)$ is **continuous** and **NEVER** zero in the interval $a \leq t \leq b$.

In other words, **a smooth curve have NO sharp corners or Cusps (尖点).**

Piecewise-smooth (区分的滑らか) Curve

A piecewise smooth curve C is **continuous** EXCEPT possibly at the points where the component smooth curves C_1, C_2, \dots, C_n are joined together.




Figure 5.16 $z'(t) = x'(t) + iy'(t)$ as a tangent vector (接ベクトル)




Figure 5.17 Curve C is not smooth because it has a cusp

6.2 Complex Integral (複素積分)

Simple (單一) Curve


A curve C in the complex plane is said to be a **simple** if $z(t_1) \neq z(t_2)$ for $t_1 \neq t_2$, except possibly for initial point $t = a$ and terminal point $t = b$.

Closed (閉) Curve


C is a closed curve if $z(a) = z(b)$.

Simple Closed Curve (單一閉曲線)

C is a **simple closed curve** if $z(t_1) \neq z(t_2)$ for $t_1 \neq t_2$ and $z(a) = z(b)$.




Smooth, simple,
not closed



Piecewise smooth,
simple, not closed



Smooth, closed,
not simple



Smooth, simple,
closed

6.2 Complex Integral (複素積分)

Contour

A piecewise smooth curve C is called a **Contour** or Path.


Direction (向き) on a Contour

- We define the **positive direction** on a contour C corresponding to increasing values of the parameter t .
- Roughly, for a simple closed curve C , the **positive direction** is the **counterclockwise** (左回りの) direction or the direction that **a person must walk on C and keep the interior (内部) of C at the left hand**.
- The **negative direction** on a contour C is the direction **opposite** (反対の) the positive direction.
- Notice:** If C has positive direction, then **its opposite curve** can be denoted by $-C$.

exterior
外部



(a) Positive direction



(b) Positive direction

Figure 5.18 Interior of each curve is at the left hand

6.2 Complex Integral (複素積分)

Note: Find more explanations in Page 247 ~ 250 of the textbook.

Definition 5.3 Complex Integral (複素積分)

An integral of a function $f(z)$ defined by

$$\int_C f(z) dz = \lim_{\|\Delta z_{max}\| \rightarrow 0} \sum_{k=1}^n f(z_k^*) \Delta z_k \quad (5.2.2)$$

is called a **complex integral**, where z is a complex number, and $f(z)$ is defined on a contour C (積分路). (Here the norm $\|\Delta z_{max}\|$ defines the length of the longest subinterval (部分区間))


- If the limit in (5.2.2) exists, then $f(z)$ is said to be **integrable** (可積分の) on C .
- The **limit exists** whenever if $f(z)$ is **continuous** at all points on C , where C is either smooth or piecewise smooth.
- The **Complex Integral** $\int_C f(z) dz$ has a more common name: **Contour Integral**.

6.2 Complex Integral (複素積分)

- Specially, we will use the notation

$$\oint_C f(z)dz$$

as a complex integral around a positively oriented closed curve C .



Positive direction

6.2 Complex Integral (複素積分)

Integral for Complex-Valued Function of a Real Variable

Example

If t represents a real variable, then the output of the function $f(t) = (2t + i)^2$ is a complex number. For $t = 2$,

$$f(2) = (2 \cdot 2 + i)^2 = 16 + 8i + i2 = 15 + 8i.$$

In general, if f_1 and f_2 are real-valued functions of a real variable t (that is, real functions), then $f(t) = f_1(t) + if_2(t)$ is a complex-valued function of a real variable t .

When we consider the interval $0 \leq t \leq 1$,

$$\int_0^1 (2t + i)^2 dt = \int_0^1 (4t^2 - 1 + i4t) dt = \int_0^1 (4t^2 - 1) dt + i \int_0^1 4t dt = \left(\frac{4}{3} \cdot t^3 - t \right) \Big|_0^1 + i \cdot 2t^2 \Big|_0^1 = \frac{1}{3} + 2i$$

Then we can define the integral of the complex-valued function $f(t) = f_1(t) + if_2(t)$ on interval $a \leq t \leq b$ as

$$\int_a^b f(t) dt = \int_a^b f_1(t) dt + i \int_a^b f_2(t) dt \quad (5.2.4)$$

6.2 Complex Integral (複素積分)

Theorem 5.1 How to Compute a Complex Integral (i.e. Complex Integral)

If f is continuous on a smooth curve C given by the parametrization (パラメータ表示) $z(t) = x(t) + iy(t)$, $a \leq t \leq b$, then

$$\int_C f(z) dz = \int_a^b f(z(t)) z'(t) dt \quad (5.2.11)$$

where

$$f(z(t)) z'(t) = [u(x(t), y(t)) + iv(x(t), y(t))][x'(t) + iy'(t)]$$

Recall $z(t)$ in Equation (5.2.1)

6.2 Complex Integral (複素積分)

EXAMPLE (例題) 5.2.1 Evaluating a Contour Integral

Evaluate $\int_C \bar{z} dz$, where C is given by $z(t) = x(t) + iy(t)$, $x(t) = 3t$, $y(t) = t^2$, $-1 \leq t \leq 4$.

Solution (解答):

Hint:

- Equation (5.2.11)
- Equation (5.2.4)

The Lecture Slides with complete solution will be uploaded with Assignment sheet after the class.

Check <https://github.com/uoaworks/ComplexAnalysisAY2018>

6.2 Complex Integral (複素積分)

EXAMPLE (例題) 5.2.2 Evaluating a Contour Integral

Evaluate $\oint_C \frac{1}{z} dz$, where C is the circle $x(t) = \cos t$, $y(t) = \sin t$, $0 \leq t \leq 2\pi$.

Solution (解答):

Hint:

- Equation (5.2.11)
- Equation (5.2.4)
- Example in Page 34 of this Lecture 6 Slides

The Lecture Slides with complete solution will be uploaded with Assignment sheet after the class.

Check <https://github.com/uoaworks/ComplexAnalysisAY2018>

6.2 Complex Integral (複素積分)

Theorem 5.2 Properties of Contour Integrals

Suppose the functions f and g are continuous in a domain D , and C is a smooth or piecewise smooth curve in D . Then

(i) $\int_C qf(z) dz = q \int_C f(z) dz$, where q is a complex constant.

(ii) $\int_C [f(z) + g(z)] dz = \int_C f(z) dz + \int_C g(z) dz$

(iii) $\int_C f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz$, where C consists of the smooth curves C_1 and C_2 joined end to end.

(iv) $\int_{-C} f(z) dz = - \int_C f(z) dz$, where $-C$ denotes the curve having the opposite Orientation (向き) of C .

6.2 Complex Integral (複素積分)

EXAMPLE (例題) 5.2.3 C is a Piecewise Smooth Curve

Evaluate $\int_C (x^2 + iy^2) dz$, where C is the contour shown in Figure 5.20.

Solution (解答):

Hint:

- Theorem 5.2 (iii)
- Equation (5.2.11)
- Equation (5.2.4)




Figure 5.20 Contour $C = C_1 \cup C_2$ is piecewise-smooth

The Lecture Slides with complete solution will be uploaded with Assignment sheet after the class.

Check <https://github.com/uoaworks/ComplexAnalysisAY2018>

6.2 Complex Integral (複素積分)

Solution (解答)(cont.):




Figure 5.20 Contour $C = C_1 \cup C_2$ is piecewise-smooth

6.2 Complex Integral (複素積分)

*Theorem 5.3 A Bounding (界) Theorem

If f is continuous on a smooth curve C and if modulus $|f(z)| \leq M$ for all z on C , Then $\left| \int_C f(z) dz \right| \leq ML$, where L is the length of C .

6.2 Complex Integral (複素積分)

*EXAMPLE (例題) 5.2.4 A Bound for a Contour Integral

Find an upper bound (上界) for the absolute value of $\oint_C \frac{e^z}{z+1} dz$, where C is the circle $|z| = 4$.

Solution (解答):

Hint:

- Theorem 5.3
- Inequality (1.2.7) $|z_1 + z_2| \geq |z_1| - |z_2|$ in Page 35 of the Lecture 1 Slides.

The Lecture Slides with complete solution will be uploaded with Assignment sheet after the class.

Check <https://github.com/uoaworks/ComplexAnalysisAY2018>

Review for Lecture 6

- Real Line Integral
- Piecewise Smooth Curve
- Simple, Closed Curve
- Complex Integral (Contour Integral)
- How to Compute Complex Integral

Assignment

Please Check <https://github.com/uoaworks/ComplexAnalysisAY2018>

Reading Materials: The Section 5.1 and 5.2 of Textbook

References

- [1] A first course in Complex Analysis with application, Dennis G. Zill and Patrick D. Shanahan, Jones and Bartlett Publishers, Inc. 2003
- [2] Calculus, 6th Edition, James Stewart, Thomas Brooks/Cole, 2009
- [3] Wikipedia

EXAMPLE (例題) Recall Real Line Integral in *Calculus II* (微積分 II)

Evaluate $\int_C (2 + x^2 y) ds$, where C is the upper half of the unit circle $x^2 + y^2 = 1$

Solution (解答):


Recall that the unit circle can be parametrized by

$$x = \cos t, y = \sin t$$

And the upper half of the circle is described by the parameter interval $0 \leq t \leq \pi$

Therefore, from the formula in Page 14 of this lecture note, we have

$$\begin{aligned} \int_C (2 + x^2 y) ds &= \int_0^\pi (2 + \cos^2 t \sin t) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt \\ &= \int_0^\pi (2 + \cos^2 t \sin t) \sqrt{(-\sin t)^2 + \cos^2 t} dt \\ &= \int_0^\pi (2 + \cos^2 t \sin t) dt = \left[2t - \frac{\cos^3 t}{3} \right]_0^\pi = 2\pi + \frac{2}{3} \end{aligned}$$



EXAMPLE (例題) Recall Real Line Integral in *Calculus II* (微積分 II)


Evaluate $\int_C 2x \, ds$, where C consists of the arc C_1 of the parabola $y = x^2$ from $(0,0)$ to $(1,1)$ followed by the vertical line segment C_2 from $(1,1)$ to $(1,2)$.

Solution (解答): ① Because from C_1 we know y is a function of x , i.e. the domain (x, y) becomes (x, x^2) , so we can use x as the parameter, then

$$x = x, \quad y = x^2, \quad 0 \leq x \leq 1$$

Therefore

$$\begin{aligned} \int_{C_1} 2x \, ds &= \int_0^1 2x \sqrt{\left(\frac{dx}{dx}\right)^2 + \left(\frac{dy}{dx}\right)^2} \, dx = \int_0^1 2x \sqrt{1 + (2x)^2} \, dx \\ &= \left[\frac{1}{4} \cdot \frac{2}{3} (1 + 4x^2)^{\frac{3}{2}} \right]_0^1 = \frac{5\sqrt{5} - 1}{6} \end{aligned}$$



Solution (解答)(cont.):


- ② Because from C_2 we see a vertical line segment, so we can use y as the parameter, then

$$x = 1, \quad y = y, \quad 1 \leq y \leq 2$$

Therefore

$$\begin{aligned} \int_{C_2} 2x ds &= \int_1^2 2 \cdot 1 \cdot \sqrt{\left(\frac{dx}{dy}\right)^2 + \left(\frac{dy}{dy}\right)^2} dy \\ &= \int_1^2 2\sqrt{0+1} dt = \int_1^2 2 dt = 2 \end{aligned}$$

$$\int_C 2x ds = \int_{C_1} 2x ds + \int_{C_2} 2x ds = \frac{5\sqrt{5}-1}{6} + 2$$



Application 2 of Complex Number :

Rotation

